Title	Zn protoporphyrin IX is formed not from heme but from protoporphyrin IX
Author(s)	Wakamatsu, Jun-ichi; Okui, Jun; Hayashi, Nobutaka; Nishimura, Takanori; Hattori, Akihito
Citation	Meat Science, 77(4), 580-586 https://doi.org/10.1016/j.meatsci.2007.05.008
Issue Date	2007-12
Doc URL	http://hdl.handle.net/2115/32300
Type	article (author version)
File Information	MEATSCI-00261.pdf

[Instructions for use](#)

Zn protoporphyrin IX is formed not from heme
but from protoporphyrin IX.

Jun-ichi WAKAMATSU*, Jun OKUI, Nobutaka HAYASHI,
Takanori NISHIMURA, Akihito HATTORI

Division of Bioresource and Product Science, Graduate School of Agriculture,
Hokkaido University, Sapporo, Hokkaido 060-8589, Japan

*Corresponding author.

Tel.: +81 11 706 2547; Fax.: +81 11 706 2547

E-mail address: jwaka@anim.agr.hokudai.ac.jp (J. Wakamatsu)

Meat Science Laboratory, Division of Bioresource and Product Science, Research
Faculty of Agriculture, Hokkaido University
N-9 W-9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

19 **Abstract**

20

21 We examined the effects of exogenous myoglobin, a bivalent chelator, and nitrite on
22 Zn protoporphyrin IX (ZPP) formation by using model systems. ZPP was formed in a
23 model solution without addition of exogenous myoglobin. After incubation, the amount
24 of ZPP in a model solution was increased but that of heme was not decreased compared
25 with the amounts before incubation. Protoporphyrin IX (PPIX) instead of ZPP also
26 accumulated in a model solution with addition of EDTA, but the amount of heme was
27 not reduced. These results suggested that ZPP was not formed by the Fe-Zn
28 substitution in heme but was formed by the insertion of Zn into PPIX, which was
29 formed independently. The fact that the effects of various factors in model systems
30 with/without addition of a bivalent chelator were similar suggested that ZPP formation
31 was strongly affected by PPIX formation. Inhibition of PPIX formation by nitrite
32 might be the reason for the low levels of ZPP in cured meats.

33

34 **Keywords:** Zn protoporphyrin IX; protoporphyrin IX; color; dry-cured meat products;
35 nitrite; nitric oxide

1. Introduction

Dry-cured meat products have been produced and consumed throughout history by a diversity of cultures in different parts of the world. Parma ham, one of the Italian dry-cured hams, is made from heavier pigs (> 150 kg in liveweight) and is processed for at least 12 months without addition of nitrite/nitrate (Toldrá, 2002). It has been reported that Parma ham has a red porphyrin derivative that is not nitrosylmyoglobin or oxymyoglobin and is extractable by water, acetone/water (75%/25%) and tetrahydrofuran (Morita, Niu, Sakata & Nagata, 1996; Møller, Adamsen & Skibsted, 2003; Parolari, Gabba & Saccani, 2003). The pigment was purified and identified to be Zn protoporphyrin IX (ZPP) by ESI-HR-MS analysis (Wakamatsu, Nishimura & Hattori, 2004). Moreover, the presence of zinc in the red pigment was directly demonstrated by scanning electron microscopy/energy dispersive X-ray microanalysis (SEM-EDX) (Wakamatsu, Ito, Nishimura & Hattori, 2007). Møller, Adamsen, Catharino, Skibsted, & Eberlin (2007) showed by using ESI-MS and TOF-MS analysis that zinc protoporphyrin IX is present not only in Parma ham but also in Iberian ham.

Effects of various factors on ZPP formation have been investigated using a model system, and it has been shown that oxygen inhibited ZPP formation and that ZPP was formed in the absence of microorganisms (Wakamatsu, Okui, Nishimura & Hattori, 2004). It has also been reported that the amount of ZPP and protoporphyrin IX (PPIX) were increased in pork and turkey during anaerobic conditions (Veberg et al., 2006). By use of autofluorescence of ZPP, it has been revealed that ZPP was distributed in not only lean meat but also fat tissue by using near-UV purple LED irradiation and an image analysis (Wakamatsu, Odagiri, Nishimura & Hattori, 2006).

On the other hand, exogenous myoglobin was added into a model solution that was established by Wakamatsu, Okui, Nishimura & Hattori (2004). The addition of exogenous myoglobin may affect ZPP formation. Moreover, the amount of ZPP formed

63 has only been measured by fluorescence intensity, and the amount of heme has not been
64 determined at all. In order to elucidate the mechanism by which ZPP is formed, these
65 porphyrins must be quantitatively determined. A recent study has shown that ZPP is
66 present not only in Parma ham but also in Iberian ham but that ZPP content in meat
67 products cured with nitrite is very low (Adamsen Møller, Laursen, Olsen & Skibsted,
68 2006). It is still not clear why ZPP content in cured meat products is much lower than
69 that in dry-cured ham without addition of nitrite.

70 In this study, to elucidate the mechanism by which ZPP is formed, we investigated
71 the effects of various factors on ZPP formation by using a simpler model system with no
72 addition of exogenous myoglobin and a new model system in which PPX was formed by
73 the addition of ethylenediaminetetraacetate (EDTA). Then we measured the contents
74 of heme, ZPP and PPIX before and after incubation in these model solutions and
75 investigated stoichiometrically the mechanism by which ZPP or PPIX was formed.
76 Additionally, we investigated the inhibitory effect on ZPP formation of nitrite, which is
77 generally used in cured meat products.

78 2. Materials and methods

79

80 2.1. *Materials*

81

82 Pork loin samples (n = 3) were purchased from various retail markets. Myoglobin
83 (from horse skeletal muscle), penicillin G potassium, streptomycin sulfate and
84 gentamicin sulfate were purchased from Wako Pure Chemical Industries, Ltd. (Osaka,
85 Japan). Acetone, acetic acid, acetate ethyl, methanol and ammonium acetate were
86 purchased from Kanto Chemical Co. Inc. (Tokyo, Japan). Methanol was of HPLC grade
87 and other chemicals were of analytical grade.

88

89 2.2. *Model system*

90

91 Model solutions were prepared as described by Wakamatsu, Okui, Nishimura &
92 Hattori (2004). Experimental designs are shown in Table 1. Pork loin was
93 homogenized with 2 volumes of distilled water using a homogenizer at 10,000 rpm for 1
94 min. Antibiotics were added to the model solutions to final concentrations of 100
95 units/ml for penicillin G potassium, 0.1 mg/ml for streptomycin sulfate and 0.05 mg/ml
96 for gentamicin sulfate. The solutions were put into gas-impermeable bags and
97 incubated at 25 °C for 5 days in darkness using an environmental chamber. An
98 anaerobic condition was obtained by using a commercial kit (Oxygen Absorbing System;
99 I.S.O. Inc., Yokohama, Japan) consisting of an oxygen absorber (A-500HS, I.S.O. Inc.)
100 and an oxygen indicator tablet. The color of the oxygen indicator tablet changes from
101 violet to pink when the oxygen concentration becomes less than 0.1%.

102

103 2.3. *Fluorescent analysis*

104

105 Extraction by acetone and fluorescent analysis were carried out as described by
106 Wakamatsu, Okui, Nishimura & Hattori (2004) with minor modification. After
107 extraction by acetone, the fluorescent spectra of the extracts were measured from 450 to
108 700 nm at 420/410 nm for excitation using a spectrofluorophotometer (RF-5300PC,
109 Shimadzu Corp., Kyoto, Japan). Fluorescence intensity at 590 nm for excitation at 420
110 nm was regarded as the amount of ZPP formed, and fluorescence intensity at 630 nm for
111 excitation at 410 nm was regarded as the amount of PPIX formed. All operations were
112 carried out in darkness as much as possible.

113

114 *2.4. Quantitative analysis of ZPP, PPIX and heme*

115

116 The contents of ZPP, PPIX and heme were determined by HPLC as described by Guo,
117 Lim & Peters (1991) with some modification. Porphyrins were extracted by acetic
118 acid/acetate ethyl (1:4, v/v) as described by Smith, Doran, Mazur & Bush (1980) with
119 some modification. One milliliter of model solution was mixed vigorously with 9 ml of
120 acetic acid/acetate ethyl (1:4, v/v), and the mixture was held on ice for 30 min. After
121 centrifugation (3,000 rpm, 15 min), the supernatant was collected. This extraction
122 operation was carried out twice. The resulting solution was mixed with an equal
123 volume of methanol/ammonium acetate (86:14, v/v, pH 5.16). The sample was filtered
124 through a 0.45-µm filter (Minisart RC4, Sartorius AG, Goettingen, Germany). An STR
125 ODS-II column (4.6 x 150 mm, Shinwa Chemical Industries Ltd., Kyoto, Japan) was
126 used for separation of porphyrin, and the separation was carried out by isocratic elution
127 using methanol/ammonium acetate (86:14, v/v, pH 5.16) at a flow rate of 0.6 ml/min at
128 35°C. Forty microliters of each sample was injected. The detection of ZPP and PPIX
129 was carried out at excitation and emission wavelengths of 420/400 and 590/630 nm,
130 respectively. Heme was monitored at 400 nm absorption. All operations of extraction

131 were carried out in darkness as much as possible.

132

133 *2.5. Statistical analysis*

134

135 Results are expressed as mean values of three independent trials in each individual.

136 Data were analyzed using one-way analysis of variance (ANOVA) with Scheffé's test.

137 Statistics were calculated using Microsoft Excel 2002 and a statistic add-in software

138 (Excel Toukei 2002 for Windows, Social Survey Research Information Co., Ltd., Tokyo,

139 Japan). A probability of $P<0.05$ was considered statistically significant.

140 3. Results and discussion

141

142 First, in order to investigate the effects of exogenous myoglobin in our established
143 experimental model, the effect of the amount of added myoglobin on the amount of ZPP
144 formed was examined (Fig. 1). When exogenous myoglobin was added at 0.1%, the
145 amount of ZPP formed was maximal. With further addition of myoglobin, the amount
146 of formed ZPP gradually decreased. The amount of ZPP formed was significantly
147 increased with the addition of 0.1% exogenous myoglobin. The cause of the increase is
148 unclear and it may be because myoglobin used in this study was metmyoglobin or
149 included impurities. However, ZPP was formed despite no addition of myoglobin.
150 Therefore, this result suggested that exogenous myoglobin is not always essential for
151 the formation of zinc protoporphyrin IX and ZPP. It is also possible that ZPP is formed
152 only from pork components. Since the addition of exogenous myoglobin in a model
153 solution might make it difficult to elucidate the mechanism by which ZPP is formed, the
154 mechanism should be investigated in a simpler model system with no addition of
155 exogenous myoglobin. However, the cause of increase in ZPP formation by the addition
156 of 0.1% exogenous myoglobin was not verified.

157 Next, we established a simpler model system composed of pork homogenate and
158 antibiotics without addition of exogenous myoglobin and investigated the effects of
159 various factors on ZPP formation (Fig. 2). Oxygen inhibited ZPP formation (Fig. 2A)
160 and the amount of ZPP formed was increased with increase in the amount of pork added
161 (Fig. 2B). The amount of ZPP formed rapidly increased during the first 3 days of
162 incubation and increased slightly after the third day (Fig. 2C). The amount of ZPP
163 formed was also increased with increase in incubation temperature within the range of
164 temperatures examined (4, 15, 25 and 35°C) (Fig. 2D). The amount of ZPP formed
165 peaked at about pH 5.5 and decreased considerably at lower or higher pH (Fig. 2E).

166 The results showing that ZPP formation depended on many factors suggested the
167 involvement of an enzyme. The most notable dependence is the pH dependence. ZPP
168 formation peaked at 5.5, which is very close to the ultimate pH of meat. The optimal
169 pH range was narrower than those for general enzymes. Although the cause is not
170 clear, this pH range might be an important clue to elucidate the mechanism by which
171 ZPP is formed.

172 On the other hand, it has been reported that little ZPP was present in meat products
173 cured with nitrite and/or nitrate (Adamsen Møller, Laursen, Olsen & Skibsted, 2006).
174 Therefore, the influence of added nitrite was investigated by using the model system
175 (Fig. 2F). The formation of ZPP was inhibited significantly by addition of nitrite of
176 more than 10 μ M (0.69 mg/L). Since the concentration at which ZPP formation was
177 inhibited was much lower than the amount of nitrite added to common meat products, it
178 was speculated that low ZPP content in meat products cured with nitrite and/or nitrate
179 is caused by the inhibition of ZPP formation by nitrite. However, the cause is still
180 unclear and future studies are needed.

181 In previous studies (Wakamatsu, Nishimura & Hattori, 2004; Wakamatsu, Okui,
182 Nishimura & Hattori, 2004), the amount of ZPP formed was determined by fluorescence
183 intensity and was not determined quantitatively. Heme content was also not
184 measured. Therefore, we determined the quantities of ZPP and heme in the model
185 solutions. As shown in Fig. 3, although the amount of ZPP had significantly increased
186 after 5-day incubation, a decrease in the amount of heme was not observed. PPIX was
187 not detected before and after incubation. Therefore, the total amount of ZPP and heme
188 was increased compared with the total amount before incubation. If ZPP was just
189 formed from heme with substitution of Zn for Fe, the sum total amount of heme and
190 ZPP would not be increased. This result therefore suggested that ZPP was not formed
191 by Fe-Zn substitution in heme but was independently formed in the model during
192 incubation. At the last step of heme biosynthesis, heme was formed by inserting Fe

193 into PPIX (Ferreira, 1999; Dailey, Dailey, Wu, Medlock, Rose & Wang, 2000; Dailey,
194 2002). It is therefore possible that ZPP is formed by insertion of Zn into PPIX
195 generated independently. Thus, the chelation of zinc by a chelator was predicted to
196 inhibit the formation of ZPP and to result in the accumulation of PPIX.

197 Next, we observed the fluorescence spectrum of acetone extract after incubation in
198 the model solution with addition of EDTA, a bivalent metal chelator (Fig. 4). A strong
199 peak at 630 nm was observed in the model solution with addition of EDTA. The
200 maximum excitation wavelength of the fluorescence peak was about 410 nm, shorter
201 than the excitation wavelength of ZPP (420 nm) in agreement with results of a previous
202 study (Shepherd & Dailey, 2005). This fluorescence peak coincided with that of PPIX
203 (Shepherd & Dailey, 2005; Veberg et al., 2006). Thus, since the chelation of zinc by
204 EDTA inhibited the formation of ZPP, ZPP seems to be formed by insertion of Zn into
205 PPIX.

206 Next, in order to determine whether PPIX was formed from the demetalation of
207 heme or independently, porphyrin contents in the model solutions with and without
208 addition of EDTA were determined after incubation (Fig. 5). Although the addition of
209 EDTA into a model solution significantly inhibited ZPP formation and the amount of
210 PPIX was increased significantly, there was no change in heme content ($P>0.05$). The
211 increase in PPIX content and the fact that the amount of heme did not change strongly
212 suggested that PPIX was formed not from heme but from another substrate. EDTA
213 was added into a model solution, but only a small amount of ZPP was present in the
214 solution. This is due to the presence of ZPP originally contained in pork (see Fig. 3).

215 We also investigated the effects of various factors on the formation of PPIX in simple
216 model systems with addition of EDTA. Oxygen tended to inhibit the formation of PPIX
217 (Fig. 6A). PPIX was increased with increase in pork content in model solutions (Fig.
218 6B). The amount of PPIX formed increased rapidly during the first 3 days of
219 incubation and increased slightly after the third day (Fig. 6C). The amount of PPIX

increased with increase in incubation temperature (Fig. 6D) and was maximal at about pH 5.5 (Fig. 6E). The addition of more than 10 μ M of nitrite inhibited ZPP formation significantly (Fig. 6F). All of these patterns of PPIX formation are very similar to those of ZPP formation (Fig. 2). Thus, these results indicated that ZPP in a model solution was formed by the insertion of Zn into PPIX, which was formed not from heme but independently. It is therefore possible that ZPP is also formed by the same pathway in Parma ham. PPIX is formed from protoporphyrinogen IX (proto'gen) by protoporphyrinogen oxidase (PPO; EC 1.3.3.4) in the heme biosynthesis system (Dailey, 2002). The enzyme activity of mammalian PPO has been shown to be maximal at pH 8.6 to 8.7 (Poulson, 1976), much higher than results obtained in the present study. On the other hand, the inhibition of PPIX formation by oxygen was reduced in model systems with addition of EDTA, but there was no significant difference between them. There are two forms of PPO, an oxygen-dependent form and an oxygen-independent form (Dailey, 2002), and they may contribute to PPIX formation. Our results revealed that nitrite inhibited the formation of not only ZPP but also PPIX. Anaerobic oxidation of proto'gen to PPIX has been demonstrated in extracts of several microorganisms by nitrite or nitrate as an electron acceptor (Jacobs & Jacobs, 1976; Klemm & Barton, 1985). Since the inhibition of PPIX formation by nitrite is not still clear, further studies are needed to elucidate the contribution of proto'gen and PPO to ZPP/PPIX formation. The patterns of ZPP and PPIX formation were similar in model systems. The pH curves with maximum peaks at about 5.5 are noteworthy, and they might serve as a clue to elucidating the mechanism by which ZPP/PPIX is formed.

Furthermore, our results revealed that nitrite inhibited the formation of not only ZPP but also PPIX. Color formation in cured meat products is caused mainly by the reaction of endogenous myoglobin in meat with nitric oxide (NO) (Sakata, 2000; Fox, 1966). NO is produced from added nitrite (or nitrate) as the curing agent. NO has a high affinity for non-heme iron proteins (Fujii & Yoshimura, 1996) and copper proteins

247 (Suzuki, 1996). NO is a paramagnetic molecule and has a high affinity for metal ions
248 other than heme (Fujii and Yoshimura, 1996). Consequently, many metalloenzymes
249 are inactivated by NO. Iron-sulfur (Fe-S) proteins have a very high affinity for NO.
250 Mammalian ferrochelatase contains a [2Fe-2S] cluster (Ferreira, Franco, Lloyd, Pereira,
251 Moura, Moura, & Huynh, 1994; Dailey, Dailey, Wu, Medlock, Rose & Wang, 2000), and
252 the cluster is labile and sensitive to destruction by NO (Sellers, Johnson, & Dailey,
253 1996). It has been reported that ferrochelatase as a zinc chelatase participate in ZPP
254 formation from myoglobin in pork loin extract and porcine heart extract (Ishikawa,
255 Yoshihara, Baba, Kawabuchi, Sato, Numata & Matsumoto, 2006a; Ishikawa, Yoshihara,
256 Baba, Kawabuchi, Sato, Numata & Matsumoto, 2006b). Although the participation of
257 ferrochelatase is still not clear, the inactivation of metalloenzyme by NO may be
258 involved in the inhibition of PPIX formation.

259 The results of this study might serve as a clue to elucidating the mechanism by which
260 ZPP/PPIX is formed. Since the formation of PPIX had a strong influence on that of ZPP,
261 it is necessary to investigate the steps prior to PPIX formation in more detail.

262

263 4. Conclusion

264

265 This study demonstrated that ZPP in a model solution was not formed by Fe-Zn
266 substitution in heme but was formed by the insertion of Zn into PPIX, which was
267 formed independently. It was shown that ZPP formation was strongly influenced by
268 PPIX formation. The fact that there is little ZPP in cured meat products is because
269 ZPP cannot be formed as a result of inhibition of PPIX formation by nitrite. A full
270 understanding of ZPP formation in Parma ham awaits elucidation of the formation of
271 PPIX or its precursor.

272

273

274 Acknowledgments

275 This study was supported partially by a Grant-in-Aid for Scientific Research
276 (16780184) from the Ministry of Education, Culture, Sports, Science and Technology of
277 Japan.

278

279 References

280

281 Adamsen, C. E., Møller, J. K. S., Laursen, K., Olsen, K. & Skibsted, L.H. (2006).

282 Zn-porphyrin formation in cured meat products: Effect of added salt and nitrite,

283 *Meat Science*, *72*, 672-679.

284 Dailey H. A. (2002). Terminal steps of haem biosynthesis, *Biochemical Society*

285 *Transactions*, *30*, 590-595.

286 Dailey, H. A., Dailey, T. A., Wu, C.-K., Medlock, A. E, Rose, J. P., & Wang, K.-F. (2000).

287 Ferrochelatase at the millennium: structures, mechanisms and [2Fe-2S] clusters,

288 *Cellular & Molecular Life Sciences*, *57*, 1909-1926.

289 Ferreira, G. C., Franco, R., Lloyd, S. G., Pereira, A. S., Moura, I. , Moura, J. J., & Huynh,

290 B. H. (1994) Mammalian ferrochelatase, a new addition to the metalloenzyme

291 family. *The Journal of Biological Chemistry*, *269*, 7062-7065.

292 Ferreira, G.C. (1999) Ferrochelatase. *The International Journal of Biochemistry & Cell*

293 *Biology*, *31*, 995-1000.

294 Fox J.B. Jr. (1966) The chemistry of meat pigments, *Journal of Agricultural and Food*

295 *Chemistry*, *14*, 207-210.

296 Fujii , S. & Yoshimura, T. (1996) (Japanese) 13 Interaction of nitric oxide with non-heme

297 iron proteins,

298 Guo, R., Lim, C. K., & Peters T. J. (1991). High-performance liquid chromatographic

299 assays for protoporphyrinogen oxidase and ferrochelatase in human leucocytes,

300 *Journal of Chromatography*, *566*, 383-396.

301 Hornsey, H. C. (1956). The colour of cooked cured pork. I. Estimation of the nitric

302 oxide-haem pigments. *Journal of the Science of Food and Agriculture*, *7*, 534-540.

303 Ishikawa, H., Yoshihara, M., Baba, A., Kawabuchi, T., Sato, M., Numata, M. &

304 Matsumoto, K. (2006a). Formation of zinc protoporphyrin IX from myoglobin with

305 pork loin extract, *Journal of The Faculty of Agriculture Kyusyu University*, *51*,

306 93-97.

307 Ishikawa, H., Yoshihara, M., Baba, A., Kawabuchi, T., Sato, M., Numata, M. &
308 Matsumoto, K. (2006b). Formation of zinc protoporphyrin IX from myoglobin in
309 porcine heart extract, *Food Science and Technology Research*, 12, 125-130.

310 Jacobs, N. J., & Jacobs, J. M. (1976). Nitrate, fumarate, and oxygen as electron
311 acceptors for a late step in microbial heme synthesis, *Biochimica et Biophysica Acta*,
312 449, 1-9.

313 Klemm, D. J., & Barton, L. L. (1985). Oxidation of protoporphyrinogen in the obligate
314 anaerobe *Desulfovibrio gigas*, *Journal of Bacteriology*, 164, 316-320.

315 Møller, J. K. S., Adamsen, C. E., & Skibsted, L. H. (2003). Spectral characterization of
316 red pigment in Italian-type dry-cured ham. Increasing lipophilicity during
317 processing and maturation. *European Food Research and Technology*, 216, 290-296.

318 Møller, J. K. S., Adamsen, C. E., Catharino, R. R., Skibsted, L. H., & Eberlin M. N.
319 (2007) Mass spectrometric evidence for a zinc-porphyrin complex as the red
320 pigment in dry-cured Iberian and Parma ham. *Meat Science*, 75, 203-210.

321 Morita, H., Niu, J., Sakata, R., & Nagata, Y. (1996). Red pigment of Parma ham and
322 bacterial influence on its formation. *Journal of Food Science*, 61, 1021-1023.

323 Okayama, T., & Nagata, Y. (1978). (Japanese) A Modification of the method for
324 measuring the color formation ability of meat products. *Japanese Journal of*
325 *Zootechnical Science*, 49, 866-871.

326 Okayama, T., & Nagata, Y. (1979). (Japanese) A method for the determination of total
327 heme pigments in meat products. *Japanese Journal of Zootechnical Science*, 50,
328 15-21.

329 Parolari, G., Gabba, L., & Saccani, G. (2003). Extraction properties and absorption
330 spectra of dry cured hams made with and without nitrate. *Meat Science*, 64,
331 483-490.

332 Poulson, R. (1976). The enzymic conversion of protoporphyrinogen IX to protoporphyrin

IX in mammalian mitochondria, *The Journal of Biological Chemistry*, 251,
3730-3733.

Sakata, R. (2000) Studies on physicochemical characteristics of red pigments in meat products. *Animal Science Journal*, 71, 1-16.

Sellers, V. M., Johnson, M. K., & Dailey, H. A. (1996) Function of the [2Fe-2S] cluster in mammalian ferrochelatase: a possible role as a nitric oxide sensor. *Biochemistry*, 35, 2699-2704.

Shepherd, M., & Dailey, H.A. (2005). A continuous fluorimetric assay for protoporphyrinogen oxidase by monitoring porphyrin accumulation, *Analytical Biochemistry*, 344, 115-121.

Smith, R. M., Doran, D., Mazur M., & Bush, B (1980). High-Performance Liquid Chromatographic Determination of Protoporphyrin and Zinc Protoporphyrin in Blood, *Journal of Chromatography*, 181, 319-327.

Toldrá, F. (2002). *Dry-cured meat products*, Connecticut, USA: Food & Nutrition Press, Inc.

Veberg, A., Sørheim, O., Moan, J., Iani, V., Juzenas, P., Nilsen, A. N., & Wold, J. P. (2006). Measurement of lipid oxidation and porphyrins in high oxygen modified atmosphere and vacuum-packed minced turkey and pork meat by fluorescence spectra and images. *Meat Science*, 73, 511-520.

Wakamatsu, J., Nishimura, T., & Hattori, A. (2004). A Zn-porphyrin complex contributes to bright red color in Parma ham. *Meat Science*, 67, 95-100.

Wakamatsu, J., Okui, J., Ikeda, Y., Nishimura, T., & Hattori, A. (2004). Establishment of a model experiment system to elucidate the mechanism by which Zn-protoporphyrin IX is formed in nitrite-free dry-cured ham. *Meat Science*, 68, 313-317.

Wakamatsu, J., Odagiri, H., Nishimura, T., & Hattori, A. (2006). Observation of the distribution of Zn protoporphyrin IX in Parma ham by using purple LED and image

360 analysis. *Meat Science.*, 74, 594-599.

361 Wakamatsu, J., Ito, T., Nishimura, T., & Hattori A. (2007). Direct demonstration of the

362 presence of zinc in the acetone-extractable red pigment from Parma ham. *Meat*

363 *Science.*, 76, 385-387.

364 Figure legends

365

366

367 Fig. 1. Effect of exogenous myoglobin addition on the formation of ZPP. Model
368 solutions with addition of various amounts of exogenous myoglobin were incubated
369 for 5 days at 25°C anaerobically. The fluorescence intensity of acetone extracts was
370 measured. Bars represent the standard deviation of the means (n=3). ^{ab}: Values
371 bearing different letters are significantly different ($P < 0.05$).

372

373 Fig. 2. Effects of oxygen (A), pork content (B), incubation day (C), temperature (D), pH
374 (E) and nitrite (F) on the formation of ZPP in model solutions with no addition of
375 exogenous myoglobin. The fluorescence intensity of acetone extracts was measured
376 after incubation. Bars represent the standard deviation of the means (n=3). ^{abcde}:
377 Values in the same examination bearing different letters are significantly different
378 ($P < 0.05$).

379

380 Fig. 3. Heme (■) and ZPP(■) concentrations in model solutions without addition of
381 exogenous myoglobin before and after incubation. Porphyrins extracted by acetic
382 acid/acetic ethyl were measured by HPLC. Bars represent the standard deviation
383 of the means (n=3). n.s.: not significant (vs before incubation).

384

385 Fig. 4. Fluorescence pattern of acetone extract of model solutions with (broken line, Ex.
386 410 nm) and without (solid line, Ex. 420 nm) addition of EDTA after 5 days at 25°C.

387

388 Fig. 5. Heme (■), ZPP (■) and PPIX (□) concentrations in model solutions with and

389 without addition of EDTA after 5 days at 25°C. Porphyrins extracted by acetic
390 acid/acetic ethyl were measured by HPLC. Bars represent the standard deviation
391 of the means (n=3). n.s.: not significant (vs with addition of EDTA).

392

393 Fig. 6. Effects of oxygen (A), pork content (B), incubation day (C), temperature (D), pH
394 (E) and nitrite (F) on the formation of ZPP in the model solution with addition of
395 EDTA. The fluorescence intensity of acetone extracts was measured after
396 incubation. Bars represent the standard deviation of the means (n=3). ^{abcde}:
397 Values in the same examination bearing different letters are significantly different
398 ($P < 0.05$). n.s.: not significant.

399

400 Table 1. Experimental designs by using a model system

401

Figure1

Fig.1

Figure2

Fluorescence intensity (Ex/Em 420/590 nm)

Fig.2

Figure3

Fig. 3

Figure4

Fig.4

Figure5

Fig. 5

Figure6

Fluorescence intensity (Ex/Em 410/630 nm)

Fig.6

Table1

Table 1. Experimental designs by using a model system

Pork	Myoglobin	EDTA	Oxygen	Temperature	Day	pH	Nitrite
Experimental model designs with addition of exogenous myoglobin							
20%	0-1% ^a	-	< 0.1%	25°C	5	-	-
Experimental model designs without addition of exogenous myoglobin							
20%	-	-	< 0.1%	25°C	5	-	-
20%	-	-	NA ^b	25°C	5	-	-
0-50% ^c	-	-	< 0.1%	25°C	5	-	-
20%	-	-	< 0.1%	25°C	0-10 ^d	-	-
20%	-	-	< 0.1%	4-37°C ^e	5	-	-
20%	-	-	< 0.1%	25°C	5	3.5-8.5	-
20%	-	-	< 0.1%	25°C	5	-	0-100 µM ^f
Experimental model designs in the chelation on bivalent metals by EDTA							
20%	-	0.5 mM	< 0.1%	25°C	5	-	-
20%	-	0.5 mM	NA ^b	25°C	5	-	-
0-50% ^c	-	0.5 mM	< 0.1%	25°C	5	-	-
20%	-	0.5 mM	< 0.1%	25°C	0-10 ^d	-	-
20%	-	0.5 mM	< 0.1%	4, 25, 37°C	5	-	-
20%	-	0.5 mM	< 0.1%	25°C	5	3.5-8.5	-
20%	-	0.5 mM	< 0.1%	25°C	5	-	0-100 µM ^f

Antibiotics were added to all model solutions to final concentrations of 100 units/ml for penicillin G potassium, 0.1 mg/ml for streptomycin sulfate and 0.05 mg/ml for gentamicin sulfate.
^a; 0, 0.05, 0.1, 0.5, 1%; ^b; no adjusted; ^c; 0, 5, 10, 25, 50%; ^d; 0, 1, 3, 5, 10 day; ^e; 4, 15, 25, 37°C; ^f; 0, 5, 10, 50, 100 µM