



Title	CORRELATION BETWEEN STIMULUS-SECRETION COUPLING AND FUROSEMIDE-SENSITIVE ION TRANSPORT IN RAT PANCREATIC ACINI : An approach monitoring digestive enzyme release and concentration of intracellular calcium ions
Author(s)	SUGIYAMA, Tomomi
Citation	Japanese Journal of Veterinary Research, 39(1), 76-76
Issue Date	1991-05-30
Doc URL	<a href="http://hdl.handle.net/2115/3265">http://hdl.handle.net/2115/3265</a>
Type	bulletin (article)
File Information	KJ00002377496.pdf



[Instructions for use](#)

CORRELATION BETWEEN STIMULUS-SECRETION COUPLING AND  
FUROSEMIDE-SENSITIVE ION TRANSPORT IN RAT PANCREATIC ACINI

— An approach monitoring digestive enzyme release and  
concentration of intracellular calcium ions —

Tomomi SUGIYAMA

*Department of Physiology,  
Faculty of Veterinary Medicine,  
Hokkaido University, Sapporo 060, Japan*

1. The present study was carried out to examine the relationship between stimulus-secretion coupling and ion transport. The cytosolic concentration of calcium ion ( $[Ca^{2+}]_c$ ) of isolated pancreatic acini from the rat loaded with fura-2 was monitored by microspectrofluorometry. Under comparable experimental conditions, acini loaded with fura-2 were perfused with modified Krebs-Henseleit bicarbonate-buffered solution and amylase release from the perfused acini was monitored.
2. During continuous stimulation with 10pM CCK-8, amylase released from acini increased rapidly to reach a plateau level three times greater than the resting level. Cessation of the stimulation resulted in a gradual decrease in amylase release, reaching the resting level within about 15 min. Stimulation with CCK-8 induced a sustained increase in  $[Ca^{2+}]_c$  with oscillatory change. During perfusion with  $Ca^{2+}$ -free solution with 1mM EGTA, amylase release induced by 10pM CCK-8 was strongly inhibited but a small and transient increase remained in the initial phase of the secretory response. The sustained increase in  $[Ca^{2+}]_c$  was abolished but the oscillation remained in the  $[Ca^{2+}]_c$ -free environment.
3. Furosemide (1mM) reduced amylase release induced by 10pM CCK-8 to the resting level. Administration of 1mM furosemide alone decreased transiently the resting level of amylase release. Furosemide (1mM) abolished the sustained increase in  $[Ca^{2+}]_c$  induced by 10pM CCK-8 and decreased oscillatory amplitude and frequency.
4. The CCK-8-induced secretory response was unchanged by perfusion with a solution containing 0.1mM ouabain, and with a  $K^+$ -free or  $Na^+$ -deficient solution to inhibit  $Na^+$ ,  $K^+$ -ATPase activity. The response was also uninfluenced by administration of 1mM amiloride or 0.1mM SITS. Replacement of  $Cl^-$  with isethionate inhibited amylase release induced by 10pM CCK-8.
5. On the basis of the present results, the following mechanism is proposed: Binding of CCK-8 to CCK-receptors in the membrane of pancreatic acinar cells may cause an influx of extracellular  $Ca^{2+}$  and, in turn, the increase in  $[Ca^{2+}]_c$  may induce exocytosis of zymogen granules.