Title	Removal of understory dwarf bamboo (<i>Sasa kurilensis</i>) induces changes in water-relations characteristics of overstory <i>Betula ermanii</i> trees
Author(s)	Ishii, Hiroaki T.; Kobayashi, Tsuyoshi; Uemura, Shigeru; Takahashi, Koichi; Hanba, Yuko T.; Sumida, Akihiro; Hara, Toshihiko
Citation	Journal of Forest Research, 13(2), 101-109 https://doi.org/10.1007/s10310-007-0058-0
Issue Date	2008-04
Doc URL	http://hdl.handle.net/2115/32745
Rights	The original publication is available at www.springerlink.com
Type	article (author version)
File Information	ishii.pdf

[Instructions for use](#)

Removal of understory dwarf bamboo (*Sasa kurilensis*) induces changes in
water-relations characteristics of overstory *Betula ermanii* trees

Hiroaki T. Ishii^{1,2}, Tsuyoshi Kobayashi^{1,3}, Shigeru Uemura⁴, Koichi Takahashi⁵, Yuko T.
Hanba⁶, Akihiro Sumida¹, Toshihiko Hara¹

¹ Biosphere Dynamics Research Group, Institute of Low Temperature Science, Hokkaido
University, Sapporo 060-0819, JAPAN

² Current address and correspondence: Division of Forest Resources, Graduate School of
Agriculture, Kobe University, Kobe 657-8501, JAPAN
Phone: +81-78-803-5826; FAX: +81-78-803-5349; hishii@alumni.washington.edu.

³ Current address: Faculty of Agriculture, Kagawa University, Kagawa 761-0795, JAPAN

⁴ Field Science Center for Northern Biosphere, Hokkaido University, Sapporo 060-0809,
JAPAN

⁵ Faculty of Science, Shinshu University, Matsumoto 390-8621, JAPAN

⁶ Center for Bioresource Field Science, Kyoto Institute of Technology, Kyoto 616-8354,
JAPAN

1 Removal of understory dwarf bamboo (*Sasa kurilensis*) induces changes in
2 water-relations characteristics of overstory *Betula ermanii* trees

3
4 **ABSTRACT**

5 We investigated how removal of understory dwarf bamboo (*Sasa kurilensis* Makino et
6 Shibata) affected growth and water-use of young *Betula ermanii* Cham. trees in a
7 secondary forest in northern Japan. We compared current-year shoot morphology, leaf
8 water-relations characteristics and whole-tree water use of *B. ermanii* trees growing with
9 dwarf bamboo in the understory (intact plot) to those of trees growing where dwarf
10 bamboo was experimentally removed (removal plot). Current-year shoot length and
11 internode frequency increased, while internode length and individual leaf area decreased in
12 response to dwarf bamboo removal. Carbon isotope composition ($\delta^{13}\text{C}$) of leaves did not
13 change indicating that leaf water-use efficiency was unaffected by the presence/absence of
14 dwarf bamboo. Pressure-volume analysis indicated that leaves in the removal plot had
15 lower water uptake ability. During mid-summer, trees in the removal plot sustained high
16 sap flow velocity after peaking at midday, while that of trees in the intact plot declined. In
17 addition, whole-tree sap flux and soil-to-leaf hydraulic conductance were both higher for
18 trees in the removal plot. We inferred that the observed changes reflect hydraulic
19 homeostasis of *B. ermanii* trees to maintain constant water-use efficiency in response to
20 belowground competition with dwarf bamboo. Removal of dwarf bamboo is, therefore, an
21 effective silvicultural prescription to enhance growth of overstory trees by relieving
22 belowground competition for soil water.

23
24 **Key Words:** hydraulic homeostasis, sap flow, shoot morphology, understory management,
25 water relations

1 Introduction

2 Dwarf bamboos (genus *Sasa*; clonal evergreen plants) are prominent understory
 3 species of many temperate forests in East Asia, including Japan (Oshima 1961; Franklin et
 4 al. 1979). Dwarf bamboo cover is especially thick in areas of heavy snowfall, where culms
 5 reach over 2 m in height. In such areas, leaf area index (LAI: total leaf area per unit ground
 6 area) of dwarf bamboo may be as high as 4 to 6 and relative light intensity on the forest
 7 floor is less than 5% (Oshima 1962; Konno 2001; Kobayashi et al. 2004). Thick dwarf
 8 bamboo cover inhibits tree regeneration by preventing seedling establishment and
 9 suppressing sapling growth (Oshima 1962; Taylor and Qin 1988; Kobayashi et al. 2004).
 10 As a result, stands with thick dwarf bamboo cover completely lack sub-canopy trees and
 11 only consist of two layers of vegetation: overstory trees and understory dwarf bamboo
 12 (Franklin et al. 1979).

13 In contrast to the many studies conducted on the negative effects of dwarf bamboo
 14 on tree regeneration (e.g., Nakashizuka 1988; Hiura et al. 1996; Takahashi 1997), its effect
 15 on the growth of overstory trees has received less attention. Dense understory vegetation
 16 competes with overstory trees for nutrients and water (Sands and Nambiar 1984; Oren et al.
 17 1987; Riegel et al. 1992; Miller et al. 1998). For example, in secondary *Pinus densiflora*
 18 Sieb et Zucc. stands in south-central Japan, maximum net photosynthetic rate and stomatal
 19 conductance of needles increased (Kume et al. 2003) and diameter growth increased
 20 (Sakamoto et al. 2004) in response to removal of understory vegetation. This suggests that
 21 stand productivity could be considerably increased through adequate control of understory
 22 vegetation.

23 In northern Japan, dwarf bamboo is the dominant understory species in many
 24 secondary forests and plantations (Noguchi and Yoshida 2005). It is likely that overstory
 25 trees and dwarf bamboo compete for belowground resources such as water and nutrients.

Takahashi et al. (2003) found that diameter-growth rate of young *Betula ermanii* Cham. in Hokkaido, northern Japan, increased in the two years following experimental removal of dwarf bamboo. This and subsequent studies in the same stand indicated that soil water and N availability both increased after dwarf bamboo removal (Takahashi et al. 2003; Tripathi et al. 2006). A separate study showed that photosynthetic rate of *B. ermanii* leaves decreased as soil water potential decreased during the growing season in the stand with understory dwarf bamboo (Takahashi et al. 2004). The fine and coarse roots of both dwarf bamboo and *B. ermanii* are concentrated in the top 20 cm of the soil (Tripathi et al. 2005). Therefore, removal of dwarf bamboo is expected to increase the growth of overstory *B. ermanii* trees by relieving belowground competition for water and nutrients.

In this study, we investigated how removal of dwarf bamboo influenced the water-relations characteristics of overstory *B. ermanii* trees in a young secondary forest in northern Japan. We compared current-year shoot morphology, leaf water-relations characteristics and whole-tree water use of *B. ermanii* trees growing with dwarf bamboo in the understory to those of trees growing where dwarf bamboo was experimentally removed. From these results, we inferred the ecophysiological mechanisms that lead to increased growth rate of *B. ermanii* trees in response to dwarf bamboo removal and evaluated the effectiveness of this silvicultural prescription for enhancing the growth of stands with dense understory dwarf bamboo cover.

Materials and Methods

Study site and sample trees

The study was conducted in an even-aged, pure stand of *B. ermanii* (ca. 25-year-old) located atop the flat ridge of Mt. Jinja (580 m above sea level), Moshiri Village, in the Uryu Experimental Forest of Hokkaido University in Hokkaido Prefecture, northern Japan

(44° 20' N, 142° 15' E). The total stand area is ca. 5 ha. Mean annual temperature in the region is 3.0°C ranging from -12.1°C mean monthly temperature in January to 18.1°C in August. Annual mean precipitation is ca. 1500 mm. Snow depth during the winter months (December–March) reaches more than 2 m.

The *B. ermanii* stand initiated after dwarf bamboo (*Sasa kurilensis* Makino et Shibata) was removed in 1973 using a bulldozer to enhance tree regeneration; (scarification cf., Umeki 2003; Yoshida et al. 2005). Following dwarf-bamboo removal, *B. ermanii* regenerated naturally from seed. In the following years, dwarf bamboo recovered from clonal growth of surviving rhizomes in the soil and formed a dense understory. According to measurements made by Takahashi et al. (2003) in 1998 (the year of experimental removal of dwarf bamboo, see below), the average height of the dwarf bamboo was 2.5 m with an LAI of 0.9. The LAI of the overstory trees was 4. Stand density and basal area of overstory trees in 1998 were: 5165 trees ha⁻¹ and 18.2 m² ha⁻¹ (Takahashi et al. 2002).

At the beginning of the growing season in 1998, all above-ground parts of dwarf bamboo were pruned in a 20 x 30 m rectangular area of the stand (here after: removal plot) and along a 5-m buffer zone on all sides to study the effects of dwarf bamboo removal on the growth and physiology of overstory *B. ermanii*. Pruning of dwarf bamboo was conducted every year for the following three years until no new shoots emerged from surviving rhizomes. Some herbaceous plants established after dwarf bamboo removal but their cover and height were small. As a control, a 20 x 30 m plot was created ca 50 m from the removal plot in the area of the stand where dwarf bamboo was left intact (hereafter: intact plot). Both plots were within the same continuous stand of *B. ermanii* with a canopy height of ca. 11 m. Because the stand is located at the top of a flat ridge, stand conditions are homogeneous and the canopies of both plots are fully exposed in all directions. There was no difference in tree size between the two plots in 1998 before removal of dwarf

bamboo (Takahashi et al. 2002).

Shoot morphology

In 2001, one scaffolding tower (13 m tall) was erected in each of the plots to study the canopy of *B. ermanii* trees. Five dominant trees near each tower were chosen as sample trees (Table 1). These trees composed the upper canopy of the stand and the crown surface of each tree was well exposed to direct sunlight. Toward the end of the growing season (September 7, 2001), we selected six current-year shoots from the crown surface of each sample tree for measurement of shoot morphology. For each current-year shoot, we measured shoot length (L_s) and number of internodes (N_i). We measured the area of each leaf (A_L) using a leaf area meter (LI-3100-C, LiCor Inc., Lincoln, NE). The leaves were then oven-dried to constant weight at 80°C to determine leaf dry mass. This was used to calculate leaf mass per area (LMA). *Betula ermanii* exhibits successive leaf emergence (Kikuzawa 1982) and early leaves had already fallen at the time of measurement. Therefore, morphological data could only be measured on late leaves.

Leaf water relations

To compare water-use efficiency of leaves, the dried leaf samples were ground to a fine powder. We analyzed carbon isotope composition of leaves using a system consisting of an elemental analyzer (EA1108, Carlo-Erba, Milan, Italy) for combustion, an interface (Finnigan MAT Confro, Bremen, Germany) and a stable isotope ratio mass spectrometer (Finnigan MAT 252). Two duplicate samples (0.1–0.2 mg) were measured for each leaf and $\delta^{13}\text{C}$ (‰) values were calculated as:

$$\delta^{13}\text{C} = \{(R_{\text{sample}}/R_{\text{standard}}) - 1\} \times 1000, \quad (3)$$

where R is the $^{13}\text{C}/^{12}\text{C}$ ratio. Relative error between duplicates from the same leaf were less

than 1.0%. Measurements from the duplicates were averaged to obtain mean $\delta^{13}\text{C}$ values for each leaf. Less negative values of $\delta^{13}\text{C}$ reflect greater integrated water-use efficiency (carbon gain per water transpired) of leaves under field conditions (Farquhar et al. 1989).

We compared leaf water relations characteristics using the pressure-volume technique (Tyree and Hammel 1972). One current-year shoot was sampled from the crown surface of each sample tree toward the end of the growing season when leaf emergence and shoot elongation had ceased (September 5, 2001). Immediately after cutting, the branches were recut under water, sealed in plastic bags with moist paper, transported to the laboratory and kept under darkness overnight. Leaf water potential measurements were made on one fully expanded, mature leaf from each current-year shoot using a Scholander-type pressure chamber (Model 600, PMS Instruments Co., Corvallis, OR). Measurements were repeated with natural dehydration in the laboratory until well after turgor loss (five to six additional data points). The leaves were then oven-dried to constant weight for measurement of dry mass.

Leaf osmotic potential at saturation ($\Psi_{\pi, \text{sat}}$), leaf water potential at turgor loss ($\Psi_{l, \text{tlp}}$) and relative water content at turgor loss (RWC_{tlp}) were obtained from the pressure-volume curve for each leaf. Bulk modulus of elasticity of leaf tissue (ϵ) was determined for each leaf from the slope of the linear relationship between leaf water potential ($\Psi_{\pi, \text{sat}}$ to $\Psi_{l, \text{tlp}}$) and free water content (Maruyama and Morikawa 1983).

Field observations of whole-tree water use

We measured whole-tree sapflow velocity in four of the five sample trees from July to September of 2001 (Table 1). Thermal Dissipation Probes (TDP-30, Dynamax Inc., Houston, TX) were inserted in opposite sides of the trunk of each tree at 1.3 m above ground level and insulated with styrofoam and reflective lining to measure sapflow velocity

by the heat dissipation technique (Granier 1987; Clearwater et al. 1999). The TDP-30 consisted of two probes, 1.2 mm in diameter and 30 mm in length. The two probes were inserted into the main stem of the tree at 40 mm vertical spacing. The upper probe received 0.2 W of current which was dissipated as heat into the sapwood. Temperature difference between the upper and lower probes (ΔT , °C) was measured every 10 seconds by a thermocouple and 30-minute averages were recorded on a data logger (CR10, Cambell Scientific, Logan, UT). We calculated sapflow velocity (U , mm s⁻¹) using the following equation proposed by Granier (1987).

$$U = 0.0119 k^{1.231} \quad (1)$$

$$k = (\Delta T_M - \Delta T) / \Delta T \quad (2)$$

where, ΔT_M is the temperature difference between the upper and lower probes during the stable period at night when there is no sap flow. Equations (1) and (2) have been applied to a wide variety of species and sites (Wullschlegel et al. 1998). However, because sapflow velocity varies with depth from the cambium, accurate estimation of sapflow requires an understanding of the radial profile of sap flux density along the length of the probe along with measurements of sapwood area to develop species-specific empirical equations (Zang et al. 1996; Wullschelger and King 2000; Nadezhina et al. 2002; Kubota et al. 2005). We failed to obtain these data and our estimates of sapflow velocity may, therefore, be inaccurate. However, assuming that the radial profile of sapflux does not change, our estimates would still be valid for comparing sap flow velocity of total sapwood area between the two stands relative to each other.

The calculated values of sapflow velocity were converted to m hr⁻¹ to facilitate comparison with daily values for whole-tree sap flux. We calculated whole-tree sap flux (m³ m⁻² day⁻¹) for clear days during the growing season by integrating the 30-min averages of sapflow velocity of each tree over a 24-hour period centered around 12:00.

We measured diurnal changes in leaf water potential at the crown surface of the sample trees in the field using the pressure chamber. Measurements were taken on five fully expanded, mature leaves on two of the five sample trees in each plot (tree numbers: 1, 3, 739, 765), six times during a 24-hour period on clear days (August 20, 30 and September 5, 2001) at predawn, morning, noon, afternoon, late afternoon and predawn of the following day. We estimated soil-to-leaf hydraulic conductance from the relationship between diurnal measurements of leaf water potential and the corresponding measurement of sapflow velocity. Assuming a steady-state water flow within the soil-plant-air continuum, we estimated soil-to-leaf hydraulic conductance (unscaled) from the slope of the relationship between leaf water potential and sapflow velocity using data from all measurements of leaf water potential in the field.

We obtained meteorological data on daily solar radiation and precipitation from the weather station at the Moshiri Field Station of Hokkaido University about 5 km from the study site as a reference for weather conditions during measurement. Takagi et al. (2001) give a detailed description of the meteorological measurement system.

Statistical analyses

For logistic reasons, the experimental unit in this study, the stand, could not be replicated. However, given homogeneous stand conditions and proximity of the two plots, we assumed that the experimental design was sufficient to infer that any observed differences between plots in tree-level characteristics would be due to the effect of dwarf bamboo removal.

For leaf water relations and shoot morphology, mean values were compared between the intact and removal plots using the two-sample *t*-test by pooling data from all sampled leaves in each plot upon finding no significant difference between sample trees.

For sapflow velocity and whole-tree sap flux, we considered measurements made on the four sample trees in each plot as replicates and compared plots using two-way ANOVA with plot and sampling time as the two main effects. We compared the slope of the relationship between leaf water potential and sapflow velocity between the plots using regression analysis.

Results

Shoot morphology

Shoot length (L_s) was longer in the removal plot than in the intact plot (Table 2, $t = 2.19$, $P = 0.03$). This was because shoots in the removal plot had more internodes ($t = 5.59$, $P < 0.01$), although internode lengths (L_i) were shorter than in the intact plot ($t = 7.95$, $P < 0.001$). Both leaf area (A_L) and leaf mass per area (LMA) were smaller in the removal plot than in the intact plot ($t = 8.36$, $P < 0.01$ and $t = 2.77$, $P < 0.01$, respectively).

Leaf water relations

Carbon isotope composition ($\delta^{13}\text{C}$) of leaves was similar for the two plots ($t = 0.482$, $P = 0.642$) indicating that integrated leaf water-use efficiency was not affected by presence/absence of dwarf bamboo. Pressure-volume measurements indicated that leaves in the removal plot lost turgor at higher leaf water potentials than leaves in the intact plot (Table 3). Both leaf water potential ($\Psi_{l, \text{tlp}}$) and relative water content at turgor loss point (RWC_{tlp}) were higher for leaves in the removal plot than for those in the intact plot ($t = 2.20$, $P = 0.03$ and $t = 2.50$, $P = 0.02$, respectively). Although not statistically significant, osmotic potential at saturation ($\Psi_{\pi, \text{sat}}$) tended to be higher for leaves in the removal plot ($t = 1.19$, $P = 0.29$). Bulk modulus of elasticity (ε) was lower for leaves in the removal plot than for leaves in the intact plot ($t = 2.71$, $P = 0.02$).

Field observations of whole-tree water use

Maximum sapflow velocity was similar between the two plots ranging from 0.118 – 0.261 m h⁻¹ in the removal plot and 0.167 – 0.247 m h⁻¹ in the intact plot during July – September ($F = 0.019$, $P = 0.089$). However, comparison of diurnal patterns of sapflow velocity indicated that, on clear days, sapflow velocity of trees in the removal plot began to increase earlier in the day and maximum sapflow velocity was sustained until later in the day than trees in the intact plot (Figure 1). For example, on a typical clear day during the early growing season (July 27), sap flow velocity began to increase in tree 739 in the removal plot at 7:00 indicating that transpiration had begun. In contrast, sapflow velocity did not start until 8:00 for tree 4 in the intact plot. On this day, sapflow velocity reached peak values (0.227–0.263 m hr⁻¹) at 12:30 in both plots. Subsequently, sapflow velocity of trees in the intact plot decreased more rapidly than that of trees in the removal plot. At 17:00, sapflow velocity of tree 4 of the intact plot had decreased to 0.265 (0.061 m hr⁻¹) of the daily maximum, whereas that of tree 739 in the removal plot had only decreased to 0.575 (0.151 m hr⁻¹). On a typical clear day during the mid-growing season (August 13), there was a one-hour difference between tree 739 and tree 4 in the time of initiation of sapflow. Sapflow velocity increased rapidly in the morning and reached peak values (0.227–0.266 m hr⁻¹) around 10:30–11:30 in both plots. Subsequently, sapflow velocity of trees in the intact plot decreased rapidly, whereas trees in the removal plot sustained relatively high sapflow velocity until about 17:00. The difference between the plots in the diurnal pattern of sapflow velocity was more marked than during the early growing season. On a typical clear day during the late growing season (September 6), the diurnal patterns of sapflow velocity was similar for trees in both plots.

On clear days during the growing season, whole-tree sap flux was higher for trees

in the removal plot than for trees in the intact plot (Figure 2, $F = 14.50$, $P < 0.01$). Average daily sap flux on clear days (the mean of all observations shown in Figure 3) was $1.37 \text{ m}^3 \text{ m}^{-2} \text{ day}^{-1}$ for the removal plot and $1.17 \text{ m}^3 \text{ m}^{-2} \text{ day}^{-1}$ for the intact plot. The largest differences between the two plots were observed on August 21 after a prolonged dry period and during September 2-3. On these dates, sap flux in the removal plot was $0.544\text{--}0.614 \text{ m}^3 \text{ m}^{-2} \text{ day}^{-1}$ greater than in the intact plot.

The relationship between sapflow velocity and leaf water potential indicated that soil-to-leaf hydraulic conductance was higher for trees in the removal plot than for those in the intact plot (Figure 3, $F = 6.57$, $P = 0.01$). Although sapflow velocity was similar between the plots at higher leaf water potentials, at leaf water potentials lower than -1.0 , sapflow velocity was higher for trees in the removal plot than for trees in the intact plot. Linear regression estimates of the slope of the relationship were -4.42 and -5.83 for trees 739 and 765 in the removal plot and -6.31 and -7.10 for trees 1 and 3 in the intact plot.

Discussion

Our results showed that removal of understory dwarf bamboo caused plastic responses in the water use of *B. ermanii* trees through changes in morphological and water-relations characteristics at the leaf, shoot and whole-tree levels. Despite similar upper-canopy light conditions, we observed differences in shoot and leaf morphology between the intact and removal plots. Removal of dwarf bamboo increased both shoot length and internode frequency of current-year shoots, suggesting that shoot elongation and leaf production decreased in the intact plot due to belowground competition with dwarf bamboo. Dwarf bamboo removal also decreased LMA. Under water-stressed conditions, both shoot elongation and leaf expansion are suppressed, resulting in shorter internodes, smaller leaf area and greater LMA (Hsiao 1973; Morgan 1984). In our study, however,

shoot length was shorter, but internode length was longer and both leaf area and LMA were greater in the intact plot. Because the study site receives moderate summer precipitation and below ground water from snow melt, the decrease in soil water availability in the intact plot may not have been so great as to cause water stress. Therefore, we inferred that the observed changes are the result of homeostatic responses to maintain constant leaf water-use efficiency under variable water availability (hydraulic homeostasis *sensu* Meinzer et al. 2004; Buckley and Roberts 2005).

Similar values of $\delta^{13}\text{C}$ in both the plots indicated that water-use efficiency of leaves during the growing season was not affected by the presence/absence of dwarf bamboo. This contradicts results obtained for *Pinus densiflora* where understory vegetation restricted water uptake by leaves and leaf water-use efficiency (inferred from $\delta^{13}\text{C}$ values) was higher than in the managed stand where understory vegetation was removed (Kume et al. 2003). The apparent homeostasis in leaf water-use efficiency of the *B. ermanii* trees in this study seems to have been realized through complementary changes in water-relations characteristics at the leaf and whole-tree levels.

In the intact plot, greater individual leaf area would increase leaf-level transpiration, while greater LMA would increase transpiration rate per leaf area. Pressure-volume measurements indicated that leaves of trees in the intact plot have greater water uptake ability than leaves in the removal plot. These changes would compensate for the reduction in leaf number (fewer internodes) to maintain constant water-use efficiency at the shoot level for trees in the intact plot. The increased water uptake ability of individual leaves in the intact plot would not be effective if leaf longevity was reduced. In general, leaf longevity is positively correlated with LMA (Kudo 1996; Wright et al. 2005). We observed that leaf longevity was not noticeably reduced in the intact plot (T. Kobayashi, unpublished data). Therefore, we inferred that increased water uptake ability of leaves in

the intact plot reflect hydraulic homeostasis to maintain constant leaf-level water-use efficiency in response to belowground competition with dwarf bamboo.

Although removal of dwarf bamboo increased the growth rate of overstory *B. ermanii*, stand-level LAI did not increase (A. Sumida, unpublished data), suggesting that tree-level leaf area might also be maintained. At the whole-tree level, trees in the removal plot maintained higher sapflow velocity than trees in the intact plot during the afternoon in early- to mid-summer when evapotranspirational demand was high. In addition, whole-tree sap flux and soil-to-leaf hydraulic conductance increased in response to dwarf bamboo removal. These results suggested that water use of trees in the intact plot was negatively affected by the presence of dwarf bamboo. Photosynthetic capacity of *B. ermanii* decreases with decreasing soil water availability due to decreasing leaf stomatal conductance (Kitao et al. 2003; Takahashi et al. 2004). Because the growing season in northern Japan is relatively short, maintaining transpiration and photosynthetic production on clear days during the early- to mid-summer may be an important factor controlling the annual growth rate of trees. Therefore, the restrictions on whole-tree water use during this period may be a determining factor for the lower growth rate of *B. ermanii* in the intact plot compared with the removal plot (Takahashi et al. 2003).

Both stomatal regulation of leaf water potential and whole-tree hydraulic conductance are important for maintaining stomatal conductance and hence photosynthetic capacity (Sperry et al. 2002; Tyree 2003). Under field conditions, stomatal conductance may be maintained by either 1) increasing the water uptake capacity of leaves by lowering the leaf water potential; or 2) increasing the water absorption capacity of the whole tree by increasing the soil-to-leaf hydraulic conductance (Ishida et al. 1992; Nardini et al. 1999). For trees in the intact plot, the greater water uptake ability of leaves may compensate for the lower soil-to-leaf hydraulic conductance and maintain leaf gas exchange capacity under

1 conditions of belowground competition with dwarf bamboo. Cienciala et al. (1994) found
2 that irrigation increased basal area increment, but water-use efficiency (dry matter
3 production per unit water transpired) was similar among irrigated, control and drought-
4 treated stands of 24-year-old *Picea abies* L. Karst. in Sweden.

5 We should also note that hydraulic sufficiency can also be maintained by
6 modifying the relationship between conducting area and leaf area (Sobrado 2003).
7 Although we did not measure sapwood area and tree-level leaf area in this study, it can be
8 conceived that trees in the intact plot had more conducting area relative to leaf area and
9 thus maintained similar water-use efficiency as trees in the removal plot. Tripathi et al.
10 (2005) found that fine-root proportion of *B. ermanii* in the soil increased in response to
11 dwarf bamboo removal, whereas fine-root biomass was not affected. This suggested that
12 relative biomass allocation to belowground was greater in the intact plot, which could have
13 also contributed to maintaining hydraulic sufficiency.

14 The allocation of water between overstory trees and understory vegetation is
15 critical to tree productivity in managed stands (Sands and Nambiar 1984; Oren et al. 1987;
16 Miller et al. 1998). Minimizing understory evapotranspiration is especially important
17 during early stages of stand growth when the potential growth rate of overstory trees is
18 high, but more open canopy conditions promote development of thick understory
19 vegetation. In addition to inhibiting regeneration and growth of overstory trees, dwarf
20 bamboo contributes a large fraction of the transpiring leaf area, biomass and photosynthetic
21 production in many Japanese forests (Sakai and Akiyama 2005). Removal of dwarf
22 bamboo in relatively young stands may greatly enhance productivity of overstory trees in
23 the long-term. The most recent study in this stand showed that removal of understory dwarf
24 bamboo increased net carbon gain and transpiration rates of overstory *B. ermanii* trees
25 (Kobayashi et al. 2006) indicating that dwarf bamboo removal is an effective silvicultural

1 prescription not only to improve the understory light environment for enhancing tree
2 seedling establishment, but also to enhance growth of overstory trees by relieving
3 belowground competition.

4
5 **Acknowledgments:** The authors thank Tabata A., Nabeshima E., Endo T. and Dolezal J.
6 for field assistance, and Tanaka Y., Moshiri Field Station technical staff for facilitating our
7 study. Many members and students of the Uryu Experimental Forest and Field Science
8 Center for Northern Biosphere, Hokkaido Univ. also supported our study. We thank the
9 Center for Ecological Research, Kyoto Univ. for use of facilities for isotope analysis. Dr.
10 Sakuta K. and Dr. Saito T. provided constructive comments and discussion during this
11 study. This research was funded by the COE project fund of the Ministry of Education,
12 Culture, Sports, Science and Technology, Japan and by the Grant for Joint Research
13 Program of ILTS, Hokkaido University.

References cited

- Buckley TN, Roberts DW (2005) How should leaf area, sapwood area and stomatal conductance vary with tree height to maximize growth? *Tree Physiol* 26:145-157
- Cienciala E, Lindroth A, Cermak J, Halgren JE, Kucera J (1994) The effects of water availability on transpiration, water potential and growth of *Picea abies* during a growing season. *J Hydrol* 155:57-71
- Clearwater MJ, Meinzer FC, Andrade JL, Goldstein G, Holbrook NM (1999) Potential errors in measurement of nonuniform sap flow using heat dissipation probes. *Tree Physiol* 19:681-687
- Farquhar GD, Ehleringer JR, Hubik KT (1989) Carbon isotope discrimination and photosynthesis. *Annu Rev Plant Physiol Plant Mol Biol* 40:503-537
- Franklin JF, Maeda T, Ohsumi Y, Matsui M, Yagi H (1979) Subalpine coniferous forests of central Honshu, Japan. *Ecol Monogr* 49:311-334
- Granier A (1987) Evaluation of transpiration in a Douglas-fir stand by means of sap flow measurements. *Tree Physiol* 3:309-319
- Hiura T, Sano J, Konno Y (1996) Age structure and response to fine-scale disturbances of *Abies sachalinensis*, *Picea jezoensis*, *Picea glehnii*, and *Betula ermanii* growing under influence of a dwarf bamboo understory in northern Japan. *Can J For Res* 26:289-297
- Hsiao TC (1973) Plant responses to water stress. *Ann Rev Plant Physiol* 24:519-570
- Ishida A, Yamamura Y, Hori Y (1992) Roles of leaf water potential and soil-to-leaf hydraulic conductance in water use by understory woody plants. *Ecol Res* 7:213-223
- Kikuzawa K (1982) Leaf survival of woody plants in deciduous broad-leaved forests. 1. Tall trees. *Can J Bot* 61:2133-2139
- Kitao M, Lei TT, Koike T, Tobita H, Maruyama Y (2003) Higher electron transport rate observed at low intercellular CO₂ concentration in long-term drought-acclimated leaves of Japanese mountain birch (*Betula ermanii*). *Physiol Plant* 118:406-413
- Kobayashi T, Miki N, Kato K, Kubo T, Nishimura S, Uemura S, Ono K, Sumida A, Ohta T, Hara T (2006) Understory removal increases carbon gain and transpiration in the overstory of birch (*Betula ermanii*) stands in northern Hokkaido, Japan: trends in leaf, shoot and canopy. In: *Proceedings of the international workshop on H₂O and CO₂ exchanges in Siberia (2nd international WS on C/H₂O/Energy balance and climate over boreal regions with special emphasis on eastern Siberia)*. Nagoya University, Nagoya, pp 19-22
- Kobayashi T, Shimano K, Muraoka H (2004) Effects of light availability on the carbon gain of beech (*Fagus crenata*) seedlings with reference to the density of dwarf bamboo (*Sasa kurilensis*) in an understory of Japan Sea type beech forest. *Plant Species Biol* 19:33-46
- Konno Y (2001) Feedback regulation of constant leaf standing crop in *Sasa tsuboiana* grasslands. *Ecol Res* 16:459-469

- 1 Kubota M, Tenhunen J, Zimmermann R, Schmidt M, Adiku S, Kakubari Y (2005)
- 2 Influences of environmental factors on the radial profiles of sap flux density in
- 3 *Fagus crenata* growing at different elevations in the Naeba Mountains, Japan. Tree
- 4 Physiol 25:537-548
- 5 Kudo G (1996) Intraspecific variation of leaf traits in several deciduous species in relation
- 6 to length of growing season. EcoScience 3:483-489
- 7 Kume A et al. (2003) Effects of understory vegetation on the ecophysiological
- 8 characteristics of an overstory pine, *Pinus densiflora*. For Ecol Manage 176:195-
- 9 203
- 10 Maruyama Y, Morikawa Y (1983) Measurement of the water relations characteristics of
- 11 leaves -the pressure-volume technique-. J Jpn For Soc 65:23-28 [In Japanese]
- 12 Meinzer FC, Woodruff DR, Shaw DC (2004) Integrated responses of hydraulic
- 13 architecture, water and carbon relations of western hemlock to dwarf mistletoe
- 14 infection. Plant Cell Environ 27:937-946
- 15 Miller BJ, Clinton PW, Buchan GD, Robson AB (1998) Transpiration rates and canopy
- 16 conductance of *Pinus radiata* growing with different pasture understories in
- 17 agroforestry systems. Tree Physiol 18:575-582
- 18 Morgan JM (1984) Osmoregulation and water stress in higher plants. Annual Review of
- 19 Plant Physiol 35:299-319
- 20 Nadezhina N, Cermak J, Ceulemans R (2002) Radial patterns of sap flow in woody stems
- 21 of dominant and understory species: scaling errors associated with positioning of
- 22 sensors. Tree Physiol 22:907-918
- 23 Nagai M, Yoshida T (2006) Variation in understory structure and plant species diversity
- 24 influenced by silvicultural treatments among 21- to 26-year-old *Picea glehnii*
- 25 plantations. J For Res 11:1-10
- 26 Nakashizuka T (1988) Regeneration of beech (*Fagus crenata*) after simultaneous death of
- 27 undergrowing dwarf bamboo (*Sasa kurilensis*). Ecol Res 3:21-35
- 28 Nardini A, Lo Guillo MA, Salleo S (1999) Competitive strategies for water availability in
- 29 two Mediterranean *Quercus* species. Plant Cell Environ 22:109-116
- 30 Noguchi M, Yoshida T (2005) Factors influencing the distribution of two co-occurring
- 31 dwarf bamboo species (*Sasa kurilensis* and *S. senanensis*) in a conifer-broadleaved
- 32 mixed stand in northern Hokkaido. Ecol Res 20:25-30
- 33 Oren R, Waring RH, Stafford SG, Barret JW (1987) Twenty-four years of Ponderosa pine
- 34 growth in relation to canopy leaf area and understory competition. For Sci 33:538-
- 35 547
- 36 Oshima Y (1961) Ecological studies of *Sasa* communities I. Productive structure of some
- 37 of the *Sasa* communities in Japan. Bot Mag Tokyo 74:199-210
- 38 Oshima Y (1962) Ecological studies of *Sasa* communities V. Influence of light intensity,
- 39 snow depth, and temperature upon the development of *Sasa kurilensis* community.
- 40 Bot Mag Tokyo 75:43-48

- Riegel GM, Miller RF, Krueger WC (1992) Competition for resources between understory vegetation and overstory *Pinus ponderosa* in northeastern Oregon. *Ecol Appl* 2:71-85
- Sakai T, Akiyama T (2005) Quantifying the spatio-temporal variability of net primary production of the understory species, *Sasa senanensis*, using multipoint measuring techniques. *Agric For Meteorol* 134:60-69
- Sakamoto K, Ishii A, Nishimoto T, Miki T, Nishikawa K (2004) Restoration of pine forests with pine wilt disease by removal of understory and A₀ horizon on the forest floor. *J Jpn Soc Reveget Tech* 30:110-115
- Samuelson LJ, McLemore PC, Somers GL (2003) Relationship between foliar $\delta^{13}\text{C}$ and hydraulic pathway length in *Pinus palustris*. *For Sci* 49:790-798
- Sands R, Nambiar EKS (1984) Water relations of *Pinus radiata* in competition with weeds. *Can J For Res* 14:233-237
- Sobrado MA (2003) Hydraulic characteristics and leaf water use efficiency in trees from tropical montane habitats. *Trees* 17:400-406
- Sperry JS, Hacke UG, Oren R, Comstock JP (2002) Water deficits and hydraulic limits to leaf water supply. *Plant Cell Environ* 25:251-263
- Takagi K et al. (2001) Meteorological characteristics of northern Hokkaido University Forests during 1995-1999. *Research Bulletin of the Hokkaido University Forests* 58:29-36 [in Japanese with English summary]
- Takahashi K (1997) Regeneration and coexistence of two subalpine conifer species in relation to dwarf bamboo in the understory. *J Veg Sci* 8:529-536
- Takahashi K, Matsuki S, Uemura S, Hara T (2004) Variations in the photosynthetic rate of *Betula ermanii* in relation to soil water potential. *Veg Sci* 21:103-108
- Takahashi K, Uemura S, Hara T (2002) Effect of understory dwarf bamboo on seasonal change in soil temperature in a *Betula ermanii* forest, northern Japan. *Eurasian J For Res* 5:49-53
- Takahashi K, Uemura S, Suzuki J-I, Hara T (2003) Effects of understory dwarf bamboo removal on soil water and the growth of overstory trees in a dense secondary *Betula ermanii* forest, northern Japan. *Ecol Res* 18:767-774
- Taylor AH, Qin Z (1988) Regeneration patterns in old-growth *Abies-Betula* forests in the Wolong Natural Reserve, Sichuan, China. *J Ecol* 76:1204-1218
- Tripathi SK et al. (2006) The effects of understorey dwarf bamboo (*Sasa kurilensis*) removal on soil fertility in a *Betula ermanii* forest of northern Japan. *Ecol Res* 21:315-320
- Tripathi SK, Sumida A, Shibata H, Uemura S, Hara T (2005) Growth and substrate quality of fine root and soil nitrogen availability in a young *Betula ermanii* forest of northern Japan: Effects of the removal of understory dwarf bamboo (*Sasa kurilensis*). *For Ecol Manage* 212:278-290
- Tyree MT (2003) Hydraulic limits on tree performance: transpiration, carbon gain and growth of trees. *Trees* 17:95-100

- 1 Tyree MT, Hammel HT (1972) The measurement of the turgor pressure and the water
2 relations of plants by the pressure-bomb technique. J Exp Bot 23:267-282
- 3 Umeki K (2003) The regeneration of natural forests in Hokkaido, northern Japan: results of
4 scarification and problems remaining to be solved. J Jpn For Soc 85:246-251 [in
5 Japanese with English abstract]
- 6 Wright IJ et al. (2005) Modulation of leaf economic traits and trait relationships by climate.
7 Global Ecol Biogeogr 14:411-421
- 8 Wullschelger SD, King AW (2000) Radial variation in sap velocity as a function of stem
9 diameter and sapwood thickness in yellow-poplar trees. Tree Physiol 20:511-518
- 10 Wullschleger SD, Meinzer FC, Vertessy RA (1998) A review of whole-plant water use
11 studies in trees. Tree Physiol 18:499-512
- 12 Yoshida T, Iga Y, Ozawa M, Noguchi M, Shibata H (2005) Factors influencing early
13 vegetation establishment following soil scarification in a mixed forest in northern
14 Japan. Can J For Res 35:175-188
- 15 Zang D, Beadle CL, White DA (1996) Variation of sapflow velocity in *Eucalyptus*
16 *globulus* with position in sapwood and use of a correction coefficient. Tree Physiol
17 16:697-703
- 18
- 19

- 1 Table 1. Structural properties of the sample trees of *Betula ermanii* growing with and
 2 without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and
 3 removal plot, respectively).

Tree	Height (m)	DBH (cm)
Intact plot		
1*	11.5	11.0
2*	10.7	10.1
3*	11.7	9.6
4*	10.9	10.1
5	11.4	8.6
Removal plot		
739*	11.2	8.6
738*	12.2	10.9
737*	12.7	12.5
765*	12.3	11.3
811	11.6	8.0

- 4 * trees used for sapflow measurement.

Table 2. Comparison of shoot and leaf morphology of *Betula ermanii* trees growing with and without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and removal plot, respectively). Mean values $\pm$ one standard deviation are shown for 30 current-year shoots sampled from six trees in each plot.

Treatment	L_s (mm)	N_i	L_i (mm)	A_L (cm ²)	LMA (g m ⁻²)
Intact plot	229.8 $\pm$ 94.1	7.1 $\pm$ 2.2	45.6 $\pm$ 10.7	42.8 $\pm$ 9.8	85.5 $\pm$ 10.5
Removal plot	275.2 $\pm$ 66.9*	9.7 $\pm$ 1.4*	36.3 $\pm$ 9.2*	34.9 $\pm$ 7.6*	80.5 $\pm$ 12.7*

L_s : length of current-year shoot

N_i : number of internodes per current-year shoot

L_i : internode length

A_L : leaf area

LMA: leaf mass per area

*: Significantly different from intact plot (Two-sample *t*-test, $P < 0.05$).

Table 3. Carbon isotope composition and leaf water-relations characteristics of *Betula ermanii* trees growing with and without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and removal plot, respectively). Mean values $\pm$ one standard deviation are shown for each plot.

Treatment	$\delta^{13}\text{C}$ (‰)	RWC_{tlp}	$\Psi_{\text{l, tlp}}$ (MPa)	$\Psi_{\pi, \text{sat}}$ (MPa)	ε (MPa)
Intact plot	-28.50 ± 0.68	0.83 ± 0.013	-2.17 ± 0.09	-1.25 ± 0.13	14.73 ± 1.36
Removal plot	-28.25 ± 0.71	$0.85 \pm 0.011^*$	$-2.02 \pm 0.10^*$	-1.16 ± 0.08	$11.33 \pm 2.11^*$

$\delta^{13}\text{C}$: $[(R_{\text{sample}} / R_{\text{standard}}) - 1] \times 1000$, where R is the $^{13}\text{C}/^{12}\text{C}$ ratio.

RWC_{tlp} : relative water content at turgor loss point,

$\Psi_{\text{l, tlp}}$: leaf water potential at turgor loss point,

$\Psi_{\pi, \text{sat}}$: osmotic potential at saturation

ε : bulk modulus of elasticity

*: Significantly different from intact plot (Two-sample t -test, $P < 0.05$).

Figure captions

Figure 1. Diurnal patterns of relative sapflow velocity (sapflow velocity relative to the daily maximum value for each tree) of *Betula ermanii* trees growing with and without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and removal plot, respectively). Typical diurnal patterns on clear days in July, August and September are shown for two of the four sample trees in each plot: intact plot (trees 2 and 4, thick lines) and removal plot (trees 739 and 765, thin lines). Diurnal changes in temperature (thin line) and solar radiation (thick line) are shown above.

Figure 2. Whole-tree sap flux ($\text{m}^3 \text{ m}^{-2} \text{ d}^{-1}$) of *Betula ermanii* trees growing with and without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and removal plot, respectively). Daily sap flux on clear days during the growing season were calculated for each plot (intact plot: filled symbols, removal plot: open symbols) based on measurements of total daily sapflow for the four sample trees in each plot. Error bars indicate one standard deviation. Measurements for the intact plot in early July were lost due to mechanical failure. Bars show daily precipitation measured at the weather station near the study site.

Figure 3. Relationship between sapflow velocity and leaf water potential of *Betula ermanii* trees growing with and without understory dwarf bamboo (*Sasa kurilensis*) in northern Japan (intact plot and removal plot, respectively). Common symbols indicate measurements from the same tree: intact plot (filled symbols and thick lines), removal plot (open symbols and thin lines)

Figure1

Figure2

Figure3