Title	Gene cloning of cold-adapted isocitrate lyase from a psychrophilic bacterium, <i>Colwellia psychrerythraea</i> , and analysis of amino acid residues involved in cold adaptation of this enzyme
Author(s)	Sato, Yuhya; Watanabe, Seiya; Yamaoka, Naoto; Takada, Yasuhiro
Citation	Extremophiles, 12(1), 107-117 https://doi.org/10.1007/s00792-007-0115-9
Issue Date	2008-01
Doc URL	http://hdl.handle.net/2115/32824
Rights	The original publication is available at www.springerlink.com
Type	article (author version)
File Information	CpICL2.pdf

[Instructions for use](#)

**Gene cloning of cold-adapted isocitrate lyase from a psychrophilic
bacterium, *Colwellia psychrerythraea*, and analysis of amino acid residues
involved in cold adaptation of this enzyme**

5

Yuhya Sato • Seiya Watanabe • Naoto Yamaoka • Yasuhiro Takada

10

15 Y. Sato · S. Watanabe · N. Yamaoka · Y. Takada (✉)
Division of Biological Sciences, Graduate School of Science, Hokkaido
University, Kita 10-jo Nishi 8-chome, Kita-ku, Sapporo 060-0810, Japan
E-mail: ytaka@sci.hokudai.ac.jp
Tel.: +81-11-706-2742
20 Fax: +81-11-706-2742

S. Watanabe

Institute of Advanced Energy, Kyoto University, Uji, Kyoto 611-0011, Japan

25 N. Yamaoka

Laboratory of Plant Pathology, Faculty of Agriculture, Ehime University,
Matsuyama 790-8566, Japan

Abstract The gene (*icl*) encoding cold-adapted isocitrate lyase (ICL) of a psychrophilic bacterium, *Colwellia psychrerythraea*, was cloned and sequenced. Open reading frame of the gene was 1,587 bp in length and corresponded to a polypeptide composed of 528 amino acids. The deduced amino acid sequence showed high homology with that of cold-adapted ICL from other psychrophilic bacterium, *Colwellia maris* (88% identity), but the sequential homology with that of the *Escherichia coli* ICL was low (28% identity). Primer extension analysis revealed that transcriptional start site for the *C. psychrerythraea icl* gene was guanine, located at 87 bases upstream of translational initiation codon. The expression of this gene in the cells of an *E. coli* mutant defective in ICL was induced by not only low temperature but also acetate. However, *cis*-acting elements for cold-inducible expression known in the several other bacterial genes were absent in the promoter region of the *C. psychrerythraea icl* gene. The substitution of Ala214 for Ser in the *C. psychrerythraea* ICL introduced by point mutation resulted in the increased thermostability and lowering of the specific activity at low temperature, indicating that Ala214 is important for psychrophilic properties of this enzyme.

Key words Psychrophilic bacterium • *Colwellia psychrerythraea* • Isocitrate lyase • Cold-adapted enzyme • Cold-inducible gene expression

Introduction

Temperature, especially low temperature, has important effects on growth of living organisms and defines their growth limits because the catalytic activities of enzymes in the cells are strongly dependent on temperature. Nevertheless, microorganisms called as psychophilic bacteria can survive and live even in permanently cold environment on Earth. The cold-adapted enzymes, which exhibit higher catalytic activity at low temperature and lower thermostability than the counterparts from mesophilic and thermophilic bacteria, are one of critical mechanisms for environmental adaptation of these bacteria because such enzymes can compensate the decrease of catalytic function at low temperature. It has been reported that these common properties of cold-adapted enzymes are attributable to their pronounced structural flexibility, which makes them possible to accomplish easy binding of the substrates to their active sites at low temperature and rapid conformational changes for the catalysis without large energetic loss (Gerday et al. 1997; Fields and Somero 1998; Geolette et al. 2003; Cavicchioli et al. 2002).

Isocitrate lyase (ICL, EC 4.1.3.1) catalyses the cleavage of isocitrate to glyoxylate and succinate, the first step in the glyoxylate cycle designated by Kornberg and Krebs (1957). Then, the second enzyme of this cycle, malate synthase, serves the conversion of glyoxylate and acetyl-CoA to malate. Thus, the glyoxylate cycle can bypass several reactions for the formation of succinate and malate from isocitrate in the TCA cycle and is known to be important for the metabolism of acetate and fatty acids in plants and microorganisms (Kornberg 1966; Vanni et al. 1990; Cozzzone 1998). Furthermore, ICL is known to be highly induced when microorganisms are grown on acetate (Vanni et al. 1990).

It has been reported that ICL (*CpICL*) of a psychophilic bacteria, *Colwellia*

psychrerythraea (D'Aoust and Kushner 1972; Deming et al. 1988), exhibits several features as a cold-adapted enzyme, as well as the corresponding enzyme (CmICL) from *Colwellia maris* (Takada et al. 1979; Yumoto et al. 1998), another psychrophilic bacterial species of the same genus (Watanabe et al. 2001, 2002a, 5 b). Namely, the CpICL showed maximum activity at 25°C and completely inactivated even by brief incubation for 2 min at 30°C (Watanabe et al. 2002b). Furthermore, the maximum catalytic efficiency of this enzyme was observed at 20°C. Preliminary experiments revealed that the expression of gene encoding this enzyme may be induced by low temperature.

10 In this study, we cloned the gene encoding the cold-adapted CpICL and examined effect of sodium acetate and temperature on the gene expression. In addition, to obtain information on the molecular mechanisms for high catalytic function of the cold-adapted ICL at low temperature, several point mutations were introduced into the *icl* gene and several properties of the mutated enzymes 15 were examined.

Materials and methods

Bacteria, plasmids and culture conditions

20 The psychrophilic *Colwellia psychrerythraea* NRC 1004 (D'Aoust and Kushner 1972; Deming et al. 1988) was grown at 15°C with vigorous shaking in a nutrient medium consisting of 1% meat extract, 40 mM MgCl₂, 3% NaCl and 25 25 mM sodium acetate as a carbon source. *E. coli* XL1-Blue (Stratagene) and TOP10 (Invitrogen) were used for the propagation of plasmids and the

production of His-tagged recombinant ICL proteins from expression vectors, respectively. A mutant defective in ICL of *E. coli*, ACA421 (Watanabe et al. 2002a), and its parent strain, KM22 (a $\Delta recBCD$ mutant; Kenan 1998), were used in the experiments for *icl* gene expression. Unless otherwise stated, these
5 *E. coli* strains were grown at 37°C with vigorous shaking in Luria-Bertani (LB) medium (Sambrook and Russell 2001). Plasmids pBluscript KS(+) (pBS; Stratagene) and pTrcHisB (Invitrogen) were used as vectors for cloning the *icl* gene and conferring N-terminal (His)₆-tag on the expressed proteins, respectively.

10

Cloning of the *C. psychrerythraea icl* gene

Genomic PCR was performed to obtain a nucleotide probe for *icl* gene encoding
15 the *CpICL*. Upstream primer, 5'-TCNAAYTAYCARAGYGCNATHGARGC-3', was designed from SNYQSAIEA, the sequence between the first and ninth amino acid residues from the N-terminal of *CpICL* (Watanabe et al. 2002b). Downstream primer, 5'-CCRTCYTGRTGNCCRCAYTGYTT-3', was designed from KQCGHQDG, the amino acid sequence of the *CmICL* corresponding to
20 highly conserved regions of the *icl* genes cloned from various organisms (Watanabe et al. 2002a). Chromosomal DNA of *C. psychrerythraea* was purified as described previously (Ishii et al. 1993). Amplification was carried out for 30 cycles in a DNA thermal cycler 2400 (Perkin-Elmer) in a 50 µl of reaction mixture containing 1.4 µg genomic DNA, 10 pmol each upstream and
25 downstream primer and 1 unit KOD-plus DNA polymerase (Toyobo) in a buffer system prepared by the manufacturer. Cycling conditions were as follows: denaturation at 94°C for 15 s, annealing at 50°C for 30 s and extension at 68°C

for 1 min, for each of 30 cycles. PCR products with a predicted length of about 700 bp were purified and ligated to the *Sma*I site of pBS. Nucleotide sequences of the PCR products were determined by using a BigDye terminator Cycle Sequencing FS Ready Reaction kit (Applied Biosystems) with an ABI
5 PRISM Genetic Analyzer 310. Plasmid carrying the 668 bp PCR product with a nucleotide sequence similar to the *C. maris icl* gene was termed pCP10. This insert DNA was amplified by PCR with pCP10 as template DNA and was used as probe for genomic Southern and colony hybridizations.

Genomic Southern hybridization was carried out as described previously
10 (Watanabe et al. 2002b) with some modifications. The chromosomal DNA of *C. psychrerythraea* digested with several restriction enzymes was separated by 1% agarose gel electrophoresis and the DNA fragments on the gel were transferred onto nylon membrane, Hybond-N⁺ (Amersham Pharmacia Biotech). Hybridization was carried out overnight at 60°C with the probe labeled with
15 [α -³²P]dCTP and a random primer labeling kit (Takara). The blotted membrane was washed successively in 2×SSC (1×SSC is composed of 15 mM sodium citrate pH 7.0 and 0.15 M NaCl) containing 0.1% (w/v) SDS for 10 min at room temperature and then twice with the same solution for 20 min at 60°C. About 3 kbp of *Spe*I fragments hybridized with the probe were cloned into pBS and the
20 resultant plasmids were transformed into *E. coli* XL1-Blue. Colony hybridization was then carried out by using the same probe as Southern hybridization. The blotted membranes were washed as described above. A colony with strongest signal was selected, and the plasmid in this colony, termed pCPB49, was propagated and purified.

25 The nucleotide sequence of the insert DNA in the pCPB49 was determined in both directions by using appropriate primers according to the method described above and was analyzed with the Genetyx computer program (Software

Development Co.). The nucleotide sequence of the *C. psychrerythraea icl* gene has been deposited in the DDBJ database under accession no. AB174852.

5 Western blot analysis

After SDS-PAGE of the purified ICLs and the cell-free extracts of the *E. coli* cells on 10% polyacrylamide gels (Laemmli 1970), the proteins were transferred onto a nitrocellulose membrane, Hybond-ECL (Amersham Bioscience).

10 Western blot analysis was carried out with the ECL Western blotting detection system (Amersham Bioscience) and rabbit antibody against the purified *CmICL* (Watanabe et al. 2002a).

15 Primer extension analysis

Primer extension analysis was performed as described by Sambrook and Russell (2001) with the following modifications. A synthetic 29-mer oligonucleotide, 5'-TTGAATTTGTTCTGTGCACGCATACGGGC-3', complementary to the
20 sequence between +91 and +119 from the translation start codon of the *C. psychrerythraea icl* gene, was used as the primer. The 5'-terminal of the primer was labeled with [γ -³²P]ATP and T4 polynucleotide kinase. Total RNA (20-50 μ g) isolated from *C. psychrerythraea* was incubated overnight at 42°C with the labeled primer in the hybridization buffer. The extension was done
25 with a reverse transcriptase (ReverTra Ace, Toyobo). The reaction products were analyzed on a 6% polyacrylamide sequencing gel with a sequencing ladder of the *C. psychrerythraea icl* gene. Sequencing was performed with the T7

Sequencing Kit (USB) with the same primer as the primer extension reaction.

Construction of expression vector for His-tagged ICL of *C. psychrerythraea* and 5 site-directed mutagenesis

By PCR, restriction sites for *Bam*HI and *Sac*I were introduced at 5'- and 3'-terminals of the ORF of *C. psychrerythraea icl* gene, respectively. For this purpose, the following two primers were synthesized: forward primer
10 5'-gcgcggatccGTCTAATTATCAGAG-3' and reverse primer 5'-gcgcgagctcTTAACTGA
AACTGA ACTGGTTAG-3'. Small letters indicate additional bases for introducing the underlined cleavage sites of the respective restriction enzymes. The amplification was carried out for 30 cycles in the DNA thermal cycler in a reaction mixture (50 µl) containing 180 ng pCPB49 as template DNA, 10 pmol
15 each forward and reverse primer and 1 unit KOD-plus DNA polymerase in a buffer system prepared by the manufacturer. Cycling conditions were as follows: denaturation at 94°C for 15 s, annealing at 50°C for 30 s, and extension at 68°C for 2 min. After the digestion with *Bam*HI and *Sac*I, the PCR product was ligated to the *Bam*HI-*Sac*I site of pTrcHisB to obtain plasmid pHis-CpWT.
20 Precise insertion of the *C. psychrerythraea icl* gene into the plasmid vector was verified by nucleotide sequencing.

Site-directed mutagenesis was carried out by a standard PCR method with synthetic primers (Table 1). The PCR was performed as described above. Resultant PCR products were ligated into pTrcHisB to obtain the mutated
25 plasmids, pHisQ47K, pHisA214S, pHisA231E, pHisF333L and pHisA341N (pHisQ47K denotes pHis-CpWT introduced the substitutional mutation of Gln47 by Lys in the CpICL). The introduced mutations were checked by

nucleotide sequencing the relevant regions of plasmids as described above.

Overexpression and purification of His-tagged ICLs

5

E. coli TOP10 transformed with pHis-*Ec*WT (plasmid for His-tagged *E. coli* ICL), pHis-*Cp*WT or the plasmids carrying the mutated *icl* genes was grown at 37°C in Super broth medium (12 g tryptone, 24 g yeast extract, 5 ml glycerol, 3.81 g KH₂PO₄ and 12.5 g K₂HPO₄ per liter (pH 7.0)) containing 50 mg/l
10 ampicillin until OD₆₀₀ of the culture reached 0.6. The cultures were rapidly cooled on ice and were further incubated for 16-24 h at 15°C after the addition of 1 mM isopropyl-β-thiogalactopyranoside (IPTG) to induce the expression of His-tagged ICL proteins. Because of the marked thermolability of *Cp*ICL (Watanabe et al. 2002b), the induction of gene expression by IPTG were carried
15 out at 15°C. As described previously (Watanabe et al. 2004), cell-free extracts were prepared and His-tagged ICLs were then purified by column chromatography of Ni-NTA agarose (Qiagen). The purified His-tagged ICLs were concentrated with polyethylene glycol #20,000 and dialyzed against 20 mM potassium phosphate (pH 6.85), containing 2 mM MgCl₂, 4 mM
20 DL-isocitrate, 10% (v/v) glycerol and 1 mM dithiothreitol (DTT). All His-tagged ICLs were stocked at -35°C until use.

Enzyme assay

25

The ICL activity was assayed at pH 6.85 by measuring the increase of absorbance at 324 nm due to the formation of glyoxyl-phenylhydrazone as

described previously (Watanabe et al. 2001). The reaction mixture contained 66.7 mM potassium phosphate buffer (pH 6.85), 5 mM MgCl₂, 3.3 mM phenylhydrazine-HCl, 2 mM cysteine-HCl, 1.67 mM sodium isocitrate and an appropriate amount of enzyme in a final volume of 3 ml. The reaction was
5 started by the addition of enzyme (50 µl) into the mixture previously equilibrated at assay temperatures. One unit of the activity was defined as the formation of 1 µmol of product per min.

10 Molecular modeling of *C. psychrerythraea* ICL

The three-dimensional structural model of CpICL was built using ICL from *E. coli* (PDB No. 1IGW) as a homology model with a homology modeling program (www.cbs.dtu.dk/services/CPHmodels/index.php).

15

Results

20 Cloning of the *C. psychrerythraea icl* gene

By genomic PCR described in Materials and Methods, the 668 bp DNA fragment was obtained and then sequenced. The deduced amino acid sequence was consistent with the partial amino acid sequence of *C. psychrerythraea* ICL
25 protein reported previously (Watanabe et al. 2002b). Genomic Southern blot analysis using this PCR product as probe revealed that about 3 kbp of single fragment of the *C. psychrerythraea* genomic DNA digested with *Spe*I hybridizes

with the probe. Therefore, for cloning of the *icl* gene, the genomic library was constructed with pBS carrying the similar size of *Spe*I-digests of the *C. psychrerythraea* genomic DNA and was screened with the same probe. A clone with strongest signal was selected and the plasmid within the clone
5 (designated as pCPB49) was purified (Fig. 1).

Nucleotide and deduced amino acid sequences of the *C. psychrerythraea icl* gene

Both strands of the insert DNA subcloned in the plasmid pCPB49 were sequenced (Fig. 2). This insert contained an ORF with full length of 1,587 bp, and a putative ribosome-binding site (Shine and Dalgarno 1974), GGAG, was found 7-10 bases upstream of the ATG codon. The ORF encodes a polypeptide
15 of 528 amino acids, and the molecular mass was calculated as 58,131 Da. This value was compatible with that of the purified ICL from this bacterium estimated by gel filtration (homotetramer of the 60 kDa subunit) (Watanabe et al. 2002b). Previously, we reported that bacterial ICLs can be classified into two phylogenetic groups, subfamilies 1 and 3, based on their amino acid sequences
20 (Watanabe et al. 2004). The deduced amino acid sequence showed 24-29% identities with subfamily 1 ICLs from many bacteria, including the *E. coli* ICL (28% identity). On the other hand, sequence identities with subfamily 3 ICLs shown in Fig. 3 were 70% or above, except for the *Hyphomicrobium methylovorum* ICL (54% identity). In particular, the sequence identity to the
25 ICL of *C. maris*, another psychrophilic bacterium of the same genus *Colwellia*, attained to 88%.

Expression of the *C. psychrerythraea icl* gene

Expression of the *C. psychrerythraea icl* gene in the *icl*-defective mutant of *E. coli*, ACA421, was examined under several conditions by Western blot analysis. As Watanabe et al. (2002b) reported previously, antibody against the purified *CmICL* cross-reacted with the ICL protein purified from the *C. psychrerythraea* cells, but not with the *E. coli* ICL. Whereas no protein cross-reactive with this antibody was present in crude extract of the *E. coli* ACA421 cells transformed with a control vector, pBS, a major cross-reactive protein band was detected at the same position as the purified *CpICL* in crude extract of the *E. coli* transformant cells carrying pCPB49 grown at 15 and 25°C on LB medium. These results demonstrate that the *C. psychrerythraea icl* gene is cloned in this plasmid. ICL is known to be highly induced in microorganisms grown on acetate (Vanni et al. 1990). When the *E. coli* transformant cells carrying pCPB49 were grown at 15 and 25°C on LB medium supplemented by 50 mM sodium acetate, the gene expression was induced. Furthermore, irrespective of the supplementation of sodium acetate in growth medium, the expression level of this gene in the *E. coli* mutant grown at 15°C was larger than that grown at 25°C, and no expression were detected when it was grown at 37°C. Similar results were obtained by the assay of ICL activity in the crude extract of *E. coli* ACA421 cells transformed with pCPB49 (Table 2). These results were in accord with those of Northern and Western blot analyses carried out using the intact cells of *C. psychrerythraea* (Watanabe et al. 2002b), indicating that the expression of *C. psychrerythraea icl* gene is not only acetate-inducible but also cold-inducible and that the insert DNA subcloned in the plasmid pCPB49 contains ORF of the *icl* gene and its complete promoter region.

Analysis of 5'-terminal region of the *C. psychrerythraea icl* mRNA

5 The 5'-end of the *C. psychrerythraea icl* mRNA was examined by primer extension analysis (Fig. 4). The result revealed that transcriptional start site of the *C. psychrerythraea icl* gene (TS site in Fig. 4) is guanine, located at 87 bases upstream of translational initiation codon. Furthermore, the putative -35 and -10 promoter elements, TTCCTA and TATTAT, were found 32 and 8 bases
10 upstream of the TS site, respectively (Fig. 2).

Site-directed mutagenesis of *CpICL* and properties of mutated ICLs

15 Cold-adapted enzymes should possess inherent characteristics on protein structure responsible for the common properties, high catalytic function at low temperature and marked thermolability, compared with mesophilic and/or thermophilic counterparts. The introduction of suitable mutation(s) to these enzymes is one of useful approaches for studying this problem. From multiple
20 alignment of amino acid sequences of subfamily 3 ICLs from several mesophilic and psychrophilic bacteria, several substitutions of amino acid residues common to the two cold-adapted ICLs, *CpICL* and *CmICL*, were found to exist in highly conserved regions of the ICL proteins (Fig. 3). Among them, Gln47 (the 47th Gln residue from N-terminal), Ala214, Ala231, Phe333 and Ala341 of the
25 *CpICL* (shown by stars in Fig. 3) were substituted for the conserved amino acid residues of other ICLs by site-directed mutagenesis as described in Materials and Methods.

To examine a possibility that His-tagging of the enzyme at the N-terminal alters its original properties, the activity of His-tagged *Cp*ICL (*Cp*-WT) was assayed at various temperatures (Fig. 5). The purified *Cp*-WT showed the maximum activity at 25°C and its *k*_{cat} value at this temperature was 28.5 s⁻¹.

5 These results were consistent with those of the native *Cp*ICL reported previously (25°C and 27.7 s⁻¹, respectively; Watanabe et al. 2002b). In addition, as well as the native *Cp*ICL and *Cm*ICL, the *Cp*-WT activity could not be measured exactly at 30°C because the enzyme was rapidly inactivated at this temperature even during the enzyme assay. These indicate that the His-tagging at the
10 N-terminal does not affect the activity and characteristics of *Cp*ICL.

The activities of recombinant *Cp*ICLs and His-tagged *E. coli* ICL (*Ec*-WT) at various temperatures were shown in Fig. 5. The ICL activity of F333L, which is *Cp*ICL substituted Phe333 by Leu, was not able to be exactly measured because of its very low activity, suggesting that Phe333 of the *Cp*ICL is
15 important for the catalytic function. Temperature dependence of ICL activities of Q47K and A341N was similar to that of *Cp*-WT. Furthermore, as well as *Cp*-WT, optimum temperature for the two enzyme activities was 25°C, and their exact activities could not be measured at 30°C because they were rapidly inactivated even during the enzyme assay at this temperature. At the optimum
20 temperature, the activity of Q47K (*k*_{cat} value, 26.6 s⁻¹) was comparable to that of *Cp*-WT, but A341N (13.3 s⁻¹) exhibited lower activity than *Cp*-WT. The ICL activities of A231E below 20°C were similar to those of A341N, but the former enzyme showed the maximum activity at 20°C. Further lowering of ICL activity at low temperatures was observed in A214S, but optimum
25 temperature for this enzyme activity increased by about 5°C, compared to that of *Cp*-WT. However, above 30°C, this enzyme was also rapidly inactivated. These results imply that Ala214 of the *Cp*ICL is involved in its high activity at

low temperatures.

To estimate whether the mutations of *CmICL* affect the thermal stability, the residual activities of wild and mutated enzymes after the incubation for various times at 25°C were examined (Fig. 6). A214S and A231E were more
5 thermostable and more thermolabile than *Cp*-WT, respectively.

Discussion

10 In this study, the *icl* gene encoding the cold-adapted ICL of *C. psychrerythraea* was cloned and sequenced. From expression analysis, DNA fragment cloned in the plasmid pCPB49 was confirmed to contain the ORF of this gene and full length of the promoter region, because the expression of this gene in the *E. coli* cells was induced by acetate and low temperature and this result was consistent
15 with that in the *C. psychrerythraea* cells reported previously (Watanabe et al. 2002b). The CCAAT sequence is present in the upstream region of the *C. maris icl* gene whose expression is induced by acetate and low temperature as well as the *C. psychrerythraea icl* gene (Watanabe et al. 2002a). This sequence has been reported to be *cis*-acting element for cold-inducible expression of
20 several genes including the *icd-II* encoding the cold-adapted isocitrate dehydrogenase isozyme II of *C. maris* (Qoronfleh et al. 1992; Sahara et al. 1999). Furthermore, several *cis*-acting elements for cold-inducible gene expression was found by promoter analysis of the genomic DNA of a Gram-negative psychrophilic bacterium, *Pseudomonas haloplanktis* strain
25 TAC125, which are about 20 bp of upstream AT-rich sequences adjacent to the -35 consensus element of promoters (Duilio et al. 2004). However, such sequences are absent in the upstream region of the *C. psychrerythraea icl* gene

despite its cold-inducible expression (Fig. 2). Therefore, it is suggested that other novel *cis*-element is involved in the cold-inducible expression of this gene or the stabilization of its mRNA at low temperature influences the gene expression as reported for *cspA* and *otsA/B* mRNA of *E. coli* (Tanabe et al. 1992; 5 Kandrór et al. 2002).

From the comparison of amino acid sequence with various bacterial ICLs, it was found that the *CpICL* belongs to subfamily 3, as well as the *CmICL* (Fig. 3). As reported previously (Watanabe et al. 2002a), six additional short insertions of 3-36 amino acids are scattered in the *CmICL*, compared to 10 subfamily 1 ICLs from *E. coli* and *Mycobacterium tuberculosis*. The same insertions were also present in the *CpICL*. In spite of such a sequential difference between the *CmICL* and *E. coli* ICL, many amino acid residues essential for catalytic function and for binding with substrates and divalent metal ion, which were identified in the *E. coli* ICL, are conserved in the 15 *CmICL* (Watanabe et al. 2002a). In this study, these amino acid residues of the *CmICL* were found to be completely conserved in the *CpICL*.

Multiple alignments of amino acid sequences of subfamily 3 ICLs and the putative ones from several bacteria, including the cold-adapted *CpICL* and *CmICL*, revealed that several substitutions of amino acid residues peculiar to 20 these two enzymes are present in the conserved regions of the ICL proteins. From the following reasons, five amino acid residues of Gln47, Ala214, Ala231, Phe333 and Ala341 were selected and were substituted by point mutations; Gln47, the corresponding amino acid residues of other ICLs are basic Lys or Arg; Ala214, in a three-dimensional model of the *CpICL* protein constructed 25 with the homology modeling program, this residue is located closely to TIM barrel structure involved in the binding of substrate (Britton et al. 2001), and the corresponding residue is more hydrophilic Ser in other ICLs; Ala231 and

Phe333, the corresponding residues are acidic Glu or Asp and less hydrophobic Leu in most of other ones, respectively; Ala341, the corresponding residues of other ones are Asn. Although the A214S mutation of *CpICL* led to lowering of the enzyme activity, the thermostability and optimum temperature for ICL activity of the mutated enzyme were elevated, compared to those of wild-type enzyme. These results indicate that Ala214 is involved in thermolability of the *CpICL* and its high activity at low temperatures. Possible hydrogen bonds of this residue predicted by a program SWISSPDB VIEWER were shown in Fig. 7. The mutated ICL, A214S, can form two hydrogen bonds between main chain of the substituted Ser214 and side chain of Gln119, and side chain of Ser214 and main chain of Gln119, whereas only one hydrogen bond is allowed between main chain of native Ala214 and side chain of Gln119 in the *Cp*-WT. Therefore, this may be responsible for the decrease of structural flexibility of the A214S and resultant increase of its thermostability. On the other hand, the decreased thermostability and lowering of catalytic activity were observed in the A231E, and Q47K and A341N, respectively. These three substitutions of amino acid residues have negative effects to thermostability and specific activity of *CpICL*, suggesting that these amino acid residues are also involved in the thermostability and catalytic activity of *CpICL*. In particular, the F333L mutation of *CpICL* resulted in almost complete loss of its catalytic function, implying that Phe333 might play an important role for catalytic function of this enzyme. Furthermore, various combinations of these amino acid substitutions may be able to bring some different effects on the psychrophilic properties of *CpICL*. On the other hand, it was reported that the surface of cold-adapted enzyme tends to have more nonpolar amino acid residue, in particular Ala, compared with the corresponding mesophilic and thermophilic ones (Siddiqui and Cavicchioli 2006; Sælensminde et al. 2007). Besides A214, A231 and

A341, several substitutions by Ala are found to be still present in the two cold-adapted *CpICL* and *CmICL* (Fig. 3). Some of such Ala residues might be responsible for the psychrophilic nature of these enzymes. Thus, further studies are planned to clarify the role of these residues of cold-adaptation and thermolability of the *CpICL*.

Acknowledgements We sincerely thank Dr. Isao Yumoto of the National Institute of Advanced Industrial Science and Technology for the donation of *C. psychrerythraea*.

References

- Britton KL, Abeysinghe IS, Baker PJ, Barynin V, Diehl P, Langridge SJ, McFadden BA, Sedelnikova SE, Stillman TJ, Weeradechapon K, Rice DW
5 (2001) The structure and domain organization of *Escherichia coli* isocitrate lyase. *Acta Cryst D* 57:1209-1218
- Cavicchioli R, Siddiqui KS, Andrews D, Sowers KR (2002) Low-temperature extremophiles and their applications. *Curr Opin Biotechnol* 13:253-261
- Cozzone AJ (1998) Regulation of acetate metabolism by protein
10 phosphorylation in *Escherichia coli*. *Annu Rev Microbiol* 52:127-164
- D'Aoust JY, Kushner DJ (1972) *Vibrio psychroerythrus* sp. n.: classification of the psychrophilic marine bacterium, NRC 1004. *J Bacteriol* 111:340-342
- Deming JW, Somers LK, Straube WL, Swartz DG, Macdonell MT (1988) Isolation of an obligately barophilic bacterium and description of a new genus,
15 *Colwellia* gen. nov. *Syst Appl Microbiol* 10:152-160
- Duilio A, Madonna S, Tutino ML, Pirozzi M, Sannia G, Marino G (2004) Promoters from a cold-adapted bacterium: definition of a consensus motif and molecular characterization of UP regulative elements. *Extremophiles* 8:125-132
- 20 Fields PA, Somero GN (1998) Hot spots in cold adaptation: localized increases in conformational flexibility in lactate dehydrogenase A4 orthologs of Antarctic notothenioid fishes. *Proc Natl Acad Sci U S A* 95:11476-11481
- Gerday C, Aittaleb M, Arpigny JL, Baise E, Chessa JP, Garsoux G, Petrescu I, Feller G (1997) Psychrophilic enzymes: a thermodynamic challenge. *Biochim*
25 *Biophys Acta* 1342:119-131
- Geolette D, Damien B, Blaise V, Depiereux E, Uversky VN, Gerday C, Feller G (2003) Structural and functional adaptation to extreme temperatures in

psychrophilic, mesophilic, and thermophilic DNA ligase. J Biol Chem 278: 37015-37023

Ishii A, Suzuki M, Sahara T, Takada Y, Sasaki S, Fukunaga N (1993) Gene encoding two isocitrate dehydrogenase isozymes of a psychrophilic bacterium, *Vibrio* sp. strain ABE-1. J Bacteriol 175:6873-6890

Kandror O, Deleon A, Goldberg AL (2002) Trehalose synthesis is induced upon exposure of *Escherichia coli* to cold and is essential for viability at low temperatures. Proc Natl Acad Sci U S A 99:9727-9732

Kenan C (1998) Use of bacteriophage lambda recombination function to promote gene replacement in *Escherichia coli*. J Bacteriol 180:2063-2071

Kornberg HL, Krebs HA (1957) Synthesis of cell constituents from C₂-units by a modified tricarboxylic acid cycle. Nature 179:988-991

Kornberg HL (1966) The role and control of the glyoxylate cycle in *Escherichia coli*. Biochem J 99:1-11

Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature 227:680-685

Qoronfleh MW, Debouck C, Keller J (1992) Identification and characterization of novel low-temperature-inducible promoters of *Escherichia coli*. J Bacteriol 174:7902-7909

Sælensminde G, Halskau Jr Ø, Helland R, Willassen NP, Jonassen I (2007) Structure-dependent relationships between growth temperature of prokaryotes and the amino acid frequency in their proteins. Extremophiles 11:585-596

Sahara T, Suzuki S, Tsuruha J, Takada Y, Fukunaga N (1999) *cis*-Acting elements responsible for low-temperature-inducible expression of the gene coding the thermolabile isocitrate dehydrogenase isozyme of a psychrophilic bacterium, *Vibrio* sp. Strain ABE-1. J Bacteriol 181:2602-2611

Sambrook J, Russell D (2001) Molecular Cloning: a laboratory manual, 3rd edn.

Cold Spring Harbor Laboratory, Cold Spring Harbor, NY

Shine J, Dalgarno L (1974) The 3'-terminal sequence of *Escherichia coli* 16S ribosomal RNA: complementarity to nonsense triplets and ribosome binding sites. *Proc Natl Acad Sci U S A* 71:1342-1346

5 Siddiqui KS, Cavicchioli R (2006) Cold-adapted enzymes. *Annu Rev Biochem* 75:403-433

Takada Y, Ochiai T, Okuyama H, Nishi K, Sasaki S (1979) An obligatory psychrophilic bacterium isolated on the Hokkaido coast. *J Gen Appl Microbiol* 25:11-19

10 Tanabe H, Goldstein J, Yang M, Inouye M (1992) Identification of the promoter region of the *Escherichia coli* major cold shock gene, *cspA*. *J Bacteriol* 174:3867-3873

Vanni P, Giachetti E, Pinzauti G, McFadden BA (1990) Comparative structure, function and regulation of isocitrate lyase, an important assimilatory enzyme.

15 *Comp Biochem Physiol* 95B:431-458

Watanabe S, Takada Y, Fukunaga N (2001) Purification and characterization of a cold-adapted isocitrate lyase and a malate synthase from *Colwellia maris*, a psychrophilic bacterium. *Biosci Biotech Biochem* 65:1095-1103

Watanabe S, Takada Y, Fukunaga N (2002a) The cold-inducible *icl* gene
20 encoding thermolabile isocitrate lyase of a psychrophilic bacterium, *Colwellia maris*. *Microbiology* 148:2579-2589

Watanabe S, Yamaoka N, Fukunaga N, Takada Y, (2002b) Purification and characterization of a cold-adapted isocitrate lyase and expression analysis of the cold-inducible isocitrate lyase gene from the psychrophilic bacterium
25 *Colwellia psychrerythraea*. *Extremophiles* 6:397-405

Watanabe S, Takada Y (2004) Amino acid residues involved in cold adaptation of isocitrate lyase from a psychrophilic bacterium, *Colwellia maris*.

Microbiology 150:3393-3403

Yumoto I, Kawasaki K, Iwata H, Matsuyama H, Okuyama H (1998)
Assignment of *Vibrio* sp. strain ABE-1 to *Colwellia maris* sp. nov., a new
psychrophilic bacterium. Int J Syst Bacteriol 48:1357-1362.

Table 1 Oligonucleotides used in site-directed mutagenesis

	Plasmid	Substitution		Primers
5	pHisQ47K	Gln47 → Lys	forward reverse	5'-GGTCTAGAGATAGCT <u>AAG</u> TACACTGCAG-3' 5'-CTGCAGTGTACTT <u>AGC</u> TATCTCTAGACC-3'
10	pHisA214S	Ala214 → Ser	forward reverse	5'-CGAAAACCAAGTAT <u>TCT</u> GATGAAAAGC-3' 5'-GCTTTTCATCAG <u>A</u> TACTTGGTTTTTCG-3'
	pHisA231E	Ala231 → Glu	forward reverse	5'-GTAACGTTCCTCATG <u>AAG</u> ATTTCCACTC-3' 5'-GAGTGGAAATC <u>TTC</u> ATGAGGAACAGTTAC-3'
15	pHisF333L	Phe333 → Leu	forward reverse	5'-GAGCGTTGTGTACTT <u>GAC</u> TGTATCGGTGC-3' 5'-GCACCGATACAGTCA <u>AGT</u> ACACAACGCTC-3'
	pHisA341N	Ala341 → Asn	forward reverse	5'-CGGTGCACTTAATA <u>AAT</u> GGCGCTGATTTAC-3' 5'-GTAAATCAGCGCC <u>ATT</u> ATTAAGTGCACCG-3'

The underlined letters indicate the codons of amino acid residues substituted by site-directed mutagenesis.

Table 2 ICL activity of *E. coli icl*-defective mutant transformed with the *C. psychrerythraea icl* gene

5	Growth medium	Growth temperature		
		37°C	25°C	15°C
	LB	1.40×10^{-3}	2.08×10^{-3}	6.07×10^{-3}
10	LB + acetate	3.72×10^{-3}	7.29×10^{-3}	1.70×10^{-2}

The values indicate ICL activity (unit/mg protein) in crude extract of *E. coli* ACA421 transformed with pCPB49. The enzyme activity was assayed at 25°C. The *E. coli* transformant was grown on LB medium supplemented with none (LB) or 50 mM sodium acetate (LB + acetate). No ICL activity was detected in the *E. coli* ACA421 cells grown at 37°C on LB or LB + acetate medium.

Legends to figures

Fig. 1 Physical and restriction map of the genomic DNA fragment containing the *C. psychrerythraea icl* gene. The black bar indicates the 668 bp DNA
5 fragment used as the probe for colony hybridization.

Fig. 2 Nucleotide and deduced amino acid sequences of the *C. psychrerythraea icl* gene. The *underlined sequences* are the putative –35 and –10 promoter elements. The *boxed nucleotides* are the putative
10 Shine-Dalgarno sequence. The guanine in black box indicates transcriptional start point (Fig. 5). The probable stem-loop structures located downstream of the translational stop codon are shown by *two opposite arrows*.

Fig. 3 Alignment of amino acid sequences of subfamily 3 ICLs from various
15 bacteria. Amino acid residues identical and similar to those of CpICL are shown as *white letters in black boxes* and *black letters in gray boxes*, respectively. Gaps in the alignments are indicated by dashes. C.m., *C. maris* (accession no. for the protein database, BAB62107), H.m., *Hyphomicrobium methylovorum* GM2 (BAA23678), and R.e., *Ralstonia. eutropha* (AAM18124).
20 Putative ICLs, R.m., *Ralstonia metallidurans* (ZP_00022497), R.p., *Rhodopseudomonas palustris* (ZP_00009157), B.j., *Bradyrhizobium japonicum* (NP_709695), N.a., *Novosphingobium aromaticivorans* (ZP_0093836), P.a., *Pseudomonas aeruginosa* PAO1 (NP_251324), A.v., *Azotobacter vinelandii* (ZP_00091825), and A.c., *Acinetobacter calcoaceticus* ADP1 (YP_045792).
25 *Stars on the sequences* indicate the positions of amino acid residues substituted by site-directed mutagenesis (see text).

Fig. 4 Primer extension analysis. Product of primer extension with a primer complementary to *icl* is shown in lane 1. Total RNA (35 µg) was isolated from the *C. psychrerythraea* cells grown at 15°C on nutrient medium containing 50 mM acetate and was used as the template. The sequence ladders are shown in lanes T, G, C and A.

Fig. 5 Effect of temperature on activities of wild and mutated ICLs. Symbols; *Cp*-WT (♦), Q47K (■), A214S (▲), A231E (×), A341N (▼) and *Ec*-WT (●). The *k_{cat}* values of ICLs were measured under standard assay condition.

Fig. 6 Thermostability of wild and mutated ICL activities. Symbols used in this figure are the same as Fig. 6. The purified enzymes were dialyzed at 4°C against 20 mM potassium phosphate (pH 6.85) containing 2 mM MgCl₂ and 1 mM DTT. After the incubation for the indicated time periods at 25°C, the dialyzed enzymes were withdrawn and immediately cooled for 10 min on ice. The ICL activities were then measured at optimum temperatures of the respective enzymes.

Fig. 7 Molecular models of *Cp*-WT and A214S. The models were built by the homology modeling program. The main chains and side chains of Gln119 and Ala214 or Ser214 are indicated by red and orange, respectively. Hydrogen bonds predicted by a program SWISSPDB VIEWER is indicated by green dashes.

Fig. 1 Sato et al

1 CTAGTAAAAATAAGTTTAACATAATTGTTTTTAAATACTTTTAAAGTAATATTTATTTGATTTTTGTTTTGGTATTTTCTGTGTGAATGGTG 90

91 GTAATCATCACTATATATTTTCTTAAATACATGGTTAGTAGACTATTATTTTTTGGCTCTCTAGTGCGAAGACTTAAAAAGTCTGTACACCA 180

-35 -10

181 ACCAAGCTAATATAGCTTTTATATACTACAAACAAAGTAAATGGAGATTGATATGTCTAATTATCAGAGTGCAATAGAAGCAGTTCAAGC 270

SD M S N Y Q S A I E A V Q A

271 GATCAAAGCAAAAGCCGGTAGCTCTTGGGATGCTATAAACCCAGAATCAATTGCCCGTATGCGTGCACAGAACAAATTCAAAACCTGGTCT 360

I K A K A G S S W D A I N P E S I A R M R A Q N K F K T G L

361 AGAGATAGCTCAGTACACTGCAGACATTATGCGTGCTGATATGGCTGCATTTGATGCAGACAAGACTCAATATACAAATCTCTAGGTTG 450

E I A Q Y T A D I M R A D M A A F D A D K T Q Y T Q S L G C

451 TTGGCATGGTTTTGTTGGCCAACAAAAAATGATTCTATCAAAAAACATTTTGGTAAACAGACCGTCGTTATTTATATCTTTTCAGG 540

W H G F V G Q Q K M I S I K K H F D G K T D R R Y L Y L S G

541 TTGGATGGTTGCTGCACTTCGCAGTGAATTTGGTCTCTTCTCTGATCAATCTATGCATGAAAAACGTCGGTAGCCTCTTTAGTTGCTGA 630

W M V A A L R S E F G P L P D Q S M H E K T S V A S L V A E

631 GCTTTATACTTTCTTACGTCAAGCTGACGCACGTGAATTAGGCGGTCTTTTCCGTGAAGTGTGAGCTGCTGAAGGTGATAAAGCTGC 720

L Y T F L R Q A D A R E L G G L F R E L D A A A E G D K A A

721 TATCCAATCACAGATTGATAACCATGTTACTCACGTTGTGCCAATTGTTGCTGATATCGATGCTGGTTTTGGTAAACGCTGAAGCGACTTA 810

I Q S Q I D N H V T H V V P I V A D I D A G F G N A E A T Y

811 CTTAATGGCTAAGCAAATGATTGAAGCAGGTGCTTGTGCGCTTCAAATCGAAAACCAAGTAGCTGATGAAAAGCAATGTGGTCATCAAGA 900

L M A K Q M I E A G A C A L Q I E N Q V A D E K Q C G H Q D

901 TGGCAAAGTAAGTGTCTCATGCTGATTTCCACTCTAAAATTCGTGCATTACGCCATGCATTTTTAGAACTAGGTATAGACAACGGTCT 990

G K V T V P H A D F H S K I R A L R H A F L E L G I D N G L

991 TATCGTTGCACGTACTGACTCTGAAGGTGCTGGTTTACTAAAGAAATCGCTGTTGTTAAAGAGCCTGGTGATTCTGGCGATATCTATAA 1080

I V A R T D S E G A G L T K E I A V V K E P G D S G D I Y N

1081 CTCTTATTTAGATGTTGAAGAAATCAGCGCAGCTGATATGGCTGAAGGCGATGTATGCTTTAATCGTGACGGTAAATTAGTTCGTCTTAA 1170

S Y L D V E E I S A A D M A E G D V C F N R D G K L V R P K

1171 GCGTTTACCTTCAGGTTTATACCAATTCCGTCAAGGTACTGGTCATGAGCGTTGTGTATTTGACTGTATCGGTGCACCTTAATGCTGGCGC 1260

R L P S G L Y Q F R Q G T G H E R C V F D C I G A L N A G A

1261 TGATTACTTTGGATTGAAACGGCAGTGCCAACGGTACACGAAATTGCTGGTATGATGGATGACGTTTCGTAAAGTTCACCCAGATGCGAA 1350

D L L W I E T A V P T V H E I A G M M D D V R K V H P D A K

1351 ACTTGTTTATAACAACTCTCCATCTTTTAACTGGACACTTAATTTCCGTGACGCAAGCATATGACGCAATGGTTGAAGCGGGTCAAGATGT 1440

L V Y N N S P S F N W T L N F R Q Q A Y D A M V E A G Q D V

1441 ATCTGCATACGTTTCGTGCTGACTTAATGAAAGCTGAATATGACGAACTGAATTGTCTGCAACAGCTGATGAACGCATTTCGTACTTTCCA 1530

S A Y V R A D L M K A E Y D E T E L S A T A D E R I R T F Q

1531 AGCTGATACTGCACGTGAAGCGAACGTATTCCATCACTTAATCACTTTACCTACTTACCATACAACTGCGTTATCAGTAGATAACTTAGC 1620

A D T A R E A N V F H H L I T L P T Y H T T A L S V D N L A

1621 TAAAGAGTACTTCGGTGAGCAAGGGATGCTTGGTTATGTTAAAGGTGTACAACGTAAAGAAATCCGTCAAGGTATTGCCTGTGTTAAACA 1710

K E Y F G E Q G M L G Y V K G V Q R K E I R Q G I A C V K H

1711 TCAAAACATGTCTGGTTCAGACATGGGTGATGATCACAAGAATACTTTGCAGGTGAAAATGCACTTAAAGCGGGTGGCGCAAAAAATAC 1800

Q N M S G S D M G D D H K E Y F A G E N A L K A G G A K N T

1801 GTCTAACCAGTTCAAGTTAATACCAATATTTTAACTCTTAAAGTGTGCTTAAAGCTGAAATATAAATACGAATCAGATTGTTACTAACG 1890

S N Q F S *

1891 TAGCAATCTGATTTTTTTTTGTTAAAAAATAGTATTGTAATTGTAAATTAGTTTACTAGAATCAAGAGAACAACTAAACAGTTTACACTT 1980

1981 TGAAACAATTGTTGCTATATTGTTTTGATCGTTCGATAAAATAAATACATGCCTGATTTTTTACTACTTTTGTACGAAGGAAATATGTTGCA 2070

2071 GTCATCTTCATCTAAACAATTGTTGCTACTTAATACCAATGCTGAAAGGCTTACTAAACTTGAAAACCTATTACTTTCTGGCACCCGCTA 2160

2161 TACTATTTTATCCGCAGCAGATACTCGCAGTGCCATAGCGATATTGAAAACAGAAAGCATCGACTTTGTGGTGAGTAATATCAATCTCGA 2250

2251 TAATTTTGATGGCTGGCGTTTGGCCCAATGGTGCCTCAGGCGTGTAAAGTGTCTCAGCAGATACACCTTTTGTGGTGGTGGCAA 2330

C.p. 1 -----MSNYQSATEAVQAIKAKAGSSWDATNPES
C.m. 1 -----MSNYQSATEAVKAIKEKAGNSWDATNPES
R.e. 1 -----MAQYODDKAVAGLKENHGSAWNAINPEY
R.m. 1 -----MAQYODDKAVAGLKEQHGSAWNAINPEY
A.v. 1 -----MSAYONEIKALAALKEKNGSAWSAINPEY
P.a. 1 -----MSAYONEIKAVAALKEKNGSSWSAINPEY
B.j. 1 MLHDITNKKQAIYTKFFSRLRRDTHFAMHCQECRWPAIAIVSERIRHMNYOPRGISTFOAFASYQSEIDAAKALLETQ-PTWNGVTAEA
R.p. 1 -----MTYHSKIVEAGKAIAPFS-QTWDGIEPES
N.a. 1 -----MTYHSKIVEAGKAIAPFS-QTWDGIEPES
A.c. 1 -----MTYQSALAEHVRAVKNKLGTTWDATRPED
H.m. 1 -----MAHKKTYSQLRSELLARYPVGLTKGGVSIDD

C.p. 30 IARMRAONKFKTGLDIAOYTADIMRADMAAFDADKTKYQTOSLGCWHGFIGOQKMSISIKKHFDGKTDRRYLYLSGWMVAALRSEFGPLPDQ
C.m. 30 VARMRAONKFKTGLDIAOYTADIMRADMAAFDADKTKYQTOSLGCWHGFIGOQKMSISIKKHFDGKTDRRYLYLSGWMVAALRSEFGPLPDQ
R.e. 30 AARMRAONKFKTGLDIAKYTAKIMRADMAAYDADSSKYQTOSLGCWHGFIGOQKMSIKMHFN-STERRYLYLSGWMVAALRSEFGPLPDQ
R.m. 30 AARMRAONKFKTGLDIAKYTAKIMRADMAAYDADSSKYQTOSLGCWHGFIGOQKMSIKKHFK-STERRYLYLSGWMVAALRSEFGPLPDQ
A.v. 30 AARMRIQNRFKTGLDIKYTAAIMRQDMAEYDADSSVYTOSLGCWHGFIGOQKMSISIKKHFK-TDKKLYLYSGWMVAALRSDFGPLPDQ
P.a. 30 AARMRIQNRFKTGLDIKYTAAIMRQDMAEYDADSSVYTOSLGCWHGFIGOQKMSISIKKHFK-TNKRYLYLSGWMVAALRSDFGPLPDQ
B.j. 90 VARMRLQNRFRFTGLDIKYTAAIMRADMAAYDADPTKYQTOSLGCWHGFIGOQKMSISIKKHFGGRTDRRYLYLSGWMVAALRSEFGPLPDQ
R.p. 42 VARMRLQNRFKTGLDIKYTAAIMRADMAAYDADPTKYQTOSLGCWHGFIGOQKMSISIKKHFGGRTDRRYLYLSGWMVAALRSEFGPLPDQ
N.a. 29 VARMRLQNRFRHTGLDIKYTAKIMRADMAAYDADPANYTOSLGCWHGFIGOQKMSISIKKHFG-TTKGRYLYLSGWMVAALRSEFGPLPDQ
A.c. 29 AARMRIQNRFRHTGLDIKYTAAIMRQDMAEYDADNSSYTOSLGCWHGFIGOQKMSIANKKFTG-TNKRYLYLSGWMVAALRSEFGPLPDQ
H.m. 32 IVQLRLQSPVESHLDVARAMASVMRADMAAYDRPTGKFTOSLGCWSGFHAQOMIKAVKRLRG-TTKGAVYLYSGWMVACLRNRWGLPDQ

C.p. 120 SMHEKTSVASLVAELYTFLROADARELGGLFRDLDAAG-----DKAAITQSQIDNHVTHVVPPIADIDAGFGNAEATYLMAKOMIEAGA
C.m. 120 SMHEKTSVASLVAELYTFLROADARELGGLFRDLDAASD-----DKAAITQDKIDNHVTHVVPPIADIDAGFGNAEATYLMAKOMIEAGA
R.e. 119 SMHEKTSVASLIRELYTFLROADARELGGLFRDLDAAGP-----AKAAITQAKIDNHVTHVVPPIADIDAGFGNAEATYLLAKOFIEAGA
R.m. 119 SMHEKTSVASLIRELYTFLROADARELGGLFRDLDAATGA-----AKAAITQEKIDNHVTHVVPPIADIDAGFGNAEATYLLAKOFIEAGA
A.v. 119 SMHEKTSVSSLIEELYTFLROADARELDLFFSALDDARNAGDK-AKEQELHKGIDGFETHVVPPIADIDAGFGNAEATYLLAKKMIIEAGA
P.a. 119 SMHEKTSVSSLIEELYTFLROADARELDLFFSALDDARNAGDK-AKEAELLAQIDNFEETHVVPPIADIDAGFGNAEATYLLAKKMIIEAGA
B.j. 180 SMHEKTSVPALIEELYTFLROADARELDLFFSALDDARNAGDK-TREKELIEKIDGFETHVVPPIADIDAGFGNAEATYLLAKKMIIEAGA
R.p. 132 SMHEKTSVPALIEELYTFLROADARELDLFFSALDDARNAGDK-AKETATIDQIDNFEETHVVPPIADIDAGFGNAEATYLLAKKMIIEAGA
N.a. 118 SMHEKTSVPALIEELYTFLROADARELDLFFSALDDARNAGDK-VNTHRLHKKIDNFEETHVVPPIADIDAGFGNAEATYLLAKKFIIEAGA
A.c. 118 SMHEKTSVPALIEELYTFLROADARELDLFFSALDDARNAGDK-AKVDTESQIDNFEETHVVPPIADIDAGFGNEEATYLLTKOMIEAGA
H.m. 121 SMHEKTSVVDLIEELYVSLROADARELDLFFSALDDARNAGDKARAKGATNKACEEISRIDGFETHVVPPIADIDAGFGNEHATYLLAKEMIKAGA

C.p. 205 CALQIENQVADAKQCGHODGKVTVPHEDFHISKIRALRHAFLELGIDNGLIVARTDSEAGLTKETIAVVKEPGDSGDIYNSYLDVEETISAA
C.m. 205 CALQIENQVADAKQCGHODGKVTVPHEDFHISKIRALRHAFLELGIDNGIIVSRDSEAGLTKETIAVVKEPGDOGDYIYNSYLDVEEIDVS
R.e. 204 CCIQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVTNPGDLGDOYNSYFLDCEELSD
R.m. 204 CCIQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVTNPGDLGDOYNSYFLDCEELSD
A.v. 208 CCIQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVTNPGDLGDKYNSYFLDCEETSPA
P.a. 208 CCIQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVTNPGDLGDLYNSYFLDCEETSES
B.j. 269 CALQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVSHKPGDIDGOYNSYFLDCEETIAE
R.p. 221 CALQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVSHKPGDIDGOYNSYFLDCEEDVAS
N.a. 207 CCIQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIAVVKEPGIDGOYNSYFLDCEEDVDA
A.c. 207 CALQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSEAGLTKQIPVSKKEDLASQYISYLDTOETIDIA
H.m. 211 CCLQIENQVADAKQCGHODGKVTVPHEDFLAKIRARHAFLELGVDGDIIVARTDSLGLTKQIPVSKKEDLASQYISYLDTOETIDIA

C.p. 295 D-MAEGDVCFNRDGLKVRPKRLPSGLYOFROGTGHERCVFDCIGALNAG-ADLLWIETAVPTVHEIAGVMDDVRKVHPDAKLVYNNSPSF
C.m. 295 D-MAEGDVCFNRDGLKVRPKRLPSGLYOFROGTGHERCVFDCIGALNAG-ADLLWIETATPNVADITAMMNEVRKEIPDAKLVYNNSPSF
R.e. 294 Q-LGNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIAQIGGMVSEIRKVIIPNAKLVYNNSPSF
R.m. 294 Q-LGNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIAQIGGMVSEIRKVIIPNAKLVYNNSPSF
A.v. 298 E-MKNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIAQIGGMVSEIRKVIIPNAKLVYNNSPSF
P.a. 298 E-LGNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIAQIGGMVSEIRKVIIPNAKLVYNNSPSF
B.j. 359 N-ARNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIEQIAKMVDRIREVIPNAKLAYNNSPSF
R.p. 311 K-IGNQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIEQIASMVDRIREVIPNAKLAYNNSPSF
N.a. 297 N-LGHQVAVITTRDGLKVRPKRLPSGLYOFRAGTGEARCVLDCVTLONG-ADLLWIETEKPHIAQIGGMVSEIRKVIIPNAKLAYNNSPSF
A.c. 297 D-AQEDFILLIKRDGLKVRPKRLPSGLYOFREGTOHDRVVLDCVTLONG-ADMTWIETPTPDVAGIAGFVNDIKKQVPNAKLAYNNSPSF
H.m. 301 NPLSEGELAIWQSGNFARPIRMPNGLFSFREGTGRARVIEDCTIASLKDGADLLWIETDTPNVDEIASMVAEIRKQVPDAKLAYNNSPSF

C.p. 383 NWTNLFROQAVDAMVEAGQDVSAVRA--DLMKAEYDETELSATADERIRTFQADTAREANVFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
C.m. 383 NWTNLFROQAVDAMVEAGQDVSAVRA--DLMNVDYDTTELAAAADDKIRSFQADTAREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
R.e. 382 NWTNLFROQAVDAMKAEAGQDVSAVRA--QLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
R.m. 382 NWTNLFROQAVDAMKAEAGQDVSAVRA--QLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
A.v. 386 NWTNLFROQAVDAMKAEAGQDVSAVRA--KLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
P.a. 386 NWTNLFROQAVDAMKAEAGQDVSAVRA--KLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
B.j. 447 NWTNLFROQAVDAMKAEAGQDVSAVRA--ELMKAEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
R.p. 399 NWTNLFROQAVDAMKAEAGQDVSAVRA--ELMKAEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
N.a. 385 NWTNLFROQAVDAMKAEAGQDVSAVRA--KLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
A.c. 385 NWTNLFROQAVDAMKAEAGQDVSAVRA--KLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ
H.m. 391 NWTNLFROQAVDAMKAEAGQDVSAVRA--KLMSVEYDQTELAKLADGKIRTFQGDASREAGIFHHLITLPTYHTTALSVDNLAKKEYFGEDQ

C.p. 471 GMLGYVKGVORKEIROGIACVKHQNMSGSDMGDDHKEYFAGEAALKAGGA-KNTSNQFS-
C.m. 471 GMLGYVKGVORKEIROGIACVKHQNMSGSDMGDDHKEYFAGEAALKAGGA-KNTSNQFS-
R.e. 470 GMLGYVAGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQF--
R.m. 470 GMLGYVAGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQF--
A.v. 474 GMLAYVKGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFH-
P.a. 474 GMLAYVKGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFH-
B.j. 535 GMLGYVKNVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFG-
R.p. 487 GMLGYVKNVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFS-
N.a. 473 GMLGYVKGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFAA
A.c. 473 GMLAYVAGVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFAH
H.m. 481 RMLAYVATVORKEIROGIACVKHQNMSGSDIGDDHKEYFAGEAALKAGG-KNTSNQFAA

Fig. 3 Sato et al

Fig. 4 Sato et al

Fig. 5 Sato et al

Fig. 6 Sato et al

Cp-WT

A214S

Fig. 7 Sato et al