



Title	FOOD HYGINE STUDIES ON PSYCHROPHILIC BACTERIA
Author(s)	HIGOSHI, Hironobu
Citation	Japanese Journal of Veterinary Research, 12(2), 33-33
Issue Date	1964-06
Doc URL	http://hdl.handle.net/2115/3287
Type	bulletin (article)
File Information	KJ00002369089.pdf



[Instructions for use](#)

INFORMATION

The Hokkaido University has granted the degree of Master of Veterinary Medicine to eight graduates on March 25, 1964. The authors' summaries of the theses excepting one are as follows:

FOOD HYGINE STUDIES ON PSYCHROPHILIC BACTERIA

Hironobu HIGOSHI

*Department of Veterinary Public Health,
Faculty of Veterinary Medicine,
Hokkaido University, Sapporo, Japan*

(Summary of Master's thesis written under direction of Dr. S. HAMADA)

Ninety samples of raw milk taken shortly after the milk arrived at 3 different milk plants, 66 bottles of pasteurized milk preserved at 5 and 20°C after bottling, 26 samples of pork and 14 samples of beef refrigerated at $-7\sim-10^{\circ}\text{C}$ for some days, and 17 groups of oysters collected directly from oyster beds, were used to examine the frequency of detection of psychrophilic bacteria. Nine hundred and sixteen strains of psychrophiles were isolated from these 213 samples by culturing at 5°C for 14 days. These strains were then examined as to growth, proteolytic and lipolytic activity, fluorescence and pigment production at 5, 20 and 35°C. The heat-resistance and the sensitivity to penicillin of all of the strains were also tested.

Among the 916 strains of several genera of psychrophilic bacteria isolated, the most common was *Pseudomonas* which was isolated primarily from the refrigerated beef and pork, and secondly from raw and pasteurized milk. Most strains of *Pseudomonas* grew well in cultures at 5 and 20°C, and were insensitive to 2.5 I. U. of penicillin. More than half of the *Pseudomonas* strains were active in gelatin liquefaction and in casein digestion, and some strains showed lipolytic activity, fluorescence and pigment production. It should be noted that 4 strains originating from refrigerated beef and pork were tolerant to heating at 60°C for 15 minutes.