



Title	Geographical variations in infection by larval <i>Anisakis simplex</i> and <i>Contracaecum osculatum</i> (Nematoda, Anisakidae) in walleye pollock <i>Theragra chalcogramma</i> stocks off Hokkaido, Japan
Author(s)	Konishi, Kenji; Sakurai, Yasunori
Citation	Fisheries Science, 68(3), 534-542 https://doi.org/10.1046/j.1444-2906.2002.00459.x
Issue Date	2002-06
Doc URL	http://hdl.handle.net/2115/35583
Rights	© 2002 公益社団法人日本水産学会; © 2002 The Japanese Society of Fisheries Science
Type	article
File Information	2002-68_p534-542.pdf



[Instructions for use](#)

Geographical variations in infection by larval *Anisakis simplex* and *Contracaecum osculatum* (Nematoda, Anisakidae) in walleye pollock *Theragra chalcogramma* stocks off Hokkaido, Japan

KENJI KONISHI^{a*} AND YASUNORI SAKURAI

Graduate School of Fisheries Science, Hokkaido University, Hakodate, Hokkaido 041-8611, Japan

ABSTRACT: Stocks of walleye pollock *Theragra chalcogramma* collected from: (i) the Sea of Japan (off Rebun Island and Kumaishi); (ii) the Pacific coast (off Shikabe and eastern Hokkaido); and (iii) Nemuro Strait off Hokkaido, northern Japan, were examined for anisakid nematodes during December 1999 to February 2000, and the prevalence and abundance of *Anisakis simplex* and *Contracaecum osculatum* larvae were compared among the various sampling sites for fish of the same size and age. *Anisakis simplex* was generally more abundant than *C. osculatum*. Infection by *A. simplex* varied between the aforementioned stocks of walleye pollock as well as within stocks, whereby fish from off Rebun Island and Nemuro Strait were infected the most, followed by those from off the Pacific coast and Kumaishi. Infection by *C. osculatum* differed between the host stocks, and *C. osculatum* was the most abundant among the fish from Nemuro Strait. The infection variations seemed to be due to differences in host growth rate, host feeding habit, and the distribution of marine mammal final hosts. The results indicate that these two larval nematodes are useful biological indicators for the population study of walleye pollock in Japanese waters.

KEY WORDS: Anisakidae, *Anisakis simplex*, *Contracaecum osculatum*, geographic variation, Nematoda, *Theragra chalcogramma*, walleye pollock.

INTRODUCTION

Walleye pollock *Theragra chalcogramma* is one of the most abundant and commercially exploited fish species around northern Japan, where it occurs at bottom or midwater depths above the continental shelf and slope.^{1,2} Because walleye pollock is distributed broadly in the North Pacific, understanding the distinctiveness between populations is important for the successful management of walleye pollock fisheries.³ Walleye pollock in Japanese waters is generally divided into several stocks (Sea of Japan, Pacific coast, Sea of Okhotsk and Nemuro Strait), based on tagging surveys,

morphometrics and meristics.^{1,4–7} However, the distribution of walleye pollock changes annually in relation to environmental factors;² therefore, more detailed information on the populations that form these stocks and on their migration routes has become increasingly important.

Many different methods should be used to increase the efficiency and accuracy of fish population studies, and parasitological analyses of marine fishes can provide useful information on the ecology of a host species.^{8–12} The advantage of such a method is that it is inexpensive and does not require the recapturing of the same individuals, unlike conventional artificial tagging.¹³ The basic principle of the method is based on the fact that the infection rate of parasites varies among their host populations depending on feeding habits, migration routes or growth rates.

Several attempts have been made to define stocks and determine the migration routes of walleye pollock in Canadian and Russian waters

*Corresponding author: Tel: 81-3-3536-6521. Fax: 81-3-3536-6522. Email: konishi@i-cetacean-r.or.jp

^aPresent address: The Institute of Cetacean Research, 4–5 Toyomi-cho, Chuo-ku, Tokyo 104-0055, Japan.

Received 13 April 2001. Accepted 7 December 2001.

using parasites.^{3,14,15} Larval anisakid nematodes, such as the whaleworm *Anisakis simplex* (Rudolphi, 1809) and *Contracaecum osculatum* (Rudolphi, 1802), are transported by crustaceans to fishes and squids as paratenic hosts and finally ingested by marine mammal definitive hosts.^{16–19} Their third-stage larvae commonly occur in the viscera of walleye pollock.^{20–24} Because the feeding habits of walleye pollock vary geographically in Japanese waters,^{25–27} differences in the abundance of anisakid larvae in walleye pollock can be used to define the stocks or their feeding locations. Previous ecologic surveys of anisakid nematodes from walleye pollock in Japanese waters have shown a broad variation of infection among sampling sites.^{20,21,28} However, the walleye pollock examined in these studies were collected throughout the year including summer, when various walleye pollock stocks sometimes mix at feeding grounds.⁷ Thus, these data can not be used to compare stocks. In addition, ecologic surveys of anisakid larvae in walleye pollock have not been carried out for more than two decades.

The aim of the present study was to elucidate the geographic variation in the prevalence and abundance of *A. simplex* and *C. osculatum* larvae in different walleye pollock stocks off Hokkaido, Japan in winter when the stocks separate into their respective spawning areas.

MATERIALS AND METHODS

Sampling

Walleye pollock were collected at five sites (Rebun Island, Kumaishi, Shikabe, Nemuro Strait, and eastern Hokkaido) off Hokkaido, Japan from December 1999 to February 2000 (Fig. 1). The collection comprised walleye pollock from three stocks: (i) Sea of Japan (Rebun Island and Kumaishi); (ii) Pacific coast (Shikabe and eastern Hokkaido); and (iii) Nemuro Strait. Fish from off Rebun Island, Kumaishi, Shikabe and in Nemuro Strait were collected from commercial catches, and fish from off eastern Hokkaido were collected

by the Hokkaido National Fisheries Research Institute (HNFRI) using bottom or midwater trawls (Table 1).

Treatment of samples

The fish samples were either frozen immediately after collection, brought to the laboratory and then thawed at room temperature for later examination, or dissected within 24 h after collection. The fish were sexed and measured to the nearest millimeter from the tip of the snout to the fork of the caudal fin, and otoliths from each specimen were removed for age determination. The otoliths were examined by the HNFRI, the Hokkaido Fisheries Experimental Station (FHES), or by KK to assess each specimen's age using the break and burn method.²⁹ The fish were then divided into both 1-year age classes and 25-mm length classes. Fish smaller than 300 mm were not included in the analyses. *Anisakis simplex* (L3), which had been




Fig. 1 Map of Hokkaido, northern Japan showing five sites where walleye pollock *Theragra chalcogramma* were collected.

Table 1 General information on walleye pollock (*Theragra chalcogramma*) collected off the coast of Hokkaido, Japan

Sampling areas	Sampling dates	N	Fishing gear	Mean length with range (mm)
Rebun Is.	Jan/2000	131	Trawl net	446.1 (371–582)
Kumaishi	Dec/1999	100	Longline	396.5 (325–491)
Shikabe	Jan/2000	100	Gill net	429.6 (372–495)
East. Hokkaido	Jan/2000	423	Trawl net	371.6 (300–538)
Nemuro Strait	Feb/2000	197	Gill net	468.0 (360–622)

described as *Anisakis* type I larva,^{30,31} and *Contracaecum osculatum* (L3), which had been described as *Contracaecum* type A³¹ or type B,³⁰ were removed from the fish mesenteries, pyloric caeca and livers, and were counted.

Numerical analyses and statistics

Nematode infections were compared among the same length classes and age classes from different stocks. 'Prevalence' was defined as the percentage of infected hosts, and 'abundance' was defined as the number of parasites per host.³² Because growth rates differ among walleye pollock stocks^{4,33} (Fig. 2), prevalence and abundance were analysed for both host length classes and age classes. Walleye pollock from each sampling site had different length distributions, probably due to the differences in fishing gears; therefore, statistical tests were conducted on data from three length classes (400–424 mm, 425–449 mm, 450–474 mm) and from fish aged 4–7 years.

As the frequency distributions of abundance data were positively skewed, these data were $\log_{10}(x+1)$ transformed to bring the data closer to normal distributions for parametric analyses.³⁴ Because of the fact that there are differences in the infection of anisakid larvae between male and female fish,^{22,35} the abundance of the two larval nematodes in males and female hosts were compared in the 400–424 mm length class and in the age 4 class using Student's *t*-test. To compare the abundances among sampling stations for the same length classes and age classes, a one-way ANOVA

and a post-hoc Tukey's test were used. The significance of the relationships between nematode abundance and host length or age classes were tested using Kendall's rank correlation test. For the correlation analyses between larval abundances and host age, walleye pollock from off Shikabe and eastern Hokkaido were not included because of their small sample sizes.

RESULTS

Comparison of larval nematode abundance between host sexes

No significant difference in the nematode abundance was found between the host sexes in the 400–424-mm length class for both *Anisakis simplex* ($t=1.15$, d.f.=105, $P=0.25$) and *Contracaecum osculatum* ($t=1.40$, d.f.=105, $P=0.16$). In contrast, nematode abundances differed significantly between age 4 male and female walleye pollock for *A. simplex* ($t=2.05$, d.f.=204, $P<0.05$), but not for *C. osculatum* ($t=-0.65$, d.f.=204, $P=0.16$). Therefore, data from male and female hosts were separated to compare *A. simplex* abundance among sampling sites according to host age. Male and female data were combined for all other analyses.

Anisakis simplex larvae in walleye pollock

The prevalence of *A. simplex* was high (>80%) in all host length classes at all sampling sites, except off Kumaishi in the 400–424-mm length class (Table 2). The abundance of *A. simplex* differed significantly among sampling sites ($F=7.79$, d.f.=4, $P=0.0001$), and the fish from Kumaishi were significantly less abundant than those from the other four sites in the 400–424-mm length class (Table 2). However, no significant difference in abundance was found among sampling sites in the 425–449-mm length class ($F=1.791$, d.f.=4, $P=0.1324$). The abundance of *A. simplex* in the 450–474-mm length class differed significantly among sampling sites ($F=5.54$, d.f.=3, $P=0.01$), and the fish from off eastern Hokkaido were significantly less abundant than those from off Rebus Island, Shikabe and Nemuro Strait (Table 2). Abundance in fish from off Kumaishi was lowest for all host age classes (Fig. 3).

The mean abundance of *A. simplex* larvae was correlated significantly with host size from off Rebus Island, Kumaishi and Nemuro Strait ($\tau=0.90$, $P=0.001$; $\tau=0.71$, $P=0.05$; $\tau=0.71$, $P=0.05$, respectively). Those were also correlated significantly with the host age only for female walleye pollock from off Rebus Island, Kumaishi and




Fig. 2 Age-length relationships for walleye pollock (*Theragra chalcogramma*) collected at five sites off Hokkaido, northern Japan.

Table 2 Prevalence and abundance of *Anisakis simplex* from walleye pollock by host length class

Host length-class (mm)	400-424				425-449				450-474			
	Reb	Kum	Shi	E.H.	N.St.	Reb	Kum	Shi	E.H.	N.St.	Shi	E.H.
Host sample size (N)	26	12	33	107	24	31	24	38	55	42	20	13
Prevalence (%)	100	50	88	93	100	97	92	97	89	100	100	85
Mean abundance	5.7 ^a	1.1 ^b	4.1 ^a	4.4 ^a	6.4 ^a	5.0	5.0	5.7	4.8	6.6	9.3 ^a	3.8 ^b

Tukey-Kramer test on abundances transformed by $\log_{10}(x+1)$ were used if results of ANOVA were significant. Abundances with the different letters in every length-class indicate significant difference at $P < 0.05$ level. Reb, off Rebun Is.; Kum, off Kumaishi; Shi, off Shikabe; E.H., off eastern Hokkaido; N.St., Nemuro Strait.




Fig. 3 Abundance of *Anisakis simplex* in relation to the age and sex of walleye pollock (*Theragra chalcogramma*) from various sampling sites off Hokkaido, northern Japan.

Nemuro Strait ($\tau = 1.00$, $P = 0.05$; $\tau = 1.00$, $P = 0.01$; $\tau = 0.93$, $P = 0.01$, respectively) (Fig. 3).

Contracaecum osculatum larvae in walleye pollock

Contracaecum osculatum was generally less abundant than *A. simplex* in walleye pollock (Tables 2,3). The prevalence of *C. osculatum* larvae in fish from Nemuro Strait was higher in all length classes compared with those from other sites, and that in fish from the Pacific coast (off Shikabe and eastern Hokkaido) was generally higher in all length classes compared with that in the Sea of Japan (off Rebun Island and Kumaishi) (Table 3).

The abundance of *C. osculatum* larvae differed significantly among sampling sites in all host length classes (400-424 mm, 425-449 mm, and 450-474 mm) ($F = 8.90$, d.f. = 4, $P = 0.001$; $F = 17.90$, d.f. = 4, $P = 0.001$; $F = 16.49$, d.f. = 3, $P = 0.001$, respectively), and that at Nemuro Strait was significantly higher than those at other sites (Table 3). In addition, the prevalences in fish from the Pacific coast of Hokkaido (off Shikabe and eastern Hokkaido) were generally higher than those in fish from the Sea of Japan coast of Hokkaido (off Rebun Island and Kumaishi) (Table 3). In hosts aged between 4 and 7 years, the abundance of *C. osculatum* larvae differed significantly among sampling sites ($F = 50.38$, d.f. = 3, $P = 0.001$), and that in Nemuro Strait was significantly higher than those in other sites (Fig. 4; Table 4).

The mean abundance of *C. osculatum* larvae was correlated significantly with host size off

Table 3 Prevalence and abundance of *Contracaecum osculatum* from walleye pollock by host length class

Host length-class (mm) Sampling site	400–424				425–449				50–474			
	Reb	Kum	Shi	E.H.	N.St.	Reb	Kum	Shi	E.H.	N.St.	Shi	N.St.
Host sample size (N)	26	12	33	107	24	31	24	38	55	42	20	40
Prevalence (%)	31	33	70	63	88	39	25	61	56	91	80.0	95.0
Mean abundance	0.4 ^a	0.5 ^{ab}	1.3 ^b	1.4 ^b	2.8 ^c	0.6 ^{ab}	0.4 ^b	2.1 ^a	2.1 ^a	5 ^d	2.5 ^a	7.7 ^b

Tukey–Kramer test on abundances transformed by $\log_{10}(x+1)$ were used if results of ANOVA were significant. Abundances with the same letter in every length-class indicate not significant difference at $P < 0.05$ level. Reb, off Rebun Is.; Kum, off Kumaishi; Shi, off Shikabe; E.H., off eastern Hokkaido; N.St., Nemuro Strait.




Fig. 4 Abundance of *Contracaecum osculatum* in relation to the age of walleye pollock (*Theragra chalcogramma*) from various sampling areas off Hokkaido, northern Japan.

Table 4 Prevalence and abundance of *Contracaecum osculatum* from walleye pollock (*Theragra chalcogramma*) in age classes 4 through 7

Areas	Reb	Kum	Shi	N.St.
Host sample size (N)	40	40	40	40
Prevalence (%)	43	13	70	95
Mean abundance	1.85 ^a	0.18	1.80 ^a	6.98

Tukey test on abundances transformed by $\log_{10}(x+1)$ were used if results of ANOVA were significant.

ANOVA were significant. Abundances with the same letters are not significantly different at $P < 0.05$ level.

Reb, off Rebun Is.; Kum, off Kumaishi; Shi, off Shikabe; N.St., Nemuro Strait.

Samples from off eastern Hokkaido were not included because a small sample size of fish older than 5 years.

Rebun Island, eastern Hokkaido and in Nemuro Strait ($\tau = 0.60$, $P = 0.05$; $\tau = 0.94$, $P = 0.001$; $\tau = 1.00$, $P = 0.001$, respectively). But the mean abundance of *C. osculatum* larvae was correlated significantly with host age only in Nemuro Strait ($\tau = 0.64$, $P = 0.05$) (Fig. 4).

DISCUSSION

Geographical distribution of *Anisakis simplex* in walleye pollock

Infection levels by *Anisakis simplex* varied widely both between walleye pollock stocks and within the Sea of Japan stock, although previous stock studies have insisted that walleye pollock in the Sea of Japan off Hokkaido form a single stock.^{7,36}

Previous studies on *A. simplex* in walleye pollock^{20,21} have shown similar results concerning such interstock variations. Walleye pollock from off Rebun Island and Nemuro Strait were generally higher in both the abundance and prevalence of *A. simplex* compared with those from the other sites, and the fish from the Pacific coast (off Shikabe and eastern Hokkaido) were generally higher than those from off Kumaishi. Furthermore, the abundance of *A. simplex* larvae in larger walleye pollock differed significantly between the two sites off the Pacific coast of Hokkaido (Shikabe and eastern Hokkaido).

Variation in the nematode abundance in walleye pollock among sampling sites was similar to that in the prey composition among and within walleye pollock stocks. Infection by *A. simplex* occurs when walleye pollock feeds on infected prey. Euphausiids are a major crustacean host of *A. simplex*,^{19,37-42} thus the intensity of predation on euphausiids may be an important factor that determines infection levels. Walleye pollock that spawn in Nemuro Strait migrate to the Sea of Okhotsk to feed,^{43,44} and euphausiids are the dominant prey of walleye pollock in the Sea of Okhotsk off Hokkaido.²⁵ Walleye pollock off the Pacific coast of Hokkaido feed on many prey items according to their development stage; that is, small walleye pollock in this area feed mainly on euphausiids, copepods and amphipods, whereas larger walleye pollock are more piscivorous and feed on larger prey (e.g. micronektonic fishes and small walleye pollock), in addition to the aforementioned crustaceans.^{25,26,45} In the Sea of Japan off northern Hokkaido, walleye pollock feed mainly on euphausiids,²⁵ whereas in the Sea of Japan off southern Hokkaido coast, they feed mainly on amphipods, chaetognaths and euphausiids.²⁷ Thus, a high consumption of euphausiids by walleye pollock is an important factor causing the high infection rates of *A. simplex*.

Geographical variation in infection by *A. simplex* within the Sea of Japan off Hokkaido (off Rebun and Kumaishi) seems to be related to the difference in the growth rate of walleye pollock. Kooka has reported that walleye pollock in the Sea of Japan off southern Hokkaido have a low growth rate;³³ hence, the low infection rate in the host from Kumaishi may be associated with the lower feeding rates in this area.

Infection in larger walleye pollock (450–474 mm) from eastern Hokkaido was also low, and this is presumably caused not by a low feeding rate, but by small-scale variations in the migration and spawning area within the Pacific stock. The Pacific stock has a large spawning ground near Shikabe, and most of the mature walleye pollock from the

Pacific coast of Hokkaido migrate to that spawning ground in winter.⁴⁶ However, some walleye pollock spawn along the coast of eastern Hokkaido.⁴⁶ Most walleye pollock in the 450–474 mm length class have reached the size of sexual maturity;¹ hence, the difference in nematode abundance in larger walleye pollock between Shikabe and eastern Hokkaido was probably more distinct in larger fish than in smaller fish, suggesting that walleye pollock from Shikabe have different feeding grounds as well as spawning areas from fish from eastern Hokkaido, which could be the cause of the different infection levels of walleye pollock between these localities.

The distribution of the definitive hosts of *A. simplex* may also influence the infection in walleye pollock. *Anisakis simplex* matures and releases eggs into seawater when it occurs in its definitive host.^{17,47} Thus, the distribution of its definitive hosts will largely determine the area where *A. simplex* larvae occur. The major definitive hosts of *A. simplex* are cetaceans,⁴⁸ and minke whale *Balaenoptera acutorostrata* is one of the major definitive host species in the north-west North Pacific.⁴⁹ Northern North Pacific minke whales occur mainly off the coast of northern Japan in summer,⁵⁰ including the areas studied in the present report. However, *A. simplex* appears to exhibit little host specificity,^{48,51} so how other cetacean species affect the infection by *A. simplex* in walleye pollock remains unknown.

The abundance of *A. simplex* in walleye pollock increased significantly with both length and age of female hosts. A similar increase in the abundance of *A. simplex* in relation to host size has been reported in rockfish *Sebastes capensis*⁵² and in Atlantic cod *Gadus morhua*.³⁵ As *A. simplex* larvae seem to live at least 2 or 3 years in host fish,^{53,54} older fish are usually more heavily infected by larvae than younger ones. Higher consumption rates by older (larger) fish also increase the chance for infection by *A. simplex*. In contrast, the abundance of larvae was not correlated with the age of male walleye pollock. Hemmingsen *et al.* have suggested that differences in infection of anisakid larvae between male and female Atlantic cod were caused by differences in their feeding habits.³⁵ However, such differences have not yet been reported in walleye pollock.

The results of the present study show that *A. simplex* can be used as a biological indicator of walleye pollock. Abundances of *A. simplex* can be used to divide some previously defined walleye pollock stocks, and used to divide the Sea of Japan and Pacific stocks into at least two respective substocks. These substocks probably have different feeding areas and migration routes.

Geographical distribution of *Contracaecum osculatum* in walleye pollock

Infection by *C. osculatum* differed among walleye pollock stocks, and the highest infection by *C. osculatum* larvae was seen in the hosts from Nemuro Strait for all length and age classes.

Contracaecum osculatum matures in the stomachs of pinniped definitive hosts and releases eggs into seawater with the pinniped feces.⁵⁵ Hatched larvae are then ingested by crustaceans and passed on to fishes, and finally transmitted to their definitive hosts.¹⁸ Larval *C. osculatum* occur in euphausiids⁵⁶ and copepods.¹⁸ Although euphausiids and copepods are both preyed upon by walleye pollock from all the sites studied,^{25-27,45} the present study's results showed heavy infection occurred only in walleye pollock from Nemuro Strait. Boily and Marcogliese have stated that the distribution of definitive hosts could strongly reflect the occurrence of *C. osculatum*.⁵⁷ In Japanese waters, the definitive hosts of *C. osculatum* include Steller sea lions *Eumetopias jubatus*, northern fur seals *Callophorus ursinus*, spotted seals *Phoca largha*, and ribbon seals *P. fasciata*.⁵⁸ All are abundant in Japanese waters, but in species other than ribbon seals, few larvae mature to adulthood⁵⁹ (Takahashi C., pers. comm., 1999). Ribbon seal occurs only in the Sea of Okhotsk,⁶⁰ where it is commonly infected by *C. osculatum*.⁶¹ Thus, the distribution of ribbon seal presumably affects the infection of *C. osculatum* in walleye pollock. As noted earlier, the Nemuro Strait walleye pollock stock feed in the Sea of Okhotsk, therefore the infection of the fish in this stock probably occurred in the Sea of Okhotsk. The abundance of *C. osculatum* was correlated significantly with walleye pollock length, but not with the host age for all stocks except the Nemuro Strait stock. This suggests that the abundance of *C. osculatum* in walleye pollock was affected only by the amount of prey consumed by fish because parasites with a shorter life span in the host will not reflect long-time fluctuations of abundance. Walleye pollock in Nemuro Strait feed mainly on euphausiids irrespective of their length; therefore, the higher consumption rates by larger fish increase the chance for infection by *C. osculatum* in Nemuro Strait.

the study's results indicate that infection by *C. osculatum* can be used to divide previously defined walleye pollock stocks, although within-stock variation was not found. The Nemuro Strait stock was characterized distinctly by the high abundance of *C. osculatum*. Thus, the present study showed different infection patterns between *A. simplex* and *C. osculatum* from some walleye pollock stocks around Hokkaido, indicating that the two larval

nematodes are both useful biological indicators for population studies of walleye pollock in Japanese waters.

ACKNOWLEDGMENTS

We thank S. Honda and A. Nishimura, Hokkaido National Fishery Research Institute, and O. Shida, T. Mutou, Y. Mihara and A. Wada, Hokkaido Fisheries Experimental Station, for providing the walleye pollock samples and age data. We also thank J. R. Bower, Hokkaido University, and K. Nagasawa, National Research Institute of Far Seas Fisheries, for their helpful comments on the manuscript, and T. Hirota, Tokyo University of Agriculture & Technology, for providing an excellent computer program for non-parametric statistics.

REFERENCES

1. Yoshida H. Walleye pollock *Theragra chalcogramma* (Pallas). In: Nagasawa K, Torisawa M (eds). *Fishes and Marine Invertebrates of Hokkaido: Biology and Fisheries*. Kita-nihon Kaiyo Center Co. Ltd, Sapporo. 1991; 120-125.
2. Maeda T, Takagi S, Kamei Y, Kajiwara Y, Meguro T, Nakatani T. History and methodology of walleye pollock studies. *Sci. Rep. Hokkaido Fish. Exp. Stn* 1993; 42: 1-14.
3. Arthur JR. A preliminary analysis of the discreteness of stocks of walleye pollock (*Theragra chalcogramma*) from the northeastern Pacific Ocean off Canada based on their parasites. *Can. Tech. Rep. Fish. Aquat. Sci.* 1983; 1184: 15.
4. Ishida A. On the age determination and morphometrical differences of the otolith of Alaska pollock in the Hokkaido coast. *Bull. Hokkaido Reg. Fish. Res. Lab.* 1954; 11: 36-49.
5. Hashimoto R, Koyachi S. Geographical variation of relative growth of walleye pollock *Theragra chalcogramma* (Pallas). *Bull. Tohoku Reg. Fish. Res. Lab.* 1977; 38: 41-74.
6. Koyachi S, Hashimoto R. Preliminary survey of variations of meristic characters of walleye pollock *Theragra chalcogramma*. *Bull. Tohoku Reg. Fish. Res. Lab.* 1977; 38: 17-40.
7. Tsuji S. General remarks on the populations of Alaska pollock (*Theragra chalcogramma* (Pallas)) in waters around Hokkaido. *J. Hokkaido Fish. Exp. Stn* 1978; 35: 1-57.
8. Beverley-Burton M. Population genetics of *Anisakis simplex* (Nematoda: Ascaridoidea) in Atlantic salmon (*Salmo salar*) and their use as biological indicators of host stocks. *Environ. Biol. Fish.* 1978; 3: 369-377.
9. McGladdery SE, Burt MD. Potential of parasites for use as biological indicators of migration, feeding, and spawning behavior of northwestern Atlantic herring (*Clupea harengus*). *Can. J. Fish. Aquat. Sci.* 1985; 42: 1957-1968.
10. Nascimento GM. Populations and assemblages of parasites in hake, *Merluccius gayi*, from the southeastern Pacific Ocean: Stock implications. *J. Fish Biol.* 1996; 48: 557-568.
11. Boje J, Riget F, Køie M. Helminth parasites as biological tags in population studies of Greenland halibut (*Reinhardtius hippoglossoides* (Walbaum)), in the north-west Atlantic. *ICES J. Mar. Sci.* 1997; 54: 886-895.

12. MacKenzie K, Abaunza P. Parasites as biological tags for stock discrimination of marine fish: A guide to procedures and methods. *Fish. Res.* 1998; **38**: 45–56.
13. Urawa S. Parasites as biological indicators contributing to salmonid biology. *Sci. Rep. Hokkaido Salmon Hatchery* 1989; **43**: 53–74.
14. Avdeev VV, Avdeev GV. A study of walleye pollock population structure and migration routes using parasitological indicators. In: *Proceedings of the International Symposium on the Biology and Management of Walleye Pollock*. Alaska Sea Grant Report No. 89-1. 1988; 569–590.
15. Avdeev GV. Infestation by helminths and redistribution of immature walleye pollock *Theragra chalcogramma* in the Sea of Okhotsk. *J. Ichthyol.* 1996; **36**: 665–673.
16. Kagei N. Euphausiids and their parasites (I). *Geiken Tsushin* 1979; **328**: 53–62.
17. Nagasawa K. The life cycle of *Anisakis simplex*: A review. In: Ishikura H, Kikuchi K (eds). *Intestinal Anisakiasis in Japan: Infected Fish, Sero-Immunological Diagnosis, and Prevention*. Springer-Verlag, Tokyo. 1990; 31–40.
18. Køie M, Fagerholm H-P. The life cycle of *Contracaecum osculatum* (Rudolphi, 1802) *sensu stricto* (Nematoda, Ascaridoidea, Anisakidae) in view of experimental infections. *Parasitol. Res.* 1995; **81**: 481–489.
19. Hays R, Measures LN, Huot J. Euphausiids as intermediate hosts of *Anisakis simplex* in the St Lawrence estuary. *Can. J. Zool.* 1998; **76**: 1226–1235.
20. Sasaki M. Survey of parasites of the Alaska pollock, *Theragra chalcogramma*. *J. Hokkaido Fish. Exp. Stn* 1973; **30**: 14–39.
21. Suzuki M, Oishi K. Parasites of *Theragra chalcogramma*. In: Japanese Society of Fisheries (ed.). *Fish and Anisakis*. Koseisha Koseikaku, Tokyo. 1974; 113–125.
22. Arthur JR, Margolis L, Whitaker DJ, McDonald TE. A quantitative study of economically important parasites of walleye pollock (*Theragra chalcogramma*) from British Columbian waters and effects of postmortem handling on their abundance in the musculature. *Can. J. Fish. Aquat. Sci.* 1982; **39**: 710–726.
23. Arthur JR. A survey of the parasites of walleye pollock (*Theragra chalcogramma*) from the northeastern Pacific Ocean off Canada and a zoogeographical analysis of the parasite fauna of this fish throughout its range. *Can. J. Zool.* 1984; **62**: 675–684.
24. Nagasawa K. Parasites of gadid fishes in Japanese waters (review and bibliography). *Sci. Rep. Hokkaido Fish. Exp. Stn* 1993; **42**: 69–89.
25. Iizuka A, Kurohagi T, Ikuta K, Imai S. Composition of the food of Alaska pollock (*Theragra chalcogramma*) in Hokkaido with special reference to its local differences. *Bull. Hokkaido Reg. Fish. Res. Lab.* 1954; **11**: 7–20.
26. Yamamura O. Ecological study on demersal fish community off Sendai Bay, northern Japan, with special reference to niche dynamics among dominant fishes. PhD thesis. Hokkaido University, Hakodate, Japan, 1994.
27. Kooka K, Takatsu T, Kamei Y, Nakatani T, Takahashi T. Vertical distribution and prey of walleye pollock in the northern Japan Sea. *Fish. Sci.* 1998; **64**: 686–693.
28. Suzuki M, Matsufuji R. Survey of parasites from marine fishes. *Res. Bull. Naylor Junior Coll.* 1975; **8**: 1–30.
29. Nishimura A. Age determination of walleye pollock based on the otoliths (Review). *Sci. Rep. Hokkaido Fish. Exp. Stn* 1993; **42**: 37–49.
30. Koyama T, Kobayashi A, Kumada M, Komiya Y, Oshima T, Kagei N, Machida M. Morphological and taxonomical studies on Anisakidae larvae found in marine fishes and squids. *Jpn. J. Parasitol.* 1969; **18**: 466–487.
31. Shiraki T. Larval nematodes of family Anisakidae (Nematoda) in the northern Sea of Japan, as a causative agent of eosinophilic phlegmone or granuloma in the human gastrointestinal tract. *Acta Med. Biol.* 1974; **22**: 57–98.
32. Bush TO, Lafferty KD, Lotz JM, Shostak AW. Parasitology meets ecology on its own terms: Margolis *et al.* revisited. *J. Parasitol.* 1997; **83**: 575–583.
33. Kooka K. Effect of feeding environment on the growth pattern of pelagic walleye pollock (*Theragra chalcogramma*) in the northern Japan Sea. PhD thesis. Hokkaido University, Hakodate, Japan, 1999.
34. Sokal RR, Rohlf JF. *Biometry: The Principles and Practice of Statistics in Biological Research*, 3rd edn. W.H. Freeman, New York. 1995.
35. Hemmingsen W, Halvorsen O, MacKenzie K. The occurrence of some metazoan parasites of Atlantic cod, *Gadus morhua* L., in relation to age and sex of the host in Balsfjord (70°N), North Norway. *Polar Biol.* 2000; **23**: 368–372.
36. Utoh H, Natsume M. Regional difference of age-composition of walleye pollock, *Theragra chalcogramma*, from the Japan Sea coast of Hokkaido and an interpretation for its cause. *Sci. Rep. Hokkaido Fish. Exp. Stn* 1993; **42**: 215–228.
37. Uspenskaya AV. *Parasitic Fauna of Benthic Crustaceans of the Barents Sea*. Izdatel'stvo Akademiiy Nauk SSSR, Moscow. 1963.
38. Oshima T. A study on the first host of *Anisakis*. *Saishin Igaku* 1969; **24**: 401–404.
39. Smith JW. *Thysanoessa inermis* and *T. longicaudata* (Euphausiidae) as first intermediate hosts of *Anisakis* sp. (Nematoda: Ascaridata) in the northern North Sea, to the north of Scotland and at Faroe. *Nature* 1971; **234**: 478.
40. Kagei N. Studies on anisakid nematoda (Anisakidae) (IV). Survey of *Anisakis* larvae in the marine crustacea. *Bull. Inst. Publ. Health* 1974; **23**: 65–71.
41. Oshima T. *Anisakis* and anisakiasis in Japan and adjacent area. In: Morishita K, Komiya Y, Matsubayashi H (eds). *Progress of Medical Parasitology in Japan*. Meguro Parasitological Museum, Tokyo. 1972; 304–393.
42. Shimazu T. Some helminth parasites of marine planktonic invertebrates. *Bull. Nagano-Ken Junior Coll.* 1982; **37**: 11–29.
43. Tsuji S. Walleye pollock survey in southwest Okhotsk Sea. *Sci. Rep. Hokkaido Fish. Exp. Stn* 1973; **30**: 1–22.
44. Ishino K. Stock structure of the southwestern Okhotsk Sea walleye pollock, *Theragra chalcogramma*, during the feeding migration based on the vertebral number. *Sci. Rep. Hokkaido Fish. Exp. Stn* 1993; **42**: 203–213.
45. Maeda T, Takahashi T, Ueno M. Behaviour in each life period of adult Alaska pollack in the adjacent waters of Funka Bay, Hokkaido. *Bull. Jpn. Soc. Sci. Fish.* 1983; **49**: 577–585.
46. Hamatsu T, Yabuki K. Spawning migration and spawning ground of walleye pollock *Theragra chalcogramma* distributed along the Pacific coast of eastern Hokkaido. *Bull. Hokkaido Natl Fish. Res. Inst.* 1995; **59**: 31–41.
47. Kagei N. Life history of nematodes of the genus *Anisakis*. *Saishin Igaku* 1969; **24**: 389–400.
48. Kagei N, Oshima T, Kobayashi A, Kumada M, Koyama T, Komiya Y, Takemura A. Survey of *Anisakis* spp. (Anisakinae, Nematoda) on marine mammals in the coast of Japan. *Jpn. J. Parasitol.* 1967; **16**: 427–435.

49. Kuramochi T, Machida M, Araki J, Uchida A, Kishiro T, Nagasawa K. Minke whales (*Balaenoptera acutorostrata*) are one of the major hosts of *Anisakis simplex* (Nematoda: Anisakidae) in the northwestern North Pacific Ocean. *Rep. Int. Whal. Comm.* 1996; **46**: 415–419.
50. Miyashita T, Hatanaka H. A note on whaling grounds for the western north Pacific minke whale. *Rep. Int. Whal. Comm.* 1997; **47**: 539–542.
51. Gibson DI, Harris EA, Bray RA, Jepson PD, Kuiken T, Baker JR, Simpson VR. A survey of the helminth parasites of cetaceans stranded on the coast of England and Wales during the period 1990–1994. *J. Zool. Soc. Lond.* 1998; **244**: 563–574.
52. González T, Acuña E. Influence of host size and sex on the endohelminth infracommunities of the red rockfish *Sebastes capensis* off northern Chile. *J. Parasitol.* 2000; **86**: 854–857.
53. Smith JW. *Anisakis simplex* (Rudolphi, 1809 det. Krabbe, 1878): Length distribution and variability of L3 of known minimum age from herring *Clupea harengus* L. *J. Helminthol.* 1984; **58**: 337–340.
54. Hemmingsen W, Lysne DA, Eidnes T, Skorping A. The occurrence of larval ascaridoid nematodes in wild-caught and in caged and artificially fed Atlantic cod, *Gadus morhua* L., in Norwegian waters. *Fish. Res.* 1993; **15**: 379–386.
55. Bratney J. Effect of temperature on egg hatching in three ascaridoid nematode species from seals. In: Bowen WD (ed.). *Population Biology of Sealworm (Pseudoterranova decipiens) in Relation to its Intermediate and Seal Hosts.* *Can. Bull. Fish. Aquat. Sci.* 1990; **222**: 27–39.
56. Shimazu T, Oshima T. Some larval nematodes from euphausiid crustaceans. In: Anraku M, Bense K, Kawamura T, Nishizawa S, Parsons TR, Takenouti Y, Tsujita T (eds). *Biological Oceanography of the Northern North Pacific.* Idemitsu Shoten, Tokyo. 1972; 403–409.
57. Boily FD, Marcogliese J. Geographical variations in abundance of larval anisakine nematodes in Atlantic cod (*Gadus morhua*) and American plaice (*Hippoglossoides platessoides*) from the Gulf of St Lawrence. *Can. J. Fish. Aquat. Sci.* 1995; **52** (Suppl. 1): 105–115.
58. Nagasawa K. Parasites of pinnipeds (Mammalia: Carnivora) in Japan: Checklist and bibliography. *Bull. Nat. Res. Inst. Far Seas Fish.* 1999; **36**: 27–32.
59. Machida M. Survey on gastric nematodes of the northern fur seal on breeding islands. *Jpn. J. Parasitol.* 1971; **20**: 371–378.
60. Burns JJ. Ribbon seal *Phoca fasciata* Zimmermann, 1783. In: Ridgway SH, Harrison RJ FRS (eds). *Handbook of Marine Mammals*, Vol. 2: *Seals*. Academic Press, London. 1981; 89–109.
61. Shultz LM, Frost KJ. Helminth parasites of ribbon seals, *Phoca fasciata*, in the Bering Sea and their intermediate hosts. *Proc. Helminthol. Soc. Wash.* 1988; **55**: 68–73.