Title	Direct synthesis of alkynyl(phenyl)iodonium salts from 1-alkynes
Author(s)	Yoshida, Masanori; Nishimura, Naoya; Hara, Shoji
Citation	Chemical Communications, 2002(9), 1014-1014 https://doi.org/10.1039/b200659f
Issue Date	2002
Doc URL	http://hdl.handle.net/2115/36648
Rights	Chem. Commun., 2002, 1014 - Reproduced by permission of The Royal Society of Chemistry (RSC)
Type	article (author version)
File Information	hara-1.pdf

[Instructions for use](#)

Direct synthesis of alkynyl(phenyl)iodonium salts from 1-alkynes

Masanori Yoshida, Naoya Nishimura, and Shoji Hara*

Division of Molecular Chemistry, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan. E-mail: hara@org-mc.eng.hokudai.ac.jp

This submission was created using the RSC Communication template (DO NOT DELETE THIS TEXT) (LINE INCLUDED FOR SPACING ONLY - DO NOT DELETE THIS TEXT)

Alkynyliodonium salts can be directly prepared from 1-alkynes by the reaction with iodosobenzene, tetrafluoroboric acid, and a catalytic amount of HgO.

Alkynyl(aryl)iodonium salts have been recently used as versatile reagents for organic synthesis.¹ They are generally prepared from 1-alkynes in two-steps via the corresponding 1-trimethylsilyl- or 1-trialkylstannyl derivatives.² Direct conversion of 1-alkynes to alkynyliodonium salts is more efficient and desirable, and has been studied by many chemists.³ However, the direct synthetic methods, which include the reaction of 1-alkynes with {hydroxy(sulfonyloxy)iodo}arenes, were applicable only for the synthesis of aryl or sterically hindered alkyl group substituted alkynyliodonium salts, and in other cases, the competitive formation of alkenyliodonium salts was a serious problem.^{3a,b} Therefore, the two-step method has been the sole way for the synthesis of the alkynyliodonium salts with a normal alkyl group. We wish to report here the direct synthesis of 1-alkynyliodonium salts from 1-alkynes of the normal alkyl chain without the formation of the alkenyliodonium salts.

To a CH₂Cl₂ solution of iodosobenzene were added an aqueous solution of HBF₄ and a catalytic amount of HgO to give two liquid phases, a clear organic phase and a clear yellow aqueous phase. When 1-alkyne was added to the mixture, the yellow color of the aqueous phase faded, and the formation of 1-alkynyl(phenyl)iodonium salt and the absence of the 1-alkyne and the alkenyliodonium salt could be confirmed from ¹HNMR of the mixture. The 1-alkynyl(phenyl)iodonium salts could be isolated in good yields as BF₄ salts (Scheme 1).

Scheme 1.

R. Caple *et al.* succeeded to prepare pentynyl(phenyl)iodonium tetrafluoroborate⁴ from 1-pentyne and iodosobenzene tetrafluoroborate.⁴ However, the synthesis of any other alkynyliodonium salts was not shown and it is a serious disadvantage of their method to use a large excess of 1-pentyne (ca. 40 times) to iodosobenzene tetrafluoroborate. According to our method, various kinds of the alkynyl(phenyl)iodonium salts can be prepared from the corresponding 1-alkynes. The alkynes having both normal alkyl group and sterically hindered one can be converted to the alkynyliodonium salts without a large excess reagents. Moreover, the introduction of functional groups such as an ester, ketone, or chloride is also possible. (Table 1).[†] It is drawback to use a harmful mercury salt but less than 0.5 mol% of HgO is enough to obtain the alkynyliodonium salts in good yield. It is also advantage of our method that PhIO, HBF₄, and HgO are all commercially available, while iodosobenzene tetrafluoroborate is not.

Notes and references

[†] Typical experimental procedure: To a CH₂Cl₂ suspension (2 ml) of iodosylbenzene (132 mg, 0.6 mmol) were added 42% aqueous solution of HBF₄ (7.2 ml, 3 mmol) and HgO (0.54 mg, 0.0025 mmol) at room temperature, and the mixture was stirred for a few minutes until the solid part dissolved completely. To the resulting mixture, 1-dodecyne (83 mg, 0.5 mmol) was added and the mixture was stirred at room temperature until the yellow color of the aqueous phase disappeared. Then the mixture was poured into a 5% aqueous solution of NaBF₄ (20 ml, 1 mmol) and the separated aqueous phase was extracted with CH₂Cl₂ three times. The combined organic phases were dried over MgSO₄ and concentrated under reduced pressure. The product was solidified by dissolving the resulting viscous liquid in a little CH₂Cl₂, followed by the addition of a large quantity of hexane. The liquid part was removed by decantation, and the remained solid was washed with hexane. Finally, the solvent was removed completely under reduced pressure to give 1-dodecynyl(phenyl)iodonium tetrafluoroborate (173 mg, 0.37 mmol). Characterization data for 1-dodecynyl(phenyl)iodonium tetrafluoroborate: white solid, M.p. 52 °C. δ_{H} (400 MHz, CDCl₃): 8.05 (2H, d, *J* = 8.1 Hz), 7.68–7.54 (3H, m), 1.64–1.56 (2H, m), 1.36–1.25 (16H, m), 0.88 (3H, t, *J* = 6.7 Hz). δ_{C} (100.4 MHz, CDCl₃): 133.82, 132.81, 132.63, 114.47, 114.02, 31.82, 29.47, 29.34, 29.32, 28.85, 28.73, 27.54, 22.63, 20.83, 15.86, 14.08. ν (KBr) cm⁻¹ 2170, 1057, 739; HRMS (FAB): *m/z* 369.1096 (M-BF₄)⁺ calcd for C₁₈H₂₆I 369.1079.

Table 1 Direct synthesis of alkynyl(phenyl)iodonium salts^a

R	Reaction time (min)	Yield(%) ^b
Bu	30	61
^t Bu	40	86
C ₁₀ H ₂₁	30	76
-CH ₂	60	78
Cl-(CH ₂) ₉	30	80
^t BuC(=O)-(CH ₂) ₉	15	79 ^c
MeOOC-(CH ₂) ₉	15	78 ^c
AcO-(CH ₂) ₉	60	54

^a The reaction was carried out as described in the note. ^b Isolated yield based on alkyne used. ^c Isolated as liquid.

- 1 P. J. Stang, *Angew. Chem. Int. Ed. Engl.*, **1992**, *31*, 274; A. Vargolis, *The Organic Chemistry of Polycordinated Iodine*, VCH Publishers, Inc. New York, 1992, pp. 267-277; P. J. Stang, V. V. Zhdankin, *Chem. Rev.*, **1996**, *96*, 1123; A. Vargolis, *Tetrahedron*,

- 1997**, 53, 1179; M. Ochiai, *Chemistry of Hypervalent Compounds*, ed. K. Akiba, Wiley-VCH, New York, 1999, pp. 359-387.
- 2 M. Ochiai, M. Kunishima, K. Sumi, Y. Nagao, E. Fujita, M. Arimoto, and H. Yamaguchi, *Tetrahedron Lett.*, **1985**, 26, 4501; T. Kitamura and P. J. Stang, *J. Org. Chem.*, **1988**, 53, 4105; P. J. Stang, A. M. Arif, and C. M. Crittall, *Angew. Chem. Int. Ed. Engl.*, **1990**, 29, 287; T. Kitamura, M. Kotani, and Y. Fujiwara, *Synthesis*, **1998**, 1416.
 - 3 a) G. F. Koser, L. Rebrovic, and R. H. Wettach, *J. Org. Chem.*, **1981**, 46, 4324. b) L. Rebrovic and G. F. Koser, *J. Org. Chem.*, **1984**, 49, 4700. c) P. J. Stang, B. W. Surber, Z.-C. Chen, K. A. Roberts, and A. G. Anderson, *J. Am. Chem. Soc.*, **1987**, 109, 228. d) T. Kitamura, R. Furuki, H. Taniguchi, and P. J. Stang, *Tetrahedron*, **1992**, 48, 7149. e) A. B. Sheremetev and E. V. Mantseva, *Tetrahedron Lett.*, **2001**, 42, 5759.
 - 4 V. V. Zhdankin, R. Tykwinski, B. Berglund, M. Mullikin, R. Caple, N. S. Zefirov, and A. S. Koz'min, *J. Org. Chem.* **1989**, 54, 2609.