



Title	Are Sea Surface Temperature Variations in the Kuroshio Extension and Subarctic Frontal Zones in the Western North Pacific Ocean Coherent?
Author(s)	Nonaka, Masami; Nakamura, Hisashi
Citation	Edited by Hisatake Okada, Shunsuke F. Mawatari, Noriyuki Suzuki, Pitambar Gautam. ISBN: 978-4-9903990-0-9, 201-205
Issue Date	2008
Doc URL	http://hdl.handle.net/2115/38466
Type	proceedings
Note	International Symposium, "The Origin and Evolution of Natural Diversity". 1-5 October 2007. Sapporo, Japan.
File Information	p201-205-origin08.pdf



[Instructions for use](#)

Are Sea Surface Temperature Variations in the Kuroshio Extension and Subarctic Frontal Zones in the Western North Pacific Ocean Coherent?

Masami Nonaka^{1,*} and Hisashi Nakamura^{1,2}

¹*Frontier Research Center for Global Change, JAMSTEC, Yokohama, Japan*

²*University of Tokyo, Tokyo, Japan*

ABSTRACT

The western North Pacific region is one of the centers of action of decadal time scale variations in the atmosphere and ocean, and also oceanic ecosystem can be strongly influenced by the variations. One of the key variable of the decadal variations is sea surface temperature (SST) that varies as a result of interaction between the atmosphere and ocean. To improve our understanding of the key element, SST variations in the two frontal zones in the western North Pacific Ocean, the Kuroshio Extension and subarctic frontal zones, are investigated on the basis of an *in situ* observational dataset. Interannual-to-decadal variations in these two frontal zones are not highly correlated, indicating that to some extent different mechanisms induce the SST variations in the frontal zones and those two frontal zone cannot be considered as a single frontal zone as has been done in most of previous studies. Meanwhile, the results are consistent with the recent studies that suggest different mechanisms for the variations in the two frontal zones on the basis of a solution to an eddy-resolving, i.e., very high horizontal resolution, ocean general circulation model.

Keywords: North Pacific, Sea surface temperature, Decadal variability

INTRODUCTION

It has been well known that populations of sardine and anchovy vary on (inter-) decadal timescale with amplitude of more than one order (e.g., [1]). For the Japanese sardine, Ref. 2 showed that sea surface temperature anomalies (SSTAs) in the western North Pacific (NP) region have crucial impact on the natural mortality coefficient for the period from the postlarval stage to age 1, and thus the population, although how the SSTAs affect the oceanic ecosystem in the region has not been clarified yet. The western NP region is also known as one of the centers of action of Pacific decadal variability ([3], among others), and several studies have suggested

that air-sea interactions in the region may have a key role to induce or intensify the decadal variations [4–6]. The corresponding decadal SSTAs in the western NP are most prominent along the so-called Kuroshio-Oyashio Extension (KOE) front, the zonal oceanic front between the NP subtropical and subarctic gyres. Thus, to improve our knowledge on decadal variations in the atmosphere, ocean and ecosystem in the NP, it is crucial to understand how the SSTAs in the KOE region are induced and if the SSTAs have feedback to force atmospheric variations.

Ref. 7 then investigated the mechanism for the SSTAs on the basis of a hindcast integration of an eddy-resolving ocean general circulation model

*Corresponding author, e-mail: nona@jamstec.go.jp

Information for use in citing this article: Okada, H., Mawatari, S.F., Suzuki, N. and Gautam, P. (eds.), *Origin and Evolution of Natural Diversity*, Proceedings of International Symposium “The Origin and Evolution of Natural Diversity”, 1–5 October 2007, Sapporo, pp. 201–205.

(OGCM), the OGCM for the Earth Simulator (OFES) [8, 9], focusing on variations in the Kuroshio Extension frontal zone (KEFZ, $\sim 35^\circ\text{N}$) and in the Oyashio Extension, the subarctic, frontal zone (SAFZ, $\sim 42^\circ\text{N}$), the two prominent frontal zones in the KOE region. Their results indicated that decadal SSTAs in the western NP region have their maxima in the two frontal zones, especially in the SAFZ, where SST has the largest meridional gradient (Fig. 1), suggesting importance of meridional migration of the frontal zones. Latitude-depth section of temperature difference apparently shows that the two frontal zones migrate meridionally in association with the decadal SSTAs, inducing temperature anomalies in the surface (subsurface) layer in the SAFZ (KEFZ) due to different vertical structures of the frontal zones (Fig. 2). Although vertical displacement of thermocline also has some contribution to subsurface temperature anomaly, influences of the associated changes in warm water advection by the Kuroshio Extension Current are not apparent in the model.

In the SAFZ, heat flux anomalies appear in association with the SST anomalies, and the former tend to damp the latter, indicating that the SST anomalies are not caused by atmospheric thermal forcing [7]. In other words, the SST anomalies induced in the frontal zone by the aforementioned oceanic processes can have feedback on the atmosphere through modifying heat flux to the atmosphere. This cool (warm) SST-downward (upward) surface heat flux anomaly relation is also shown by Ref. 10 in the SAFZ on the basis of an *in situ* observational dataset.

The results of the previous studies indicate that the SAFZ has higher SST variability and higher possibility to affect the atmosphere than the KEFZ, suggesting that the two frontal zones have different properties. Then, do the interannual SSTAs in the SAFZ and KEFZ vary coherently? In this study, we investigate this based on a dataset of observed SST compiled by the Japan Meteorological Agency (JMA-SST).

RESULTS

Figure 3a indicates that wintertime SST variance is large in the SAFZ, in the Oyashio frontal zone (the western end of the SAFZ with southwest-northeast extending strong SST gradient), and in the upstream region of the KEFZ. To investigate if these SST variations are temporally coherent or not, we plot a simultaneous correlation map (Fig. 3b) of wintertime SST on the area mean SSTAs in a part of the SAFZ (black rectangular in Fig. 3a, b). High correlation region extends along the SAFZ and also to the Oyashio frontal zone. However, correlations gradually decrease southward to the KEFZ, especially in the upstream region (around $140\text{--}150^\circ\text{E}$), and negative correlations appear to the south of 35°N , the southern recirculation region of the Kuroshio Extension Current. Consequently simultaneous correlations between the SAFZ and the KEFZ are less than 0.5, and lower in the upstream region.

Time series of area mean SSTAs in the SAFZ and in the upstream KEFZ ($32^\circ\text{--}38^\circ\text{N}$, $141^\circ\text{--}153^\circ\text{E}$, following Ref. 11) indicate that they are not coherent




Fig. 1 Wintertime (January-March) mean SST fields over the North Pacific based on the OFES hindcast simulation. Contours indicate the five-winter mean for 1984–88 (every 1°C), and shading indicates the difference of that mean field from another five-winter mean for 1968–72, as indicated to the right of the panel.




Fig. 2 Latitude-depth sections of five-winter (January–March) mean temperature fields for the (top) 1968–72, and (bottom) 1984–88 periods averaged between 145° and 160°E based on the OFES hindcast simulation. Contour intervals are 1°C, and the temperature anomalies from the 1950–2003 mean field are plotted with shading, as indicated below the bottom panel.

on interannual and also decadal time scales (for example, from the end of 1960s to the early 1970s, SSTAs tend to be positive in the SAFZ but they tend to be slightly negative in the KEFZ, and negative peak in the 1980s appeared several years earlier in the KEFZ than in the SAFZ). As a result, their correlation coefficient (r) is only 0.37, and if we see area mean SSTA in [32°–36°N, 141°–153°E], excluding the Oyashio frontal zone from the upstream KEFZ in Ref. 11, their correlation becomes lower further to $r = 0.15$ (Fig. 3c, thin line). These results indicate that SST variations in the SAFZ and in the KEFZ are not highly correlative, and it can be misleading to consider them as one single frontal zone as has been done in most of previous studies on variations in the western NP Ocean, owing to rather coarse horizontal resolution of the models or datasets used.

SUMMARY AND DISCUSSION

In this study, we investigate whether interannual variations in the two frontal zones in the so-called KOE region, the KEFZ and the SAFZ, are coherent on the basis of an observational SST data product, JMA-SST. The result of the correlation analysis indicates that the interannual-to-decadal SST variations in the SAFZ and KEFZ are not highly correlative, indicating that to some extent those variations are governed by different mechanisms.

Previous studies have suggested that decadal timescale variations in the KOE region are strongly influenced by westward propagation of Rossby waves driven by wind-stress curl anomalies over the central portion of the North Pacific ([12, 13] among others). Ref. 7 confirmed this process in the KEFZ on the basis of an eddy-resolving OGCM, while Ref. 14 further pointed out a probable role of some

JMA-SST correlation based on SAF SST (1950-1999)




Fig. 3 Winter (January-March) mean SST based on the JMA-SST for 1950-1999. (a) Long-year mean (contours with intervals of 1°C) and standard deviations of interannual variations (shadings as indicated to the right of the panel). (b) Simultaneous correlation map of winter mean SST on the area mean SSTA in [38.5°-43.5°N, 155.5°-175.5°E] in the SAFZ (black rectangular in the panels a and b). (c) Time series of area mean SSTAs standardized by their standard deviations for [38.5°-43.5°N, 155.5°-175.5°E] (thick solid line), [32°-38°N, 141°-153°E] (thick dashed line), and [32°-36°N, 141°-153°E] (thin dashed line).

nonlinear oceanic processes to induce prominent anomalies strongly trapped to the narrow oceanic front. In contrast to the KEFZ, westward propagations of anomalies are sometimes not clear in the SAFZ, and some eastward propagating signals appear in the western portion of the basin [7; Fig. 12 there], suggesting the mechanisms that induce interannual-to-decadal variations in the SAFZ to be different than those in the KEFZ. Indeed, using the solution to the same eddy-resolving OGCM, Ref. 15 indicated that variations in the Oyashio Current tend to induce SST anomalies that develop eastward in the SAFZ from off Hokkaido Island. That these different mechanisms are suggested by an OGCM for variations in each of the KEFZ and the SAFZ is consistent with the SST variations, with not very high degree of correlation, in those frontal zones indicated in this study.

ACKNOWLEDGMENTS

We thank Prof. S. Minobe who gave one of us (M. N.) an opportunity to present results, described in this paper, at the international symposium “The Origin and Evolution of Natural Diversity”. This study was supported by the Frontier Research Center for Global Change. A part of this study was carried out under a research project of the Agriculture, Forestry and Fisheries Research Council.

REFERENCES

1. Chavez, F.P., Ryan, J., Kluch-Cota, S.E. and Niquen, M., 2003. From anchovies to sardines and back: multidecadal change in the Pacific Ocean. *Science*, 299, 217–221.
2. Noto, M. and Yasuda, I. 1999. Population decline of the Japanese sardine, *Sardinops melanostictus*, in relation to sea surface temperature in the Kuroshio Extension. *Can. J. Fish. Aquat. Sci.*, 56, 973–983.
3. Nakamura, H., Lin, G. and Yamagata, T., 1997. Decadal climate variability in the North Pacific during the recent decades. *Bull. Amer. Meteor. Soc.*, 78, 2215–2225.
4. Latif, M. and Barnett, T.P., 1994. Causes of decadal climate variability over the North Pacific and North America. *Science*, 266, 634–637.
5. Pierce, D.W., Barnett, T.P., Schneider, N., Saravanan, R., Dommengat, D. and Latif, M., 2001. The role of ocean dynamics in producing decadal climate variability in the North Pacific. *Clim. Dyn.* 18, 51–70.
6. Schneider, N. and Cornuelle, B.D., 2005. The forcing of the Pacific Decadal Oscillation. *J. Climate*, 18, 4355–4373.
7. Nonaka, M., Nakamura, H., Tanimoto, Y., Kagimoto, T. and Sasaki, H., 2006. Decadal variability in the Kuroshio-Oyashio Extension simulated in an eddy-resolving OGCM. *J. Climate*, 19, 1970–1989.
8. Masumoto, Y., Sasaki, H., Kagimoto, T., Komori, N., Ishida, A., Sasai, Y., Miyama, T., Motoi, T., Mitsudera, H., Takahashi, K., Sakuma, H. and Yamagata, T. 2004. A fifty-year eddy-resolving simulation of the World Ocean -preliminary outcomes of OFES (OGCM for the Earth Simulator)-, *The Journal of the Earth Simulator*, 1, 31–52.
9. Sasaki, H., Nonaka, M., Masumoto, Y., Sasai, Y., Uehara, H. and Sakuma, H., 2007. An eddy-resolving hindcast simulation of the quasi-global ocean from 1950 to 2003 on the Earth Simulator, In: W. Ohfuchi and K. Hamilton (eds.), *High resolution numerical modelling of the atmosphere and ocean*, Springer, New York., 157–186.
10. Tanimoto, Y., Nakamura, H., Kagimoto, T. and Yamane, S., 2003. An active role of extratropical sea surface temperature anomalies in determining anomalous turbulent heat flux. *J. Geophys. Res.*, 108 (C10), 3304, doi: 10.1029/2002JC001750.
11. Qiu, B. and Chen, S., 2005. Variability of the Kuroshio Extension Jet, recirculation gyre, and mesoscale eddies on decadal time scales. *J. Phys. Oceanogr.*, 35, 2090–2103.
12. Miller, A.J. and Schneider, N., 2000. Interdecadal climate regime dynamics in the North Pacific Ocean: theories, observations and ecosystem impacts. *Prog. Oceanogr.*, 47, 355–379.
13. Seager, R., Kushnir, Y., Naik, N.H., Cane M.A. and Miller, J., 2001. Wind-driven shifts in the latitude of the Kuroshio-Oyashio extension and generation of SST anomalies on decadal timescales. *J. Climate*, 14, 4149–4165.
14. Taguchi, B., Xie, S.-P., Schneider, N., Nonaka, M., Sasaki, H. and Sasai, Y. 2007. Decadal variability of the Kuroshio Extension: Observations and an eddy-resolving model hindcast. *J. Climate*, 20, 2357–2377.
15. Nonaka, M., Nakamura, H., Tanimoto, Y., Kagimoto, T. and Sasaki, H., 2007. Interannual-to-decadal variability in the Oyashio Current and its influence on temperature in the subarctic frontal zone: An eddy-resolving OGCM simulation. *J. Climate*, submitted.