



Title	Biomarker Compositions of Dinoflagellates and Their Applications for Paleoenvironmental Proxies
Author(s)	Amo, Miki; Suzuki, Noriyuki; Kawamura, Hiroshi; Yamaguchi, Aika; Takano, Yoshihito; Horiguchi, Takeo
Citation	Edited by Hisatake Okada, Shunsuke F. Mawatari, Noriyuki Suzuki, Pitambar Gautam. ISBN: 978-4-9903990-0-9, 223-226
Issue Date	2008
Doc URL	http://hdl.handle.net/2115/38469
Type	proceedings
Note	International Symposium, "The Origin and Evolution of Natural Diversity". 1-5 October 2007. Sapporo, Japan.
File Information	p223-226-origin08.pdf



[Instructions for use](#)

Biomarker Compositions of Dinoflagellates and Their Applications for Paleoenvironmental Proxies

Miki Amo^{1,*}, Noriyuki Suzuki¹, Hiroshi Kawamura¹, Aika Yamaguchi¹, Yoshihito Takano² and Takeo Horiguchi¹

¹*Department of Natural History Sciences, Graduate School of Science, Hokkaido University,
N10 W8, Kita-ku, Sapporo 060-0810, Japan*

²*Harmful Phytoplankton Section, Harmful Algal Bloom Division, National Research Institute of Fisheries and Environment of Inland
Sea, Fisheries Research Agency of Japan, 2-17-5 Maruishi, Hatsukaichi, Hiroshima 739-0452, Japan*

ABSTRACT

Four motile cells of autotrophic dinoflagellates, *Peridinium umbonatum* var. *inaequale*, *Akashiwo sanguinea*, *Scrippsiella tinctoria*, and *Prorocentrum micans*, commonly contain five major sterols: cholesterol, 4-methylcholestan-3-ol, 4, 24-dimethylcholestan-3-ol, dinosterol and dinostanol. A motile cell of heterotrophic dinoflagellate, *Proto-peridinium crassipes*, contains cholesterol, 4, 24-dimethylcholestan-3-ol, dinosterol, dinostanol and 4-tetramethylcholestan-3-ol as major free sterols. Dinosterol concentration of heterotrophic dinoflagellate is about 4–12 times higher than those of autotrophic species, suggesting that the heterotrophic dinoflagellate is important source of dinosterol in some sediments. 4-Methylcholestan-3-ol and 4-tetramethylcholestan-3-ol occur respectively in autotrophic and heterotrophic dinoflagellates are believed to serve as potential biomarkers in respective types. Resting cyst of *P. inaequale* contains 4-methylcholestan-3 β -ol, 4, 24-dimethyl-5 α -cholestan-3 β -ol, dinosterol, dinostanol, and unknown C₃₁ sterol as major free sterols. The unknown C₃₁ sterol detected only in resting cyst could have been produced during resting stage. This compound may serve as a potential biomarker for resting cysts of dinoflagellate.

Keywords: Autotrophic dinoflagellate, Heterotrophic dinoflagellate, Resting cyst, 4 α -Methyl sterol

INTRODUCTION

Dinoflagellate is one of the major primary producers in the ocean since the Mesozoic. These microalgae occur throughout the world's oceans but are often more abundant in coastal areas. About half species of dinoflagellate are autotrophic ones, and others are heterotrophic. Some species are able to force themselves into a dormant or resting stage as part of their relatively complicated life cycle. These dormant stages, called resting cysts, are typically characterized by a thick and highly specialized cell

covering. The motile stage of dinoflagellate is hardly recorded in sediments since the motile cell of dinoflagellate is labile against bio- and chemical degradations during the settling and the early diagenesis. The resting cysts of some dinoflagellates species are composed of resistant biomacromolecules, which can be preserved in sediments and sedimentary rocks. Geologic record of dinoflagellate evolution, therefore, is based on their resting cyst fossils in sedimentary rocks.

The sterol compositions of dinoflagellates are generally dominated by 4 α -methyl sterols including

*Corresponding author, e-mail: mamo@nature.sci.hokudai.ac.jp

Information for use in citing this article: Okada, H., Mawatari, S.F., Suzuki, N. and Gautam, P. (eds.), *Origin and Evolution of Natural Diversity*, Proceedings of International Symposium "The Origin and Evolution of Natural Diversity", 1–5 October 2007, Sapporo, pp. 223–226.

the C₃₀ sterol called dinosterol (4 α , 23, 24-trimethyl-5 α -cholest-22E-en-3 β -ol). This sterol is rarely found in other algae and hence has been often used as an indicator of dinoflagellate contribution to the marine sediments. However, sedimentary dinosterols do not necessarily provide sufficient information about the motile stage of dinoflagellate and the relative contribution of heterotrophic and autotrophic dinoflagellates. Sterol compositions vary quite considerably between different species of dinoflagellates [1]. Some particular sterols can be potential biomarkers for motile cells of dinoflagellates. The objective of the present study is to search for characteristic sterols of motile cell and resting cyst of autotrophic and heterotrophic dinoflagellates.

MATERIALS AND METHODS

Five motile cells of dinoflagellates (*Peridinium umbonatum* var. *inaequale*, *Akashiwo sanguinea*, *Scrippsiella tinctoria*, *Prorocentrum micans*, *Proto-peridinium crassipes*) and a resting cyst of *P. inaequale* were cultured. After the cultivation, the samples were collected at GF/F filters. Lipids were extracted by ultrasonication with methanol/dichloromethane. The combined lipid extract was saponified with 0.5 mol KOH/methanol. The neutral fraction was extracted with diethylether/*n*-hexane (1 : 9), and was fractionated into four fractions by silica gel column chromatography. The sterol fraction was analyzed using a Hewlett Packard 6890 series gas chromatograph (GC). The compounds were identified by mass spectral analysis using a gas chromatograph-mass spectrometer (GC/MS) (Hewlett Packard GC HP6890 and MS HP5973).

RESULTS AND DISCUSSION

Four motile cells of autotrophic dinoflagellates, *Peridinium umbonatum* var. *inaequale*, *Akashiwo sanguinea*, *Scrippsiella tinctoria*, and *Prorocentrum micans*, commonly contain five major sterols in the free sterol fraction (Fig. 1). These major sterols are cholesterol, 4-methylcholestan-3-ol, 4, 24-dimethylcholestan-3-ol, dinosterol and dinostanol (4, 23, 24-trimethylcholestan-3-ol). A motile cell of heterotrophic dinoflagellate, *Proto-peridinium crassipes*, contains cholesterol, 4, 24-dimethylcholestan-3-ol, dinosterol, dinostanol and 4-tetramethylcholestan-3-ol as major free sterols (Fig. 1). The dinosterol concentrations of autotrophic dinoflagellates vary within a range of 0.18 to 0.52 pg/cell. The dinosterol concentration of heterotrophic dinoflagellate *P. crassipes*

is 2.3 pg/cell, which is about 4–12 times higher than the range for autotrophic species. The genus *Proto-peridinium* currently includes more than 200 spe-




Fig. 1 Total ion chromatograms of sterol fractions from all motile cells of dinoflagellates. 1: Cholesterol, 2: 4 α -Methylcholestan-3 β -ol, 3: Campesterol, 4: β -Sitosterol, 5: 4 α , 24-Dimethyl-5 α -cholestan-3 β -ol, 6: Dinosterol, 7: 4 α -23, 24-Trimethylcholest-17 (20)-en-3 β -ol, 8: 4 α , 23R, 24R-Trimethyl-5 α -cholestan-3 β -ol, 9: 4 α -Tetramethylcholestan-3 β -ol.

cies, and its members are some of the commonest organisms in the neritic plankton [2]. These observations suggest that the heterotrophic dinoflagellate can be an important source of dinosterol in some sediments. The dinosterol flux is not always applicable as the proxy of primary production by dinoflagellates in those sediments, in which heterotrophic dinoflagellates are significant contributors of organic matter.

4-Methylcholestan-3-ol was commonly detected in autotrophic dinoflagellates, but not in heterotrophic dinoflagellate in the present study (Fig. 1). On the contrary, 4-tetramethylcholestan-3-ol was detected only in heterotrophic dinoflagellate (Fig. 1). The sterol composition of dinoflagellates is dominated by 4-methyl sterols, including dinosterol, which is found in many dinoflagellate species [e.g. 3]. Haptophyte algae of the genus *Pavlova* contain 4, 24-dimethyl-5 α -cholest-22E-en-3 β -ol, 4, 24-dimethyl-

5 α -cholestan-3 β -ol, 4-methyl-24-ethyl-5 α -cholest-22E-en-3 β -ol and 4-methyl-24-ethyl-5 α -cholestan-3 β -ol [4]. Bacteria have ever been proposed as one of the sources of 4-methyl sterols in sediments [5], but only a few bacteria contain 4-methyl sterols [6, 7]. Some higher plants also contain C₃₀ 4-methyl sterols. However, they occur generally in small amounts since the sterols are biosynthetic intermediates to other sterols [8]. These observations thus lead to the conclusion that dinoflagellates are major sources of 4-methylcholestan-3-ol and 4-tetramethylcholestan-3-ol. 4-Methylcholestan-3-ol and 4-tetramethylcholestan-3-ol may be useful biomarkers of autotrophic and heterotrophic dinoflagellates, respectively.

Resting cyst of *P. inaequale* contains 4-methylcholestan-3 β -ol, 4, 24-dimethyl-5 α -cholestan-3 β -ol, dinosterol, dinostanol, and unknown C₃₁ sterol as major free sterols (Fig. 2). Although sterol distribution of resting cyst is nearly similar to that of motile cell, the unknown C₃₁ sterol occurred only in resting cyst (Fig. 2). None of motile cells contains this unknown C₃₁ sterol. 4 α -Methylgorgostanol is known as typical C₃₁ sterol produced by dinoflagellate [8]. 4 α -Methylgorgostanol was originally detected in zoxanthellae of *Briareum asbestinum* and dinoflagellate, *Kryptoperidinium* (= *Glenodinium*) *foliaceum* [9, 10]. However, mass spectral study shows that the unknown C₃₁ sterol in the present study is clearly different from 4 α -methylgorgostanol. The unknown C₃₁ sterol has not been detected in the microalgal culture samples. This compound could have been produced during resting stage. Further study on this unknown C₃₁ sterol will provide clues for biomarkers of dinoflagellate resting cysts.




Fig. 2 Total ion chromatograms of sterol fractions from motile cell and resting cyst of *P. inaequale* and mass spectra of C₃₁ sterol. 1: Cholesterol, 2: 4 α -Methylcholestan-3 β -ol, 5: 4 α , 24-Dimethyl-5 α -cholestan-3 β -ol, 6: Dinosterol, 8: 4 α , 23R, 24R-Trimethyl-5 α -cholestan-3 β -ol, 9: 4 α -Tetramethylcholestan-3 β -ol.

CONCLUSIONS

1. The dinosterol concentration of heterotrophic dinoflagellate is about 4–12 times higher than those of autotrophic species, suggesting that the heterotrophic dinoflagellate can be an important source of dinosterol in some sediments.

2. 4-Methylcholestan-3-ol was detected only in autotrophic dinoflagellates, whereas 4-tetramethylcholestan-3-ol was confined to the heterotrophic dinoflagellates. These compounds may serve as potential biomarkers in the respective types of dinoflagellates.

3. The unknown C₃₁ sterol detected in resting cyst of *P. inaequale* could be a potential biomarker of dinoflagellate resting cysts.

ACKNOWLEDGMENTS

This work was supported in part by the 21st Century COE Program for the “Neo-Science of Natural History” Program (Leader: Prof. H. Okada) under a grant from the Japanese Ministry of Education, Science, Sports and Culture.

REFERENCES

1. Volkman, J.K., 1999. Sterols of four dinoflagellates from the genus *Prorocentrum*. *Phytochem.*, 52, 659–668.
2. Yamaguchi, A. and Horiguchi, T., 2005. Molecular phylogenetic study of the heterotrophic dinoflagellate genus *Protoperidinium* (Dinophyceae) inferred from small subunit rRNA gene sequences. *Phycol. Res.*, 53, 30–42.
3. Pirretti, M.V., Giampiero, P., Boni, L., Pistocchi, R., Diamante, M. and Gazzotti, T., 1997. Investigation of 4-methyl sterols from cultured dinoflagellate algal strains. *Journal of Phycology*, 33, 61–67.
4. Volkman, J.K., Kearney, P. and Jeffrey, S.W., 1990. A new source of 4-methyl sterols and 5 α (H)-stanols in sediments: prmnesiophyte microalgae of the genus *Pavlova*. *Org. Geochem.*, 15, 489–497.
5. Mermoud, F., Glucar, F.O., Siles, S., Chassaing, B. and Buchs, A., 1982. 4-Methylsterols in recent lacustrine sediments: terrestrial, planktonic or some other origin? *Chemosphere*, 11, 557–567.
6. Bird, C.W., Lynch, J.M., Pirt, F.J., Reid, W.W., Brooks, C.J. W. and Middleditch, B.S., 1971. Steroids and squalene in *Methylococcus capsulatus* grown on methane. *Nature*, 230, 473–474.
7. Bouvier, P., Rohmer, M., Benveniste, P. and Ourisson, G., 1976. $\Delta^{8(14)}$ -steroids in the bacterium *Methylococcus capsulatus*. *Biochem. J.*, 159, 267–271.
8. Goad, L. and Goodwin, T.W., 1972. The biosynthesis of plant sterols. *Progr. Phytochem.*, 3, 113–198.
9. Alam, M., Martin, G.E., Ray, S.M., 1979. Dinoflagellate sterols. 2. Isolation and structure of 4-methyl gorgostanol from the dinoflagellate *Glenodinium foliaceum*. *J. Org. Chem.*, 44, 4466–4467.
10. Stuedler, P.A., Schmitz, F.J. and Ciereszko, L.S., 1977. Chemistry of coelenterates. Sterol composition of some predatory-prey pairs on coral reefs. *Comp. Biochem. Physiol.*, 56B, 385–392.