Title	Geometry-driven shift in the Tomonaga-Luttinger exponent of deformed cylinders
Author(s)	Shima, Hiroyuki; Yoshioka, Hideo; Onoe, Jun
Citation	Physical Review B, 79(20), 201401 https://doi.org/10.1103/PhysRevB.79.201401
Issue Date	2009-05
Doc URL	http://hdl.handle.net/2115/38684
Rights	© 2009 The American Physical Society
Type	article
File Information	79-20_201401.pdf

[Instructions for use](#)

Geometry-driven shift in the Tomonaga-Luttinger exponent of deformed cylinders

Hiroyuki Shima,^{1,*} Hideo Yoshioka,² and Jun Onoe³

¹*Department of Applied Physics, Graduate School of Engineering, Hokkaido University, Sapporo 060-8628, Japan*

²*Department of Physics, Nara Women's University, Nara 630-8506, Japan*

³*Department of Nuclear Engineering and Research Laboratory for Nuclear Reactors, Tokyo Institute of Technology, 2-12-1 Ookayama, Meguro, Tokyo 152-8550, Japan*

(Received 10 March 2009; published 15 May 2009)

We demonstrate the effects of geometric perturbation on the Tomonaga-Luttinger liquid (TLL) states in a long, thin, and hollow cylinder whose radius varies periodically. The variation in the surface curvature inherent to the system gives rise to a significant increase in the power-law exponent of the single-particle density of states. The increase in the TLL exponent is caused by a curvature-induced potential that attracts low-energy electrons to region that has large curvature.

DOI: 10.1103/PhysRevB.79.201401

PACS number(s): 73.21.Hb, 02.40.-k, 03.65.Ge, 71.10.Pm

Studying the quantum mechanics of a particle confined to curved surfaces has been a problem for more than 50 years. The difficulty arises from operator-ordering ambiguities,¹ which permit multiple consistent quantizations for a curved system. The conventional method used to resolve the ambiguities is the confining-potential approach.^{2,3} In this approach, the motion of a particle on a curved surface (or, more generally, a curved space) is regarded as being confined by a strong force acting normal to the surface. Because of the confinement, quantum excitation energies in the normal direction are raised far beyond those in the tangential direction. Hence, we can safely ignore the particle motion normal to the surface, which leads to an effective Hamiltonian for propagation along the curved surface with no ambiguity.

It is well known that the effective Hamiltonian involves an effective scalar potential whose magnitude depends on the local surface curvature.²⁻⁵ As a result, quantum particles confined to a thin curved layer behave differently from those on a flat plane, even in the absence of any external field (except for the confining force). Such curvature effects have gained renewed attention in the last decade, mainly, because of the technological progress that has enabled the fabrication of low-dimensional nanostructures with complex geometry.^{6,8-11,7,12-14} From the theoretical perspective, many intriguing phenomena pertinent to electronic states,¹⁵⁻²² electron diffusion,²³ and electron transport²⁴⁻²⁷ have been suggested. In particular, the correlation between surface curvature and spin-orbit interaction^{28,29} as well as with the external magnetic field³⁰⁻³² has been recently considered as a fascinating subject.

Most of the previous works focused on noninteracting electron systems, though few have focused on interacting electrons³³ and their collective excitations. However, in a low-dimensional system, Coulombic interactions may drastically change the quantum nature of the system. Particularly noteworthy are one-dimensional systems, where the Fermi-liquid theory breaks down so that the system is in a Tomonaga-Luttinger liquid (TLL) state.³⁴ In a TLL state, many physical quantities exhibit a power-law dependence stemming from the absence of single-particle excitations near the Fermi energy; this situation naturally raises the question as to how geometric perturbation affects the TLL behaviors of quasi-one-dimensional curved systems. Peanut-

shaped C₆₀ polymers^{35,36} and MoS₂ hollow nanotubes³⁷ are exemplary materials to be considered for studying TLL behaviors; they are thin, long, and hollow tubules whose radius is periodically modulated along the tube axis. Hence, the periodic surface curvature intrinsic to the systems will produce sizable effects on their TLL properties if they exhibit one-dimensional metallic properties.

In this Rapid Communication, we examined the curvature effects on the TLL states in quantum hollow cylinders with a periodically varying radius. We demonstrate that the presence of a curvature-induced potential can yield a significant increase in the power-law exponent α of the single-particle density of states $n(\omega)$ near the Fermi energy E_F ; i.e., $n(\omega) \propto |\hbar\omega - E_F|^\alpha$.^{38,39} The geometric conditions required for the shift in α to be observable are within the realm of laboratory experiments, which implies that our predictions can be verified with existing materials.

We first considered noninteracting spinless electrons confined to a general two-dimensional curved surface S embedded in a three-dimensional Euclidean space. A point $\mathbf{p}$ on S is represented by $\mathbf{p} = [x(u^1, u^2), y(u^1, u^2), z(u^1, u^2)]$, where (u^1, u^2) is a curvilinear coordinate spanning the surface and (x, y, z) are the Cartesian coordinates in the embedding space. Using the notation $\mathbf{p}_i \equiv \partial \mathbf{p} / \partial u^i$ ($i = 1, 2$), we introduced the following quantities $g_{ij} = \mathbf{p}_i \cdot \mathbf{p}_j$, $h_{ij} = \mathbf{p}_{ij} \cdot \mathbf{n}$, and $\mathbf{n} = (\mathbf{p}_i \times \mathbf{p}_j) / \|\mathbf{p}_i \times \mathbf{p}_j\|$, where $\mathbf{n}$ is the unit vector normal to the surface. Using the confining-potential approach,^{2,3} we obtained the Schrödinger equation for noninteracting electron systems on curved surfaces as follows:

FIG. 1. (Color online) Schematic illustration of a quantum hollow cylinder with periodic radius modulation.

FIG. 2. (Color online) Profiles of the curvature-induced effective potential $U_n(\xi)$ for one period $[0, \Lambda]$. Geometric parameters $r_0=4.0$ and $\lambda=8.0$ in units of a are fixed. Integers n represent the angular momentum of eigenstates in the circumferential direction of a hollow tube.

$$-\frac{\hbar^2}{2m^*} \left[\frac{1}{\sqrt{g}} \sum_{i,j=1}^2 \frac{\partial}{\partial u^i} \sqrt{g} g^{ij} \frac{\partial}{\partial u^j} + (\mathcal{H}^2 - \mathcal{K}) \right] \Psi = E \Psi, \quad (1)$$

where $g = \det(g_{ij})$, $g^{ij} = g_{ij}^{-1}$,⁴⁰ and m^* is the effective mass of electrons. The quantities $\mathcal{K} = (h_{11}h_{22} - h_{12}^2)/g$ and $\mathcal{H} = (g_{11}h_{22} + g_{22}h_{11} - 2g_{12}h_{12})/(2g)$ are the so-called Gaussian curvature and mean curvature, respectively, both of which are functions of (u^1, u^2) . The term proportional to $\mathcal{H}^2 - \mathcal{K}$ in Eq. (1) is the effective scalar potential induced by the surface curvature.

We next focused on a hollow tube with a periodically varying radius represented by $\mathbf{p} = [r(z)\cos\theta, r(z)\sin\theta, z]$ (see Fig. 1). The tube radius $r(z)$ is periodically modulated in the axial z direction as $r(z) = r_0 - \frac{\delta r}{2} + \frac{\delta r}{2} \cos(\frac{2\pi}{\lambda}z)$, where the parameters r_0 and δr are introduced to express the maximum and minimum of $r(z)$ as r_0 and $r_0 - \delta r$, respectively. Because of the rotational symmetry, the eigenfunctions of the system have the form of $\Psi(z, \theta) = e^{in\theta} \psi_n(z)$. Thus, the problem reduces to the one-dimensional Schrödinger equation

$$-\frac{\hbar^2}{2m^*} \left[\mathcal{D} - \frac{n^2}{r^2} + (\mathcal{H}^2 - \mathcal{K}) \right] \psi_n(z) = E \psi_n(z), \quad (2)$$

where $\mathcal{D} = \frac{1}{r} \frac{d}{dz} r \frac{d}{dz}$, $f(z) = \sqrt{1 + r'^2}$, $\mathcal{K} = -r''/(rf^2)$, and $\mathcal{H} = (f^2 - rr'')/(2rf^3)$ with $r' \equiv dr/dz$.

Equation (2) is simplified by using a new variable $\xi = \xi(z) = \int_0^z f(\eta) d\eta$, which corresponds to the line length along the curve on the surface with a fixed θ . Straightforward calculation yields¹⁹

$$\left[-a^2 \frac{d^2}{d\xi^2} + U_n(\xi) \right] \psi_n(\xi) = \varepsilon \psi_n(\xi), \quad \varepsilon = \frac{2m^* a^2 E}{\hbar^2} \quad (3)$$

with $U_n(\xi) = (n^2 - \frac{1}{4})a^2/r^2 - r''^2 a^2/(4f^6)$, where r , r'' , and f are regarded as the functions of ξ using the inverse relation $z = z^{-1}(\xi)$. In order to derive Eq. (3), we introduced the length scale a and then multiplied both sides of Eq. (2) with $2m^* a^2/\hbar^2$ to make the units of U_n and ε dimensionless. Notice that by the definition of $\xi(z)$, U_n is periodic with a period $\Lambda = \xi(\lambda)$ depending on r_0 and δr (as well as λ). Figure 2

FIG. 3. (Color online) Energy-band structure of a sinusoidal hollow tubule with $r_0=4.0$ and $\lambda=8.0$ in units of a .

shows the spatial profile of U_n within one period; throughout the present work, we fixed $r_0=4.0$ and $\lambda=8.0$ in units of a by simulating the geometry of actual peanut-shaped C_{60} polymers whose geometry is reproduced by imposing $a=1$ Å. Mapping the discrete atomic structure of one-dimensional C_{60} polymers to a continuum curved surface is based on the result of first-principle calculations⁴¹ which indicated that π electrons on the polymers are almost free from its atomic configurations. We found in Fig. 2 that U_n takes extrema at $\xi=0$ (or Λ) and $\xi=\Lambda/2$, where r takes the maximum ($r=r_0$) and the minimum ($r=r_0 - \delta r$) values, respectively.

To solve Eq. (3), we use the Fourier series expansions $U_n(\xi) = \sum_G U_G^{(n)} e^{iG\xi}$ and $\psi_n(\xi) = \sum_k c_k^{(n)} e^{ik\xi}$, where $G = \pm 2\pi j/\Lambda$ ($j=0, 1, 2, \dots$). Substituting the expansions into Eq. (3), we obtain the secular equation $(\frac{\hbar^2}{2m^*} k^2 - \varepsilon) c_k^{(n)} + \sum_{G=-G_c}^{G_c} U_G^{(n)} c_{k-G}^{(n)} = 0$ that holds for all possible k 's and n 's. The summation has been truncated by $G_c = 20\pi/\Lambda$ because of the rapid decay of U_G with $|G|$. We then numerically calculated eigenvalues for $0 \leq k \leq \pi/\Lambda$ and evaluated the low-energy-band structure for several different δr , as depicted in Fig. 3. In all the cases, there is some energy gap at the Brillouin-zone boundary $G_0 \equiv \pi/\Lambda$, where a wider energy gap occurs for a larger δr , as expected from the large amplitude of $|U_n(\xi)|$ with increasing δr (see Fig. 2).

We now consider the Coulombic interactions between spinless electrons. The interactions make the electron-hole pairs share the ground state of the noninteracting electron system, wherein the most strongly affected states are those lying in the vicinity of E_F . As a consequence, the single-particle density of states $n(\omega)$ near E_F exhibits a power-law singularity of the following form³⁴:

$$n(\omega) \propto |\hbar\omega - E_F|^\alpha, \quad \alpha = \frac{K + K^{-1}}{2} - 1. \quad (4)$$

The explicit form of K is derived using the bosonization procedure³⁴ as follows:

$$K = \lim_{q \rightarrow 0} \sqrt{\frac{2\pi\hbar v_F + g_4(q) - g_2(q)}{2\pi\hbar v_F + g_4(q) + g_2(q)}}. \quad (5)$$

Here, $v_F = \hbar^{-1} dE/dk|_{k=k_F}$ is the Fermi velocity, and $g_4(q) = V(q, m)$ and $g_2(q) = V(q, m) - V(2k_F, m)$ are q -dependent

FIG. 4. (Color online) (a) k dependence of the derivative $d\epsilon/dk$ for the lowest-energy band. (b) δr dependence of the dimensionless Fermi velocity $\tilde{v}_F = d\epsilon/d(ka)|_{k=k_F}$ for different $k_F a$ with increments $\Delta(k_F a)=0.02$.

coupling constants. $V(q, m)$ is the Fourier transform of the screened interaction $V(\mathbf{r}) = -e^2 e^{-\kappa|\mathbf{r}|}/(4\pi\epsilon|\mathbf{r}|)$, where ϵ is the dielectric constant and κ is the screening length. The transformation is performed in terms of the curvilinear coordinates (ξ, θ) , and thus, the resulting $V(q, m)$ becomes a function of both the momentum and angular momentum transfers q and m , respectively. To make concise arguments, E_F was assumed to lie in the lowest-energy band ($n=0$). This allows us to eliminate the index m from $V(q, m)$, which leads to $V(q) = -\frac{e^2}{4\pi\epsilon} \log[(q^2 + \kappa^2)r_0^2]$ for $qr_0 \ll 1$, in which r_0 serves as the short-length-scale cutoff.

The aim of the present study is to examine the δr dependence of K and α , which requires quantification of k_F and v_F . Among the many alternatives, we set $k_F a = 2.0\ell \times 10^{-2}$ ($\ell = 1, \dots, 8$), and evaluate the dimensionless Fermi velocity defined by $\tilde{v}_F \equiv d\epsilon/d(ka)|_{k=k_F}$ for each k_F ; the original v_F is recovered by the relation $v_F = \hbar\tilde{v}_F/(2m^*a)$. We chose the maximum value of k_F such that it does not exceed the δr -dependent zone boundary G_0 for all δr . Figure 4(a) shows the plot of $d\epsilon/dk$ as a function of ka , from which $\tilde{v}_F$ for various k_F and δr are deduced, as shown in Fig. 4(b). Figure 4(b) shows that $\tilde{v}_F$ for all k_F remains constant for $\delta r/a < 2.0$ but decreases remarkably with increasing δr for $\delta r/a > 2.5$. The decrease in $\tilde{v}_F$ is caused by the decrease in the slope of dispersion curves $\epsilon = \epsilon(k)$ with δr (see Fig. 3). The results shown in Fig. 4(b) imply that a change in δr leads to a quantitative alteration in K and α for $\delta r/a > 2.5$, and furthermore, the degree of alteration increases for larger k_F , as will be demonstrated later.

Figure 5 shows the δr dependence of both K and α for different k_F values. According to the bosonization procedure,³⁴ we set the screening parameter κ to be $\kappa a = 1.0 \times 10^{-3}$, which is smaller than all $k_F a$ values that we have chosen. We also set the interaction-energy scale $e^2/(4\pi\epsilon a)$ to be 1.1 in units of $\hbar^2/(2m^*a^2)$ by simulating that of C₆₀-related materials.^{42,43} The insets in Fig. 5 show the k_F dependence of K and α at $\delta r/a = 2.0$; each K and α takes the minimum and maximum values, respectively, at the specific k_F satisfying the relation $(d/dk_F) \log[(2\pi\hbar v_F + g_4)/g_2] = 0$, which is equivalent to $dK/dk_F = 0$. This result is the same for all δr at $\delta r/a < 2.0$.

FIG. 5. (Color online) δr dependences of K and α defined by Eqs. (4) and (5), respectively. The screening parameter κ is set to be $\kappa a = 1.0 \times 10^{-3}$. Insets: nonmonotonic behaviors of K and α as a function of k_F at $\delta r/a = 2.0$.

The salient features of Fig. 5 are the significant decrease in K and increase in α with an increase in δr for $\delta r/a > 2.5$, as predicted earlier. Such δr -driven shifts in K and α are attributed to the effects of geometric curvature on the nature of TLL states. In fact, an increase in δr amplifies the curvature-induced effective potential $U_n(\xi)$, thus yielding a monotonic decrease in $\tilde{v}_F$ at $\delta r/a > 2.5$ (see Fig. 4). The decrease in $\tilde{v}_F$ plays a dominant role in the numerator of the expression in Eq. (5) and eventually leads to the systematic shifts in K and α . We have confirmed that a change in the value of κ does not substantially affect the behaviors of K and α qualitatively, while the absolute values of K and α moderately depends on the choice of κ . It is noted that Eq. (3) can formally deal with an uncurved one-dimensional system subject to a periodically electrostatic potential. In a similar manner to the present curved system, therefore, the α and K are expected to be shifted even for an uncurved quantum wire by the electric-field modulation.

An important consequence of the results shown in Fig. 5 is that nonzero surface curvature yields diverse alterations in the TLL behaviors of deformed cylinders. This is because various kinds of power-law exponents observed in TLL states are related to the quantity K ; some such exponents are the exponent of power-law decay in Friedel oscillation⁴⁴ and that of temperature (or voltage)-dependent conductance.⁴⁵ The present theoretical predictions need to be confirmed experimentally, thus opening a field of science that deals with quantum electron systems on curved surfaces.

In conclusion, we reveal that the power-law exponent α of the TLL states in deformed hollow nanocylinders shows a monotonic increase with an increase in the degree of surface curvature. The increase in α is attributed to the curvature-driven effective potential $U_n(\xi)$ that acts on electrons moving along the curved surface. The present results suggest that there are shifts in the power-law exponents of TLL states of real low-dimensional materials such as the peanut-shaped C₆₀ polymers and MoS₂.

We acknowledge K. Yakubo, Y. Toda, A. Tokuno, and T. Nishii for stimulating discussions. This study was supported

by a Grant-in-Aid for Scientific Research from the MEXT, Japan. H.S. is thankful for the financial support from the Executive Office of Research Strategy in Hokkaido Uni-

versity. Numerical simulations were carried out using the facilities of the Supercomputer Center, ISSP, University of Tokyo.

*Corresponding author; shima@eng.hokudai.ac.jp

- ¹B. De Witt, Phys. Rev. **85**, 653 (1952); Rev. Mod. Phys. **29**, 377 (1957).
- ²H. Jensen and H. Koppe, Ann. Phys. **63**, 586 (1971).
- ³R. C. T. da Costa, Phys. Rev. A **23**, 1982 (1981).
- ⁴L. Kaplan, N. T. Maitra, and E. J. Heller, Phys. Rev. A **56**, 2592 (1997).
- ⁵P. C. Schuster and R. L. Jaffe, Ann. Phys. **307**, 132 (2003).
- ⁶O. G. Schmidt and K. Eberl, Nature (London) **410**, 168 (2001).
- ⁷M. Sano, A. Kamino, J. Okamura, and S. Shinkai, Science **293**, 1299 (2001).
- ⁸S. Tanda, T. Tsuneta, Y. Okajima, K. Inagaki, K. Yamaya, and N. Hatakenaka, Nature (London) **417**, 397 (2002).
- ⁹Y. Oshima, A. Onga, and K. Takayanagi, Phys. Rev. Lett. **91**, 205503 (2003).
- ¹⁰A. Lorke, S. Bohm, and W. Wegscheider, Superlattices Microstruct. **33**, 347 (2003).
- ¹¹D. N. McIlroy, A. Alkhateeb, D. Zhang, D. E. Aston, A. C. Marcy, and M. G. Norton, J. Phys.: Condens. Matter **16**, R415 (2004), and references therein.
- ¹²H. Shima and M. Sato, Nanotechnology **19**, 495705 (2008).
- ¹³T. Fujita, H. Okada, K. Koyama, K. Watanabe, S. Maekawa, and M. W. Chen, Appl. Phys. Lett. **92**, 251902 (2008).
- ¹⁴I. Arias and M. Arroyo, Phys. Rev. Lett. **100**, 085503 (2008).
- ¹⁵A. Mostafazadeh, Phys. Rev. A **54**, 1165 (1996).
- ¹⁶G. Cantele, D. Ninno, and G. Iadonisi, Phys. Rev. B **61**, 13730 (2000).
- ¹⁷H. Aoki, M. Koshino, D. Takeda, H. Morise, and K. Kuroki, Phys. Rev. B **65**, 035102 (2001); M. Koshino and H. Aoki, *ibid.* **71**, 073405 (2005).
- ¹⁸M. V. Entin and L. I. Magarill, Phys. Rev. B **66**, 205308 (2002).
- ¹⁹N. Fujita, J. Phys. Soc. Jpn. **73**, 3115 (2004).
- ²⁰J. Gravesen and M. Willatzen, Phys. Rev. A **72**, 032108 (2005).
- ²¹H. Taira and H. Shima, Surf. Sci. **601**, 5270 (2007).
- ²²V. Atanasov and R. Dandoloff, Phys. Lett. A **372**, 6141 (2008).
- ²³S. K. Baek, S. D. Yi, and B. J. Kim, Phys. Rev. E **77**, 022104 (2008).
- ²⁴D. V. Bulaev, V. A. Geyler, and V. A. Margulis, Phys. Rev. B **69**, 195313 (2004).
- ²⁵A. V. Chaplik and R. H. Blick, New J. Phys. **6**, 33 (2004).
- ²⁶A. Marchi, S. Reggiani, M. Rudan, and A. Bertoni, Phys. Rev. B **72**, 035403 (2005).
- ²⁷G. Cuoghi, G. Ferrari, and A. Bertoni, Phys. Rev. B **79**, 073410 (2009).
- ²⁸M. V. Entin and L. I. Magarill, Phys. Rev. B **64**, 085330 (2001); Europhys. Lett. **68**, 853 (2004).
- ²⁹E. Zhang, S. Zhang, and Q. Wang, Phys. Rev. B **75**, 085308 (2007).
- ³⁰M. Encinosa, Phys. Rev. A **73**, 012102 (2006).
- ³¹O. Olendski and L. Mikhailovska, Phys. Rev. B **72**, 235314 (2005); **77**, 174405 (2008).
- ³²G. Ferrari and G. Cuoghi, Phys. Rev. Lett. **100**, 230403 (2008); G. Ferrari, A. Bertoni, G. Goldoni, and E. Molinari, Phys. Rev. B **78**, 115326 (2008).
- ³³G. Parascandolo, G. Cantele, D. Ninno, and G. Iadonisi, Phys. Rev. B **68**, 245318 (2003).
- ³⁴J. Voit, Rep. Prog. Phys. **57**, 977 (1994).
- ³⁵J. Onoe, T. Nakayama, M. Aono, and T. Hara, Appl. Phys. Lett. **82**, 595 (2003); J. Onoe, T. Ito, S. I. Kimura, K. Ohno, Y. Noguchi, and S. Ueda, Phys. Rev. B **75**, 233410 (2007).
- ³⁶Y. Toda, S. Ryuzaki, and J. Onoe, Appl. Phys. Lett. **92**, 094102 (2008).
- ³⁷P. Santiago, J. A. Ascencio, D. Mendoza, M. Pérez-Alvarez, A. Espinosa, C. Reza-sangermán, P. Schabes-Retchkiman, G. A. Camacho-bragado, and M. José-Yacamán, Appl. Phys. A **78**, 513 (2004).
- ³⁸H. Yoshioka, Physica E **18**, 212 (2003).
- ³⁹H. Ishii, H. Kataura, H. Shiozawa, H. Yoshioka, H. Otsubo, Y. Takayama, T. Miyahara, S. Suzuki, Y. Achiba, M. Nakatake, T. Narimura, M. Higashiguchi, K. Shimada, H. Namatame, and M. Taniguchi, Nature (London) **426**, 540 (2003).
- ⁴⁰H. Shima and T. Nakayama, *Higher Mathematics for Physics and Engineering* (Springer-Verlag, Berlin, 2009).
- ⁴¹T. A. Beu, J. Onoe, and A. Hida, Phys. Rev. B **72**, 155416 (2005).
- ⁴²A. F. Hebard, R. C. Haddon, R. M. Fleming, and A. R. Kortan, Appl. Phys. Lett. **59**, 2109 (1991).
- ⁴³A. Oshiyama, S. Saito, N. Hamada, and Y. Miyamoto, J. Phys. Chem. Solids **53**, 1457 (1992).
- ⁴⁴R. Egger and H. Grabert, Phys. Rev. Lett. **75**, 3505 (1995).
- ⁴⁵C. L. Kane and M. P. A. Fisher, Phys. Rev. B **46**, 15233 (1992).