



Title	Development of an adaptive marine ecosystem management and co-management plan at the Shiretoko World Natural Heritage Site
Author(s)	Matsuda, Hiroyuki; Makino, Mitsutaku; Sakurai, Yasunori
Citation	Biological Conservation, 142(9), 1937-1942 https://doi.org/10.1016/j.biocon.2009.03.017
Issue Date	2009-09
Doc URL	http://hdl.handle.net/2115/39325
Type	article (author version)
File Information	BC142-9_p1937-1942.pdf



[Instructions for use](#)

1 **Development of an adaptive marine ecosystem management and**
2 **co-management plan at the Shiretoko World Natural Heritage Site**

3

4 Hiroyuki MATSUDA ^{1*} (Corresponding author)

5 Mitsutaku MAKINO ²

6 Yasunori SAKURAI ³

7

8 ¹ Faculty of Environment and Information Sciences, Yokohama National University, 79-7

9 Tokiwadai, Hodogaya-ku Yokohama 240-8501, Japan, matsuda@ynu.ac.jp

10 Tel:+81-45-339-4362; Fax:+81-45-339-4373

11

12 ² National Research Institute of Fisheries Science, Fisheries Research Agency, 2-12-4, Fukuura, Kanazawa-ku,

13 Yokohama 236-8648, Japan

14 E-mail: mmakino@affrc.go.jp

15

16 ³ Graduate School of Fisheries Sciences, Hokkaido University, Mitano-cho, Hakodate, Hokkaido

17 041-8611, Japan

18 E-mail: sakurai@fish.hokudai.ac.jp

19

20

21 Word count = 4610 (including cover page)

22

23 **Development of an adaptive marine ecosystem management and**
24 **co-management plan at the Shiretoko World Natural Heritage Site**

25

26 The Marine Management Plan for the Shiretoko World Natural Heritage Site, Japan, provides a
27 case study for adaptive marine ecosystem management and co-management of coastal fisheries.
28 Shiretoko was the third World Natural Heritage Site registered in Japan and earned this title because
29 of its (i) formation of seasonal sea ice at some of the lowest latitudes in the world, (ii) high
30 biodiversity, and (iii) many globally threatened species. The natural resource management plan of
31 the Shiretoko site is characterized by transparency and consensus building, because (i) UNESCO
32 and IUCN require that the plan be sustainable; and (ii) the Government of Japan has guaranteed
33 local fisheries that there will be no additional regulations included in the plan. The Marine
34 Management Plan describes which species and factors are monitored, how these data are evaluated,
35 and how the benchmarks specified by ecosystem management are determined. The Plan will
36 provide a valuable example for the establishment of “environment-friendly fisheries” in Japan and
37 other countries, because it includes voluntary activities by resource users that are suitable for use in
38 a local context, flexible to ecological/social fluctuations, and efficiently implemented through
39 increased legitimacy and compliance. This approach is appropriate for coastal communities where a
40 large number of small-scale fishers catch a variety of species using various types of gear. We
41 develop a method to evaluate fisheries integrity by catch and yield data.

42 Keywords: adaptive management, co-management, scientific council, Steller sea lions, walleye
43 pollock.

44

45 **Introduction**

46 Adaptive management predicts and monitors changes in the ecosystem and subsequently
47 reviews and adjusts the management and use of natural resources (Walters 1986). Such predictions
48 and monitoring are best accompanied by feedback controls, such as the verification of hypotheses
49 based on the results of monitoring in order to review and modify management activities.

50 Marine management in Japan is characterized by a long history of coastal co-management
51 of fishers' organizations (Makino et al., 2008). Co-management is defined as the sharing of
52 responsibilities between governmental institutions and groups of resource users (Persoon et al.,
53 2005). In many countries environmental management is reformulated from exclusive state control
54 to various kinds of joint management in which local communities, indigenous peoples, and
55 nongovernmental organizations share authority and benefits with governmental institutions.
56 Fisheries in Japan face several important challenges, e.g., (i) exclusive use by fisherman with
57 fishery rights/licenses (there are few exceptions for free-fisheries and recreational angling), (ii) lack
58 of full transparency in management procedures, and (iii) lack of objective benchmarks or numerical
59 goals in management plans. Here, we elaborate on the characteristics and issues facing the
60 management of marine ecosystems in Japan. We use the "Multiple-Use Integrated Marine
61 Management Plan for the Shiretoko World Natural Heritage Site, Japan" (Ministry of the
62 Environment, the Government of Japan, and Hokkaido Prefectural Government, 2007), as a case
63 study of adaptive marine ecosystem management and co-management of coastal fisheries.

64 The natural resource management plan of the Shiretoko site is characterized by
65 transparency and consensus building, because (i) UNESCO and IUCN require that the plan be
66 sustainable; and (ii) the Government of Japan has guaranteed local fisheries that there will be no
67 additional regulations included in the plan. These "constraints" seemed to be incompatible with
68 each other. In this paper, we show the solution of these constraints. The solution will characterize
69 Japanese fisheries. In addition, we will present a method to analyze the sustainability from fisheries
70 catch and yield data, which is to be available in the Shiretoko site. Autonomous management by

71 fishers is often based on fisheries data. We use catch and yield data in Shiretoko site (Fig. 1) to
72 evaluate stock and economic status of each resource.

73

74 **Role and function of the Scientific Council for the Shiretoko World Natural Heritage Site**

75 *Marine ecosystems with coastal fisheries*

76 The objective of the Marine Management Plan for the Shiretoko site is to ensure a balance
77 between the conservation of the marine ecosystem and stable fisheries through the sustainable use
78 of fisheries resources in the marine component of the heritage area. According to the IUCN's
79 request, the marine component extends up to 3 km from the coastline, which needed the revision of
80 Natural Park Law in Japan.

81 The management plan defines measures to conserve the marine ecosystem, strategies to
82 maintain major fisheries resources, monitoring methods for those resources, and policies for marine
83 recreation. The Oyashio shelf region and the seasonally ice-covered areas north of Hokkaido are
84 highly productive and support a wide range of species, including marine mammals, seabirds, and
85 commercially important species in the western subarctic Pacific (Sakurai, 2007). The management
86 plan identifies many indicator species, such as salmonids (e.g., chum, pink, and masu salmon,
87 *Oncorhynchus keta*, *O. gorbuscha*, and *O. masou masou*, respectively), walleye pollock (*Theragra*
88 *chalcogramma*), Pacific cod (*Gadus macrocephalus*), and Steller sea lions (*Eumetopias jubatus*),
89 and also outlines the monitoring and conservation of these important species. They are selected
90 from keystone species, predators at higher trophic levels that probably have a great impact on
91 ecosystems, and endangered species in the waters surrounding Shiretoko.

92 In addition, the management plan details the vast food web structure of the Shiretoko site
93 (Makino et al. 2009) and includes catch statistics for ten categories of major fisheries resources (Fig.
94 1). Adaptive management plans usually determine criteria and feedback control measures for
95 indicator species. For example, management plans monitor and enforce conservation actions to
96 satisfy numerical goals within a limited amount of time. Management plans usually devise action
97 plans to achieve these numerical goals or to maintain thresholds for indicator species. However, the

98 present Marine Management Plan for the Shiretoko site (Ministry of the Environment, the
99 Government of Japan, and Hokkaido Prefectural Government, 2007) does not include any
100 thresholds or numerical goals for its indicator species, which are currently only monitored. A
101 crucial short-term goal will be to establish such thresholds and/or numerical goals for these
102 indicator species.

103 The optimal fisheries policy of maximizing sustainable yield from the entire ecosystem
104 does not guarantee the coexistence of all species (Matsuda and Abrams, 2006). Therefore, the goal
105 of the management plan is twofold: sustainable fisheries and biodiversity conservation. Here, we
106 focus on our monitoring efforts of sea lions and walleye pollock as well as the catch statistics for ten
107 categories of fisheries resources (Fig. 1). We consider two periods 1985-89 and 1998-2002, denoted
108 by period I and II, respectively. To evaluate fisheries integrity of the Shiretoko site, we calculate the
109 average yield and catch of resource i during period i (denoted by C_i^i and Y_i^i) and calculate the
110 average price $P_i^i = Y_i^i / C_i^i$. We also calculate the coefficient of variation of catch and yield of each
111 resource throughout 1985-2002. We compare C_i^i , Y_i^i and P_i^i between the two periods. If the fish
112 price per unit weight is positively correlated with the fish body size, the fish price is a useful
113 indicator of fisheries integrity. We also calculate the mean trophic level (MTL, Pauly et al. 1998).
114 The mean trophic level of each fish is given by FISHBASE (<http://www.fishbase.org/>). Trophic
115 level of kelp, common squids, scallop and octopus are set to be 1, 3.6, 2 and 4, respectively.

116 In addition to fisheries resources, we need to conserve species that are not utilized by
117 fisheries. Sea lions are threatened species and important from conservation viewpoints. Walleye
118 pollock is a target species of fisheries management and is also a prey of sea lions. These species are
119 controllable by several conservation measures. A flow diagram of the adaptive management
120 procedure is presented in Fig. 2.

121 If some categories of fisheries resources decrease in stock and/or catch, we recommend
122 target switching from the decreased resource to another resource that is temporally abundant
123 (Katsukawa and Matsuda 2003). If all major fisheries resources decrease, the extension or
124 improvement of marine protected areas (MPAs) is effective for fisheries resource management. In

125 Shiretoko site, fishers autonomously introduced seasonally closed fishing areas to protect spawning
126 fish of walleye pollock in 1995, as well as they decreased the number of fishing boats that fish
127 walleye pollock from 177 in 1997 to 86 in 2004. There are many autonomous closed fishing areas in
128 Japan because the area of closed fishing zone is often flexible from year to year, depending on the
129 stock status (Tomiyama et al. 2005). In this paper, we use the term MPA defined as an area where
130 fisheries are closed or regulated by either government or fishers.

131 If the stock of walleye pollock decreases, the extension of MPAs, the reduction of fishing
132 capacity, and/or revision of the total allowable catch for this species should be implemented. If the
133 population size of the Steller sea lions decreases, the cull limit, as determined by the potential
134 biological removal (defined below), should be revised, or explosives should be used to scare sea
135 lions away.

136 We will need to establish clear benchmarks for the catch statistics of major fisheries
137 resources and numerical goals for stock recovery of walleye pollock and Steller sea lions. Such
138 benchmarks should be used to monitor the feasibility of the management plan, and if the plan
139 includes any unfeasible or unrealistic goals, it should be revised (Fig. 2).

140

141 **Trends in catch and price of fisheries resources**

142 Fig. 1 shows catch amount of the top 10 largest average catch and the mean trophic level
143 (MTL) of Shiretoko fisheries during 1985 to 2002. Table 1 shows some fisheries characteristics of
144 the top 10 taxa of long-term yield and those of long-term catch amount. The coefficient of variation
145 (C.V.) of these resources were relatively stable, ranged from 100% (common squid) to 27% (kelp).

146 The mean trophic level (MTL) of catch in Shiretoko site was stable throughout 1985-2002.
147 The MTL has slightly increased since 1992 mainly because of increasing catch of salmons. The
148 integrity of marine ecosystem is not characterized by MTL of catch in Shiretoko site.

149 Among 54 taxa, 26 and 38 taxa changed their catch amount more than 100 and 10 folds,
150 respectively. Among resources in Table 1, catch of walleye pollock and rock fish by 86% and 61%
151 from period I to II, while Price of rock fish increased by 83%. Catch of sardine, anchovy, red king

152 crab (*Paralithodes camtschaticus*), *Sebastes* and herring (*Clupea pallasii*) substantially decreased
153 by 100%, 99.4%, 99%, 97% and 96%, respectively. Sardine (Watanabe et al. 1995), anchovy and
154 herring decreased probably because of natural fluctuation in Japanese waters, while we do not
155 know the reason why red crab king and *Sebastes* decreased. Greenling (*Hexagrammos otakii*)
156 decreased their catch by 70% and the fish price by 64% from period I to II.

157 There are two major resources in Shiretoko site, chum salmon and walleye pollock. Those
158 were two of largest-yield resource throughout 1985-2002, except in 1996, 97 and 2001 while the 2nd
159 largest yield resource was common squid in 1996, 97 and kelp in 2001 . The largest yield resource
160 was walleye pollock during 1985-92 and chum salmon during 1993-2002. Common squid, kelp,
161 rock fish, *Sebastes* and Pacific cod have at least once been the 3rd largest yield throughout
162 1985-2002. Sum of the top three largest yield resource ranged from 81% in 1985 to 53% in 1996 of
163 the total annual yield. Since walleye pollock decreased in 1991, Shiretoko fisheries now depend on
164 salmon fisheries, which is supported by **release of hatching stock**. The yield of common squid, sea
165 cucumber, octopus, pink salmon and Pacific cod increased from period I to II by 38-fold, 367%,
166 64%, 48% and 37%, respectively.

167

168 **Role of co-management in coastal fisheries**

169 Unlike fisheries in modern countries, there is no centralized top-down management in
170 traditional fisheries. When Japan was modernized during the second half of the 19th century, the
171 country attempted to centralize the fisheries institution. However, these attempts resulted in a great
172 deal of confusion and disturbance within fisheries societies; thus, Japan still has a decentralized
173 co-management system involving fishers and the government. The transaction costs for fisheries
174 management constitute one of the strongest arguments against top-down management systems. In
175 the co-managed system, the costs for monitoring, enforcement, and compliance are shared between
176 the government and local fishers and are remarkably lower than in systems with top-down
177 regulation (Makino and Matsuda, 2005).

178 FCAs collected data on the amount of catch, catch area, and body size of catch, and they

179 provided these data to prefectural research stations. Consequently, the local fishers voluntarily
180 enlarged the mesh size of pollock gillnets from 91 to 95 mm in the 1990s. In 1995, the local fishers
181 divided their fishing grounds into 34 areas based on local knowledge and then introduced a
182 temporal fishing ban in seven of these areas. FCAs could pay these efforts to maintain their
183 sustainable fisheries. After the stock of walleye pollock decreased, the fishers voluntarily added six
184 more areas to the fishing ban in 2005. This is probably because the FCAs expected benefit of the
185 World Heritage. These fishers annually re-examine the protected areas based on the scientific
186 advice of the local research station. Clearly, local fishers often contribute considerable efforts
187 toward consensus building in regard to voluntary regulations. Because the Nemuro stock of
188 walleye pollock is also utilized by Russian fisheries, Rausu FCA needs to cooperate with Russian
189 fishers and scientists.

190 Although the voluntary management procedure of the Rausu FCA is not well defined, they
191 regulate the impact of fisheries in terms of recent stock conditions. The marine management plan
192 for the Shiretoko site (Ministry of the Environment, the Government of Japan, and Hokkaido
193 Prefectural Government, 2007) recognizes this feedback control as adaptive management. However,
194 adaptive management must determine how to change its policy prior to the implementation of
195 management.

196 Even in coastal fisheries, resources are not used by a single fisheries organization. Japanese
197 coastal and offshore fishers and Russian trawl fishers exploit the walleye pollock in the Nemuro
198 stock. Therefore, international cooperation is indispensable for the effective resource management
199 of this species. As we mention below, the expansion of the Shiretoko World Heritage Site may be a
200 good opportunity for the international management of walleye pollock.

201

202 *Cull limit of Steller sea lions based on potential biological removal*

203 The Okhotsk and Kurile populations of Steller sea lions migrate from their breeding and
204 landing grounds in Russian waters to the waters surrounding Shiretoko for over-wintering and
205 foraging. Because the Asian population of Steller sea lions sharply declined until the 1980s, this

206 species is classified as endangered on the IUCN Red List. Fortunately, it has been gradually
207 increasing at a rate of 1.2% per year since the early 1990s (Burkanov and Loughlin, 2005). The
208 entire population size, which is spread throughout the Sea of Okhotsk, the western part of the
209 Bering Sea, and the Komandorskie Islands, was estimated at 15,676 in 2005 by enumerating the
210 reproductive colonies. In 2007, the JME ranked the sea lion as vulnerable (the third rank of
211 threatened species).

212 Despite the threatened status of this species, sea lions are still culled by Japanese fishers
213 because of the extensive damage caused to fishing nets, the extent of which increased during the
214 1980s (Fig. 4). When an international movement for the conservation of marine mammals began in
215 the 1990s, the Hokkaido Fishing Coordination Commission established a cull limit of 116 sea lions
216 per year. In 2007, the Fisheries Agency of Japan (JFA) recommended a revised cull limit based on
217 the potential biological removal (PBR, Wade, 1998). The PBR is 120 determined by the number of
218 migrant sea lions to Japanese waters and life history parameters used for the eastern Aleutian
219 population (Angliss and Outlaw, 2007). An additional problem facing Steller sea lions is by-catch.
220 For example, sea lions are often by-catch in bottom-set-net, gillnet, and set-net fisheries of
221 Hokkaido Island. Unfortunately, there have been no reports of sea lion by-catch by these fisheries.
222 According to informal interviews, the total number of by-catch was estimated between 55 and 107,
223 the highest of which (107) was assumed for management purposes. The cull limit was revised based
224 on the inferred number of by-catch from the PBR; thus, the revised cull limit is 120 (227 less 107)
225 sea lions, which, until 2006, was nearly the same as the original cull limit. If the true number of
226 by-catch is reported, the cull limit could be revised. Because we used the highest estimate for
227 by-catch, the cull limit would likely increase, thus creating an incentive to compile a by-catch
228 report.

229

230 *Importance of cooperation with Russia in fisheries management*

231 Sheppard (2005) documented the clear and apparent similarities between the environment
232 and ecology of the Shiretoko Peninsula and the Kunashiri and Itrup Islands (Fig. 5). He also

233 addressed the possibility for the future development of these regions as a more broad-scale
234 “World Heritage Peace Park.” Coastal fishers in the Shiretoko area are also concerned about the
235 effects of Russian fisheries on the Nemuro stock of walleye pollock. Japan and Russia have been in
236 conflict over the national boundary between the two countries since World War II. Russia (formerly
237 the Soviet Union) actually occupied Habomai Shikotan, Kunashiri, and Itrup Islands, whereas
238 Japan argued for inherent sovereignty of these islands. Despite these disagreements, UNESCO can
239 register a world heritage site that is multi-national and includes a boundary under international
240 dispute in accordance with the Convention on World Heritage. After the registration of the
241 Shiretoko World Heritage Site, Japanese Prime Minister Shinzo Abe and Russian President
242 Vladimir Putin agreed to organize scientific meetings for a cooperative effort in ecosystem
243 conservation within the Sea of Okhotsk. Furthermore, if Japan and Russia agree to expand the
244 Shiretoko World Heritage Site to southern Kurile and Urup Islands, the site will become an
245 “international peace park” recommended by Sheppard (2005) based on mutual understanding and
246 peace. In the first step, we share ecological data and knowledge between Japan and Russia.

247

248 **Discussion**

249 Scientists played a very important role during the registration process of the Shiretoko
250 World Heritage Site by interpreting the evaluation and criticism of the IUCN to Japanese society.
251 We propose a general procedure for environmental risk management (Rossberg et al., 2006), the
252 key point of which is to devise a scientific procedure using consensus building among stakeholders.
253 The purpose of management depends in part on all involved stakeholders (excluding the scientists).
254 After a consensus concerning the objectives of management is reached, scientists can propose an
255 action plan and numerical targets to achieve these goals.

256 The marine management plan at the Shiretoko site (Ministry of the Environment, the
257 Government of Japan, and Hokkaido Prefectural Government, 2007) was accepted in 2007 by the
258 Shiretoko World Natural Heritage Site Scientific Council and the Shiretoko World Natural Heritage
259 Site Regional Liaison Committee. An English version was translated and published before the

260 UNESCO and IUCN inspection in February 2008. Coastal fisheries persist and exploit many
261 species in the food web (Fig. 1) at the Shieretoko site and may function in this region like a top
262 predator or an umbrella species. The catch statistics of fisheries (e.g., Fig. 2) can be used to
263 determine the current status of marine food webs. However, it is too costly and virtually impossible
264 for the government to monitor all details within an ecosystem. Therefore, the knowledge of fishers
265 and data from fisheries activities should be fully utilized.

266 Matsuda et al. (2008) found that an economic rent-maximizing policy may lead to the
267 convergence of target species and gear types and could provide information about very limited
268 aspects of the ecosystem. These authors also used simple mathematical models of a multi-trophic
269 level ecosystem to demonstrate that economically efficient fisheries would result in the loss of a
270 significant fraction of the species in the ecosystem (Matsuda et al., 2008). In other words,
271 economically efficient fisheries make sense for high fishery rents but not for sustaining total
272 ecosystem services for society as a whole. One reasonable alternative to single species fisheries
273 management is to conduct responsible fisheries that target a wide range of species using a variety of
274 gear (Katsukawa and Matsuda 2003). As the Shiretoko case demonstrates, responsible and diverse
275 fisheries operations can significantly contribute to the sustainability of ecosystem monitoring
276 (Makino et al., 2008).

277 JFA surveys the stock status of walleye pollock and sea lions. In addition, Government of
278 Japan and Hokkaido Prefectural Government (HPG) pay some additional effort for the world
279 heritage, e.g., HPG supports scientists to survey upstream run of chum and pink salmon in two
280 rivers and scientists evaluate the nutrient contribution from salmon to the terrestrial ecosystems.
281 Taxon-specific catch and yield data are useful to evaluate the status of coastal fisheries and marine
282 ecosystem. If both catch and price of some resource decrease, fishers pay attention to the status of
283 these resources, such as greenling and sailfin sandfish. The Scientific Council recommended more
284 conservation than persistence of the major salmon population, and a sufficient contribution of
285 salmon from the sea.

286 Throughout the establishment of the marine management plan, fishers, scientists and

287 environmental groups built trust and understood their sense of values. Scientists analyze and
288 publish the status of coastal fisheries. This paper clarified some resources, walleye pollock, red
289 king crab, rock fish (*Sebastes*) and greenling, under bad stock or economic conditions. In
290 co-management, fishers will decide a conservation plan. Scientists can advise the fishers if
291 scientists analyze both ecology and economy in the fisheries resources in the case of sand lance
292 fisheries in Aichi Prefecture, Japan (Tomiyama et al. 2005).

293 Biodiversity may support the robustness of ecosystem processes against climate change and
294 disasters. We typically investigate the value of ecosystem services under normal conditions;
295 however, the value of biodiversity may be effective against unusual disasters. Japan has a rich
296 biodiversity in comparison to other developed countries, potentially because it has geographic
297 characteristics that have been resilient to past climate change. In future studies, we will further
298 examine the value of biodiversity in Japan.

299

300 **Acknowledgments**

301 We thank Drs. H. Homma, Y. Kagami, M. Kobayashi, Y. Maruyama, Y. Ohtaishi, S. Oyamada, M.
302 Sano, anonymous reviewers, Professros R. Primack and H. Higuchi for their valuable comments on
303 the earlier version of the manuscript. We also thank the members, secretary, and observers of
304 Marine Working Group of the Shiretoko World Heritage Site. This work was supported in part by
305 the Pew Fellowship grant and the Global COE Program “Eco-Risk Asia” at Yokohama National
306 University to H.M.

307

308 **References**

309 Angliss, R.P., Outlaw, R.B., 2007. Alaska Marine Mammal Stock Assessments, 2006. NOAA Tech.
310 Memo. NMFS-AFSC-168, pp.1-16.

311 Burkanov, N.V., Loughlin, T.R., 2005. Distribution and abundance of Steller sea lions, *Eumetopias*
312 *jubatus*, on the Asian Coast, 1720's-2005. Mar. Fish. Rev. 67, 1-62.

313 Japan Fisheries Agency 2007. Stock assessment for Nemuro stock of walleye pollock, Stock
314 Assessment Report in Fiscal year 2006, Japan Fisheries Agency, Yokohama. (in Japanese)

- 315 <http://www.jfa.maff.go.jp/release/19/081001.pdf>
- 316 Katsukawa, T., Matsuda, H., 2003. Simulated effects of target switching on yield and sustainability of
317 fish stocks. *Fish. Res.* 60, 515-525.
- 318 Makino, M., Matsuda, H., 2005. Co-management in Japanese Coastal Fishery: Its institutional
319 features and transaction cost. *Mar. Policy* 29, 441-450.
- 320 Makino, M., Matsuda, H., Sakurai, Y., 2008. Expanding Fisheries Co-management to
321 Ecosystem-based management: A case in the Shiretoko World Natural Heritage area, Japan.
322 *Mar. Policy* 33, 207-214
- 323 Matsuda, H., Abrams, P.A., 2006. Maximal yields from multi-species fisheries systems: rules for
324 systems with multiple trophic levels. *Ecol. Appl.* 16, 225-237.
- 325 Matsuda, H., Makino, M., Kotani, K., 2008. Optimal fishing policies that maximize sustainable
326 ecosystem services. (Tsukamoto K, Kawamura T eds) *The Memorial Book for the fifth World*
327 *Fisheries Congress*, 359-369.
- 328 Ministry of the Environment, the Government of Japan, and Hokkaido Prefectural Government
329 (2007) *The Multiple Use Integrated Marine Management Plan and Explanatory Material for*
330 *Shiretoko World Natural Heritage Site*, December 2007.
331 http://shiretoko.env.gr.jp/uploads/kaigi/iucn/seawg/seawg_iucn_english.pdf
- 332 Ohtaishi, N., Wada, K., eds. 1999. *Migrative ecology and conservation of Steller sea lions*, Tokai
333 University Press (in Japanese).
- 334 Pauly, D., V. Christensen, D. Johannes, F. Rainer and T. Francisco Jr. 1998. Fishing Down Marine
335 Food Webs. *Science* 279: 860-863.
- 336 Persoon GA, van Est DME, Sajise PE, eds. 2005. *Co-Management of Natural Resources in Asia: A*
337 *Comparative Perspective*. Copenhagen, NIAS Press. 320pp.
- 338 Rossberg, A.G., Matsuda, H., Koike, F., Amemiya, T., Makino, M., Morino, M., Kubo, T., Shimoide,
339 S., Nakai, S., Katoh, M., Shigeoka, T., Urano, K., 2005. A guideline for ecological risk
340 management procedures. *Landscape Ecol. Eng.* 1, 221-228.
- 341 Sakurai Y., 2007. An overview of the Oyashio ecosystem. *Deep-Sea Res. II* 54, 2526-2542.

- 342 Sheppard, D., 2005. World heritage nomination –IUCN technical evaluation, Shiretoko, Japan, ID
343 No: 1193. IUCN (The World Conservation Union).
- 344 Tomiyama, T., Lesage, C-M., Komatsu, T., 2005. Practice of Sandeel Fisheries Management in Ise
345 Bay toward Responsible and Sustainable Fisheries. *Gl. Env. Res.* 9, 139-150.
- 346 Wade, P.R., 1998. Calculating limits to the allowable human-caused mortality of cetaceans and
347 pinnipeds. *Mar. Mammal Sci.* 14, 1-37.
- 348 Walters, C.J., 1986. Adaptive management of renewable resources. McMillan, NY, USA.
- 349 Watanabe, Y., Zenitani, H. and Kimura R., 1995. Population decline of the Japanese sardine
350 *Sardinops melanostictus* owing to recruitment failures. *Can J Fish Aq Sci* 52, 1609-1616.
- 351
- 352
- 353

354 No tables in this manuscript

355

356 Figure Legends

357

358

359 Fig. 1. Catch statistics for ten major exploited taxa and total catch in the Shiretoko-daiichi, Utoro,
360 and Rausu Fisheries Cooperative Associations (Ministry of the Environment, the Government of
361 Japan, and Hokkaido Prefectural Government, 2007). Resource number is the order in the
362 long-term average catch in Shiretoko site, as shown in Table 1 (12th resource is sardine).

363

364 Fig. 2. Proposed flow diagram for the Marine Management Plan of the Shiretoko World Heritage
365 Site.

366

367 Fig. 3. Catch statistics for Steller sea lions and damage to fisheries caused by sea lions in Hokkaido,
368 Japan (Ohtaishi and Wada, 1999; Japan Fisheries Agency, 2007). Numbers of by-catch are not
369 included or are not known.

370

371 Fig. 4. Present protected areas in the Shiretoko Peninsula of Hokkaido Island and Habomai,
372 Shikotan, Kunashiri, Itrup, and Urup Islands, originally drawn by Mari Kobayashi. A national
373 boundary between Japan and Russia exists between Hokkaido and Kunashiri and Habomai (as
374 claimed by Russia) and between the Itrup and Urup Islands (as claimed by Japan).

375

376 Table 1. Annual catch and yield of major resources in Shiretoko site (Rausu, Shari and Utoro Fisheries Cooperative Associations) based on the long-term
 377 annual (Ministry of the Environment, the Government of Japan, and Hokkaido Prefectural Government, 2007). Orders and coefficient of variations in catch and
 378 yield are based on the long-term average throughout 1985-2002.

common name	academic name	Order in		Catch amount (ton)			Yield (thosand yen)			price (yen/kg)	
		catch	yield	1985-1989	1998-2002	C.V.	1985-1989	1998-2002	C.V.	1985-2002	1998-2002
chum salmon	<i>Oncorhynchus keta</i>	2	1	12,509	33,459	47%	6,660	8,466	23%	532	253
walleye pollock	<i>Theragra chalcogramma</i>	1	2	88,580	12,433	84%	11,063	1,846	78%	125	148
kelp	<i>Laminaria japonica</i>	9	3	648	586	27%	1,274	1,486	26%	1,967	2,534
common squid	<i>Todarodes pacificus</i>	3	4	225	13,182	100%	38	1,518	93%	170	115
rock fish	<i>Sebastes sp</i>	8	5	935	367	53%	1,318	947	35%	1,410	2,580
Pacific cod	<i>Gadus macrocephalus</i>	6	6	4,637	3,443	41%	662	905	38%	143	263
Okhostk atka mackerel	<i>Pleurogrammus azonus</i>	4	7	6,299	6,187	34%	810	805	23%	129	130
pink salmon	<i>Oncorhynchus gorbuscha</i>	5	8	1,060	5,362	80%	429	637	50%	405	119
sea urchin	<i>Strongylocentrotus intermedius</i>	26	9	48	35	34%	508	372	29%	10,591	10,711
scallop	<i>Patinopecten yessoensis</i>	7	10	1,767	1,665	53%	302	140	45%	171	84
octopus	<i>Octopus dofleini</i>	10	11	313	471	34%	120	198	25%	384	420

Fig. 1




Fig. 2




Fig. 3




Fig. 4

