



<b>Title</b>	Arctic sea ice reduction under global change : in association with deterioration of Greenland and Antarctic ice sheets
<b>Author(s)</b>	Ikeda, Motoyoshi
<b>Citation</b>	Intenational Symposium "Toward a Sustainable Low Carbon Society -Green New Deal and Global Change-". 4-5 November 2009. Sapporo, Japan.
<b>Issue Date</b>	2009-11-05
<b>Doc URL</b>	<a href="http://hdl.handle.net/2115/39961">http://hdl.handle.net/2115/39961</a>
<b>Type</b>	conference presentation
<b>Note</b>	Intenational Symposium "Toward a Sustainable Low Carbon Society -Green New Deal and Global Change-" PartII Global Change. 5 November 2009. Sapporo, Japan.
<b>File Information</b>	ikeda1105.pdf




[Instructions for use](#)

**Arctic sea ice reduction  
under global change:**  
in association with deterioration of  
Greenland and Antarctic ice sheets

Motoyoshi Ikeda

Graduate School of Environmental Science,  
Hokkaido University,  
Sapporo, Japan

# Sea-level rise addition due to ice sheet melting



Thermal expansion is main, while ice sheet melting is also important.

Deep ocean (Antonov et al., GRL, 2005; Kohl et al., JPO, 2007).


Ice sheets (Lemke et al., IPCC, 2007)

Glacier and small-ice caps (Dyurgerov and Meier, 2004).

Terrestrial storage (Ngo-Duc et al., GRL, 2005).

# IPCC projection as the lower limit

Ice sheet melting is additional.



# Distinct Indicator

## Arctic sea ice

Indigenous people



Polar bear



# How soon will the Arctic sea ice change to a seasonal one?

- IPCC Report prediction:  
the ice cover in summer will disappear near the end of the 21st century.
- The anomalously small ice cover in 2007 summer  
>>some specialists believe disappearance by 2020
- Significant year-to-year variability  
>>such an early disappearance is exaggeration!?

# Ice cover trend and decadal variability

Wang and Ikeda (GRL, 2001) +  $\alpha$

- Winter (DJF) and summer (JJA)
- Beaufort and Chukchi
- East Siberian and Laptev
- Barents and Kara





# Significant trend and decadal variability

- Sea ice has been reducing in last several decades.
- Decadal variability are clear and related with NAM/AO.
- We should be cautious about change in a few years.






# EOF-1 of winter Atmos. Pressure at sea level Northern Annular Mode/Arctic Oscillation



Decadal variability  
in sea ice cover produced  
by NAM/AO  
(Ikeda, 1990)

Sea ice cover  
anomaly propagates  
in three regions




# Arctic Dipole Mode (ADM)

- Sea ice has been reducing more rapidly in the Pacific sector after 1990.
- NAM/AO is not so significant after 1990.
- EOF-2 has been proposed to be important by Wu et al. (2006), Wang et al. (2009), etc.

# 2nd EOF of sea level pressure in winter Arctic Dipole Mode (ADM)

Beaufort-  
Chukchi


Winter



CONTOUR INTERVAL = 5.000E-02

Barents-  
Kara

Positive ADM




1950


2000

Sea ice anomalies in  
(Barents & Kara) minus  
(Beaufort & Chukchi)  
(1 year lagged)  
vs.  
Winter ADM

Correlation is 0.5  
after trend removal.  
This mode has become  
significant after 1980.



# Summary of shift from NAM/AO to ADM



# Remark and open questions?

- We should separate interannual fluctuations from the trend! Our projection should not be disturbed.
- Precipitation increase under global warming?
- More intense snow and rain also?
- Hydrology cycle important for ice cover?

# Global citizens should

- watch the transportation sector to reduce total energy consumption by using the Arctic routes (higher efficiency often leads to more business activities rather than saving energy)
- more importantly, prevent the petroleum industry from developing offshore oil on the Arctic Shelves (our descendants would criticize the present human as we take advantage of the consequence of global warming/**ice-free Arctic** for acceleration of global warming/**more CO2 emission**)