Title	Roles of specialized DNA polymerases in mutagenesis by 8-hydroxyguanine in human cells
Author(s)	Kamiya, Hiroyuki; Yamaguchi, Ayaka; Suzuki, Tetsuya; Harashima, Hideyoshi
Citation	Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis, 686(1-2), 90-95 https://doi.org/10.1016/j.mrfmmm.2010.02.001
Issue Date	2010-04-01
Doc URL	http://hdl.handle.net/2115/42891
Type	article (author version)
File Information	MRFMMM686-1-2_90-95.pdf

[Instructions for use](#)

Roles of specialized DNA polymerases in mutagenesis by 8-hydroxyguanine in human cells

Hiroyuki Kamiya^{a,*}, Ayaka Yamaguchi^a, Tetsuya Suzuki^{a,b}, and Hideyoshi Harashima^a

^aFaculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

^bPresent address: Division of Genetics and Mutagenesis, National Institute of Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

*Corresponding author: Tel +81-11-706-3733, Fax +81-11-706-4879, E-mail hirokam@pharm.hokudai.ac.jp (Hiroyuki Kamiya)

Abstract

The formation of 8-hydroxyguanine (8-OH-Gua, 7,8-dihydro-8-oxoguanine) in DNA and in the nucleotide pool results in G:C→T:A and A:T→C:G substitution mutations, respectively, due to the ability of 8-OH-Gua to pair with both C and A. In this study, shuttle plasmid DNAs containing 8-OH-Gua paired with C and A in the *supF* gene were transfected into human 293T cells, in which specialized DNA polymerases were knocked-down. The DNAs replicated in the cells were recovered and then introduced into an indicator strain of *Escherichia coli*. Mutation analysis indicated that the knock-downs of DNA polymerases η and ζ by siRNAs enhanced the G:C→T:A mutations caused by 8-OH-Gua:C. The 8-OH-Gua:C-induced mutation frequency was not further increased by double knock-downs of DNA polymerases η and ζ , suggesting that the two DNA polymerases work in the same pathway. In addition, the reduction of DNA polymerase η slightly decreased the A:T→C:G substitutions caused by 8-OH-Gua:A. These results suggest that DNA polymerases η and ζ are involved in the bypass of 8-OH-Gua in human cells.

Keywords: 8-Hydroxyguanine; specialized DNA polymerases; oxidative mutagenesis

Abbreviations: 8-OH-dGTP, 8-hydroxy-2'-deoxyguanosine 5'-triphosphate; 8-OH-Gua, 8-hydroxyguanine; MBL; methylene blue plus light; pol, polymerase; ROS, reactive oxygen species; RT, reverse transcription; TLS, translesion synthesis.

1. Introduction

DNA is continuously oxidized by reactive oxygen species (ROS) that are produced endogenously by normal oxygen metabolism as well as by many environmental mutagens and carcinogens. 8-Hydroxyguanine (8-OH-Gua, 7,8-dihydro-8-oxoguanine) is one of the major oxidized nucleobases produced by ROS, and 100–500 residues are generated per cell per day [1-3]. This oxidized G base is miscoding, due to its ability to form base pairs with A in addition to C [4-7]. Mutagenesis experiments with DNA containing 8-OH-Gua at a predetermined site indicated that 8-OH-Gua is highly mutagenic and induces G:C→T:A transversion mutations in *Escherichia coli* and mammalian cells [8-16]. ROS oxidize DNA precursors (2'-deoxyribonucleotides) as well as DNA. 8-Hydroxy-2'-deoxyguanosine 5'-triphosphate (8-OH-dGTP, 7,8-dihydro-8-oxo-dGTP) is the major oxidation product of dGTP in *in vitro* oxidation reactions [17]. Pursell et al. demonstrated that the concentration of 8-OH-dGTP was 1–10% of that of unmodified dGTP in the mitochondrial nucleotide pool [18]. 8-OH-dGTP is mutagenic and specifically induced A:T→C:G transversions in living bacterial and mammalian cells [19-21]. The A:T→C:G mutations seem to be triggered by the incorporation of 8-OH-dGTP opposite A in template DNA, resulting in the formation of 8-OH-Gua:A base pairs, and the subsequent incorporation of dCTP opposite 8-OH-Gua during the next round of replication would cause the transversion [22].

Mammalian cells contain specialized DNA polymerases (pols) that play important roles in translesion synthesis (TLS) [23,24]. It has been shown that TLS DNA pols bypassed 8-OH-Gua *in vitro* [25-34]. Thus, to understand the mutagenesis and carcinogenesis processes by 8-OH-Gua, it is important to examine the contributions of the polymerases to the induction and suppression of the G:C→T:A mutations caused by this lesion in living mammalian cells. Previously, Lee and Pfeifer

transfected plasmid DNA after a treatment with methylene blue plus light (MBL), which produces predominantly 8-OH-Gua in DNA, into human cells in which DNA pol η was knocked-down and mutated [35]. However, the MBL treatment could potentially produces other types of DNA damages [36]. Therefore, we surmised that the transfection of plasmid DNA containing 8-OH-Gua at a specific position would provide more direct evidence. In addition, we attempted to examine the involvement of other TLS DNA pols in the 8-OH-Gua-induced mutagenesis.

We recently found that the knock-downs of DNA pols η and ζ , and REV1 reduced the frequency of A:T $\rightarrow$ C:G mutations induced by exogenous 8-OH-dGTP [21], suggesting that these proteins are involved in the mutagenesis induced by 8-OH-dGTP. This result raised the question of whether these DNA pols contribute to the incorporation of 8-OH-dGTP and/or to the insertion of dCTP opposite 8-OH-Gua. The discrimination would be possible if we introduced plasmid DNA containing an 8-OH-Gua:A pair, an intermediate in the mutagenic process of 8-OH-dGTP [22], into cells with knocked-down TLS DNA pols.

In this study, double-stranded plasmid DNA containing an 8-OH-Gua:C pair was transfected into human cells in which DNA pols η , ι , and ζ , and REV1 were knocked-down. Moreover, plasmid DNA containing an 8-OH-Gua:A pair was also introduced into the knocked-down cells, as the intermediate in the mutagenic process of 8-OH-dGTP. We found that the knock-downs of DNA pols η and ζ enhanced the G:C $\rightarrow$ T:A transversion mutation induced by the 8-OH-Gua:C pair. Moreover, the knock-down of DNA pol η reduced the A:T $\rightarrow$ C:G substitution mutation by the 8-OH-Gua:A pair. These results suggest that some TLS DNA pols indeed play important roles in the mutagenesis processes of 8-OH-Gua.

2. Materials and methods

2.1. Materials

Oligodeoxyribonucleotides containing 8-OH-Gua and their control oligodeoxyribonucleotides were purchased from Japan Bio Services (Asaka, Japan) and purified by reverse phase and anion exchange HPLC, as described previously [37]. Other oligodeoxyribonucleotides were obtained from Hokkaido System Science (Sapporo, Japan) and Sigma Genosys Japan (Ishikari, Japan) in purified forms. The siRNAs (“stealth RNAi”, Invitrogen, Carlsbad, California, USA) were the same ones used in the previous studies [21]. The control siRNA was that for low GC content (Invitrogen). The *E. coli* strain KS40/pOF105 [38] was provided by Dr. Tatsuo Nunoshiba, of International Christian University, and was used as an indicator strain of the *supF* mutants.

2.2. Plasmid DNAs containing 8-OH-Gua

The oligodeoxyribonucleotides indicated below were synthesized and chemically phosphorylated at their 5'-end on the support during synthesis.

T-96, 5'-dGCAGACTCTAAATCTGCCGTCAT;

ohG-96, 5'-dGCAGACTCG*AAATCTGCCGTCAT;

G-122, 5'-dCGACTTCGAAGGTTCTGAATCC;

ohG-122, 5'-dCGACTTCGAAGG*TTCGAATC; where G* represents 8-OH-Gua.

Double-stranded plasmid DNAs containing 8-OH-Gua were constructed as described [39] from the single-stranded forms of pZ189-StuI [21] and the oligodeoxyribonucleotides listed above, by annealing and treatments with T4 DNA pol and T4 DNA ligase, and with Dam methylase.

2.3. *Mutagenesis experiments*

For the experiments using the G:C and 8-OH-Gua:C plasmid DNAs, 293T cells (3×10^4 cells) were plated into 24-well dishes and were cultured in Dulbecco's modified Eagle medium supplemented with 10% fetal calf serum, at 37°C under a 5% CO₂ atmosphere for 24 h. The siRNA (7.2 pmol each) and "carrier DNA", the pZeo plasmid (100 ng), which was constructed by ligation of the EM7-Zeo and pUC *ori* fragments of pCMV/Zeo (Invitrogen), were mixed with Lipofectamine (Invitrogen) and introduced into the cultured 293T cells according to the supplier's recommendations. Note that the pZeo plasmid lacks the ampicillin resistance gene and thus does not affect the mutagenesis assay. After 24 h, the plasmids containing G:C and 8-OH-Gua:C (100 ng, 29 fmol) was transfected with Lipofectamine. After 48 h of culture, the plasmid amplified in the cells was recovered by the method of Stary and Sarasin [40]. The recovered DNA was treated with DpnI, to digest the unreplicated plasmids, and was introduced into *E. coli* KS40/pOF105 cells by electroporation. The *supF* mutant frequency was determined as described previously [21,38]. For the T:A and 8-OH-Gua:A plasmids, the mutagenesis experiments were performed similarly, using 293T cells (6×10^4 cells) in 12-well dishes.

2.4. *Quantitative reverse transcription (RT)-PCR analysis of mRNA*

Total RNA was extracted from 293T cells using an RNeasy Mini Kit (Qiagen, Hilden, Germany) combined with RNase-free DNase I (Takara, Otsu, Japan) for the degradation of genomic DNA in total RNA samples. First-strand cDNA synthesis was performed on 500 ng of total RNA using an oligo dT primer and an RNA PCR Kit (AMV) (Takara), according to the manufacturer's instructions. Each of the mRNA transcripts was measured by the quantitative PCR method with an ABI 7500 Real Time PCR System and SYBR-Green chemistry (Applied Biosystems, Foster City, CA), as described previously [21]. Data are expressed as the ratio to the GAPDH mRNA.

2.5. Statistical analysis

Statistical significance was examined by the Student's *t*-test. Levels of $P < 0.05$ were considered to be significant.

3. Results

3.1. Design of plasmid DNAs involving 8-OH-Gua

We constructed plasmid DNAs containing 8-OH-Gua:C and 8-OH-Gua:A base pairs (Fig. 1). The parental plasmid, pZ189-StuI, contains the *supF* gene, the SV40 large T antigen gene, and the SV40 origin of replication [21]. 8-OH-Gua was introduced into positions 122 and 96 of the sense strand of the *supF* gene for 8-OH-Gua:C and 8-OH-Gua:A, respectively. Position 96 was one of the

A:T→C:G hot spots in the *supF* gene when 8-OH-dGTP was introduced [20,21]. Plasmid DNAs containing G:C and T:A at positions 122 and 96, respectively, were used as control plasmids.

3.2. Knock-downs of DNA polymerases η and ζ affect the induced mutation frequency of the 8-OH-Gua:C pair

We first examined the effects of the knock-downs of specialized DNA pols by siRNAs on mutations induced by 8-OH-Gua:C. DNA pols η , ι , and ζ (REV3), and REV1 were selected as the knock-down targets, as in the previous study [21]. These knock-downs were measured by quantitative RT-PCR (Table 1). The treatment of 293T cells with siRNA reduced the amounts of the mRNAs encoding these DNA pols. Since we did not observe the effective knock-down of DNA pol κ , this pol was excluded from the analysis.

The treatment with siRNAs against DNA pols η , ι , and ζ , and REV1 resulted in *supF* mutant frequencies similar to the case for the control siRNA in the experiments using the G:C plasmid ($\sim 1 \times 10^{-3}$, Fig. 2, open columns), indicating that the reduction in these DNA pols did not affect the background mutant frequency in this experimental system. When the plasmid with the 8-OH-Gua:C pair at position 122 was introduced into 293T cells treated with the control siRNA, the *supF* mutant frequency was increased to $\sim 4 \times 10^{-3}$ (Fig. 2), indicating the mutagenicity of 8-OH-Gua in DNA. The *supF* mutant frequencies were enhanced to $\sim 6.5 \times 10^{-3}$ in the cells where DNA pols η and ζ were knocked-down (Fig. 2). This result suggests that the two specialized DNA pols are involved in “high-fidelity” bypass of the 8-OH-Gua lesion.

We next examined the effects of the double knock-downs of DNA pols η and ζ . These knock-downs could further increase the *supF* mutant frequency when the two DNA pols act independently. However, we did not observe the expected additive effect upon the double knock-downs (Fig. 3). This result may suggest that DNA pols η and ζ work in the same pathway in the mutation process of 8-OH-Gua, although the knock-down efficiencies for these DNA pols were lower than those in the single knock-down experiments at 24 h, at the time point of transfection of the *supF* plasmid (Table 1).

We analyzed the sequences of the *supF* genes in colonies on the selection agar plates. As expected, the G:C→T:A transversion was the mutations found most frequently in the colonies obtained from the control siRNA-treated and knocked-down cells (Table 2). The total *supF* mutant frequencies were multiplied by the percentages of the targeted G:C→T:A transversion, and the targeted G:C→T:A frequencies were calculated to be 2.6, 5.0, and 4.8 × 10⁻³ in the control, pol η , and pol ζ experiments, respectively. Thus, we concluded that the G:C→T:A mutation was increased by the knock-down of the two TLS DNA pols, and that they play suppressive roles in the mutagenesis caused by 8-OH-Gua:C. In contrast, no G:C→T:A mutation at position 122 was observed for the transfection of the control G:C plasmid (Supplementary Table 1).

Interestingly, the G:C→A:T transition mutation was found in five of the 57 colonies obtained by transfection of the 8-OH-Gua:C plasmid into the cells where pol ι was knocked-down (Table 2). The G:C→C:G mutation was detected in all cases, except for the DNA pol η experiment.

3.3. *Effects of knock-downs of specialized DNA polymerases on the induced mutation frequency of the 8-OH-Gua:A pair*

Recently, we found that the knock-downs of DNA pols η and ζ , and REV1 decreased the A:T $\rightarrow$ C:G mutations induced by exogenous 8-OH-dGTP [21]. In the 8-OH-dGTP experiment, knock-down of a certain DNA pol could affect (i) incorporation of 8-OH-dGTP opposite A during the first round of replication and (ii) that of dCTP opposite 8-OH-Gua after 8-OH-dGTP was incorporated into DNA. We next examined the effects of the knock-downs of specialized DNA pols on the mutations induced by 8-OH-Gua:A, to assess their contributions to the insertion of dCTP opposite 8-OH-Gua. The plasmid DNA containing an 8-OH-Gua:A pair at position 96 of the *supF* gene was prepared. This position was one of the hot spots of the A:T $\rightarrow$ C:G mutations induced by 8-OH-dGTP [20,21].

The treatment with siRNAs against the TLS DNA pols did not affect the *supF* mutant frequencies in the experiments using the T:A plasmid ($\sim 1 \times 10^{-3}$, Fig. 4A), although a slightly higher frequency was observed for the knock-down of DNA pol ζ ($P=0.08$). When the plasmid with the 8-OH-Gua:A pair at position 96 was introduced into 293T cells treated with the control siRNA, the *supF* mutant frequency was 88×10^{-2} (88%), which is quite high (Fig. 4B). This high mutant frequency was due to the MUTYH protein, which removes A opposite 8-OH-Gua [41]. The *supF* mutant frequencies were decreased to 81×10^{-2} (81%) in the cells where DNA pol η was knocked-down ($P<0.05$). This result suggests that this DNA pol is involved in the incorporation of dCTP opposite the 8-OH-Gua base, although the effect of the knock-down was minimal.

We analyzed the sequences of the *supF* genes in colonies on the selection agar plates (Table 3 and Supplementary Table 2). As expected, the A:T→C:G transversion was found in most cases for the 8-OH-Gua:A plasmid. In the cases of the control and pol η siRNAs, all colonies examined contained the A:T→C:G mutation. Thus, the knock-down of DNA pol η reduced the occurrence of the A:T→C:G mutation induced by 8-OH-Gua:A by 8%, while the knock-down decreased the A:T→C:G mutations caused by exogenous 8-OH-dGTP by 32% [21]. Mutations other than A:T→C:G at position 96 were found in three and one colonies for the knock-downs of DNA pols ι and ζ , respectively (Table 3).

4. Discussion

In this study, we examined the involvement of TLS DNA pols in the mutagenesis by 8-OH-Gua. As shown in Fig. 2, the reduction of DNA pol η , a Y-family DNA pol, significantly increased the mutations induced by 8-OH-Gua paired with C in DNA. This result suggests that DNA pol η plays a suppressive role in the mutagenesis by 8-OH-Gua in human cells. The incorporation of 2'-deoxyribonucleotides opposite 8-OH-Gua by human DNA pol η was analyzed *in vitro*. Human DNA pol η incorporates dATP and dCTP opposite the lesion with comparable efficiencies, but proliferating cell nuclear antigen (PCNA) and replication protein A (RP-A) enhance the correct 8-OH-Gua bypass [25-30]. The mammalian replicative DNA pols α and δ are error-prone against 8-OH-Gua [27,28,42-47]. Thus, the knock-down of DNA pol η might force the “error-prone” replicative DNA pols to bypass 8-OH-Gua.

In addition, the knock-down of DNA pol ζ , a B-family DNA pol, increased the mutagenesis caused by 8-OH-Gua paired with C (Fig. 2). This result suggests that DNA pol ζ also plays a suppressive role in the 8-OH-Gua-induced mutations in human cells. DNA pol ζ is a highly specialized pol, and it extends primer termini opposite from a diverse array of DNA lesions, including 8-OH-Gua [24,48]. Thus, DNA pol ζ might extend a nascent strand from C (correct 2'-deoxyribonucleotide) opposite 8-OH-Gua that was incorporated by other DNA pols.

The double knock-downs of DNA pols η and ζ did not further increase the mutant frequency (Fig. 3) and this result may suggest that these DNA pols work in the same pathway. Their properties described above prompted us to speculate that DNA pol η incorporated dCTP opposite 8-OH-Gua and that DNA pol ζ extended the C nucleotide opposite 8-OH-Gua. However, this interpretation should be reviewed carefully, since the knock-down efficiencies for these DNA pols in the double knock-down experiment were lower than those in the single knock-down experiments at 24 h, at the time point of transfection of the *supF* plasmid (Table 1). Moreover, this result was in contrast to the previous finding that the double knock-downs of DNA pols η and ζ affected additively when mutations induced by 8-OH-dGTP were analyzed [21].

In contrast, the suppression of a Y-family DNA pol, DNA pol ι , did not affect the mutant frequency of the plasmid containing 8-OH-Gua:C (Fig. 2). Although the incorporation of 2'-deoxyribonucleotides opposite 8-OH-Gua in template DNA by DNA pol ι *in vitro* was reported [31-33], the result obtained in this study suggests that the contribution of the DNA pol to 8-OH-Gua bypass is minimal, at best, at least in this human cell line. REV1 is a member of the Y-family DNA pols, but it acts as deoxycytidyl transferase opposite a range of damaged bases [23,24]. Human

REV1 inserts dCTP opposite 8-OH-Gua *in vitro* [34]. However, the knock-down of REV1 did not affect the mutation caused by 8-OH-Gua (Fig. 2), suggesting that this error-free function of the protein is not important in the case of 8-OH-Gua:C.

We recently reported the A:T→C:G mutations induced by 8-OH-dGTP in living simian and human cells [20,21]. In this study, we introduced plasmid DNA containing 8-OH-Gua:A as an intermediate in the mutagenic process of 8-OH-dGTP (Fig. 1). Unexpectedly, the total *supF* mutant frequency was 88×10^{-2} in the case of 293T cells treated with the control siRNA (Fig. 4B). Since all colonies among the 50 colonies obtained in the experiment contained the targeted A:T→C:G mutation (Table 3), the mutation frequency of 8-OH-Gua:A was 88×10^{-2} in 293T cells. Le Page et al. reported that the mutation frequencies of 8-OH-Gua:A were 36 and 32×10^{-2} in simian COS-7 and human MRC5V1 cells, respectively [49]. This discrepancy would be due to the difference in the MUTYH activity, which removes an A base opposite 8-OH-Gua. The knock-down of MUTYH in 293T cells reduced the mutagenicity of the 8-OH-Gua:A base pair (to $\sim 40 \times 10^{-2}$) [41].

As shown in Fig. 4B, a reduction in the amount of DNA pol η significantly decreased the mutation induced by 8-OH-Gua paired with A (88×10^{-2} to 81×10^{-2}). This $\sim 8\%$ reduction in the mutant frequency was in contrast to the $\sim 32\%$ decrease in mutant frequency by the knock-down of DNA pol η in the previous 8-OH-dGTP introduction experiment [21]. Since the knock-down efficiency of the pol η mRNA (79% at 24 h, Table 1) was higher than that in the previous study (56% at 24 h), the slight effect observed in this study was not due to low knock-down efficiency. As mentioned above, the A:T→C:G mutation induced by 8-OH-Gua:A was MUTYH-dependent. Based on the quantitatively different effects of the knock-down of DNA pol η on the mutations

induced by 8-OH-dGTP and 8-OH-Gua:A, we speculated that DNA pol η did not play the major role in gap-filling after the removal of A opposite 8-OH-Gua by MUTYH. DNA pol λ was suggested to be involved in the bypass of 8-OH-Gua in the gap-filling reactions [28,50]. Taken together with the increased mutation by 8-OH-Gua:C and the decreased mutation by 8-OH-Gua:A in the cells with knocked-down pol η , this DNA pol would bypass 8-OH-Gua during DNA replication, with a higher dCTP/dATP incorporation ratio than those of the replicative DNA pols. Moreover, the decrease in the A:T $\rightarrow$ C:G mutations by 8-OH-dGTP in the pol η -knocked-down cells, observed in the previous study [21], was suggested to be mainly due to reduced 8-OH-dGTP incorporation into the nascent strand. This speculation is consistent with the highly erroneous incorporation of 8-OH-dGTP opposite A by this DNA pol *in vitro* [51,52].

In contrast to the previous study [21], no obvious effects were observed when DNA pol ζ and REV1 were knocked-down (Fig. 4B). Thus, DNA pol ζ and REV1 seemed to play important roles in the incorporation of 8-OH-dGTP but not in the insertion of dCTP opposite 8-OH-Gua.

Lee and Pfeifer introduced MBL-treated plasmid DNA into human cells in which DNA pol η was knocked-down and mutated [35]. They observed an increase in the total mutant frequency by the reduction/loss of DNA pol η , as we have shown in this report (Fig. 2). However, the ratios of G:C $\rightarrow$ T:A transversions in the knocked-down/knock-out cells were similar to (or less than) that in the control cells. Since the MBL treatment produces other damaged bases besides 8-OH-Gua, such as 2,5-diamino-4*H*-imidazol-4-one [36], their results might also partly reflect the effects of the reduction/loss on mutations caused by several damaged bases,.

Avkin and Livneh introduced a gapped plasmid carrying 8-OH-Gua in the single-stranded DNA region into DNA pol η -proficient and -deficient human cells [53]. They reported that the ratio of dATP insertion (corresponding to the G:C $\rightarrow$ T:A mutation) was nearly identical in both cell lines. Their result was in contrast to our present observation that the knock-down of DNA pol η increased the mutation frequency and the ratio of the G:C $\rightarrow$ T:A mutation (Fig. 2 and Table 2). This discrepancy is a consequence of the gapped plasmid that Avkin and Livneh used. DNA pol η might not be recruited to the gap, and thus may not participate in the 8-OH-Gua bypass.

In conclusion, the knock-downs of DNA pols η and ξ enhanced the mutagenicity of 8-OH-Gua paired with C in human 293T cells. This result suggests that these DNA pols are involved in the error-free bypass of the oxidized G base. Moreover, the knock-downs of DNA pol η slightly reduced the mutations induced by 8-OH-Gua paired with A. In addition to DNA pols, DNA repair enzymes and nucleotide pool sanitization enzymes would play important roles in the mutagenesis processes of damaged DNA and DNA precursors [54]. Experiments to reveal their roles are currently in progress.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

This work was supported in part by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan, and the Takeda Science Foundation to H.K.

References

- [1] T. Lindahl, Instability and decay of the primary structure of DNA, *Nature* 362 (1993) 709-715.
- [2] H. Kasai, Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis, *Mutat. Res.* 387 (1997) 147-163.
- [3] H. Kamiya, Mutagenicities of 8-hydroxyguanine and 2-hydroxyadenine produced by reactive oxygen species, *Biol. Pharm. Bull.* 27 (2004) 475-479.
- [4] Y. Oda, S. Uesugi, M. Ikehara, S. Nishimura, Y. Kawase, H. Ishikawa, H. Inoue, E. Ohtsuka, NMR studies of a DNA containing 8-hydroxydeoxyguanosine, *Nucleic Acids Res.* 19 (1991) 1407-1412.
- [5] M. Kouchakdjian, V. Bodepudi, S. Shibutani, M. Eisenberg, F. Johnson, A.P. Grollman, D.J. Patel, NMR structural studies of the ionizing radiation adduct 7-hydro-8-oxodeoxyguanosine (8-oxo-7H-dG) opposite deoxyadenosine in a DNA duplex. 8-Oxo-7H-dG(syn)•dA(anti) alignment at lesion site, *Biochemistry* 30 (1991) 1403-1412.
- [6] G.E. Plum, A.P. Grollman, F. Johnson, K.J. Breslauer, Influence of the oxidatively damaged adduct 8-oxodeoxyguanosine on the conformation, energetics, and thermodynamic stability of a DNA duplex, *Biochemistry* 34 (1995) 16148-16160.
- [7] S. Koizume, H. Kamiya, H. Inoue, E. Ohtsuka, Synthesis and thermodynamic stabilities of damaged DNA involving 8-hydroxyguanine (7,8-dihydro-8-oxoguanine) in a *ras* gene fragment, *Nucleos. Nucleot.* 13 (1994) 1517-1534.

- [8] M.L. Wood, M. Dizdaroglu, E. Gajewski, J.M. Essigmann, Mechanistic studies of ionizing radiation and oxidative mutagenesis: genetic effects of a single 8-hydroxyguanine (7-hydro-8-oxoguanine) residue inserted at a unique site in a viral genome, *Biochemistry* 29 (1990) 7024-7032.
- [9] K.C. Cheng, D.S. Cahill, H. Kasai, S. Nishimura, L.A. Loeb, 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G→T and A→C substitutions, *J. Biol. Chem.* 267 (1992) 166-172.
- [10] M. Moriya, A.P. Grollman, Mutations in the *mutY* gene of *Escherichia coli* enhance the frequency of targeted G:C→T:A transversions induced by a single 8-oxoguanine residue in single-stranded DNA, *Mol. Gen. Genet.* 239 (1993) 72-76.
- [11] J. Wagner, H. Kamiya, R.P.P. Fuchs, Leading versus lagging strand mutagenesis induced by 7,8-dihydro-8-oxo-2'-deoxyguanosine in *E. coli*, *J. Mol. Biol.* 265 (1997) 302-309.
- [12] H. Kamiya, K. Miura, H. Ishikawa, H. Inoue, S. Nishimura, E. Ohtsuka, c-Ha-*ras* containing 8-hydroxyguanine at codon 12 induces point mutations at the modified and adjacent positions, *Cancer Res.* 52 (1992) 3483-3485.
- [13] M. Moriya, Single-stranded shuttle phagemid for mutagenesis studies in mammalian cells: 8-oxoguanine in DNA induces targeted G•C→T•A transversions in simian kidney cells, *Proc. Natl. Acad. Sci. U.S.A.* 90 (1993) 1122-1126.
- [14] H. Kamiya, N. Murata-Kamiya, S. Koizume, H. Inoue, S. Nishimura, E. Ohtsuka, 8-Hydroxyguanine (7,8-dihydro-8-oxoguanine) in hot spots of the c-Ha-*ras* gene: effects of sequence contexts on mutation spectra, *Carcinogenesis* 16 (1995) 883-889.

- [15] F. Le Page, A. Margot, A.P. Grollman, A. Sarasin, A. Gentil, Mutagenicity of a unique 8-oxoguanine in a human *Ha-ras* sequence in mammalian cells, *Carcinogenesis* 16 (1995) 2779-2784.
- [16] X. Tan, A.P. Grollman, S. Shibutani, Comparison of the mutagenic properties of 8-oxo-7,8-dihydro-2'-deoxyadenosine and 8-oxo-7,8-dihydro-2'-deoxyguanosine DNA lesions in mammalian cells, *Carcinogenesis* 20 (1999) 2287-2292.
- [17] N. Murata-Kamiya, H. Kamiya, M. Muraoka, H. Kaji, H. Kasai. Comparison of oxidation products from DNA components by γ -irradiation and Fenton-type reactions, *J. Radiat. Res.* 38 (1997) 121-131.
- [18] Z.F. Pursell, J.T. McDonald, C.K. Mathews, T.A. Kunkel. Trace amounts of 8-oxo-dGTP in mitochondrial dNTP pools reduce DNA polymerase γ replication fidelity, *Nucleic Acids Res.* 36 (2008) 2174-2181.
- [19] M. Inoue, H. Kamiya, K. Fujikawa, Y. Ootsuyama, N. Murata-Kamiya, T. Osaki, K. Yasumoto, H. Kasai, Induction of chromosomal gene mutations in *Escherichia coli* by direct incorporation of oxidatively damaged nucleotides, *J. Biol. Chem.* 273 (1998) 11069-11074.
- [20] K. Satou, K. Kawai, H. Kasai, H. Harashima, H. Kamiya, Mutagenic effects of 8-hydroxy-dGTP in live mammalian cells, *Free Radic. Biol. Med.* 42 (2007) 1552-1560.
- [21] K. Satou, M. Hori, K. Kawai, H. Kasai, H. Harashima, H. Kamiya, Involvement of specialized DNA polymerases in mutagenesis by 8-hydroxy-dGTP in human cells, *DNA Repair (Amst.)* 8 (2009) 637-642.

- [22] H. Kamiya, Mutations induced by oxidized DNA precursors and their prevention by nucleotide pool sanitization enzymes, *Genes Environ.* 29 (2007) 133-140.
- [23] A.R. Lehmann, A. Niimi, T. Ogi, S. Brown, S. Sabbioneda, J.F. Wing, P.L. Kannouche, C.M. Green, Translesion synthesis: Y-family polymerases and the polymerase switch, *DNA Repair (Amst.)* 6 (2007) 891-899.
- [24] S. Prakash, R.E. Johnson, L. Prakash, Eukaryotic translesion synthesis DNA polymerases: specificity of structure and function, *Annu. Rev. Biochem.* 74 (2005) 317-353.
- [25] L. Haracska, S.-L. Yu, R.E. Johnson, L. Prakash S. Prakash, Efficient and accurate replication in the presence of 7,8-dihydro-8-oxoguanine by DNA polymerase η , *Nat. Genet.* 25 (2000) 458-461.
- [26] Y. Zhang, F. Yuan, X. Wu, O. Rechkoblit, J.-S. Taylor, N.E. Geacintov, Z. Wang, Error-prone lesion bypass by human DNA polymerase η , *Nucleic Acids Res.* 28 (2000) 4717-4724.
- [27] P. Jałoszyński, C. Masutani, F. Hanaoka, A.B. Perez, S. Nishimura, 8-Hydroxyguanine in a mutational hotspot of the *c-Ha-ras* gene causes misreplication, 'action-at-a-distance' mutagenesis and inhibition of replication, *Nucleic Acids Res.* 31 (2003) 6085-6095.
- [28] G. Maga, G. Villani, E. Crespan, U. Wimmer, E. Ferrari, B. Bertocci, U. Hübscher, 8-Oxo-guanine bypass by human DNA polymerases in the presence of auxiliary proteins, *Nature* 447 (2007) 606-608.

- [29] C.-W. Yung, Y. Okugawa, C. Otsuka, K. Okamoto, S. Arimoto, D. Loakes, K. Negishi, T. Negishi, Influence of neighbouring base sequences on the mutagenesis induced by 7,8-dihydro-8-oxoguanine in yeast, *Mutagenesis* 23 (2008) 509-513.
- [30] S.D. McCulloch, R.J. Kokoska, P. Garg, P.M. Burgers, T.A. Kunkel, The efficiency and fidelity of 8-oxo-guanine bypass by DNA polymerases δ and η , *Nucleic Acids Res.* 37 (2009) 2830-2840.
- [31] Y. Zhang, F. Yuan, X. Wu, J.-S. Taylor, Z. Wang, Response of human DNA polymerase ι to DNA lesions, *Nucleic Acids Res.* 29 (2001) 928-935.
- [32] A. Vaisman, R. Woodgate, Unique misinsertion specificity of pol ι may decrease the mutagenic potential of deaminated cytosines, *EMBO J.* 20 (2001) 6520-6529.
- [33] R.E. Johnson, L. Haracska, L. Prakash, S. Prakash, Role of Hoogsteen edge hydrogen bonding at template purines in nucleotide incorporation by human DNA polymerase ι , *Mol. Cell. Biol.* 26 (2006) 6435-6441.
- [34] Y. Zhang, X. Wu, O. Rechkoblit, N.E. Geacintov, J.-S. Taylor, Z. Wang, Response of human REV1 to different DNA damage: preferential dCMP insertion opposite the lesion, *Nucleic Acids Res.* 30 (2002) 1630-1638.
- [35] D.-H. Lee, G.P. Pfeifer, Translesion synthesis of 7,8-dihydro-8-oxo-2'-deoxyguanosine by DNA polymerase η in vivo, *Mutat. Res.* 641 (2008) 19-26.
- [36] G.W. Buchko, J. Cadet, B. Morin, M. Weinfeld, Photooxidation of d(TpG) by riboflavin and methylene blue. Isolation and characterization of thymidylyl-(3',5')-2-amino-5-[(2-deoxy- β -D-*erythro*-pentofuranosyl)amino]-4*H*-imidazol-4-

one and its primary decomposition product thymidyl-(3',5')-2,2-diamino-4-[(2-deoxy- β -D-*erythro*-pentofuranosyl)amino]-5(2*H*)-oxazalone, Nucleic Acids Res. 23 (1995) 3954-3961.

- [37] H. Kamiya, H. Kasai, Substitution and deletion mutations induced by 2-hydroxyadenine in *Escherichia coli*: effects of sequence contexts in leading and lagging strands, Nucleic Acids Res. 25 (1997) 304-310.
- [38] F. Obata, T. Nunoshiba, T. Hashimoto-Gotoh, K. Yamamoto, An improved system for selection of forward mutations in an *Escherichia coli supF* gene carried by plasmids, J. Radiat. Res. 39 (1998) 263-270.
- [39] S. Tornaletti, L.S. Maeda, R.D. Kolodner, P.C. Hanawalt, Effect of 8-oxoguanine on transcription elongation by T7 RNA polymerase and mammalian RNA polymerase II, DNA Repair (Amst.) 3 (2004) 483-494.
- [40] A. Strydom, A. Sarasin, Simian virus 40 (SV40) large T antigen-dependent amplification of an Epstein-Barr virus-SV40 hybrid shuttle vector integrated into the human HeLa cell genome, J. Gen. Virol. 73 (1992) 1679-1685.
- [41] T. Suzuki, H. Harashima, H. Kamiya, Effects of base excision repair proteins on mutagenesis by 8-oxo-7,8-dihydroguanine (8-hydroxyguanine) paired with cytosine and adenine, DNA Repair (Amst.) in press. doi:10.1016/j.dnarep.2010.02.004
- [42] S. Shibutani, M. Takeshita, A. P. Grollman, Insertion of specific bases during DNA synthesis past the oxidation-damaged base 8-oxodG, Nature 349 (1991) 431-434.

- [43] H. Kamiya, T. Sakaguchi, N. Murata, M. Fujimuro, H. Miura, H. Ishikawa, M. Shimizu, H. Inoue, S. Nishimura, A. Matsukage, C. Masutani, F. Hanaoka, E. Ohtsuka, *In vitro* replication study of modified bases in *ras* sequences, *Chem. Pharm. Bull.* 40 (1992) 2792-2795.
- [44] H. Kamiya, N. Murata-Kamiya, M. Fujimuro, K. Kido, H. Inoue, S. Nishimura, C. Masutani, F. Hanaoka, E. Ohtsuka, Comparison of incorporation and extension of nucleotides *in vitro* opposite 8-hydroxyguanine (7,8-dihydro-8-oxoguanine) in hot spots of the c-Ha-*ras* gene, *Jpn. J. Cancer Res.* 86 (1995) 270-276.
- [45] E. Efrati, G. Tocco, R. Eritja, S.H. Wilson, M.F. Goodman, "Action-at-a-distance" mutagenesis. 8-oxo-7,8-dihydro-2'-deoxyguanosine causes base substitution errors at neighboring template sites when copied by DNA polymerase β , *J. Biol. Chem.* 274 (1999) 15920-15926.
- [46] R. Fazlieva, C.S. Spittle, D. Morrissey, H. Hayashi, H. Yan, Y. Matsumoto, Proofreading exonuclease activity of human DNA polymerase δ and its effects on lesion-bypass DNA synthesis, *Nucleic Acids Res.* 37 (2009) 2854-2866.
- [47] H.J. Einolf, F.P. Guengerich, Fidelity of nucleotide insertion at 8-oxo-7,8-dihydroguanine by mammalian DNA polymerase δ : steady-state and pre-steady-state kinetic analysis, *J. Biol. Chem.* 276 (2001) 3764-3771.
- [48] L. Haracska, S. Prakash, L. Prakash, Yeast DNA polymerase zeta is an efficient extender of primer ends opposite from 7,8-dihydro-8-oxoguanine and *O*⁶-methylguanine, *Mol. Cell. Biol.* 23 (2003) 1453-1459.

- [49] F. Le Page, A. Guy, J. Cadet, A. Sarasin, A. Gentil, Repair and mutagenic potency of 8-oxoG:A and 8-oxoG:C base pairs in mammalian cells, *Nucleic Acids Res.* 26 (1998) 1276-1281.
- [50] G. Maga, E. Crespan, U. Wimmer, B. van Loon, A. Amoroso, C. Mondello, C. Belgiovine, E. Ferrari, G. Locatelli, G. Villani, U. Hübscher, Replication protein A and proliferating cell nuclear antigen coordinate DNA polymerase selection in 8-oxo-guanine repair, *Proc. Natl. Acad. Sci. U.S.A.* 105 (2008) 20689-20694.
- [51] M. Shimizu, P. Gruz, H. Kamiya, S.-R. Kim, F.M. Pisani, C. Masutani, Y. Kanke, H. Harashima, F. Hanaoka, T. Nohmi. Erroneous incorporation of oxidized DNA precursors by Y-family DNA polymerases, *EMBO Rep.* 4 (2003) 269-273.
- [52] M. Shimizu, P. Gruz, H. Kamiya, C. Masutani, Y. Xu, Y. Usui, H. Sugiyama, H. Harashima, F. Hanaoka, T. Nohmi, Efficient and erroneous incorporation of oxidized DNA precursors by human DNA polymerase η , *Biochemistry* 46 (2007) 5515-5522.
- [53] S. Avkin, Z. Livneh. Efficiency, specificity and DNA polymerase-dependence of translesion replication across the oxidative DNA lesion 8-oxoguanine in human cells, *Mutat. Res.* 510 (2002) 81-90.
- [54] H. Kamiya, Mutagenic potentials of damaged nucleic acids produced by reactive oxygen/nitrogen species: approaches using synthetic oligonucleotides and nucleotides, *Nucleic Acids Res.* 31 (2003) 517-531.

Figure legends

Fig. 1 Structures of plasmid DNAs containing 8-OH-Gua, used in this study. 8-OH-Gua was introduced to either positions 96 (8-OH-Gua:A) or 122 (8-OH-Gua:C) in the *supF* gene. The X's indicate the positions where 8-OH-Gua was incorporated.

Fig. 2 Effects of knock-downs of specialized DNA pols in 293T cells on the mutant frequencies induced by 8-OH-Gua:C. Open columns, control plasmid containing G:C at position 122; closed columns, plasmid containing 8-OH-Gua:C at position 122. Data are expressed as means $\pm$ SEM (n=4 for control siRNA (G:C and 8-OH-Gua:C), pol η siRNA, (8-OH-Gua:C), and pol ι siRNA (G:C), and n=3 for others). (* P <0.05 vs control).

Fig. 3 Effects of double knock-downs of DNA pols η and ξ in 293T cells on the mutant frequencies induced by 8-OH-Gua:C. Open columns, control plasmid containing G:C at position 122; closed columns, plasmid containing 8-OH-Gua:C at position 122. Data are expressed as means $\pm$ SEM (n=3). (* P <0.05 vs control).

Fig. 4 Effects of knock-downs of specialized DNA pols in 293T cells on mutant frequencies induced by 8-OH-Gua:A. (A) Mutant frequencies upon transfection of the control plasmid containing T:A at position 96. (B) Mutant frequencies upon transfection of the plasmid containing

8-OH-Gua:A at position 96. Data are expressed as means $\pm$ SEM (n=4 for REV3 siRNA (8-OH-Gua:A), n=6 for control siRNA (8-OH-Gua:A), and n=3 for others). (* P <0.05 vs control).

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Table 1

Amounts of mRNAs in 293T cells with knocked-down specialized DNA pols.

DNA pol	24 h	48 h	72 h
Single knock-down experiment			
Pol η	0.21 ± 0.03^a	0.27^b	0.21 ± 0.03^a
Pol ϵ	0.21 ± 0.01^a	0.20^b	0.13^b
REV3	0.41 ± 0.04^a	0.54 ± 0.06^c	0.85 ± 0.12^d
REV1	0.22 ± 0.15^a	0.27 ± 0.00^d	0.37 ± 0.07^a
Double knock-down experiment			
Pol η	0.43 ± 0.03^a	0.51 ± 0.12^a	0.50 ± 0.03^a
REV3	0.54 ± 0.08^a	0.54 ± 0.02^a	0.62 ± 0.07^a

The amount of mRNA was measured by quantitative RT-PCR at 24, 48, and 72 h after siRNA introduction according to the methods described in Section 2. Relative values to that in 293T cells treated with the control siRNA are shown. Data are expressed as means ($\pm$ SEM).

^a $n=3$.

^b $n=2$.

^c $n=5$.

^d $n=4$.

Table 2. Targeted mutations at position 122 found in progeny plasmid of 8-OH-Gua:C
plasmid, replicated in 293T cells with knocked-down DNA pols.

Mutation	Control	Pol h	Poli	REV3 (Pol z)	REV1
Single base substitutions					
Transition					
G:C-->A:T	0 (0)	1 (2)	5 (9)	1 (2)	0 (0)
Transversion					
G:C-->T:A	38 (68)	41 (76)	39 (68)	37 (71)	34 (68)
G:C-->C:G	3 (5)	0 (0)	1 (2)	1 (2)	1 (2)
Single base deletion	0 (0)	2 (4)	1 (2)	0 (0)	0 (0)
Others	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
No mutation	15 (27)	10 (19)	11 (19)	13 (25)	15 (30)
Total	56 (100)	54 (100)	57 (100)	52 (100)	50 (100)

All data are represented as cases found (%).

Table 3. Targeted mutations at position 96 found in progeny plasmid of 8-OH-Gua:A
plasmid, replicated in 293T cells with knocked-down DNA pols.

Mutation	Control	Pol h	Pol i	REV3 (Pol z)	REV1
Single base substitutions					
Transition					
A:T-->G:C	0 (0)	0 (0)	1 (2)	0 (0)	0 (0)
Transversion					
A:T-->T:A	0 (0)	0 (0)	1 (2)	1 (2)	0 (0)
A:T-->C:G	50 (100)	46 (100)	47 (94)	53 (98)	58 (100)
Single base deletion	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Others	0 (0)	0 (0)	1 (2) ^a	0 (0)	0 (0)
No mutation	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Total	50 (100)	46 (100)	50 (100)	54 (100)	58 (100)

All data are represented as cases found (%).

^a8-OH-Gua --> AG.