Title	Aligned 18S for Zoraptera (Insecta) : Phylogenetic position and molecular evolution
Author(s)	Yoshizawa, Kazunori; Johnson, Kevin P.
Citation	Molecular Phylogenetics and Evolution, 37(2), 572-580 https://doi.org/10.1016/j.ympev.2005.05.008
Issue Date	2005-11
Doc URL	http://hdl.handle.net/2115/43133
Type	article (author version)
File Information	2005mpe-1.pdf

[Instructions for use](#)

ELSEVIER

Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

Molecular Phylogenetics and Evolution xxx (2005) xxx–xxx

MOLECULAR
PHYLOGENETICS
AND
EVOLUTIONwww.elsevier.com/locate/ympev

Aligned 18S for Zoraptera (Insecta): Phylogenetic position and molecular evolution

Kazunori Yoshizawa^{a,*}, Kevin P. Johnson^b^a Systematic Entomology, Graduate School of Agriculture, Hokkaido University, Sapporo 060-8589, Japan^b Illinois Natural History Survey, 607 East Peabody Drive, Champaign, IL 61820, USA

Received 11 January 2005; revised 6 May 2005

Abstract

The order Zoraptera (angel insects) is one of the least known insect groups, containing only 32 extant species. The phylogenetic position of Zoraptera is poorly understood, but it is generally thought to be closely related to either Paraneoptera (hemipteroid orders: booklice, lice, thrips, and bugs), Dictyoptera (blattoid orders: cockroaches, termites, and mantis), or Embioptera (web spinners). We inferred the phylogenetic position of Zoraptera by analyzing nuclear 18S rDNA sequences, which we aligned according to a secondary structure model. Maximum likelihood and Bayesian analyses both supported a close relationship between Zoraptera and Dictyoptera with relatively high posterior probability. The 18S sequences of Zoraptera exhibited several unusual properties: (1) a dramatically increased substitution rate, which resulted in very long branches; (2) long insertions at helix E23; and (3) modifications of secondary structures at helices 12 and 18.

© 2005 Published by Elsevier Inc.

Keywords: Zoraptera; 18S rDNA; Secondary structure based alignment; Phylogeny; Molecular evolution

1. Introduction

Zoraptera (angel insects) is one of the least diverse and poorly known insect orders. To date, only 38 species (which includes six fossil species) are described, and all extant species are classified under a single genus, *Zorotypus* (Engel and Grimaldi, 2002). Some other genera have been proposed for extant species (Chao and Chen, 2000; Kukalová-Peck and Peck, 1993), but more the conservative taxonomic system is adopted here, as suggested by Engel and Grimaldi (2000) and New (2000). All species of Zoraptera live under the bark of rotting wood (Smithers, 1991).

Based on morphological characters, the order Zoraptera is thought to be closely related to either Paraneoptera (= hemipteroid orders: bugs, thrips, booklice, and

lice: Hennig, 1981; Kristensen, 1975, 1981; Wheeler et al., 2001), Dictyoptera (= blattoid orders: cockroaches, termites, and mantis: Boudreaux, 1979; Kukalová-Peck and Peck, 1993; Smithers, 1991) or Embioptera (= web spinners: Engel and Grimaldi, 2000; Minet and Bourgoin, 1986). Combined morphological and molecular analysis by Wheeler et al. (2001) supported a close relationship between Zoraptera and Dictyoptera. However, separate analysis of molecular data (18S rDNA) placed Zoraptera as the sister taxon of Psocodea (booklice and parasitic lice), conflicting with combined tree (Wheeler et al., 2001).

Separate analyses of 18S data (Wheeler et al., 2001) resulted in tree with a very unconventional placement of some insect orders (e.g., Diplura and Grylloblattodea were imbedded within Holometabola). Wheeler et al. (2001) used direct optimization of morphological and molecular data, which minimizes incongruence between two data partitions. Kjer (2004) pointed out that, when

* Corresponding author. Fax: +81 11 706 4939.

E-mail address: psocid@res.agr.hokudai.ac.jp (K. Yoshizawa).

support from molecular data for nodes is small, conclusions from molecular data by direct optimization would be highly dependent on some combination of (1) morphological data, (2) noise from the homoplastic data, and (3) arbitrarily optimized homology of unalignable data (see also Kjer, 1995 and Simmons, 2004).

To address the problems of direct optimization, Kjer (2004) conducted phylogenetic analyses of insect orders based on 18S sequences aligned manually according to secondary structure. The resulting tree matched traditional insect classification reasonably well. However, Kjer's (2004) study lacked a sequence of Zoraptera and thus could not address the phylogenetic position of this order. Part of the reason for the exclusion of Zoraptera by Kjer was that he concluded that the 18S of Zoraptera presented in Wheeler et al. (2001) was either contaminated in part by mite (Acari) DNA sequences, because of a homologous unique sequence shared by the Zoraptera and mites, or that if the zorapteran sequence was not a contaminant, it was highly autapomorphic and problematic.

Thus, a more detailed molecular test of the phylogenetic position of Zoraptera is needed. The 18S rDNA gene has played an important role in resolving the deep phylogeny of insects (Campbell et al., 1995; Johnson et al., 2004; Kjer, 2004; Whiting et al., 1997). However, a correct 18S sequence of Zoraptera may not be available to date. In the present study, we amplified and analyzed the 18S rDNA of Zoraptera using samples collected in the USA, Malaysia, and Vietnam. These sequences of Zoraptera plus additional sequences of Blattodea (cockroaches), Phasmatodea (stick insects), Embioptera, and Paraneoptera were aligned with the 18S data provided by Kjer (2004). We address two questions: (1) is the 18S sequence of Zoraptera used by Wheeler et al. (2001) really a contaminant and (2) what is the closest relative of Zoraptera?

2. Materials and methods

We sequenced four species of Zoraptera, *Zorotypus hubbardi* from the USA, *Z. sp.*MY1 and *Z. sp.*MY2 from Malaysia, and *Z. sp.*VN from Vietnam (the latter three species are currently being described). Methods of total DNA extraction and 18S amplification and sequencing followed Johnson et al. (2004). Primer sets used were Ns1-Ns2a (Barker et al., 2003), 18Sai-18Sbi (Whiting et al., 1997), and Ns5aP2-Ns8P (Johnson et al., 2004). The 18S sequence of *Z. snyderi* was obtained from GenBank and was only used to check whether the 18S sequence of the species was contaminant or not. The sequence was not used for phylogenetic analyses because only a short piece of the 18S sequence was available for this species. Additional 18S sequences of Blattodea, Phasmatodea, Embioptera, Psocodea,

Thysanoptera, and Hemiptera were obtained from GenBank (Appendix A). These sequences were manually aligned to the data matrix provided by Kjer (2004) according to the secondary structure model presented on his website. When we detected a modification of the secondary structure in the new sequences, the secondary structure of the region was estimated using GeneBee (Brodsky et al., 1995). Except for these additional samples, the taxon set was largely unchanged from Kjer (2004). However, we replaced sequences of Ectopsocidae Gen. sp. and *Pthirus pubis* with *Ectopsocus perkinsi* and *Pedicinus* sp., respectively, because only a short piece of 18S sequence was available for the former two species (Appendix A). Unalignable regions were excluded from the analyses, and the exclusion set followed Kjer (2004). Aligned data is available at <http://insect3.agr.hokudai.ac.jp/psoco-web/data/>.

Preliminary parsimony (MP) and neighbor-joining (NJ) analyses using PAUP* (Swofford, 2002) placed Zoraptera as the sister taxon of Diptera (flies). Diptera is a holometabolus order (insects with pupal stage) whereas Zoraptera is hemimetabolous (insects without pupal stage), so this result seems unlikely. As mentioned below, the basal branch leading to Zoraptera was very long as was the case for Diptera, and thus this result appeared to be an artifact of long-branch attraction (Felsenstein, 1978). Kjer (2004) also suggested that long-branch attraction was problematic for his MP analysis, with Diptera grouping outside of insect, as the sister taxon of Crustacea. In contrast to MP, the Bayesian tree recovered by Kjer (2004) was more reasonable. Likelihood analysis is thought to be less affected by long-branch attraction (Huelsenbeck, 1997; Huelsenbeck and Hillis, 1993). Therefore, we conducted further phylogenetic analyses using maximum likelihood (ML) in PAUP* (Swofford, 2002) and Bayesian ML in MrBayes (Huelsenbeck and Ronquist, 2001). The simplest model for ML analyses was determined by a hierarchic likelihood ratio test using Modeltest (Posada and Crandall, 1998). The GTR + I + G model was selected (unequal base frequencies: A = 0.2496, C = 0.2210, G = 0.2781, T = 0.2513; six substitution categories: A-C = 1.5445, A-G = 3.5713, A-T = 1.5224, C-G = 0.7884, C-T = 5.0195, G-T = 1; gamma distributions shape parameter = 0.6195; proportion of invariant sites = 0.1861). For ML analysis, the NJ tree was used as a starting tree and TBR branch swapping option was selected. For Bayesian analysis, we ran four chains for 10 million generations, and the tree was sampled every 1000 generations. By analyzing the change in likelihood score during the chain using Tracer (Rambaut and Drummond, 2004), we identified a suitable burn-in of 600,000 generations (Fig. 5). Therefore, the first 600 trees were excluded as burn-in, and we computed a 50% majority consensus tree of the remaining 9400

Table 1

Parameters from the Bayesian likelihood analysis under GTR + I + G model

	Kjer (2004)	Present
$r(A \rightleftharpoons C)$	1.52 ± 0.16	1.56 ± 0.27
$r(A \rightleftharpoons G)$	3.45 ± 0.27	3.57 ± 0.49
$r(A \rightleftharpoons T)$	1.23 ± 0.11	1.49 ± 0.22
$r(C \rightleftharpoons G)$	0.75 ± 0.08	0.69 ± 0.12
$r(C \rightleftharpoons T)$	5.17 ± 0.49	5.27 ± 0.67
$r(G \rightleftharpoons T)$	1.00 ± 0.00	1.00 ± 0.00
Alpha	0.58 ± 0.04	0.60 ± 0.05
Pinvar	0.17 ± 0.02	0.16 ± 0.04

Parameters obtained by Kjer (2004) are also indicated for comparison. Following abbreviations are used: r, substitution rates between the listed nucleotides; alpha, shape parameter of the gamma distribution; and pinvar, proportion of invariable sites.

164 trees to estimate posterior probabilities of branches in
165 the tree Table 1.

166 For ML bootstrapping, the NJ tree was used as a
167 starting tree, and the NNI branch swapping option was
168 selected with 100 replicates. TBR branch swapping was
169 not performed because it was computationally infeasible.
170 However, as mentioned above, the tree obtained by NJ
171 method was problematic, and preliminary analysis indi-
172 cated that NNI branch swapping was not sufficient to es-
173 cape from long-branch attraction caused by a number of
174 problematic taxa: Diplura + Protura (Entognatha),
175 Zoraptera (“Hemimetabola”), and Diptera (Holometab-
176 bola) (Fig. 1). Therefore, to avoid long-branch attraction
177 of these distantly related orders, monophyly of Insecta,
178 Neoptera, and Holometabola were given as three con-
179 straints for ML bootstrapping. Monophyly of those
180 higher level groups have previously received very strong
181 support from morphological and molecular studies and
182 are not controversial (Kjer, 2004; Kristensen, 1975,
183 1981; Wheeler et al., 2001). No constraints were given
184 for ML and Bayesian tree searches.

185 3. Results

186 3.1. Sequences and data evaluation

187 We successfully amplified and sequenced the 18S
188 rDNA gene from four species of Zoraptera. As men-
189 tioned below, the 18S of Zoraptera had large insertions
190 (E23 sensu Wuyts et al., 2000) and modifications of
191 secondary structure. However, all four *Zorotypus* 18S
192 sequences obtained here, as well as sequences of
193 *Z. hubbardi* obtained by Vawter (1991) and *Z. snyderi*
194 obtained by Wheeler et al. (2001), could be readily
195 aligned according to the secondary structure model for
196 insect 18S (Kjer, 2004). As mentioned by Kjer (2004),
197 our preliminary MP and NJ analyses (trees not shown)
198 indicated that the 18S of *Z. hubbardi* analyzed by
199 Vawter (1991) was not close to the other *Zorotypus*

sequences, but was imbedded within an odonate (drag-
onflies) clade composed of the genera *Leucorrhinia*,
Sympetrum, and *Celithemis* (Fig. 1). However, the 18S
of *Z. snyderi* analyzed by Wheeler et al. (2001) was very
similar to all the *Zorotypus* sequences obtained in the
present study. For example, using MP and NJ analyses
based only on the middle segment of 18S available for *Z.*
snyderi (i.e., no missing data), a sister group relationship
between *Z. snyderi* and *Z. hubbardi* and monophyly of
Zoraptera were always recovered with 100% bootstrap
support (trees not shown). The sequence AAAACTTA
CCCGGCC, which appeared in the 18S of *Z. snyderi*
near helix 36 and was considered by Kjer (2004) to be
evidence of acarine contamination, was also detected
in the newly sequenced samples, although the underlined
bases were not A, T, and C but A,T, and A in other
zorapterans (Fig. 4). As mentioned by Kjer (2004), when
this sequence and the neighboring region was subjected
to a BLAST search, the 18S of some Arachnida and
Annelida were returned as the top three matches (Sep-
tember 23, 2004: Fig. 4). However, when the middle por-
tion of these 18S sequences were aligned to the data set
and analyzed by MP and NJ methods (trees not shown),
the arachnid and annelid sequences were distant from
Zoraptera and placed near the root of the tree.

In addition to this unusual short fragment, other un-
ique characteristics were observed in the 18S sequences
of Zoraptera. For example, helix 18 of Zoraptera could
not be aligned to the other insect sequences, although
the region was otherwise very conservative throughout
insects. Analysis of secondary structure indicated that
the shape of helix 18 in Zoraptera differed from that of
other insects by having a longer stem and a very small
hairpin loop (Fig. 3). Modifications of secondary struc-
tures were also identified in helix 12. Although the region
was well aligned throughout insects including Zoraptera,
the estimated secondary structures of helix 12 in Zorap-
tera were greatly modified from other insects (Fig. 2).
Rather long insertions, ranging about 90–160 bp, were
observed between helices E23-2 and E23-8 in Zoraptera.
Such insertions were not observed in any other polyne-
opteran (orthopteroid insects: i.e., Neoptera excluding
Paraneoptera and Holometabola) nor holometabolous
orders, but were observed in some species of Paraneop-
tera (e.g., *Pedicinus* sp. had an 800 bp insertion). A very
large insertion at E23 has also been reported for holome-
tabolous Strepsiptera (twisted wings) by Gillespie et al.
(in press), but sequences from this order are not analyzed
in our study. Finally, Zoraptera was on a very long
branch in the ML tree (Fig. 1), indicating an accelerated
substitution rate of the 18S of Zoraptera.

506 3.2. Phylogenetic analyses

The trees obtained from our data set (Fig. 1) were
generally in agreement with the tree obtained by Kjer

ZORAPTERA

Fig. 1. Tree obtained by ML analysis of the 18S rDNA data ($-\ln L = 32543.23325$). The tree is rooted on *Limulus polyphemus* (horseshoe crab). Branch lengths are proportional to ML estimated branch lengths. The numbers associated with the nodes are bootstrap values or posterior probabilities obtained by ML/Bayes analyses. Bootstrap values higher than 50% and/or Bayesian posterior probabilities higher than 90% are indicated. Monophyly of Insecta, Neoptera and Holometabola are constrained for ML bootstrapping (indicated by CON). No constraints are given for tree searches.

Fig. 2. Estimated secondary structure of helix 12 for selected samples. Stems are shaded, and complementary regions are connected by solid line.

Fig. 3. Estimated secondary structure of helix 18 for selected samples. Stems are shaded. Secondary structure model follows Gutell (1993, 1994) and Gillespie et al. (in press). Their model employs some non-canonical pairs (such as G-A pairs), but Kjer (2004) did not follow Gutell model and employ canonical pairs only (A-T, G-C).

Fig. 4. Sequences near helix 36 for selected samples. Based on the character states at highlighted positions, Kjer (2004) concluded that the 18S sequence of *Z. snyderi* was an acarine contaminant. When this and neighboring regions of *Z. snyderi* were subjected to a BLAST search, the top three sequences (harvestmen spiders and an annelid) were identified as the closest matches.

(2004), which showed high congruence with the traditional classification of insect orders. However, four results from our analyses differed from Kjer's (2004) tree (outlined below).

(1) Monophyly of Pterygota (winged insects) was recovered by our ML analyses; in contrast, the tree obtained by Kjer (2004) placed wingless apterygote *Tricholepidon gertschi* (Thysanura: silver fish) as a sister taxon of Odonata (dragon flies). The ML bootstrap value for monophyly of Pterygota was low (<50%), and the present Bayesian consensus placed *T. gertschi* as a sister of Odonata with 54% posterior probability.

(2) Monophyly of Paraneoptera was not recovered by either ML or Bayesian analyses. Monophyly of Paraneoptera was recovered by Kjer (2004), but with low posterior probabilities (64–87%). The present results indicated a sister group relationship between Holometabola and Hemiptera + Thysanoptera. The posterior probability for non-monophyly of Paraneoptera was 92%, but the ML bootstrap value was lower than 50%.

(3) A close relationships between Holometabola and Paraneoptera and between Orthoptera (grasshoppers) and Paraneoptera + Holometabola were recovered by the present analyses as well as by Kjer (2004). Although the posterior probabilities

for these relationships obtained by Kjer (2004) were low (48–63% for Holometabola + Paraneoptera and 38–76% for Holometabola + Paraneoptera + Orthoptera), results from the present Bayesian analysis provided relatively strong support for these clades (97 and 100% posterior probability, respectively). In contrast, ML bootstrap support for the clades was lower than 50%.

(4) Two orders, Zoraptera and Thysanoptera (thrips), were not analyzed by Kjer (2004) but are newly added by our analyses. Zoraptera was always placed as the sister taxon of Dictyoptera (54% bootstrap and 99% posterior probability). Thysanoptera was placed as a sister taxon of Hemiptera (bugs, aphids, cicadas, etc.) which suggested monophyletic Condylgnatha (Yoshizawa and Saigusa, 2001). However, support for the clade was low (<50% ML bootstrap and 72% Bayesian posterior probability).

300

301 4. Discussion

302 4.1. 18S of Zoraptera

Kjer (2004) suggested that the 18S sequence of *Zorotypus snyderi* analyzed by Wheeler et al. (2001) might be an acarine contaminant, at least in part. However, new obtained 18S sequences for four species of Zoraptera show a close match with the 18S of *Z. snyderi*. MP and NJ analyses based on the smaller fragment available for previously published sequences indicate that the 18S sequences from five zorapteran species compose a monophyletic group (100% bootstrap supports), and they are divided into two well supported clades: Oriental (*Z. sp.MY1*, *Z. sp.MY2* and *Z. sp.VN*: 100% support) and North American species (*Z. snyderi* and *Z. hubbardi*: 100% support). In addition to the close match of the nucleotide sequences, all zorapteran sequences including *Z. snyderi* have indels at the same position of helix 18 (Fig. 3).

It is very unlikely that extractions from five species extracted in three different laboratories contain the same contaminant (*Z. snyderi* at lab of Wheeler and colleagues, USA, *Z. hubbardi* at Illinois Natural History Survey, USA and *Z. sp.MY1*, *Z. sp.MY2* and *Z. sp.VN* at Hokkaido University, Japan). In addition, one of three base positions (Fig. 4), which was thought to be evidence of acarine contaminant by Kjer (2004), is variable within Zoraptera, and the nucleotide in that position in some zorapterans agrees with that of the other insects. The phylogenetic trees based on these 18S sequences placed Zoraptera within Neoptera, which is reasonable in light of morphological evidence (e.g., Kristensen, 1975, 1981). The intra-ordinal relationships

of Zoraptera based on these sequences are also very reasonable, agreeing with morphological observations (the three Oriental species have an ovoid coil on the phallosome, which is lacking in the New World species; Engel and Grimaldi, 2000; New, 1978, 2000; Yoshizawa, pers. obs.). Therefore, we conclude that the zorapteran 18S sequence analyzed by Wheeler et al. (2001) is not a contaminant, and that the fragment near helix 36 in the 18S of Zoraptera is highly variable, which causes convergence with the acarine sequences, a possibility also considered by Kjer (2004).

In addition to the region near helix 36 (Fig. 4), the 18S of Zoraptera shows other unique characteristics, including an accelerated substitution rate (Fig. 1), modification of secondary structures (Figs. 2 and 3), and long insertions. These phenomena are uniquely and uniformly observed in all the 18S sequences of Zoraptera, and thus a correlated origin of these phenomena is likely. A correlation of unique molecular evolutionary trends is detected in the mitochondrial genomes of lice, which includes accelerated substitution rates, modifications of rRNA secondary structures, long insertions/deletions, increased GC contents, and genome rearrangements (Johnson et al., 2003; Page et al., 2002; Shao et al., 2003; Yoshizawa and Johnson, 2003). However, the forces that cause these trends in molecular evolution is less understood.

4.2. Phylogenetic analyses

Kjer (2004) showed that 18S sequences aligned according to a secondary structure model provide reasonable results for the phylogeny of insects. The results obtained by the present analyses are basically in agreement with Kjer (2004), so here we focus only on some novel findings or incongruence between our trees and Kjer (2004).

In the present analyses, a sister group relationship between Zoraptera and Dictyoptera is recovered by both ML and Bayesian methods. Zoraptera + Dictyoptera received 99% posterior probability. Morphologically, Zoraptera and Dictyoptera share a reduced pterothoracic phragmata and dorsolongitudinal muscles (Boudreaux, 1979) and a derived wing venation (Kukalová-Peck and Peck, 1993). Therefore, there is also some morphological support for this placement of Zoraptera.

Additional sequences of 18S for Thysanoptera are newly available for our broader study (Johnson et al., 2004). Morphologically, a close relationship between Thysanoptera and Hemiptera has been suggested (Kristensen, 1975, 1981; Yoshizawa and Saigusa, 2001, 2003). In contrast, the combined data set produced by direct optimization (Wheeler et al., 2001) recovered a sister relationship between Thysanoptera and Psocodea. Wheeler et al. (2001) mentioned that

there were no morphological apomorphies supporting Thysanoptera + Psocodea, but that the result had strong molecular support (10 transitions and 4 transversions in 18S). The 18S alignment analyzed here recovers a sister relationship between Thysanoptera and Hemiptera, whereas a sister relationship between Thysanoptera and Psocodea receives only 1% bootstrap support and 0.8% posterior probability. In addition, a sister relationship between Thysanoptera and Psocodea is not recovered by MP analysis. Therefore, it is evident that 18S has little phylogenetic signal supporting Thysanoptera + Psocodea, and the present results are congruent with morphological data.

While several of our results agree with morphological characters, incongruence between the present results and morphological characters occurs with respect to Paraneoptera. Monophyly of Paraneoptera is strongly supported by morphological autapomorphies such as modified mouth parts, derived wing base structures, a single abdominal ganglion, and the absence of cerci (Kristensen, 1975, 1981; Yoshizawa and Saigusa, 2001, 2003). However, the present results suggest a paraphyletic grade of Paraneoptera (i.e., Hemiptera + Thysanoptera sister to Holometabola) with posterior probability 92%. ML bootstrap support for the non-monophyly of Paraneoptera is very low (<50%). There is no morphological evidence published supporting Holometabola + (Hemiptera + Thysanoptera). Therefore, further evidence is needed to resolve whether Paraneoptera is monophyletic.

A close relationship between Orthoptera and Paraneoptera + Holometabola was recovered by our analyses as well as by Kjer (2004). However, this relationship is also unexpected from the morphological point of view. Although the support for this relationship obtained by Kjer (2004) was very weak (38–76% posterior probability), the present Bayesian analysis provides strong support for the clade (100% posterior probability). However, ML bootstrap support for the clade is very low (<50%). As far as we are aware, no one has ever suggested a close relationship between Orthoptera and Paraneoptera + Holometabola based on morphology.

Remarkable differences between bootstrap support and posterior probability are frequent at deep and short nodes as mentioned above. Recent analyses of empirical and simulated data sets revealed that posterior probability is excessively high and can provide erroneous conclusions more often (Cummings et al., 2003; Erixon et al., 2003; Simmons et al., 2004). Inflation of posterior probability is known to be especially frequent for short nodes (Alfaro et al., 2003; Lewis et al., in press). Nodes supporting Condylgnatha + Holometabola and Orthoptera + Paraneoptera + Holometabola are very short, and bootstrap supports for these clades are very low (<50%) compared to high posterior probabilities (>92%). Therefore, further morphological and molecu-

lar data sets are required to test these clades. ML bootstrap support for Zoraptera + Dictyoptera is also relatively low (54%) in comparison to a high posterior probability (99%). However, the basal node supporting this sister relationship is not short, and is almost as long as, or even longer than the basal nodes of very well supported groups, such as Archaeognatha (91% bs, 100% pp), Neoptera (constrained for ML bootstrap, 100% pp), Holometabola (constrained for ML bootstrap, 100% pp) and Mecoptera + Siphonaptera (96% bs and 99% pp). Therefore, a different explanation may be required for the low ML bootstrap support for Zoraptera + Dictyoptera clade in comparison to a high posterior probability. One possible explanation is that NNI branch swapping was used for ML bootstrapping. As mentioned previously, a neighbor-joining starting tree for ML estimation was problematic especially for the placement of Zoraptera, and NNI branch swapping was not sufficient to escape from long-branch attraction problems. Because more thorough searches should more readily converge on the most likely tree, bootstrap support for some clades should be improved by the use of TRB branch swapping for ML bootstrapping. However, TBR is computationally infeasible for the present data set and processor power available.

The present result is based only on a single gene and thus represents gene tree. In addition, unusual characteristics of the zorapteran 18S sequences might be problematic for resolving the placement of this order. Thus, although the present result provides a very reasonable phylogenetic tree for insect orders and relatively strong support for a Zoraptera + Dictyoptera clade, it will be important to test the present results with additional morphological and molecular data.

4.3. Concluding comment

The 18S sequences aligned according to secondary structure model provide very reasonable insect phylogeny. It is especially notable that both Kjer (2004) and present analyses provided well resolved and very reasonable phylogenetic hypotheses among deep hexapod lineage (e.g., monophyly of Hexapoda), even though previous analyses of molecular data failed to provide a reasonable result (Bitsch et al., 2004). Samples of Zoraptera are newly analyzed, and a close relationship between Zoraptera and Dictyoptera is recovered by molecular data for the first time. In addition, examination of “molecular morphology” indicate unique evolutionary trends in the zorapteran 18S. Such interesting findings have also been provided for some insect ribosomal RNA by examinations of secondary structure (Ouvrard et al., 2000; Page et al., 2002; Yoshizawa and Johnson, 2003). Secondary structure-based manual alignment is valuable for both phylogenetic analyses and examinations of molecular morphology (Gillespie et al., in press; Kjer, 2004).

497 **5. Uncited reference**

498 Brodsky et al. (1992).

499 **Acknowledgments**

500 We thank the following individuals: A B. Idris, H.
501 Kojima, D. Morris, S. Nomura, R. Rakitov, and

N. Takahashi for providing specimens; G. Ito for infor- 502
mation on orthopteroid phylogeny; J. Gillespie for allow- 503
ing to access his unpublished paper; K. Kjer and an 504
anonymous referee for valuable comments. KY thanks 505
Y. Saito and M. Ôhara for allowing the use of molecular 506
facilities and laboratories. KY's collecting trip to Malay- 507
sia was supported by JSPS grant (142550161 to O. Yata). 508
This study was partly supported by JSPS grant (1577052) 509
to K.Y. and NSF grant (DEB-0107891) to K.P.J. 510

Appendix A.

Additional taxa included in the present study. Species not listed here are from Kjer (2004). *Ectopsocus perkinsi* [*Ectopsocus* sp. of Johnson et al. (2004): re-identification based on DNA voucher by KY] and *Pedicinus* sp. are replacement samples of Ectopsocidae Gen. sp. and *Pediculus humanus* of Kjer (2004), respectively. *Zorotypus snyderi* and *Drosophila melanogaster* were included only for the analysis of 18S secondary structure and/or preliminary phylogenetic inference.

Order	Family	Species	GenBank accession number
Zoraptera	Zorotypidae	<i>Zorotypus snyderi</i>	AF372432
Zoraptera	Zorotypidae	<i>Zorotypus hubbardi</i>	DQ013288
Zoraptera	Zorotypidae	<i>Zorotypus</i> sp.MY1	DQ013289
Zoraptera	Zorotypidae	<i>Zorotypus</i> sp.MY2	DQ013291
Zoraptera	Zorotypidae	<i>Zorotypus</i> sp.VN	DQ013290
Phasmatodea	Timematidae	<i>Timema knulli</i>	AF423806
Blattodea	Polyphagidae	<i>Polyphaga aegyptiaca</i>	AF220575
Blattodea	Blattellidae	<i>Blattella germanica</i>	AF220573
Blattodea	Blattellidae	<i>Supella longipalpa</i>	AY491149
Blattodea	Blaberidae	<i>Gromphadrhina laevigata</i>	AY210820
Embioptera	Notoligotomidae	<i>Notoligotoma</i> sp.	AY338693
Psocodea	Prionoglarididae	<i>Prionoglaris</i> sp.	AY630456
Psocodea	Lepidopsocidae	<i>Echmepteryx madagascariensis</i>	AY630447
Psocodea	Troctopsocidae	<i>Selenopsocus</i> sp.	AY630457
Psocodea	Archipsocidae	<i>Archipsocus</i> sp.	AY630479
Psocodea	Epipsocidae	<i>Epipsocus</i> sp.	AY630539
Psocodea	Ectopsocidae	<i>Ectopsocus perkinsi</i>	AY630510
Psocodea	Mesopsocidae	<i>Mesopsocus hongkongensis</i>	AY630516
Psocodea	Pachytroctidae	<i>Tapinella</i> sp.	AY630466
Psocodea	Pedicinidae	<i>Pedicinus</i> sp.	AY077777
Thysanoptera	Thripidae	<i>Frankliniella</i> sp.	AY630445
Thysanoptera	Phlaeothripidae	<i>Idolothrips spectrum</i>	AY630443
Diptera	Drosophilidae	<i>Drosophila melanogaster</i>	M21017

References

- Alfaro, M.E., Zoller, S., Lutzoni, F., 2003. Bayes or bootstrap? A simulation study comparing the performance of Bayesian Markov chain Monte Carlo sampling and bootstrapping in assessing phylogenetic confidence. *Mol. Biol. Evol.* 20, 255–266.
- Barker, S.C., Whiting, M., Johnson, K.P., Murrell, A., 2003. Phylogeny of the lice (Insecta, Phthiraptera) inferred from small subunit rRNA. *Zool. Scr.* 32, 407–414.
- Bitsch, J., Bitsch, C., Bourgoin, T., d'Haese, C., 2004. The phylogenetic position of early hexapod lineages: morphological data contradict molecular data. *Syst. Entomol.* 29, 433–440.
- Boudreaux, H.B., 1979. *Arthropod Phylogeny, with Special Reference to Insects*. Wiley, New York.
- Brodsky, L.I., Vasiliev, A.V., Kalaidzidis, Y.L., Osipov, Y.S., Tatuzov, R.L., Feranchuk, S.I., 1992. GeneBee: the program package for biopolymer structure analysis. *Dimacs* 8, 127–139.
- Brodsky, L.I., Ivanov, V.V., Kalaidzidis, Y.L., Leontovich, A.M., Nikolaev, V.K., Feranchuk, S.I., Drachev, V.A., 1995. GeneBee-NET: Internet-based server for analyzing biopolymers structure. *Biochemistry* 60, 923–928.
- Campbell, B.C., Steffen-Campbell, J.D., Sorensen, J.T., Gill, R.J., 1995. Paraphyly of Homoptera and Auchenorrhyncha inferred from 18S rDNA nucleotide sequences. *Syst. Entomol.* 20, 175–194.

- 511 Chao, R.F., Chen, C.S., 2000. *Formosozoros newi*, a new genus and
512 species of Zoraptera (Insecta) from Taiwan. Pan-Pacific Entomol.
513 76, 24–27.
- 514 Cummings, M.P., Handley, S.A., Myers, D.S., Reed, D.L., Rokas, A.,
515 Winka, K., 2003. Comparing bootstrap and posterior probability
516 values in the four-taxon case. Syst. Biol. 52, 187–477.
- 517 Engel, M.S., Grimaldi, D.A., 2000. A winged *Zorotypus* in Miocene
518 amber from the Dominican Republic (Zoraptera: Zorotypidae),
519 with discussion on relationships of and within the order. Acta
520 Geol. Hispanica 35, 149–164.
- 521 Engel, M.S., Grimaldi, D.A., 2002. The first Mesozoic Zoraptera
522 (Insecta). Am. Mus. Novitates 3362, 1–20.
- 523 Erixon, P., Svennblom, B., Britton, T., Oxelman, B., 2003. Reliability
524 of Bayesian posterior probabilities and bootstrap frequencies in
525 phylogenetics. Syst. Biol. 52, 665–673.
- 526 Felsenstein, J., 1978. Cases in which parsimony and compatibility
527 methods will be positively misleading. Syst. Zool. 27, 401–410.
- 528 Gillespie, J.J., McKenna, C.H., Gutell, R.R., Johnston, J.S., Kath-
529 irithamby, J., Cognato, A.I., in press. Assessing the odd secondary
530 structural properties of nuclear small subunit ribosomal RNA
531 sequences (18S) of the twisted-wing parasites (Insecta: Strepsiptera).
532 Insect. Mol. Biol.
- 533 Hennig, W., 1981. Insect Phylogeny (ed. and translated by A. C. Pont).
534 Wiley, New York.
- 535 Huelsenbeck, J.P., 1997. Is the Felsenstein zone a fly trap? Syst. Biol.
536 46, 69–74.
- 537 Huelsenbeck, J.P., Hillis, D.M., 1993. Success of phylogenetic methods
538 in the four-taxon case. Syst. Biol. 42, 247–265.
- 539 Huelsenbeck, J.P., Ronquist, F., 2001. MrBayes: Bayesian inference on
540 a phylogeny. Bioinformatics 17, 754–755.
- 541 Johnson, K.P., Cruickshank, R.H., Adams, R.J., Smith, V.S., Page,
542 R.D.M., Clayton, D.H., 2003. Dramatically elevated rate of
543 mitochondrial substitution in lice (Insecta: Phthiraptera). Mol.
544 Phylogenet. Evol. 26, 231–242.
- 545 Johnson, K.P., Yoshizawa, K., Smith, V.S., 2004. Multiple origins of
546 parasitism in lice. Proc. R. Soc. Lond. B 271, 1771–1776.
- 547 Kjer, K.M., 1995. Use of rRNA secondary structure in phylogenetic
548 studies to identify homologous positions: an example of alignment
549 and data presentation from the frogs. Mol. Biol. Evol. 4, 314–330.
- 550 Kjer, K.M., 2004. Aligned 18S and insect phylogeny. Syst. Biol. 53,
551 506–514.
- 552 Kristensen, N.P., 1975. The phylogeny of hexapod orders. A critical
553 review of recent accounts. Zeitschrift für zoologische Systematik
554 und Evolutionsforschung 13, 1–44.
- 555 Kristensen, N.P., 1981. Phylogeny of insect orders. Ann. Rev.
556 Entomol. 26, 135–157.
- 557 Kukalová-Peck, J., Peck, S.B., 1993. Zoraptera wing structures:
558 evidence for new genera and relationship with the blattoid orders
559 (Insecta: Blattellodea). Syst. Entomol. 18, 333–350.
- 560 Lewis, P.O., Holder, M.T., Holsinger, K.E., in press. Polytomies and
561 Bayesian phylogenetic inference. Syst. Biol.
- 562 Minet, J., Bourgoin, T., 1986. Phylogénie et classification des
563 Hexapodes (Arthropoda). Cah. Liaison, O.P.I.E. 20 (4), 23–28.
- New, T.R., 1978. Notes on Neotropical Zoraptera, with descriptions of
two new species. Syst. Entomol. 3, 361–370.
- New, T.R., 2000. Zoraptera (Insecta) in east Malaysia: notes on
Zorotypus caudelli Karny. Orient. Insects 34, 7782.
- Ouvrard, D., Campbell, B.C., Bourgoin, T., Chan, K.L., 2000. 18S
rRNA secondary structure and phylogenetic position of Pelori-
diidae (Insecta, Hemiptera). Mol. Phylogenet. Evol. 16, 403–
417.
- Page, R.D.M., Cruickshank, R., Johnson, K.P., 2002. Louse (Insecta:
Phthiraptera) mitochondrial 12S rDNA secondary structure is
highly variable. Insect. Mol. Biol. 11, 361–369.
- Posada, D., Crandall, K.A., 1998. MODELTEST: testing the model of
DNA substitution. Bioinformatics 14, 817–818.
- Rambaut, A., Drummond, A., 2004. TRACER v1.2. Distributed by
authors from URL at <http://evolve.zoo.ox.ac.uk/beer/>.
- Shao, R., Dowton, M., Murrell, A., Barker, S.C., 2003. Rates of gene
rearrangement and nucleotide substitution are correlated in the
mitochondrial genomes of insects. Mol. Biol. Evol. 20, 1612–1619.
- Simmons, M.P., 2004. Independence of alignment and tree search.
Mol. Phylogenet. Evol. 31, 874–879.
- Simmons, M.P., Pickett, K.M., Miya, M., 2004. How meaningful are
Bayesian support values? Mol. Biol. Evol. 21, 188–199.
- Smithers, C.N., 1991. Zoraptera. In: Naumann et al. (Eds.), Insect of
Australia. Cornell University Press, New York, pp. 410–411.
- Swofford, D.L., 2002. PAUP*: phylogenetic analysis using parsimony
(*and other methods), version 4 beta. Sinauer, MA.
- Vawter, L., 1991. Evolution of blattoid insects and of the small subunit
ribosomal RNA gene. Unpublished PhD Thesis, University of
Michigan.
- Wheeler, W.C., Whiting, M., Wheeler, Q.D., Carpenter, J.M., 2001.
The phylogeny of the extant hexapod orders. Cladistics 17, 113–
169.
- Whiting, M.F., Carpenter, J.C., Wheeler, Q.D., Wheeler, W.C., 1997.
The Strepsiptera problem: phylogeny of the holometabolous insect
orders inferred from 18S and 28S ribosomal DNA sequences and
morphology. Syst. Biol. 46, 1–68.
- Wuyts, J., De Rijk, P., Van de Peer, Y., Pison, G., Rousseeuw, P., De
Wachter, R., 2000. Comparative analysis of more than 3000
sequences reveals the existence of two pseudoknots in area V4 of
eukaryotic small subunit ribosomal RNA. Nucleic Acids Res. 28,
4698–4708.
- Yoshizawa, K., Johnson, K.P., 2003. Phylogenetic position of Phthi-
raptera (Insecta: Paraneoptera) and elevated rate of evolution in
mitochondrial 12S and 16S rDNA. Mol. Phylogenet. Evol. 29, 102–
114.
- Yoshizawa, K., Saigusa, T., 2001. Phylogenetic analysis of paraneopt-
eran orders (Insecta: Neoptera) based on forewing base structure,
with comments on monophyly of Auchenorrhyncha (Hemiptera).
Syst. Entomol. 26, 1–13.
- Yoshizawa, K., Saigusa, T., 2003. Reinterpretations of clypeus and
maxilla in Psocoptera, and their significance in phylogeny of
Paraneoptera (Insecta: Neoptera). Acta Zool. (Stockholm) 84, 33–
40.