



Title	Discharge use of angiotensin receptor blockers provides comparable effects with angiotensin-converting enzyme inhibitors on outcomes in patients hospitalized for heart failure
Author(s)	Tsuchihashi-Makaya, Miyuki; Furumoto, Tomoo; Kinugawa, Shintaro; Hamaguchi, Sanae; Goto, Kazutomo; Goto, Daisuke; Yamada, Satoshi; Yokoshiki, Hisashi; Takeshita, Akira; Tsutsui, Hiroyuki
Citation	Hypertension Research, 33(3), 197-202 <a href="https://doi.org/10.1038/hr.2009.199">https://doi.org/10.1038/hr.2009.199</a>
Issue Date	2010-03
Doc URL	<a href="http://hdl.handle.net/2115/43817">http://hdl.handle.net/2115/43817</a>
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	HR33-3_197-202.pdf



[Instructions for use](#)

**Discharge use of angiotensin receptor blockers provides comparable effects with angiotensin-converting enzyme inhibitors on outcomes in patients hospitalized for heart failure**

Miyuki Tsuchihashi-Makaya<sup>1</sup>, Tomoo Furumoto<sup>1</sup>, Shintaro Kinugawa<sup>1</sup>, Sanae Hamaguchi<sup>1</sup>, Kazutomo Goto<sup>1</sup>, Daisuke Goto<sup>1</sup>, Satoshi Yamada<sup>1</sup>, Hisashi Yokoshiki<sup>1</sup>, Akira Takeshita<sup>2</sup>, Hiroyuki Tsutsui<sup>1</sup>, and for the JCARE-CARD Investigators

*1. Department of Cardiovascular Medicine, Hokkaido University Graduate School of Medicine, Sapporo 060-8638 (M-TM, TF, SK, SH, KG, DG, SY, HY, HT),*

*2. Futsukaichi Saiseikai Hospital, Futsukaichi 820-0018 (AT), Japan*

Dr. Akira Takeshita deceased on March 15 2009.

Running title: ARB and heart failure

Word count of abstract: 238 words

Total word count: 4590 words

Total number of tables and figures: 4 tables, 1 figure, and 3 supplemental tables

Address for correspondence:

Hiroyuki Tsutsui, M.D., Ph.D.

Department of Cardiovascular Medicine,

Hokkaido University Graduate School of Medicine,

Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

Phone: +81-11-706-6970, FAX: +81-11-706-7874

e-mail: [htsutsui@med.hokudai.ac.jp](mailto:htsutsui@med.hokudai.ac.jp)

This work is supported by the grants from the Japanese the Ministry of Health, Labour and Welfare, the Japan Heart Foundation, and Japan Arteriosclerosis Prevention Fund.

## **Abstract**

Large-scale, placebo-controlled, randomized clinical trials have demonstrated that angiotensin-converting enzyme (ACE) inhibitors or angiotensin receptor blockers (ARBs) reduce mortality and hospitalization in patients with heart failure (HF) caused by left ventricular systolic dysfunction (LVSD). However, it is unknown whether ACE inhibitors and ARBs have similar effects on the long-term outcomes in HF patients encountered in routine clinical practice. The Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD) enrolled HF patients hospitalized with worsening symptoms and they were followed during an average of 2.2 years. The outcome data were compared in patients with LVSD by echocardiography (ejection fraction <40%) according to the predischARGE use of ACE inhibitors (n=356) or ARBs (n=372). The clinical characteristics were similar between patients with ACE inhibitor and ARB use except for higher prevalence of hypertensive etiology and diabetes mellitus. There was no significant difference between ACE inhibitor and ARB use in all cause death (adjusted hazard ratio [HR] 0.958, 95%CI 0.601-1.527,  $P=0.858$ ) and rehospitalization (adjusted HR0.964, 95% CI 0.683-1.362,  $P=0.836$ ). The effects of ACE inhibitor and ARB use on the outcomes were generally consistent across all clinically relevant subgroups examined, including age, sex, etiology, EF, hypertension, diabetes mellitus, and  $\beta$ -blocker use. Discharge use of ARBs provided comparable effects with ACE inhibitors on outcomes in patients hospitalized for HF. These findings provide further support for guideline recommendations that ARBs can be used in patients with HF and LVSD as an alternative of ACE inhibitors.

**Key Words:** ACE inhibitor; ARB; heart failure; outcome; survival

## **INTRODUCTION**

Heart failure (HF) is a leading cause of morbidity and mortality in industrialized countries and is also a growing public health problem, mainly because of aging of the population and the increase in the prevalence of HF in the elderly. Large-scale, placebo-controlled, randomized clinical trials such as CONSENSUS and SOLVD have demonstrated that angiotensin-converting enzyme (ACE) inhibitors significantly improved clinical outcomes in patients with chronic HF with left ventricular systolic dysfunction (LVSD).<sup>1, 2)</sup>

Angiotensin receptor blockers (ARBs) can directly block angiotensin II at the AT<sub>1</sub> receptor with no accumulation of bradykinin. They should provide similar benefits to ACE inhibitors in blocking the harmful effects of angiotensin II with fewer side effects. ARBs are highly effective in reducing blood pressure and preventing cardiovascular events in patients with hypertension. They are generally considered to be appropriate specifically in hypertensive patients with cardiovascular diseases including HF. In the ELITE II study, losartan was not superior to captopril in improving survival in elderly HF patients, but was significantly better tolerated.<sup>3)</sup> However, ELITE II study was designed as a superiority trial and not to address equivalence between losartan and captopril and thus could not provide any direct information in the difference in efficacy between these two drugs. In the CHARM-alternative study, candesartan reduced the risk of cardiovascular death or hospital admission for HF in patients with LVSD not currently treated with an ACE inhibitor because of previous intolerance.<sup>4)</sup> Based on these results, recent treatment guidelines for HF in the United States and Europe recommended that ACE inhibitors should remain the treatment of the first choice in HF and ARB can be a useful alternative agent in a limited group of patients in whom ACE inhibitors are not tolerated.<sup>5, 6)</sup> Therefore, it still remains to be uncertain whether ARBs are a fully effective substitute for ACE inhibitors in HF. Moreover, it has not been determined whether ACE inhibitors and ARBs have similar effects on the long-term survival in HF patients encountered in routine

clinical practice. It should be confirmed in another setting whether there is a benefit associated with ARB use in an unselected population of patients with HF.

The Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD) is designed to study prospectively the characteristics, treatments, and outcomes in a broad sample of patients hospitalized with HF at teaching hospitals in Japan between January 2004 and June 2005 and the outcomes including death and hospitalization due to HF were followed through 2007.<sup>7)</sup> The JCARE-CARD program enrolled 2675 patients admitted with HF in a web-based registry with an average follow-up of 2.2 years.

The present study was aimed to investigate the hypothesis that the long-term outcomes including all-cause mortality and hospitalization due to the worsening of HF were comparable between ACE inhibitor and ARB use at discharge among a nonselected population of patients hospitalized for HF and LVSD.

## **METHODS**

### **Patients**

The details of the JCARE-CARD program have been described previously.<sup>7-11)</sup> Briefly, eligible patients were those hospitalized with worsening HF as the primary cause of admission. JCARE-CARD enrolled a total of 2675 patients hospitalized for HF at 164 participating hospitals. The diagnosis of HF was established by the simultaneous presence of at least two major criteria or one major criterion in conjunction with two minor criteria by use of the Framingham criteria. LV ejection fraction (LVEF) was assessed in 1692 patients, and 728 were determined to have LVSD who had documented LVEF <40% by echocardiography and without valvular etiology. Mean postdischarge follow-up within this group was 801±300 days (2.2±0.8 years).

### **Data collection and processing**

For each patient, baseline data recorded on the form included (1) demography; (2) causes of HF; (3) precipitating causes; (4) comorbidities; (5) complications; (6) clinical status; (7) electrocardiographic and echocardiographic findings; (8) plasma BNP; and (9) treatments including discharge medications.

The status of all patients was surveyed after discharge and the following information was obtained; (1) survival, (2) cause of death, and (3) the hospital readmission due to an exacerbation of HF that required more than continuation of their usual therapy on prior admission.

### **Statistical analysis**

Patient characteristics and treatments were compared using Pearson  $\chi^2$  test for categorical variables and unpaired t-test for continuous variables. Only patients

who survived the initial hospitalization were included in the follow-up analysis. The relationship between medication use at discharge and outcomes was evaluated among patients with multivariable adjustment. Baseline clinical variables and treatment factors including other HF medications as shown in Table 1 and 2 were used in developing the 2 postdischarge Cox proportional hazards models. A *P* value of .05 was used for criteria for variables to stay in the model. SPSS version 14.0 (SAS Institute, Cary, NC) was used for all statistical analyses.

## **RESULTS**

### **Patient characteristics**

**Table 1** provides a comparison of clinical characteristics for the entire cohort (n=728) and according to the treatment groups; ACE inhibitor (n=356) and ARB (n=372) use at the time of hospital discharge. The mean age was 66±14 years and 73% were male. HF etiology was ischemic in 39%, cardiomyopathic in 39%, and hypertensive in 21% of the patients. The mean LVEF was 27%. The majority of patients taking ACE inhibitors had enalapril (66%) and some had other ACE inhibitors including temocapril (10%), imidapril (7%), and others. ARBs included candesartan (38%), valsartan (30%), losartan (22%), and others.

The characteristics were similar between patients with ACE inhibitor and ARB use except for higher prevalence of hypertensive etiology and diabetes mellitus in ARB use (**Table 1**). Serum creatinine, body mass index (BMI), and systolic blood pressure were significantly higher in patients with ARB. NYHA functional class was lower in ARB group. However, echocardiographic findings including LV diameters and EF as well as plasma BNP levels did not differ between ACE inhibitor and ARB. Importantly, concurrent cardiovascular medications other than ACE inhibitors or ARBs were similar between groups except for higher use of warfarin in ACE inhibitor group (**Table 2**). Clinical characteristics were comparable among different ARB subgroups except for higher prevalence of hypertension, high blood pressure values, and greater wall thickness by echocardiography in patients treated with valsartan or other ARBs (**Supplemental Table 1**). Medication use was also comparable among different ARB subgroups except for higher use of calcium channel blocker in patients with valsartan or others, which might be due to higher prevalence of hypertension in these groups (**Supplemental Table 2**).

## Postdischarge long-term outcomes

The long-term follow-up data could be obtained in 652 (322 and 330 for ACE inhibitors and ARBs, respectively) out of 728 registered patients (90.0%). Mean post-discharge follow-up was  $801 \pm 300$  days ( $2.2 \pm 0.8$  years);  $803 \pm 281$  for ACE inhibitors and  $813 \pm 286$  days for ARBs ( $P=0.629$ ).

In the overall cohort of patients with HF and LVSD, mortality at 2.4 year after hospital discharge was 15.6%. Rehospitalization during the same follow-up period was 30.7%. In the group of patients discharged on ARBs, there were 51 deaths from any cause (15.5%) compared with 51 (15.8%) in ACE inhibitors ( $P=0.807$ ) (**Table 3**). The rate of cardiac death was also comparable between ACE inhibitor and ARB groups (10.2% vs 11.2%;  $P=0.775$ ). The rate of rehospitalization due to the worsening of HF did not differ between groups (31.4% vs 30.0%;  $P=0.612$ ). These postdischarge outcomes were comparable among different ARB subgroups (**Supplemental Table 3**).

## Multivariable analysis

The effect of ACE inhibitor and ARB use on long-term (2.4 years) outcomes was tested in risk-adjusted models. After adjustment for multiple variables predictive of postdischarge mortality, there was no significant difference in a Kaplan-Meier plot for all-cause mortality after hospital discharge between ACE inhibitor and ARB (adjusted hazard ratio [HR] 0.958, 95% CI 0.601-1.527,  $P=0.858$ ) (**Table 3** and **Figure 1**). For the combined endpoint of all cause mortality or hospitalization, ARB use was also associated with comparable risk (adjusted HR 0.951, 95% CI 0.694-1.302,  $P=0.752$ ) (**Table 3** and **Figure 1**).

In the risk-adjusted model for all-cause mortality, the effects of ACE inhibitor and ARB use were generally consistent across all clinically relevant subgroups examined, including age (<65 vs  $\geq 65$  years), sex (male vs female), etiology (ischemic vs non-ischcemic), LVEF (<25 vs  $\geq 25\%$ ), systolic blood pressure

(<110 vs  $\geq$ 110mmHg), diastolic blood pressure (<65 vs  $\geq$ 65mmHg), hypertension, diabetes, and  $\beta$ -blocker use (**Table 4**).

## **DISCUSSION**

The present study demonstrated that there was no significant difference in outcomes including all-cause mortality, cardiac mortality, and hospitalization due to the worsening of HF between the discharge use of ACE inhibitors and ARBs among patients hospitalized with HF and LVSD during the long-term (2.2 year) follow-up.

Large-scale, placebo-controlled, randomized clinical trials such as CONSENSUS and SOLVD demonstrated that ACE inhibitors significantly improved clinical outcomes in patients with chronic HF with LVSD.<sup>1, 2)</sup> ARBs can offer an alternative approach to the inhibition of the renin–angiotensin system. CHARM-alternative study studied candesartan compared with placebo among patients with HF who were receiving no background ACE inhibitor treatment.<sup>4)</sup> In addition, post-hoc subgroup analysis of Val-Heft trial in a subset of 366 patients not taking ACE-inhibitor treatment also demonstrated the large reduction in mortality and morbidity with valsartan.<sup>12)</sup> Based on these results from clinical trials, current HF management guidelines in United States, Europe, and Japan have recommend that ACE inhibitors should remain the treatment of the first choice in HF with LVSD. In patients in whom ACE inhibitors are not tolerated, an ARB might be a useful alternative agent to block the renin-angiotensin-aldosterone system.<sup>5, 6)</sup>

The ELITE II study demonstrated that losartan was not superior to captopril in improving survival in elderly HF patients.<sup>3)</sup> Therefore, the present study has extended the results shown by ELITE II, which, however, was a superiority trial and was not designed to address equivalence between losartan and captopril and thus could not provide any direct information in the difference in efficacy between these two drugs. Moreover, both CHARM and Val-Heft have suggested that ARBs may similar beneficial effects on outcomes as ACE inhibitors in patients with HF.<sup>4,</sup>

<sup>12)</sup> However, these trials used placebo, not ACE inhibitors, as a comparator.

Therefore, the present study provided the first demonstration that ARBs could exert comparable effects with ACE inhibitors on outcomes in HF patients. The survival curves of ACE inhibitors and ARBS were almost completely overlapped during the average follow-up of 2.2 years (**Figure 1**). These findings should reassure clinicians that ARB is as effective as ACE inhibitors, and provides a similar opportunity to improve outcomes for patients with HF as ACE inhibitors. Moreover, our results parallel the findings of OPTIMAAL and VALIANT, which established the noninferiority of ARBs as compared to captopril in patients with LVSD or HF after acute myocardial infarction. <sup>13, 14)</sup>

Although evidence from randomized clinical trials shows a significant survival benefit for ARB therapy in systolic HF, <sup>3, 4, 15, 16)</sup> such trials are recognized as unrepresentative of the general HF population encountered in routine clinical practice. Therefore, uncertainty pertaining to the applicability of these findings to the population of patients with HF at large persists. Therefore, it is of critical importance to analyze the registry data of HF patients. For this purpose, JCARE-CARD was designed to focus on the demographic and clinical characteristics, treatment strategies, and outcomes in HF patients in Japan. <sup>7)</sup> The results from representative HF population in JCARE-CARD extended the results of ELITE II conducted in selected outpatients with systolic HF to a diverse cohort of patients with HF and confirmed the findings from large-scale clinical trials in that ARB treatment could be associated with survival benefit to the same extent as ACE inhibitors. Therefore, ARBs, one of major classes of antihypertensive drugs, are useful in reducing adverse events not only in hypertensive patients but also in those with HF.

Most patients with HF have hypertension. <sup>17)</sup> Hypertension is not only an important comorbidity of HF, but it also contributes to the pathogenesis of systolic and diastolic HF by inducing cardiac hypertrophy. In addition, hypertension is a major risk factor for ischemic heart disease and can lead to the development of HF

via impaired cardiac contractility, remodeling, and eventual systolic and diastolic dysfunction.<sup>18)</sup> Target blood pressure values in HF have not been firmly established, but, in most successful clinical trials, systolic blood pressure was lowered to the range of 110 to 130 mm Hg.<sup>19)</sup> On this basis, the American Heart Association Council for High Blood Pressure Research and the Councils on Clinical Cardiology and Epidemiology and Prevention have made the recommendation that the target blood pressure in patients with HF should be <130/80 mmHg, and also suggest that consideration should be given to lowering the BP even further to <120/80 mmHg.<sup>20)</sup> The present study demonstrated that the comparable effects of ACE inhibitors and ARBs on the long-term outcomes were consistent irrespective of the blood pressure levels (**Table 4**).

Several crucial limitations inherent in the design of the present study should be considered. First, documentation of ACE inhibitor or ARB prescription at hospital discharge might not accurately reflect the level of drug use after discharge or adherence to ACE inhibitor or ARB use over time. Second, JCARE-CARD is not a prospective randomized trial and, despite covariate adjustment, other measured and unmeasured factors might have influenced outcomes. We thus could not completely exclude other unmeasured factors that might also affect outcomes. Third, the number of the study patients was not sufficient to avoid type 2 (beta) statistical error in this study. Forth, as the dose of medication was not recorded, this study could not confirm that ACE inhibitors and ARBs exerted comparable inhibitory effects against angiotensin II in the studied patients. Finally, data were dependent on the accuracy of documentation and abstraction by individual medical centers that participated in the program. However, it was not the objective of this survey to restrict enrollment to the narrowly defined population of HF patients usually included in clinical trials but rather to include a broad range of patients reflecting the current reality of clinical practice rather than trials.

In conclusion, discharge use of ARBs provided comparable effects with ACE inhibitors on outcomes in patients hospitalized for HF in routine clinical

practice. These findings provide further support for guideline recommendations that ARBs can be used in patients with HF and LVSD as an alternative of ACE inhibitors.

## **CONFLICT OF INTEREST**

The authors declare no conflict of interest.

## **ACKNOWLEDGEMENTS**

The JCARE-CARD investigators and participating cardiologists are listed in the Appendix of our previous publication.<sup>7)</sup> This study could not have been carried out without the help, cooperation and support of the cardiologists in the survey institutions. We thank them for allowing us to obtain the data. The JCARE-CARD was supported by the Japanese Circulation Society and the Japanese Society of Heart Failure and by grants from Health Sciences Research Grants from the Japanese Ministry of Health, Labor and Welfare (Comprehensive Research on Cardiovascular Diseases), the Japan Heart Foundation, and Japan Arteriosclerosis Prevention Fund.

Supplementary information is available at Hypertension Research's website.

## References

1. Effects of enalapril on mortality in severe congestive heart failure. Results of the Cooperative North Scandinavian Enalapril Survival Study (CONSENSUS). The CONSENSUS Trial Study Group. *N Engl J Med* 1987; 316:1429-1435.
2. Effect of enalapril on survival in patients with reduced left ventricular ejection fractions and congestive heart failure. The SOLVD Investigators. *N Engl J Med* 1991; 325:293-302.
3. Pitt B, Poole-Wilson PA, Segal R, Martinez FA, Dickstein K, Camm AJ, Konstam MA, Riegger G, Klingler GH, Neaton J, Sharma D, Thiyagarajan B. Effect of losartan compared with captopril on mortality in patients with symptomatic heart failure: randomised trial--the Losartan Heart Failure Survival Study ELITE II. *Lancet* 2000; 355:1582-1587.
4. Granger CB, McMurray JJ, Yusuf S, Held P, Michelson EL, Olofsson B, Ostergren J, Pfeffer MA, Swedberg K. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function intolerant to angiotensin-converting-enzyme inhibitors: the CHARM-Alternative trial. *Lancet* 2003; 362:772-776.
5. Jessup M, Abraham WT, Casey DE, Feldman AM, Francis GS, Ganiats TG, Konstam MA, Mancini DM, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 focused update: ACCF/AHA Guidelines for the Diagnosis and Management of Heart Failure in Adults: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Practice Guidelines: developed in collaboration with the International Society for Heart and Lung Transplantation. *Circulation* 2009; 119:1977-2016.
6. Dickstein K, Cohen-Solal A, Filippatos G, McMurray JJ, Ponikowski P, Poole-Wilson PA, Stromberg A, van Veldhuisen DJ, Atar D, Hoes AW, Keren A, Mebazaa A, Nieminen M, Priori SG, Swedberg K, Vahanian A,

- Camm J, De Caterina R, Dean V, Funck-Brentano C, Hellemans I, Kristensen SD, McGregor K, Sechtem U, Silber S, Tendera M, Widimsky P, Zamorano JL. ESC Guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J* 2008; 29:2388-2442.
7. Tsutsui H, Tsuchihashi-Makaya M, Kinugawa S, Goto D, Takeshita A. Clinical characteristics and outcome of hospitalized patients with heart failure in Japan -rationale and design of Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2006; 70:1617-1623.
  8. Hamaguchi S, Tsuchihashi-Makaya M, Kinugawa S, Yokota T, Ide T, Takeshita A, Tsutsui H. Chronic kidney disease as an independent risk for long-term adverse outcomes in patients hospitalized with heart failure in Japan. report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2009; 73:1442-1447.
  9. Tsuchihashi-Makaya M, Hamaguchi S, Kinugawa S, Yokota T, Goto D, Yokoshiki H, Kato N, Takeshita A, Tsutsui H. Characteristics and outcomes of hospitalized patients with heart failure and reduced vs preserved ejection fraction - a report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2009; 73: 1893-1900.
  10. Hamaguchi S, Tsuchihashi-Makaya M, Kinugawa S, Yokota T, Takeshita A, Yokoshiki H, Tsutsui H. Anemia is an independent predictor of long-term adverse outcomes in patients hospitalized with heart failure in Japan. a report from the Japanese Cardiac Registry of Heart Failure in Cardiology (JCARE-CARD). *Circ J* 2009; 73:1901-1908.
  11. Hamaguchi S, Yokoshiki H, Kinugawa S, Tsuchihashi-Makaya M, Yokota T, Takeshita A, Tsutsui H. Effects of atrial fibrillation on long-term outcomes in patients hospitalized for heart failure in Japan. *Circ J* (in

- press).
12. Maggioni AP, Anand I, Gottlieb SO, Latini R, Tognoni G, Cohn JN. Effects of valsartan on morbidity and mortality in patients with heart failure not receiving angiotensin-converting enzyme inhibitors. *J Am Coll Cardiol* 2002; 40:1414-1421.
  13. Dickstein K, Kjekshus J. Effects of losartan and captopril on mortality and morbidity in high-risk patients after acute myocardial infarction: the OPTIMAAL randomised trial. Optimal Trial in Myocardial Infarction with Angiotensin II Antagonist Losartan. *Lancet* 2002; 360:752-760.
  14. Pfeffer MA, McMurray JJ, Velazquez EJ, Rouleau JL, Kober L, Maggioni AP, Solomon SD, Swedberg K, Van de Werf F, White H, Leimberger JD, Henis M, Edwards S, Zelenkofske S, Sellers MA, Califf RM. Valsartan, captopril, or both in myocardial infarction complicated by heart failure, left ventricular dysfunction, or both. *N Engl J Med* 2003; 349:1893-1906.
  15. Cohn JN, Tognoni G. A randomized trial of the angiotensin-receptor blocker valsartan in chronic heart failure. *N Engl J Med* 2001; 345:1667-1675.
  16. McMurray JJ, Ostergren J, Swedberg K, Granger CB, Held P, Michelson EL, Olofsson B, Yusuf S, Pfeffer MA. Effects of candesartan in patients with chronic heart failure and reduced left-ventricular systolic function taking angiotensin-converting-enzyme inhibitors: the CHARM-Added trial. *Lancet* 2003; 362:767-771.
  17. Kannel WB, Belanger AJ. Epidemiology of heart failure. *Am Heart J* 1991; 121:951-957.
  18. McKee PA, Castelli WP, McNamara PM, Kannel WB. The natural history of congestive heart failure: the Framingham study. *N Engl J Med* 1971; 285:1441-1446.
  19. Packer M, Fowler MB, Roecker EB, Coats AJ, Katus HA, Krum H, Mohacsi P, Rouleau JL, Tendera M, Staiger C, Holcslaw TL, Amann-Zalan I, DeMets DL. Effect of carvedilol on the morbidity of patients with severe

chronic heart failure: results of the carvedilol prospective randomized cumulative survival (COPERNICUS) study. *Circulation* 2002; 106:2194-2199.

20. Rosendorff C, Black HR, Cannon CP, Gersh BJ, Gore J, Izzo JL, Jr., Kaplan NM, O'Connor CM, O'Gara PT, Oparil S. Treatment of hypertension in the prevention and management of ischemic heart disease: a scientific statement from the American Heart Association Council for High Blood Pressure Research and the Councils on Clinical Cardiology and Epidemiology and Prevention. *Circulation* 2007; 115:2761-2788.

## **Figure Legends**

**Figure 1** Event-free curves from (A) all-cause death, (B) cardiac death, (C) rehospitalization due to HF, and (D) all-cause death or rehospitalization due to HF in patients using ACE inhibitors (black lines; n=322) compared with ARBs (red lines; n=330). The data were adjusted for differences in baseline variables, including age, gender, BMI, hypertension, diabetes mellitus, serum creatinine level, NYHA class at discharge, systolic blood pressure, and warfarin.

**Table 1 Patient characteristics**

	All (n=728)	ACE inhibitor (n=356)	ARB (n=372)	P value
<b>Demographic</b>				
Age, years	65.9±13.8	65.6±13.9	66.2±13.7	0.622
Older than 65 yr, %	59.2	56.5	61.8	0.141
Male, %	73.1	76.1	70.2	0.070
<b>Cause of HF, %</b>				
Ischemic	39.3	38.2	40.3	0.558
Hypertensive	21.2	16.9	25.3	0.005
Cardiomyopathic	38.7	41.9	35.8	0.091
Undetermined	13.5	15.2	11.8	0.187
<b>History</b>				
Hypertension, %	50.3	46.7	53.6	0.064
Diabetes mellitus, %	32.3	28.4	36.1	0.026
Hyperlipidemia, %	30.1	30.4	29.8	0.864
Chronic renal failure, %	7.0	5.6	8.4	0.151
Serum creatinine, mg/dL	1.25±1.20	1.14±0.65	1.35±1.54	0.041
Hyperuricemia, %	50.3	50.7	49.9	0.817
Stroke, %	13.3	14.2	12.4	0.483
Anemia, %	10.6	10.5	10.8	0.895
Hemoglobin, g/dL	13.0±2.3	12.9±2.3	13.1±2.3	0.403
COPD, %	5.7	4.8	6.6	0.310
Atrial fibrillation, %	23.6	25.9	21.4	0.148
Sustained VT/Vf, %	9.0	8.8	9.1	0.894
PCI, %	20.9	21.8	20.0	0.559
CABG, %	11.1	12.6	9.8	0.235
Smoking, %	46.8	46.0	47.6	0.665
<b>Physical findings</b>				
Body mass index, kg/m <sup>2</sup>	22.9±4.2	22.5±3.9	22.3±4.5	0.011
Heart rate, bpm	70.4±11.8	70.7±12.4	70.1±11.3	0.445

Discharge SBP, mmHg	112.5±16.7	110.0±15.4	113.9±17.8	0.018
Discharge DBP, mmHg	65.9±11.4	65.1±11.1	66.7±11.6	0.063
NYHA class at discharge				
I	36.8	30.9	42.5	
II	57.4	62.9	52.2	0.005
III	5.8	6.2	5.4	
Echocardiographic parameters				
LV EDD, mm	62.3±9.3	62.5±9.5	62.1±9.1	0.576
LV ESD, mm	53.7±9.4	53.7±9.7	53.6±9.1	0.902
LV EF, %	26.9±7.4	26.6±7.5	27.2±7.3	0.283
IVS thickness, mm	9.7±2.9	9.6±3.3	9.7±2.3	0.909
LV PW, mm	10.0±2.5	9.8±2.8	10.1±2.1	0.139
Admission plasma BNP, pg/ml	951.1±991.5	926.3±974.5	974.5±1027.4	0.540
Discharge plasma BNP, pg/ml	352.7±495.8	359.9±541.6	345.2±441.4	0.761

HF, Heart failure; COPD, chronic obstructive pulmonary disease; VT/Vf, ventricular tachycardia/fibrillation; PCI, percutaneous coronary intervention; CABG, coronary artery bypass grafting; SBP, systolic blood pressure; DBP, diastolic blood pressure; NYHA, New York Heart Association; LV, left ventricular; EDD, end-diastolic dimension; ESD, end-systolic dimension; EF, ejection fraction; IVS, inter-ventricular septum; PW, left ventricular posterior wall; BNP, B-type natriuretic peptide.

**Table 2 Medication use other than ACE inhibitors and ARBs**

	All (n=728)	ACE inhibitor (n=356)	ARB (n=372)	P value
β-blocker	68.3	69.4	67.2	0.528
Diuretics	88.5	88.8	88.2	0.803
Spironolactone	48.5	47.8	49.2	0.697
Digitalis	29.5	32.0	27.2	0.150
Calcium-channel blocker	15.1	13.5	16.7	0.231
Antiarrhythmic	20.5	22.5	18.5	0.190
Aspirin	49.2	47.8	50.5	0.452
Warfarin	44.0	48.3	39.8	0.020
Statin	24.6	26.1	23.1	0.347

ACE, angiotensin converting enzyme; ARB, angiotensin receptor blocker

**Table 3 Unadjusted and adjusted hazard risk of long-term outcomes for patients with ACE inhibitors versus ARBs**

Outcomes	Number (%)		Unadjusted HR (95% CI) P-value	Adjusted HR* (95% CI) P-value
	ACE inhibitors (n=322)	ARBs (n=330)		
All cause death	51 (15.8%)	51 (15.5%)	0.958 (0.646-1.405) P=0.807	0.958 (0.601-1.527) P=0.858
Cardiac death	33 (10.2%)	37 (11.2%)	1.071 (0.669-1.713) P=0.775	1.186 (0.680-2.067) P=0.548
Sudden death	9 (2.8%)	11 (3.3%)	1.180 (0.489-2.848) P=0.712	0.770 (0.270-2.199) P=0.626
Hospitalization	101 (31.4%)	99 (30.0%)	0.931 (0.705-1.228) P=0.612	0.964 (0.683-1.362) P=0.836
All cause death or hospitalization	121 (37.6%)	118 (35.8%)	0.924 (0.717-1.191) P=0.540	0.951 (0.694-1.302) P=0.752

HR, hazard risk; CI, confidence interval.

Use of ACE inhibitors was used as a reference against ARBs when the hazard ratios were calculated.

\*Adjusted for age, sex, BMI, hypertension, diabetes mellitus, serum creatinine, NYHA class at discharge, systolic blood pressure, and warfarin.

**Table 4 Subgroup analysis of risk-adjusted all-cause death for HF patients with ACE inhibitors versus ARBs**

Subgroup	n	HR for mortality ACE inhibitors vs ARBs	95% CI	P value
Age				
≥65 yr	386	0.802	0.467-1.378	0.425
<65 yr	266	0.871	0.282-2.690	0.810
Gender				
Male	473	0.979	0.573-1.674	0.940
Female	179	0.506	0.165-0.155	0.233
Etiology				
Ischemic	255	0.747	0.397-1.405	0.365
Non-ischemic	397	1.024	0.491-2.135	0.950
LVEF				
≥25%	422	0.825	0.445-1.529	0.541
<25%	230	1.022	0.459-2.275	0.958
Systolic blood pressure at discharge				
≥110mmHg	388	0.756	0.398-1.437	0.394
<110mmHg	264	1.003	0.498-2.131	0.937
Diastolic blood pressure at discharge				
≥65mmHg	318	0.556	0.250-1.236	0.150
<65mmHg	334	1.156	0.645-2.074	0.626
Hypertension*				
Yes	326	0.966	0.422-2.215	0.935
No	322	1.178	0.645-2.150	0.593
Diabetes				
Diabetes	206	1.413	0.590-3.386	0.438
No diabetes	446	0.760	0.417-1.384	0.369
β-blocker				
β-blocker use	446	0.928	0.492-1.748	0.816
No β-blocker use	206	0.757	0.353-1.623	0.475

HR, hazard risk; CI, confidence interval.

Use of ACE inhibitors was used as a reference against ARBs when the hazard ratios were calculated.

\*Four subjects with insufficient information are not included in the analysis.

**Figure 1**

