

HOKKAIDO UNIVERSITY

Title	Neuroendocrine Control of Anuran Anterior Preoptic Neurons and Initiation of Mating Behavior
Author(s)	Urano, Akihisa
Citation	Zoological Science, 5(5), 925-937
Issue Date	1988-10-15
Doc URL	http://hdl.handle.net/2115/43963
Туре	article
File Information	Z\$5-5_925-937.pdf



REVIEW

Neuroendocrine Control of Anuran Anterior Preoptic Neurons and Initiation of Mating Behavior

Akihisa Urano¹

Department of Regulation Biology, Faculty of Science, Saitama University, Urawa, Saitama 338, Japan

INTRODUCTION

The preoptic area plays an important role in the evocation of sexual behavior in many vertebrate species [1-6]. In the anuran brain, the anterior part of the preoptic nucleus (APON) is considered to be the center for triggering male mate calling behavior. After careful re-examination of the lesion study by Aronson and Noble [7] on the brain of the leopard frog Rana pipiens, Schmidt [2, 8] proposed that the APON initiates an organizing activity of mating call patterns in the brain stem "call center" which is composed of the pretrigeminal nucleus and the larvngeal motor neurons. The same system may function in the brain of the Japanese toad, Bufo japonicus [9, 10]. The APON is further concerned with female orientation to conspecific mating calls in R. pipiens [11].

Anuran mating behavior is controlled not only by neural input signals but also by various hormonal signals. Testosterone seems to be not a sole, but a crucial hormonal factor in the initiation of male mate calling behavior [12–14]. In gravid female *Rana pipiens*, vasotocin, an anuran neurohypophysial hormone, suppressed female release calling behavior, and in turn elevated sexual receptivity [15]. Luteinizing hormone-releasing hormone (LHRH) also increased sexual receptivity in female *Xenopus laevis* [16]. The amphibian brain shows a fundamental organizational pattern common to the structure of the vertebrate brain [17]. Clarification of neuroendocrine control of mating behavior in amphibians is thus important for understanding phylogenetically fundamental control mehanisms of sexual behavior. In this paper, I first describe the morphological characteristics of the preoptic nucleus (PON), and then review recent studies concerning the neural and hormonal control mechanisms for the initiation of seasonal reproductive behavior in anuran amphibians.

CYTOARCHITECTURE OF THE PON

Subnuclear Organization

The PON is a neuronal cell mass which surrounds the preoptic recess, and is considered to be an important neuroendocrine center. It is easily divisible into anterior and posterior halves by a relatively cell-poor zone along the sulcus preopticus (Fig. 1), and each half is composed of several subnuclei, some of which include many neurosecretory neurons. Recent physiological studies indicate that these subnuclei are functionally distinctive [18].

In the Japanese toad, the PON can be divided into seven subnuclei: the anterior part of the PON (APON), the dorsal and the ventral periventricular parts, the dorsal and the ventral magnocellular parts (*dmc* and *vmc*), the suprachiasmatic part, and the posterior part of the PON (PPON) [18]. The APON is composed of a rather compact cell

Received May 27, 1988

Present address: Ocean Research Institute, University of Tokyo, Minamidai, Nakano-ku, Tokyo 164, Japan.



FIG. 1. Diagram of the parasagittal midplain of the toad diencephalon. Note that the preoptic nucleus which locates antero-dorsal to the optic chiasma (OC) is divisible into the anterior part (APON) and the posterior part (PPON). AC, anterior commissure; EP, epiphysis; LT, lamina terminalis; ME, median eminence; NID, nucleus infundibularis dorsalis; NIV, nucleus infundibularis ventralis; OL, optic lobe; PD, pars distalis of hypophysis; PI, pars intermedia of hypophysis; PN, pars nervosa of hypophysis; TEG, tegmentum mesencephali; TEL, telencephalon; IIIv, third ventricle. (from Urano and Ishihara [110])

mass surrounding the anterior portion of the preoptic recess in the toad brain, while, in many ranid species, APON neurons are well organized in a laminar pattern [19–21]. The *dmc* and the *vmc* consist of magnocellular neurosecretory neurons which contain immunoreactive (ir) vasotocin [18, 22], mesotocin [22], somatostatin [23] and other peptides [24]. The toad suprachiasmatic part in-

cludes small ir-vasotocin neurons and receives direct retinal innervation (Shimotoso and Urano, unpublished). Synapses between the optic terminals and the PON neurons were observed in *Rana temporaria* [25]. The PPON, the name of which follows nomenclatorial usage in the brain atlas of *Rana pipiens* [21], is a common preoptic structure in many anuran species.

Sexual Dimorphism

Differences in patterns of sexual behavior between males and females are partly due to differences between male and female neuronal circuitries, i. e., sexual dimorphism of nervous systems. As in songbirds [26], the Japanese quail [27], and some mammalian species [28-31], the brain loci concerning male mate calling behavior showed sexual dimorphism in Xenopus [32], Rana pipiens [33] and Bufo japonicus [34]. As for the toad PON, the nuclear volume of the APON in the male was significantly larger than that in the female (male/female, 1.25-1.39). The mean of the total cell number in the male APON was 1.20 times that of the female, while the cell sizes of male and female APON neurons were in the same range [35]. These results suggest that the sexual difference in the APON volumes was caused by a greater cell number in the male APON. In addition, the amygdala pars medialis (Am), which forms a morphological and probably functional complex with the APON [36], showed a similar

TABLE 1. Volumes $(\times 10^{-3} \text{ mm}^3, \text{mean} \pm \text{S.E.})$ of the anterior part of the preoptic nucleus (APON) and the amygdala pars medialis (Am) in 1-year old toadlets, and hibernating and post-breeding adult toads

Subnuclei	Male	Female	M/F
APON			
1-year old toad	103.6 ± 4.5	98.1±3.2	1.06
hibernating	311.5 ± 23.0	224.1 ± 14.2	1.39*
post-breeding	250.0 ± 12.3	199.3 ± 9.7	1.25**
Am			
1-year old toad	45.9 ± 2.1	35.5 ± 3.1	1.30*
hibernating	153.8 ± 9.5	90.1 ± 4.2	1.71**
post-breeding	110.1 ± 8.4	58.9 ± 3.2	1.87**

*, P<0.02; ** P<0.005 by the t-test.

M/F, the ratio of the nuclear volume in the male to the female.

926

sexual difference. Ontogenetically, the sexual difference in the Am volumes appeared in yearling toadlets, preceding the development of the APON (Table 1).

The sexual dimorphism in the APON seems to be a fundamental part of vertebrate brain structure in general, since the preoptic area is sexually dimorphic in the quail [27], the rat [30] and the macaque monkey [29].

MATE CALLING TRIGGER CENTER

Location in the APON

Mate calling can be induced by electrical stimulation of the APON. In freely moving *Rana pipiens*, the effective sites for electrical evocation of calling were located in the rostral extreme of the PON [37]. Using isolated *pipiens* brain preparations in which neural correlates of mate calling can be recorded, the area most sensitive to electrical stimulation was localized in the ventral half of the APON [38]. The ventrolateral border of the preoptic gray was much more sensitive than the ventricular surface. Stimulation of the same locus was also effective in inducing electrical activity correlating to mate calling in *in situ* perfused brains of Japanese toads (Fig. 2).

Scanning Electron Microscopy of the Ventricular Surface

In the anuran brain, the ventricular wall of the dorsal part of the APON is ciliated as are the surfaces of the ventricular walls in many other brain loci. However, the wall of the ventral part is only sparsely ciliated; instead, it is studded with numerous large bulbous protrusions which are large cytoplasmic extensions of ependymal cells and intraventricular dendritic end bulbs of secretory neurons [36, 39–41]. This area includes almost all the portions of the ventricular wall of the rostroventral part of the APON, and corresponds topographically to the area mentioned above that is most sensitive to the initiation of calling behavior by electrical stimulation.

Steroid Hormone Accumulating Neurons

The APON has been considered to be a sex steroid-sensitive center that triggers sexual behavior in anurans. Autoradiographic studies of the brains in *Xenopus laevis* [42, 43] and *Rana pipiens* [44] showed that many APON neurons can



FIG. 2. Neural correlates of mate calling induced by electric stimulation (2 ms biphasic pulse, 50 Hz for 1 sec) of the APON through a fine bipolar electrode in a perfused toad brain. Electrical activity correlating mate calling was recorded at the root of the vagus nerve (n.X, upper trace) and the hyoglossal muscle (m. hg, lower trace). Magnitudes of the neural correlates increased when the intensity of stimulation was elevated from 0.2 mA to 0.6 and 1.0 mA.

accumulate sex steroid hormones. Such neurons were localized throughout the dorsal-ventral extent in the rostral part of the APON, but were found ventrally in the caudal part. This distributional pattern of sex steroid-accumulating APON neurons corresponds to the localization pattern of the ventricular bulbous protrusions. In mammals, the medial preoptic nucleus, which is the homologue of the APON, includes many sex steroidaccumulating neurons [45, 46].

Intracranial implantation of testosterone into or near the APON enhanced the incidence of mate calling which was evoked acoustically by play-back of tape-recorded conspecific mating calls in *Rana pipiens* [14]. In mammals, steroid hormones are reported to excite electrical activity of central neurons [47–50]. At present, it is not clear whether testosterone can modulate electrical activity of APON neurons in the amphibian brain due to the lack of any experimental data.

Projection of the APON to the Brain Stem Call Center

Mate calling behavior and its neural correlates could be evoked even after massive ablations or lesions to brain areas including almost all of the telencephalon, the dorsal thalamus, the torus semicircularis, the dorsal part of the isthmic nucleus and the infundibulum [38, 51]. These results indicate the presence of pathways at the ventrolateral border of the central gray from the APON to the isthmo-trigeminal tegmentum.

The retrograde axonal transport study in which afferents to laryngeal motor neurons were traced in the brain of *Xenopus laevis* showed the presence of direct projection of APON neurons to the pre-trigeminal nucleus of the dorsal tegmental area (DTAM). DTAM neurons send efferent fibers to laryngeal motor neurons [52]. The projection from the APON to the DTAM is less flourishing in the female than in the male.

HUMORAL SIGNAL RECEPTIVITY

As is described above, the APON from which efferent fibers project to the brain stem call center is believed to be the androgen-sensitive center for male mate calling behavior. An electrophysiological study of this locus in *Rana pipiens* showed the presence of neurons responsive to auditory stimulation by playback of conspecific mating calls, and injections of pituitary homogenate significantly increased the percentage of these units excited by the calls [53]. Activity of APON neurons thus can be modulated by humoral signals. A Golgielectron microscopic study revealed that a portion of the APON neurons have the proper anatomical features for detecting humoral signals (Fig. 3) [36, 54].



FIG. 3. Diagram showing CSF-contacting and BCcontacting neurons in the APON. These neurons have the proper morphological features to receive both humoral and neuronal synaptic, either chemical or electric, input signals. bc, blood capillary; EP, ependymal layer; Med, medial gray of the APON; Lat, lateral white of the APON; POR, preoptic recess. (from Urano and Ishihara [110])

Cerebrospinal Fluid (CSF)-Contacting Neurons

In many anuran species, the APON, especially its ventral half, contains CSF-contacting neurons [36, 39-41]. These neurons were rapid-Golgi stained and gold-toned, and then were examined by scanning electron microscopy after removal of surrounding tissues with hydrochloric acid and collagenase. The somata of stripped neurons bear debris of nerve terminals on their surfaces, suggesting that the neuronal activity of these cells can be modulated by various synaptic input signals.

Processes of CSF-contacting neurons, probably dendritic, protrude into the preoptic recess. The dendrites projecting into the preoptic recess from preoptic neurosecretory cells can be equipped to serve both secretory and sensory functions [55]. The CSF-ventricular system is thought to distribute biologically-active hormonal substances within the brain, since many researchers have found various hormones in the CSF, such as LHRH and thyrotropin-releasing hormone (TRH) [56], oxytocin and vasopressin [57], and melatonin [58]. The concentrations of these hormones varied according to various physiological statuses. Further, single intraventricular injections of LHRH and TRH increased the amplitudes and frequency of electroencephalographic (EEG) activity recorded from the brain of a hibernating Japanese toad (Fig. 4) [59]. The effective dose of $1 \mu g$ needed for enhancement of EEG activity through a single intraventricular injection of LHRH or TRH was much less than that needed for systemic injections. It is thus possible that the CSF-contacting neurons



FIG. 4. Effects of intraventricularly injected LHRH and TRH on EEG activity of a hibernating Japanese toad. According to amplitudes and frequency, EEG activity was categorized into 4 levels: 1, resting; 11, awake; 111, active; and IV, very active. Each level corresponds well to a behavioral state. Note that both LHRH and TRH induced dual fast and slow enhancements of EEG activity. Although the EEG record is not shown, the pattern of LHRH-induced fast response includes highly synchronized bursting waves that were not observed upon injection of control saline. (Fujita, thesis, Saitama University)

whose dendritic processes protrude into the preoptic recess detect changes in ventricular hormonal status and motivate the neuronal circuitry in preparation for mating behavior in pre-breeding anurans.

Blood Capillary (BC)-Contacting Neurons

The presence of BC-contacting neurons is incompatible with the general concept of the relations between brain neurons and capillaries. Blood capillaries in the vertebrate brain are generally surrounded by astrocytic endfeet with an intervening basement membrane, so that brain neurons, even fish hypothalamic neurosecretory cells, are separated from the vascular endothelium Nonetheless, neurosecretory cells which [60]. directly come into contact with blood capillaries were shown in the toad preoptic nucleus [61]. Recently, it was found that a considerable number of peptidergic neurons come into contact with blood capillaries with only an intervening basement membrane in the APON of both the bullfrog and the Japanese toad [36].

BC-contacting neurons send their dendrites laterad toward the preoptic white matter. Although arborization is rather poor, the dendrites usually bifurcate several times and form dendritic fields. There, many axon terminals form synapses on the dendritic spines of these neurons. It is highly probable that APON neurons receive the input signals of the afferent fibers mainly through the dendritic synapses in the preoptic white matter along the border of APON cell mass, since Halpern [62] noted that terminal degeneration by the telencephalic lesions was located along the lateral edges of cell masses in the frog hypothalamus. The single BC-contacting neurons thus detect changes in titers of blood-born hormones, preferably sex steroid hormones which have activational effects on APON neurons, and further receive neuronal input signals through dendritic synapses to integrate hormonal and neural signals concerned with the initiation of sex behavior.

AFFERENTS OF THE APON

Retrograde Horseradish Peroxidase (HRP) Study

It is important to know what neuroanatomic afferent relations the APON has with other parts of the brain. Such information is requisite for better understanding of the sensory modalities and activating or inhibiting pathways that might trigger or modulate sexual behavior through the PON. Thus, the afferents of the APON were examined in *Rana pipiens* (Urano and Gorbman, unpublished) and *Bufo japonicus* [63] using the retrograde HRP method, which is a particularly useful tool in studies of neural connections.

Evidence of retrogradely transported enzymatic activity was observed in perikarya and neuropil in the following brain regions: the ventro-medial limbic cortex, the posterior part of the preoptic nucleus including the magnocellular part, the infundibular nuclei, the thalamic area, the subtectal and tegmental regions including the reticular formation, and the rhombencephalic central gray. Neurons in these regions appear to send their axons to the APON mainly via the medial and lateral forebrain bundles. Localization of some HRP-labeled perikarya and fibers coincides with that of immunoreactive perikarya and fibers containing either LHRH, vasotocin or TRH which have been considered to project to the APON [64, 65].

Particular HRP-labeled loci in the ventro-medial limbic cortex included the nucleus medialis septi, the nucleus lateralis septi, the nucleus accumbens septi, the amygdala pars medialis and the nucleus of the diagonal band of Broca. The amygdalapreoptic tract may exist in all vertebrate classes from cyclostomes to mammals [66]. The septal projection to the preoptic area in the leopard frog and the Japanese toad has an apparently homologous relationship to a similar pattern in the lizard [67] and the rat [68, 69]. Although in anurans, the physiological significance of amygdaloid and septal projections to the APON is not clear at present, it is possible that these projections are concerned with the control of sexual behavior as has been claimed in mammals [70, 71].

HRP-labeled structures in the subtectal and

tegmental regions were the nucleus anterodorsalis tegmenti mesencephali, the torus semicircularis, the nucleus posteroventralis tegmenti mesencephali, the nucleus isthmi and the mesencephalic reticular nuclei. Mesencephalic projections to the anterior hypothalamus are well known in amphibian brains [17, 19, 66, 72] as well as in other vertebrate classes [66, 73, 74]. The mammalian preoptic area is directly continuous with a vast nonspecific neuronal apparatus of the brain stem reticular formation [75]. In frog brains, the mesencephalic reticular system receives afferents from various parts of the brain, such as the telencephalon [62, 76], the optic tectum [77], and the superior olivary nucleus [78]. The presence of multimodal inputs suggests a nonspecific or generalized character of function of the anuran reticular formation as a possible activating or inhibitory regulatory system which may influence the neural substrate for mating behavior.

Chemical Neuroanatomy of the APON Afferents

Information on the chemical nature of APON afferents is important for the examination of control mechanisms of APON neuronal activity at the cellular and molecular levels.

The HRP study mentioned above showed the presence of HRP-labeled neurons in the magnocellular part of the PON, and in the nuclei infundibularis dorsalis and ventralis in the toad brain. These regions are rich in vasotocinergic and mesotocinergic neurosecretory neurons [22], and TRH neurons [65, 79], respectively. Jokura and Urano [64] verified that varicose ir-vasotocin fibers are found in the ventrolateral region of the APON where the APON neurons have their dendritic fields. Some ir-vasotocin fibers from the vmc protrude into the APON cell mass, and appeared to come into contact with somata of APON neurons. In Japanese toads, ir-TRH neurons were localized mainly in the nucleus infundibularis ventralis (NIV) [65]. Ir-TRH fibers arising from the NIV neurons project to the median eminence to form the hypothalamo-hypophysial tract. In addition, a considerable number of ir-TRH fibers innervate into the APON. In the APON, varicose ir-TRH fibers are scattered widely among the neuronal cell mass and the white matter.

Other important loci in the toad brain where HRP-labeled neurons were found include the nucleus medialis septi and the nucleus of the diagonal band of Broca. These loci contained many ir-LHRH neurons which project to the APON [64]. Most ir-LHRH fibers emanating from the nucleus medialis septi form a loose fiber bundle with those arising from the diagonal band of Broca. These ir-LHRH fibers, which have typical beaded features, project to the ventrolateral border of the preoptic gray.

In mammalian brains, peptidergic axonic processes form ordinary synapses [80] and en passant synapses with dendritic profiles [81]. Therefore, it is highly probable that varicose ir-vasotocin, ir-TRH and ir-LHRH fibers form ordinary or en passant synapses in the dendritic fields of APON neuron in the toad brain.

Functional Significance of the APON Afferents

The retrograde HRP study indicates that there are multimodal inputs to the APON from various regions of the brain. The septal nuclei, which send ir-LHRH fibers to the APON, receive olfactory inputs through the medial olfactory tract [82, 83], and the amygdala is innervated by projections from the accessory olfactory bulb [84]. These limbic nuclei, from which afferents to the APON arise, may relay olfactory signals to the APON neurons. In addition, the terminal nerve, which may function in odor processing, sends an ir-LHRH-ergic projection to the preoptic region in the tiger salamander and the bullfrog [85].

Visual cues can be conveyed through direct retinal projection to the suprachiasmatic part of the PON. This was clarified in the Japanese toad by use of a cobaltic lysine method (Shimotoso and Urano, unpublished). The presence of direct retino-preoptic projetion has also been supposed in the brain of *Rana temporaria* [25]. Acoustic signals which excite APON neurons may reach the preoptic region through at least two ascending pathways in the brain stem [78]. One is the pathway relayed through the nucleus oliva superior and the nucleus profundus mesencephali; the other is that relayed through the nucleus oliva superior and the torus semicircularis. The thalamo-preoptic connection is a possible pathway for transmission of tactile signals. Thus, the APON neurons may be influenced by various kinds of sensory inputs, although almost all sensory modalities are relayed and may be regulated either by sex steroid hormones or by neurohormones released from extrahypothalamic terminals of neurosecretory neurons [86, 87]. Since the electrical activity of many APON units was excited by iontophoretically applied LHRH, TRH and vasotocin (Fujita and Urano, in preparation), the APON neurons probably integrate various sensory inputs under the influence of peptidergic neurosecretory neurons, and then generate neural signals for the initiation of mate calling behavior.

SEASONAL VARIATIONS

Many anurans, especially those in the temperate zone, are typical seasonal breeders which spawn in spring or early summer. The neuroendocrine systems associated with reproductive behavior also show seasonal changes in their synthetic and secretory activities. In bullfrogs, the plasma level of luteinizing hormone (LH), which can increase androgen secretion from the testes [88], was elevated during the breeding season [89]. In the Xenopus hypothalamus, the contents of LHRH, which can stimulate pituitary gonadotropin release in bullfrogs [90] and plasma androgen levels in Japanese toads [91], varied seasonally in correspondence to reproductive physiological states [92]. Ishii and his collaborators measured circannual changes in plasma levels of various pituitary hormones [93, 94], thyroid hormones [95], adrenal steroids [96, 97] and sex steroids [98]. Most of these hormones showed marked increases in their plasma titers prior to or during the breeding season. Further, classical histochemical and morphometric studies showed seasonal morphological changes in the hypothalamo-neurohypophysial neurosecretory system in Rana temporaria [99, 100]. The results of these studies suggest the possibility that the APON neurons do show some seasonal changes in their morphological and functional features, since the activity of APON neurons was modulated by administrations of testosterone [14] and pituitary homogenates [53].

Seasonal Changes in the Volumes of PON Subnuclei

Seasonal variations in the volumes of several preoptic and amygdala subnuclei were found between hibernating and postbreeding Japanese toads [34]. The APON in the hibernating males was 125% larger than that in the post-breeding animals. The seasonal difference in the female APON was not statistically significant. In both sexes, the hibernating animals had larger ventral magnocellular parts of the PON, amygdala medialis and amygdala lateralis than the postbreeding animals did. The seasonal variation in the volumes of several subnuclei mentioned above may be due to hypertrophy of the neurons in these loci, since cell nulear volumes of PON neurons increase prior to the breeding season [100]. Morphological changes in the APON and the amygdala thus precede physiological and behavioral changes in the breeding season. A similar result was observed in the song control nucleus in the brain of the male canary which is larger during breeding season than in the fall when the animal is sexually inactive [101], probably because sex steroids induce dendritic growth in this nucleus [102].

Immunoreactivity of Neuroendocrine Cells

It is highly probable that neuronal input signals to APON neurons differ seasonally. Therefore, seasonal variations in LHRH, TRH and vasotocin that were localized in varicose afferent fibers to the APON were examined immunohistochemically in toad forebrains and neurohypophyses [65, 103]. The immunohistochemical technique utilized was the avidin-biotin-peroxidase complex (ABC) method, which is superior to the peroxidaseantiperoxidase method in a quantitative study. LHRH immunoreactivity (ir) was strong in both perikarya and fibers in animals captured in spring and autumn, while in summer animals, LHRH-ir was weak. Seasonal changes in TRH-ir were similar to those in LHRH-ir, while significant seasonal variations were not found in vasotocin-ir. The circannual changes in LHRH-ir appear to correspond with seasonal variations in plasma steroid levels reported by Inoue et al. [98]. This coincidence implies that LHRH and sex steroids

can have synergistic effects on the control of APON neurons.

Effects of Castration

As is described above, testosterone may modulate neural activity of the APON to initiate reproductive behavior. Structures of the nervous system are modified by sex steroids during both fetal and adult periods in many vertebrate species [101, 102, 104]. In Japanese toads, the volumes of the APON and the amygdala in the male are larger than those in the female. Furthermore, the volumes of these nuclei and LHRH-ir also changed seasonally. These changes appear to correlate with the annual variation of plasma testosterone levels. Castration experiments, in which the role of testosterone in the control of the phenomena mentioned above was examined, showed that the effects of castration differ seasonally [105]. The volume of the amygdala medialis in autumn toads was significantly reduced by castration; however, the reduction in spring animals was not statistically signif-Meanwhile, castration did not modify icant. LHRH-ir in the median eminence in either spring or autumn toads, although dense ir-LHRH fibers were observed in the mesencephalic tegmental region in the castrated spring toads but not in the autumn toads either intact or castrated. These results suggest that seasonal influences on the effects of castration were not uniform among the different brain loci.

COORDINATION OF NEURAL AND ENDOCRINE ACTIVITY

Temporal coordination of neural and endocrine events is a cruial requisite for successful reproduction. Plausible candidates for coordinating the brain and endocrine functions are LHRH-ergic and vasotocinergic neurosecretory systems, both of which send fine varicose fibers to various extrahypothalamic brain loci other than the median eminence and the pars nervosa [86, 87]. Both LHRH-ergic and vasotocinergic fibers innervate either sensory or motor centers concerned with reproductive behavior.

LHRH applied by microiontophoresis increased discharge rates of individual neurons in the septal

preoptic area of the rat [106], and in the APON of the Japanese toad (Fujita and Urano, in preparation). An intracellular study showed that LHRH can mimic slow excitatory postsynaptic potentials when applied to the sympathetic neurons in the bullfrog [107]. LHRH at the synaptic level may play a role in increasing neuronal excitability in the loci where LHRH fibers innervate. On the other hand, LHRH applied systemically or intraventricularly can stimulate the pituitary-gonadal axis to elevate plasma androgen levels in male Japanese toads (Fig. 5). This evidence suggests that LHRH simultaneously affects both the neuronal activity of the APON neurons as an excitatory neurotransmitter or neuromodulator and the endocrine events of the pituitary-gonadal axis as a hypothalamic releasing hormone.

The endocrine functions of vasotocin in amphib-

ians are well documented in many endocrine textbooks. In addition, vasotocin and its homologues can excite unit-spike activity of neurons in the rat supraoptic and paraventricular nuclei [106, 108], the eel preoptic nucleus [109] and the toad APON (Fujita and Urano, in preparation). Vasotocin thus may facilitate the activity of many central neurons as a neuromodulator or a local hormone. The latter possibility is supported by the fact that 10^{-9} M vasopressin, comparable to the effective dose of vasopressin necessary for peripheral targets, can excite rat paraventricular neurons [108].

At present, it is difficult to account for the temporal discrepancy between LHRH-induced neural events (Fig. 4) and endocrine events (Fig. 5). When LHRH functions as a neurotransmitter or a neuromodulator, its influence on target



FIG. 5. Changes in the plasma testosterone levels after administrations of intravenous and intraventricular LHRH. Each curve represents a change in testosterone levels in an individual male toad. a, effects of intravenous saline as a control; b, single intravenous injection of 10 μg LHRH; c, single intravenous injection of 100 μg LHRH; d, continuous infusion of LHRH at a dose of 100 μg/hour for 4 hours; and e, intracerebroventricular administration of 100 ng LHRH. Note that the dose of intracranial LHRH which markedly elevated plasma testosterone was much less than that of intravenous LHRH. (Fujita, thesis, Saitama University) neurons lasted for within the order of seconds or minutes. However, endocrine events, e.g., the secretion of androgen, take a much longer time. Since the APON neurons are sex steroid-sensitive and are excited by LHRH, some unknown intrinsic cellular mechanisms within the APON neurons and neurons having the same characteristics may regulate the above temporal discrepancy in order to complete seasonal breeding successfully.

REFERENCES

- Demski, L. S. and Knigge, K. M. (1971) The telencephalon and hypothalamus of the bluegill (*Lepomis macrochirus*): evoked feeding, aggressive and reproductive behavior with representative frontal sections. J. Comp. Neurol., 143: 1-16.
- 2 Schmidt, R. S. (1968) Preoptic activation of frog mating behavior. Behaviour, 30: 239-257.
- 3 Wheeler, J. M. and Crews, D. (1978) The role of the anterior hypothalamus-preoptic area in the regulation of male reproductive behavior in the lizard, *Anolis carolinensis*: lesion studies. Horm. Behav., 11: 42-60.
- 4 Barfield, R. J. (1969) Activation of copulatory behavior by androgen implanted into the preoptic area of the male fowl. Horm. Behav., 1: 37–52.
- 5 Morali, G. and Beyer, C. (1979) Neuroendocrine control of mammalian estrous behavior. In "Endocrine Control of Sexual Behavior". Ed. by C. Beyer, Raven Press, New York, pp. 33-175.
- 6 Larssen, K. (1979) Features of the neuroendocrine regulation of masculine sexual behavior. In "Endocrine Control of Sexual Behavior". Ed. by C. Beyer, Raven Press, New York, pp. 77-163.
- 7 Aronson, L. R. and Noble, G. K. (1945) The sexual behavior of anura. II. Neural mechanisms controlling mating in the male leopard frog, *Rana pipiens*. Bull. Am. Mus. Nat. Hist., 86: 83-140.
- 8 Schmidt, R. S. (1976) Neural correlates of frog calling: isolated brainstem. J. Comp. Physiol., 108: 99-113.
- 9 Urano, A. (1983) Neural correlates of mating behavior in perfused toad brains. Zool. Mag., 92: 601.
- 10 Satou, M., Oka, Y. and Ueda, K. (1987) The nervous system. In "Biology of Toad". Ed. by A. Urano and K. Ishihara, Shokabo, Tokyo, pp. 62– 90.
- Schmidt, R. S. (1969) Preoptic activation of mating call orientation in female anurans. Behaviour, 35: 114-127.
- 12 Schmidt, R. S. (1966) Hormonal mechanisms of frog mating calling. Copeia, 1966: 637–644.

- 13 Palka, Y. S. and Gorbman, A. (1973) Pituitary and testicular influenced sexual behavior in male frogs, *Rana pipiens*. Gen. Comp. Endocrinol., 21: 148-151.
- 14 Wada, M. and Gorbman, A. (1977) Relation of mode of administration of testosterone to evocation of male sex behavior in frogs. Horm. Behav., 8: 310-319.
- 15 Diakow, C. and Nemiroff, A. (1981) Vasotocin, prostaglandin, and female reproductive behavior in the frog, *Rana pipiens*. Horm. Behav., 15: 86-93.
- 16 Kelley, D. B. (1982) Female sex behaviors in the South African clawed frog, *Xenopus laevis*: gonadotropin-releasing, gonadotropic, and steroid hormones. Horm. Behav., 16: 158-174.
- 17 Herrick, C. J. (1948) The brain of tiger salamander, *Ambystoma tigrinum*. Univ. Chicago Press, Chicago.
- 18 Takami, S., Jokura, Y. and Urano, A. (1984) Subnuclear organization of the preoptic nucleus in the toad, *Bufo japonicus*. Zool. Sci., 1: 759–770.
- 19 Frontera, J. G. (1952) A study of anuran diencephalon. J. Comp. Neurol., 96: 1-69.
- 20 Hoffman, H. H. (1963) The olfactory bulb, accessory olfactory bulb and hemisphere of some anurans. J. Comp. Neurol., 120: 317–368.
- 21 Wada, M., Urano, A. and Gorbman, A. (1980) A stereotaxic atlas for diencephalic nuclei of the frog, *Rana pipiens*. Arch. Histol. Jpn., 43: 157–173.
- 22 Vandesande, F. and Dierickx, K. (1976) Immunocytochemical demonstration of separate vasotocinergic and mesotocinergic neurons in the amphibian hypothalamic magnocellular neurosecretory system. Cell Tissue Res., 175: 289– 296.
- 23 Vandesande, F. and Dierickx, K. (1980) Immunocytochemical localization of somatostatincontaining neurons in the brain of *Rana temporaria*. Cell Tissue Res., 205: 45–53.

ł

- Yui, R. (1983) Immunohistochemical studies on peptide neurons in the hypothalamus of the bullfrog *Rana catesbeiana*. Gen. Comp. Endocrinol., 49: 195-209.
- 25 Vullings, H. B. G. and Heussen, A. M. A. (1975) Electron microscopic observations on the retinopreoptic pathway of *Rana temporaria*. Cell Tissue Res., 161: 177-182.
- 26 Arnold, A. P. (1980) Sexual differences in the brain. Am. Sci., 68: 165-173.
- 27 Panzica, G. C., Viglietti-Panzica, C., Calacagni, M., Anselmetti, G. C., Schumacher, M. and Balthazart, J. (1987) Sexual differentiation and hormonal control of the sexually dimorphic medial preoptic nucleus in the quail. Brain Res., 416: 59-68.
- 28 Arimatsu, Y., Seto, W. and Amano, T. (1981) Sexual dimorphism in alpha-bungarotoxin binding

capacity in the mouse amygdala. Brain Res., 213: 432-437.

- 29 Aynob, D. M. and Greenough, W. T. (1983) Sex differences in dendritic structure in the preoptic area of the juvenile macaque monkey brain. Science, 219: 197-198.
- 30 Gorski, R. A., Gordon, J. H., Shryne, J. E. and Southam, A. M. (1978) Evidence for a morphological sex difference within the medial preoptic area of the rat brain. Brain Res., 148: 333-346.
- 31 Matsumoto, A. and Arai, Y. (1986) Morphological evidence for sexual dimorphism in wiring pattern in the neuroendocrine brain. In "Pars Distalis of the Pituitary Gland-Structure, Function and Regulation". Ed. by F. Yoshimura and A. Gorbman, Elsevier, Amsterdam, pp. 239-245.
- 32 Hannigan, P. and Kelley, D. B. (1981) Male and female laryngeal motor neurons in *Xenopus laevis*. Soc. Neurosci. Abstr., 7: 269.
- 33 Schmidt, R. S. (1982) Sexual dimorphism in succinic dehydrogenase staining of toad pretrigeminal nucleus. Exp. Brain Res., 45: 447-450.
- 34 Takami, S. and Urano, A. (1984) The volume of the toad medial amygdala-anterior preoptic complex is sexually dimorphic and seasonally variable. Neurosci. Lett., 44: 253–258.
- 35 Takami, S., Higashiyama, T. and Urano, A. (1984) A quantitative analysis of sexual dimorphism in the anterior part of the preoptic nucleus in the toad. Neurosci. Lett., S17: S35.
- 36 Urano, A. (1984) A Golgi-electron microscopic study of anterior preoptic neurons in the bullfrog and the toad. Zool. Sci., 1: 89–99.
- 37 Wada, M. and Gorbman, A. (1977) Mate calling induced by electrical stimulation in freely moving leopard frogs, *Rana pipiens*. Horm. Behav., 9: 141-149.
- 38 Schmidt, R. S. (1984) Neural correlates of frog calling: preoptic area trigger of 'mating calling'. J. Comp. Physiol., 154: 847-853.
- 39 Nakai, Y., Ochiai, H., Shioda, S. and Ochi, J. (1977) Cytological evidence for different types of cerebrospinal fluid-contacting subependymal cells in the preoptic and infundibular recesses of the frog. Cell Tissue Res., 176: 317-334.
- 40 Fasolo, A. and Franzoni, M. F. (1977) A Golgi study on the hypothalamus of amphibia. The neuronal typology. Cell Tissue Res., 178: 341-354.
- 41 De Waele, G. and Dierickx, K. (1979) Scanning electron microscopy of the wall of the third ventricle in the brain of *Rana temporaria*. Part IV. Cell Tissue Res., 203: 53-64.
- 42 Kelley, D. B., Morrell, J. I. and Pfaff, D. W. (1975) Autoradiographic localization of hormoneconcentrating cells in the brain of an amphibian, *Xenopus laevis.* I. Testosterone. J. Comp. Neurol.,

164: 47-62.

- 43 Morrell, J. I., Kelley, D. B. and Pfaff, D. W. (1975) Autoradiographic localization of hormoneconcentrating cells in the brain of an amphibian, *Xenopus laevis*. II. Estradiol. J. Comp. Neurol., 164: 63-78.
- 44 Kelley, D. B., Lieberburg, I., McEwen, B. S. and Pfaff, D. W. (1978) Autoradiographic and biochemical studies of steroid hormone-concentrating cells in the brain of *Rana pipiens*. Brain Res., 140: 287– 305.
- 45 Sar, M. and Stumpf, W. E. (1975) Distribution of androgen-concentrating neurons in rat brain. In "Anatomical Neuroendocrinology". Ed. by W. E. Stumpf and L. D. Grant, S. Karger, Basel, pp. 120-133.
- 46 Pfaff, D. W. (1980) Estrogen and Brain Function. Springer-Verlag, New York, Heidelberg, Berlin.
- 47 Yagi, K. (1973) Changes in firing rates of single preoptic and hypothalamic units following an intravenous administration of estrogen in the castrated female rat. Brain Res., 53: 343–352.
- 48 Bueno, J. and Pfaff, D. W. (1976) Single unit recording in hypothalamus and preoptic area of estrogen-treated and untreated ovariectomized female rats. Brain Res., 101: 67–78.
- 49 Kow, L.-M. and Pfaff, D. W. (1985) Estrogen effects on neuronal responsiveness to electrical and neurotransmitter stimulation: an in vitro study on the ventromedial nucleus of the hypothalamus. Brain Res., 347: 1-10.
- 50 Nabekura, J., Oomura, Y., Minami, T., Mizuno, Y. and Fukuda, A. (1986) Mechanism of the rapid effect of 17-estradiol on medial amygdala neurons. Science, 233: 226-228.
- 51 Schmidt, R. S. (1966) Central mechanisms of frog calling. Behaviour, 26: 251-285.
- 52 Wetzel, D. M., Haerter, U. L. and Kelley, D. B. (1985) A proposed neural pathway for vocalization in South African clawed frogs, *Xenopus laevis*. J. Comp. Physiol. A, 157: 749-761.
- 53 Urano, A. and Gorbman, A. (1981) Effects of pituitary hormonal treatment on responsiveness of anterior preoptic neurons in male leopard frogs, *Rana pipiens. J. Comp. Physiol.*, 141: 163-171.
- 54 Urano, A. (1985) The anterior preoptic neurons of the frog, *Rana catesbeiana* and toad, *Bufo bufo japonicus*: control of mating behavior. In "Current Trends in Comparative Endocrinology". Ed. by B. Lofts and W. N. Holmes, Hong Kong Univ. Press, Hong Kong. pp. 1125–1126.
- 55 Smoller, C. G. (1965) Neurosecretory processes extending into third ventricle: secretory or sensory? Science, 147: 882-884.
- 56 Joseph, S. A., Sorrentino, S., Jr. and Sundberg, D. K. (1975) Releasing hormones, LRF and TRF,

in the cerebrospinal fluid of the third ventricle. In "Brain-Endocrine Interaction II. The Ventricular System in Neuroendocrine Mechanisms". Ed. by K. M. Knigge, D. E. Scott, H. Kobayashi and S. Ishii, Karger, Basel, pp. 306–312.

- 57 Robinson, I. C. A. F. and Jones, P. M. (1982) Neurohypophysial peptides in cerebrospinal fluid: recent studies. In "Neuroendocrinology of Vasopressin, Corticoliberin and Opiomelanocortins". Ed. by A. J. Baertchi and J. J. Dreifuss, Academic Press, London, pp. 21–31.
- 58 Klein, D. C. (1979) Circadian rhythms in the pineal gland. In "Endocrine Rhythms". Ed. by D. T. Krieger, Raven Press, New York, pp. 202–223.
- 59 Fujita, Y. and Urano, A. (1985) Enhancement of EEG activity by injection of LHRH and TRH in hibernating toad, *Bufo japonicus*. Zool. Sci., 2: 979.
- 60 Reaves, T. A., Cumming, R. and Hayward, J. N. (1982) Light- and electron-microscopic characterization of electrophysiologically identified, horseradish peoxidase-injected magnocellular neuroendocrine cells in goldfish preoptic nucleus. Neuroscience, 7: 1545-1557.
- 61 Murakami, M. (1964) Electronenmikroskopisch Untersuchungen am Nucleus Praeopticus der Krote (Bufo vulgaris formosus). Z. Zellforsch., 63: 208-225.
- 62 Halpern, M. (1972) Some connections of the telencephalon of the frog, *Rana pipiens*. An experimental study. Brain Behav. Evol., 6: 42-68.
- 63 Fujita, Y. and Urano, A. (1986) The afferent projection to the anterior part of the preoptic nucleus in Japanese toads, *Bufo japonicus*. Zool. Sci., 3: 677-686.
- 64 Jokura, Y. and Urano, A. (1985) Projections of luteinizing hormone-releasing hormone and vasotocin fibers to the anterior part of the preoptic nucleus in the toad, *Bufo japonicus*. Gen. Comp. Endocrinol., 60: 390-397.
- 65 Fujita, Y. and Urano, A. (1986) Immunohistochemical localization of thyrotropin-releasing hormone (TRH) in the hypothalamus of the toad, *Bufo japonicus*. Proc. Jpn. Soc. Com. Endocrinol., 1: 23.
- 66 Crosby, E. C. and Showers, M. J. C. (1969) Comparative anatomy of the preoptic and hypothalamic areas. In "The Hypothalamus". Ed. by W. Haymaker, E. Anderson and W. J. H. Nauta, Charles, C. Thomas, Springfield, pp. 61–135.
- 67 Hoogland, P. V., Ten Donkelaar, H. J. and Cruce, J. A. F. (1978) Efferent connections of the septal area in a lizard. Neurosci. Lett., 7: 61-65.
- 68 Raisman, G. (1966) The connections of the septum. Brain, 89: 317-348.
- 69 Swanson, L. W. and Cowan, W. M. (1976) Auto-

radiographic studies of the development and connections of the septal area in the rat. In "The Septal Nuclei. Adv. Behav. Biol., Vol. 20". Ed. by J. E. DeFrance, Plenum, New York, London, pp. 37-64.

- 70 Lisk, R. D. (1967) Sexual behavior: hormonal control. In "Neuroendocrinology. Vol.II". Ed. by L. Martini and W. F. Ganong, Academic Press, New York, pp. 197–239.
- 71 Malsbury, C. W. and Pfaff, D. W. (1974) Neural and hormonal determinants of mating behavior in adult male rats. A review. In "Limbic and Autonomic Nervous Systems Research". Ed. by L. V. DiCara, Plenum, New York, London, pp. 85-136.
- 72 Kicliter, E. and Northcutt, R. (1975) Ascending afferents to the telencephalon of Ranid frogs: an anterograde degeneration study. J. Comp. Neurol., 161: 239-254.
- 73 Ariens Kappers, C. U., Huber, G. C. and Crosby, E. C. (1960) The Comparative Anatomy of the Nervous System of Vertebrates, including Man. Vol. 2. Hafner, New York.
- 74 Zyo, K., Oki, T. and Ban, T. (1963) Experimental studies on the medial forebrain bundle, medial longitudial fasciculus and supraoptic decussations in the rabbit. Med. J. Osaka Univ., 13: 193-239.
- 75 Nauta, W. J. H. and Haymaker, W. (1969) Hypothalamic nuclei and fiber connections. In "The Hypothalamus". Ed. by W. Haymaker, E. Anderson and W. J. H. Nauta, Charles C. Thomas, Springfield, pp. 136–209.
- 76 Kokoros, J. J. (1972) Amphibian telencephalic efferents: an experimental study. Am. Zool., 12: 727.
- 77 Rubinson, K. (1968) Projection of the tectum opticum of the frog. Brain Behav. Evol., 1: 529– 561.
- 78 Rubinson, K. and Skiles, M. (1973) Efferent projections from the superior olivary nucleus in *Rana catesbeiana*. Anat. Rec., 175: 431–432.
- 79 Seki, T., Nakai, Y., Shioda, S., Mitsuma, T. and Kikuyama, S. (1983) Distribution of immunoreactive thyrotropin-releasing hormone in the forebrain and hypophysis of the bullfrog, *Rana catesbeiana*. Cell Tissue Res., 233: 507-516.
- 80 Buijs, R. M. (1985) Extrahypothalamic pathways of a neurosecretory system: their role in neurotransmission. In "Neurosecretion and the Biology of Neuropeptides". Ed. by H. Kobayashi, H. A. Bern and A. Urano, Jpn. Sci. Soc. Press, Tokyo, pp. 279-286.
- 81 Silverman, A. J. (1983) Luteinizing hormone releasing hormone (LHRH) synapses in the diagonal band (DBB) and preoptic area (POA) of the guinea pig. Soc. Neurosci. Abstr., 9: 1182.
- 82 Scalia, F., Halpern, M., Knapp, H. and Riss, W.

(1968) The efferent connections of the olfactory bulb in the frog: a study of degenerating unmyelinated fibers. J. Anat., 103: 245-262.

- 83 Northcutt, R. G. and Royce, G. J. (1975) Olfactory bulb projections in the bullfrog, *R. catesbeiana* Shaw. J. Morphol., 145: 251–268.
- 84 Scalia, F. (1972) The projection of the accessory olfactory bulb in the frog. Brain Res., 36: 409-411.
- 85 Wirsig, C. R. and Getchell, T. (1986) Amphibian terminal nerve: distribution revealed by LHRH and AChE markers. Brain Res., **385**: 10–21.
- 86 Jokura, Y. and Urano, A. (1986) Extrahypothalamic projection of luteinizing hormone-releasing hormone fibers in the brain of the toad, *Bufo japonicus*. Gen. Comp. Endocrinol., 62: 80-88.
- Jokura, Y. and Urano, A. (1987) Extrahypothalamic projection of immunoreactive vasotocin fibers in the brain of the toad, *Bufo japonicus*. Zool. Sci., 4: 675–681.
- 88 Muller, C. H. and Licht, P. (1980) Gonadotropin specificity of androgen secretion by amphibian testes. Gen. Comp. Endocrinol., 42: 365–377.
- 89 Licht, P., McCreery, B. R., Barnes, R. and Pang, R. (1983) Seasonal and stress related changes in plasma gonadotropins, sex steroids, and corticosterone in the bullfrog, *Rana catesbeiana*. Gen. Comp. Endocrinol., 50: 124-145.
- 90 Daniels, E. and Licht, P. (1980) Effects of gonadotropin-releasing hormone on the levels of plasma gonadotropins (FSH and LH) in the bullfrog, *Rana catesbeiana*. Gen. Comp. Endocrinol., 42: 455– 463.
- 91 Fujita, Y. and Urano, A. (1987) Effects of LHRH on EEG activity and plasma androgen levels in Japanese toads. Zool. Sci., 4: 1079.
- 92 King, J. A. and Millar, R. P. (1979) Hypothalamic luteinizing hormone-releasing hormone content in relation to the seasonal reproductive cycle of *Xenopus laevis*. Gen. Comp. Endocrinol., 39: 309–312.
- 93 Yoneyama, H., Ishii, S., Yamamoto, K. and Kikuyama, S. (1984) Plasma prolactin levels of *Bufo japonicus* before, during and after breeding in the pond. Zool. Sci., 1 :969.
- 94 Itoh, M. and Ishii, S. (1986) Establishment of radioimmunoassays of toad (*Bufo japonicus*) gonadotropins and annual cycle of plasma FSH and LH levels in a wild toad population. Zool. Sci., 3: 1078.
- 95 Tasaki, Y., Inoue, M. and Ishii, S. (1986) Annual cycle of plasma thyroid hormone levels in the toad, *Bufo japonicus*. Gen. Comp. Endocrinol., 62: 404– 410.
- 96 Jolivet-Jaudet, G., Inoue, M., Takada, K. and Ishii, S. (1984) Circannual changes in corticosterone plasma levels and binding of corticosterone to plasma in *Bufo japonicus formosus*. Zool. Sci., 1: 317-325.

- 97 Jolivet-Jaudet, G., Inoue, M., Takada, K. and Ishii, S. (1984) Circannual changes in plasma aldosterone levels in *Bufo japonicus formosus*. Gen. Comp. Endocrinol., 53: 163-167.
- 98 Inoue, M., Takada, K. and Ishii, S. (1982) Seasonal changes in plasma androgen concentrations and reproductive behavior in toads. Zool. Mag., 91: 568.
- 99 Dierickx, K. and Van Meirvenne, N. (1961) Karyometric studies of the preoptic nucleus of *Rana temporaria*. Gen. Comp. Endocrinol., 1: 51-58.
- 100 Dierickx, K. and Vandesande, F. (1965) The magnocellular preoptic nuclei and the reproduction in *Rana temporaria*. Z. Zellforsch., 68: 190–193.
- 101 Nottebohm, F. (1981) A brain for all seasons: Cyclical anatomical changes in song control nuclei of the canary brain. Science, 214: 1368–1370.
- 102 DeVoogd, T. and Nottebohm, F. (1981) Gonadal hormones induce dendritic growth in the adult avian brain. Science, 214: 202-204.
- 103 Jokura, Y. and Urano, A. (1985) An Immunohistochemical study of seasonal changes in luteinizing hormone-releasing hormone and vasotocin in the forebrain and the neurohypophysis of the toad, *Bufo japonicus*. Gen. Comp. Endocrinol., 59: 238-245.
- 104 Kurz, E. M., Sengelaub, D. R. and Arnold, A. P. (1986) Androgens regulate the dendritic length of mammalian motoneurons in adult-hood. Science, 232: 395-398.
- 105 Fujita, Y., Jokura, Y., Takami, S. and Urano, A. (1987) Effects of castration on volumes of the preoptic nucleus and the amygdala and on immunoreactivity of LHRH fibers in the brain of the toad *Bufo japonicus*. Gen. Comp. Endocrinol., 68: 278-285.
- 106 Baker, J. L. (1977) Physiological roles of peptides in the nervous system. In "Peptides in Neurobiology". Ed. by H. Gainer, Plenum Press, New York, pp. 295-343.
- 107 Jan, L. Y., Jan, Y. N. and Brownfield, M. S. (1980) Peptidergic transmitters in synaptic boutons of sympathetic ganglia. Nature, 288: 380–382.
- 108 Inenaga, K. and Yamashita, H. (1986) Excitation of neurones in the rat paraventricular nucleus in vitro by vasopressin and oxytocin. J. Physiol., 370: 165-180.
- 109 Sugita, R. and Urano, A. (1986) Responses of magnocellular neurons in *in vitro* eel preoptic nucleus (PONmg) to acetylcholine, catecholamines, vasotocin, isotocin, angiotensin, and Na⁺. Zool. Sci., 3: 1081.
- 110 Urano, A. and Ishihara, K. [eds] (1987) Biology of Toad. Shokabo, Tokyo.