



Title	Relation between excessive CO2 expiration and performance in high-intensity exercise
Author(s)	Yano, T.; Yunoki, T.; Ogata, H.; Mastuura, R.
Citation	Biology of Sport, 22(2), 117-124
Issue Date	2005-06
Doc URL	http://hdl.handle.net/2115/44020
Type	article
File Information	BS22-2_117-124.pdf



[Instructions for use](#)

RELATION BETWEEN EXCESSIVE CO₂ EXPIRATION AND PERFORMANCE IN HIGH-INTENSITY EXERCISE

T.Yano, T.Yunoki, H.Ogata, R.Matsuura

Dept. of Exercise Physiology, Graduate School of Education, Hokkaido University, Kita-ku Sapporo, Japan

Abstract. This study examined the relation between excessive CO₂ expiration and exercise performance in exercise exhausted for 1-2 min. First, the Wingate test was conducted to determine the exercise intensity of the performance test. The exercise intensity was set at 80% of the power exerted during the last 5 s of the Wingate test so that subjects could continue for 1-2 min. Total work, which was the value of the work rate times the performed duration was used as an index of the performance level. The excessive CO₂ expiration rate ($\dot{V}CO_{2\text{excess}}$) was defined as the difference between CO₂ output and O₂ uptake. Integration of $\dot{V}CO_{2\text{excess}}$ from the start of exercise to the zero level of $\dot{V}CO_{2\text{excess}}$ observed at around 10 min in recovery was defined as the amount of excessive CO₂ expiration (CO₂excess). The performance was associated with the peak value of blood lactate (peak La) obtained after exercise ($r=0.744$). The performance was also associated with CO₂excess ($r=0.848$). It is likely that CO₂excess is a better physiological index for assessing the performance in exercise exhausted for 1-2 min than peak La.

(Biol.Sport 22:117-124, 2005)

Key words: CO₂ excess-intensive exercise-blood lactate-performance

Introduction

Performance in exercise exhausted for 1-2 min is associated with peak blood lactate after exercise [4,7]. This was interpreted as relating to the amount of energy derived from lactic formation (lactic capacity). However, a recent study demonstrated that a peak blood lactate after exercise does not reflect the amount of lactate formation [8]. Therefore, the relation between the performance and amount of lactate formation has not been verified.

The lactic acid formed in exercise is hydrolyzed. Within the physiological range

Reprint request to: Prof. Tokuo Yano, Dept. of Exercise Physiology, Graduate School of Education, Hokkaido University, Kita-ku Sapporo, Japan

Tel.: 011-706-5090; Fax: 011-552-1347; E-mail: yano@edu.hokudai.ac.jp

in pH, the produced hydrogen ion is equivalent to the formed lactate ion but is buffered in the body. Therefore, the actual increase of hydrogen ion is very slight. This buffer action in the active muscle is mainly made by a non-bicarbonate buffer system and absorption of hydrogen ion accompanied by the resolution of phosphocreatine (CrP) [2]. After stopping exercise, lactic acid is spread throughout the entire body and consequently is buffered by the entire body rather than active muscle. Since CrP is recovered after exercise, hydrogen ion absorbed by the resolution of CrP is released. Therefore, the action of CrP is invalid when buffer action from exercise to recovery is considered. As the buffer action in the extracellular fluid is mainly owing to the bicarbonate buffer system, buffer action after exercise is mainly dependent on the bicarbonate buffer system, which can be observed as excessive CO₂ expiration [9]. Thus, through the shift of lactic acid from the muscle in exercise to the entire body during recovery, buffer action in exercise could be assessed by the excessive CO₂ expiration (CO₂excess).

Lactic acid produced during intense exercise is involved in the development of fatigue [3]. A high ability of the active muscles to release lactic acid and the high buffer capacity of muscle could prevent muscle functionality and delay the onset of fatigue. Lactic acid released from the active muscle in exercise induces CO₂excess. Lactic acid accumulated in the muscle in exercise gradually spreads throughout the entire body during recovery. This also causes the CO₂ excess. As the buffer capacity of active muscle and the release ability of lactic acid from active muscle have an advantage in the performance in exercise exhausted for 1-2 min, the amount of CO₂excess in exercise and during recovery could be used to assess performance.

In the present study, therefore, we examined whether the performance in the exercise exhausted for 1-2 min is associated with the amount of CO₂ excess.

Material and Methods

Table 1

Characteristics of the subjects and results of the Wingate test

	Age (years)	Height (cm)	Weight (kg)	Peak power (watts)	Average power (watts)
Mean	21	172	62	695	538
SD	0.9	4.1	6.5	130	82

Thirteen men belonging to a sports club in university participated in the present experiments. Physical characteristics of the subjects are listed in Table 1. Each subject was informed of the purpose of the study, the experimental procedure, and the risks associated with the experiments before consent was obtained. Furthermore, they were instructed to abstain from heavy training on the day before the experiments and rest for one hour prior to the start of the experiments.

A cycle ergometer (Powermax-VII, Combi) was used in the experiments. On the first day, each subject performed a 30-s maximal exercise test at a load (kp) corresponding to 7.5% x body weight (Wingate test). During the Wingate test, the power output (watts) was calculated from the imposed load and cycling rate (rpm) averaged for every 5 s. A few days later, a short-term intensive exercise test (SIET) was performed at a work load corresponding to 80% of the power exerted during the last 5 s of the Wingate test [11]. Each subject, after resting on the cycle ergometer for 5 min, performed SIET until he could no longer maintain 90 rpm. After the end of SIET, each subject sat on a chair for a 30-min recovery period.

Ventilation (\dot{V}_E), O₂ uptake (\dot{V}_{O_2}), CO₂ output (\dot{V}_{CO_2}) and end tidal CO₂ pressure (PETCO₂) were measured breath-by-breath using a respiratory gas analyzer (Aeromonitor AE-280S, Minato) throughout the SIET. These data were averaged and were outputted for each 20-s period. \dot{V}_E was measured by a hot-wire flow meter that was fixed to the front of the subject's gas mask. The flow meter was calibrated with a syringe of known volume (2.0 l). Sample gas was drawn continuously from the front of the gas mask (220 ml·min⁻¹) for determination of the fractional concentrations of O₂ and CO₂ by a zirconium sensor and an infrared absorption analyzer, respectively. The gas analyzer was calibrated by set standard gases. Twenty-five microliters of blood were sampled from a fingertip using a capillary tube at rest before exercise, at 0, 3, 6, 10, 20, and 30 min after the end of exercise. The lactate concentration (La) in the sampled blood was determined by an automatic lactate analyzer (1500 sport, YSI). The analyzer was calibrated with a standard liquid (5 mM of lactate, YSI).

Excessive CO₂ expiration per unit time ($\dot{V}_{CO_2\text{excess}}$) was calculated by subtracting the \dot{V}_{O_2} values from the \dot{V}_{CO_2} values during the SIET [10,11]. The $\dot{V}_{CO_2\text{excess}}$ was integrated from the start of exercise to the zero level of $\dot{V}_{O_2\text{excess}}$, and this was defined as the total excessive CO₂ expiration (CO₂excess).

The strength of the relation between dependent and independent variables was expressed by a single correlation coefficient of Pearson. A P value of 0.05 or less was significant. Results were expressed as mean ± standard deviation (SD).

Results

Table 2

Blood lactate (mM) at rest and during recovery

	Rest	Recovery					
		0 min	3 min	6 min	10 min	20 min	30 min
Mean	0.97	8.03	9.69	9.55	8.46	6.54	4.69
SD	0.22	1.96	1.55	1.77	1.82	1.78	1.56

Table 2 shows La at rest and during recovery. La immediately after the SIET increased by 77% of peak La. The peak La was observed at 3 or 6 min after SIET. Then La decreased. The work rate given for the performance test of SIET ranged from 279 watts to 453 watts. The duration of SIET was almost the same (70 to 110 s). Total work, which was obtained by multiplying the duration and a given work rate for SIET ranged from 22320 to 37510 joule. There was a significant relation between total work and peak La after SIET (Fig. 1, $r=0.744$).






Fig. 1
Relation between peak blood lactate (La) after exercise and total work performed in exercise

**Fig. 2**

Relation between amount of excessive CO₂ expiration (CO₂excess) and total work performed in exercise

**Fig. 3**

Change in excessive CO₂ expiration per unit time ($\dot{V}CO_2$ excess) from the start of

exercise to recovery phase

There was a significant relation between total work and CO_2 excess (Fig. 2, $r=0.848$). As shown in Fig. 3, $\ddagger\text{CO}_2$ excess did not increase but decreased at the onset of SIET. Then $\ddagger\text{CO}_2$ excess started to increase before the end of SIET and showed peak at around at 1.5 min after SIET. $\ddagger\text{CO}_2$ excess became zero level at around 10 min. PETCO_2 increased in SIET. PETCO_2 started to decrease during recovery and dropped below resting level (Fig. 4).




Fig. 4

Change in end tidal CO_2 pressure (PETCO_2) from the start of exercise to recovery phase

Discussion

$\ddagger\text{O}_2$ excess showed negative values in SIET. This could be due to the rise of CO_2 pressure. As venous CO_2 pressure increases in intensive exercise [6], this would result in the increase of bicarbonate ion not only in the active muscle but also in venous blood. As a result, $\ddagger\text{CO}_2$ excess can be depressed. At the end of exercise, lactic acid is thought to move from the muscle to extra cellular fluid, since the La immediately after SIET increased. This suggests an increase in $\ddagger\text{CO}_2$ excess.

However, the rise of CO₂ pressure could mask the effect of the lactic acid shift. The mask was released after exercise due to the decrease in CO₂ pressure. Therefore, even if lactate is not shifted into the extra cellular fluid, ‡CO₂excess is increased. Afterwards, PETCO₂ reached below the resting level. As this is induced by the hyperventilation, ‡ CO₂excess continues. However, the hyperventilation leads to overestimation of CO₂excess.

CrP is synthesized during recovery. This causes the increase of hydrogen ion in the muscle. Although lactic acid is spread throughout the entire body after exercise, this does not always mean that all hydrogen ion spreads throughout the entire body. However, there is a mechanism to exchange Na⁺ and H⁺ [2,5]. This may well work to help the buffer action in the muscle by spreading the hydrogen throughout the entire body.

Entire body buffer action consists of a non-bicarbonate buffer system and bicarbonate buffer system [2,9]. The former system does not induce CO₂excess. This leads to underestimation of the amount of lactate formation by CO₂excess. However, this underestimation may be counteracted by the overestimation due to hyperventilation.

As mentioned in Introduction, peak La is an index of lactic capacity. This index was associated with the performance. Since it is considered that the performance exhausted for 1-2 min is mainly determined of lactic capacity, the obtained relation between the performance and peak La is well understandable. We postulated that CO₂excess should be a new physiological index for assessing the performance exhausted for 1-2 min. The CO₂excess was also associated with the performance. Furthermore, correlation coefficient of this relation was higher than that obtained in peak La. Although there is a limitation in the use of the CO₂excess, it is likely that CO₂excess is a better physiological index for assessing the performance in exercise exhausted for 1-2 min than peak La.

References

1. Bret C., L.Messonier, J.M.Nouck, H.Freund, A.B.Dufour, J.R.Lacour (2003) Differences in lactate exchange and removal abilities in athletes specialized in different track running events (100 to 1500 m). *Int.J.Sports Med.* 24:108-113
2. Cerretelli P., M.Samaja (2003) Acid-base balance at exercise normoxia and in chronic hypoxia. Revisiting the "lactate paradox". *Eur.J.Appl.Physiol.* 90:431-448
3. Fitts R.H. (1994) Cellular mechanisms of muscle fatigue. *Physiol.Rev.* 74:49-94
4. Fujitsuka N., T.Yamamoto, T.Ohkuwa, M.Saito, M.Miyamura (1982) Peak blood lactate after short periods of maximal treadmill running. *Eur.J.Appl.Physiol.* 48:289-296
5. Juel C., C.Klarskov, J.J.Nielsen, P.Krustrup, M.Mohr, J.Bangasbo (2004) Effect of

high-intensity intermittent training on lactate and H⁺ release from human skeletal muscle. *Am.J.Physiol.Endocrinol.Metab.* 286:E245-E251

6. Kowalchuk J.M., G.J.F.Heigenhauser, M.I.Lindinger, J.R.Satton, N.L.Jones (1988) Factors influencing hydrogen ion concentration in muscle after intense exercise. *J.Appl.Physiol.* 65:2080-2089

7. Lacour J.R., E.Bouvat, J.C.Barthelemy (1990) Post-competition blood lactate concentrations as indicators of anaerobic energy expenditure during 400-m and 800-m races. *Eur.J.Appl.Physiol.* 61:172-176

8. Medbo J.I., K.Toska (2001) Lactate release, concentration in blood, and apparent distribution volume after intense bicycling. *Jpn.J.Physiol.* 51:303-312

9. Yano T (1987) A theoretical approach to excessive CO₂ expiration due to lactate production in exercise. *Jpn.J.Physiol.* 37:937-940

10. Yunoki T., M.Horiuchi, T.Yano (1999) Kinetics of excess CO₂ output during and after intensive exercise. *Jpn.J.Physiol.* 49:139-144

11. Yunoki T., M.Horiuch, T.Yano (2000) Kinetics of excess CO₂ output during and after intensive exercise in sprinters and long distance runners. *Jpn.J.Physiol.* 50:199-205

Accepted for publication 10.12.2004