



Title	Water-soluble organic compounds in PM2.5 and size-segregated aerosols over Mount Tai in North China Plain
Author(s)	Wang, Gehui; Kawamura, Kimitaka; Umemoto, Nobuhiko; Xie, Mingjie; Hu, Shuyuan; Wang, Zifa
Citation	Journal of Geophysical Research, Atmospheres, 114, D19208 https://doi.org/10.1029/2008JD011390
Issue Date	2009-10-10
Doc URL	http://hdl.handle.net/2115/44585
Rights	Wang, G., K. Kawamura, N. Umemoto, M. Xie, S. Hu, and Z. Wang, Water-soluble organic compounds in PM2.5 and size-segregated aerosols over Mount Tai in North China Plain, J. Geophys. Res., 114, D19208, 2009. Copyright 2009 American Geophysical Union. Reproduced/modified by permission of American Geophysical Union.
Type	article
File Information	JGRA114_D19208.pdf



[Instructions for use](#)



Water-soluble organic compounds in PM_{2.5} and size-segregated aerosols over Mount Tai in North China Plain

Gehui Wang,^{1,2,3} Kimitaka Kawamura,¹ Nobuhiko Umemoto,¹ Mingjie Xie,³ Shuyuan Hu,³ and Zifa Wang⁴

Received 31 October 2008; revised 14 May 2009; accepted 6 July 2009; published 10 October 2009.

[1] Daytime and nighttime PM_{2.5} samples were collected at the summit of Mount Tai (1534 m) located in North China Plain during a week in 2006 summer. Size-segregated aerosol particles were also collected using an eight-stage impactor during the same period. Samples were analyzed for various water-soluble organic compounds using GC/FID and GC/MS techniques. Among the species identified in PM_{2.5} samples, dicarboxylic acids (C₂–C₁₁) were found as the most abundant compound class, followed by ketocarboxylic acids, saccharides, polyols and polyacids, and dicarbonyls. Daytime concentrations of most compounds were found to be 2–3 times higher than in nighttime. Such a diurnal variation was first interpreted by the depressed transport of pollutants in nighttime from the lowlands to the mountaintop owing to the decreased heights of planetary boundary layer, and second by the photochemical production in daytime. The diurnal variation trends of secondary organic aerosols (SOA) such as diacids at the mountain site are the same as those on lowlands, but the diurnal patterns of primary organic aerosols (POA) on the mountaintop are in contrast to those on lowlands, where POA such as saccharides and polyols are usually higher in nighttime owing to the accumulation within inversion layer developed. The eight-stage impactor samples showed bimodal distributions of diacids and related compounds peaking at size ranges of 0.70–1.1 μm and 5.8–9.0 μm. In the present study, water-soluble organics in the fine mode are largely originated from biomass burning and/or photooxidation of gaseous precursors and the subsequent adsorption on the preexisting particles, whereas those in the coarse mode are mainly derived from suspended soil particles and pollens and in part via the hygroscopic growth of fine particles and formation of cloud/fog droplets.

Citation: Wang, G., K. Kawamura, N. Umemoto, M. Xie, S. Hu, and Z. Wang (2009), Water-soluble organic compounds in PM_{2.5} and size-segregated aerosols over Mount Tai in North China Plain, *J. Geophys. Res.*, 114, D19208, doi:10.1029/2008JD011390.

1. Introduction

[2] Tropospheric aerosols are deeply involved in global radiative balance [Seinfeld and Pandis, 1998] and pollution transport [Akimoto, 2003; Wilkening *et al.*, 2000]. Radiative effects of fine particles in the troposphere are much stronger than coarse particles owing to their more active response to radiation [Seinfeld and Pandis, 1998]. Because water-soluble organic compounds comprise a substantial fraction of the tropospheric organic aerosols, they significantly alter the hygroscopic properties of particles and thus affect the radiative forcing of the atmosphere and climate by acting as

cloud condensation nuclei (CCN) [Hemming and Seinfeld, 2001]. In particular, some water-soluble organics such as carboxylic acids and alcohols can act as surfactants in aerosols [Latif and Brimblecombe, 2004], and have an influence on hygroscopic properties of the particles. They also increase the solubility of toxic pollutants in aerosol particles and fog/rain droplets.

[3] Dicarboxylic acids (DCA) are a major water-soluble organic compound class in the tropospheric aerosols from continental urban sites [Kawamura and Yasui, 2005; Wang *et al.*, 2002; Yao *et al.*, 2004] to remote ocean [Kawamura and Gagosian, 1987; Kawamura and Sakaguchi, 1999]. They can lower the surface tension of water in aerosols [Hyvarinen *et al.*, 2006], and as such enhance the hygroscopic growth of the particles [Zhang *et al.*, 2004]. Although DCAs can be directly emitted from combustion sources [Kawamura and Kaplan, 1987], photochemical oxidation of gaseous organic precursors is more important [Claeys *et al.*, 2004; Kawamura *et al.*, 2005]. Lower molecular weight DCAs ($\leq C_6$) are produced by photooxidation of anthropogenic volatile organic compounds (VOC) whereas higher molecular weight DCAs ($> C_6$) such as azelaic acid (C₉) are

¹Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan.

²State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an, China.

³School of the Environment, State Key Laboratory of Pollution Control and Resources Reuse, Nanjing University, Nanjing, China.

⁴Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing, China.




Figure 1. A map of eastern China and surroundings with a sampling location at Mount Tai (36.26°N , 117.11°E).

produced by oxidation of unsaturated fatty acids from marine and terrestrial plants [Kawamura and Gagosian, 1987; Nepotchaykh and Ariya, 2002] and human cooking activities [Robinson et al., 2006].

[4] Saccharides are another important class of water-soluble organic compounds in atmospheric aerosols and are reported from many geographic regions including Amazonia, Brazil [Graham et al., 2002], the western North Pacific [Mochida et al., 2003], and Santiago, Chile [Simoneit et al., 1993]. The dominant saccharides include levoglucosan, α - and β -glucose, α - and β -fructose, sucrose, and trehalose, with lesser amount of inositol, α - and β -mannose, α - and β -xylose, and α - and β -galactose. Levoglucosan has been used as a specific biomass burning tracer, which is produced by the pyrolysis of cellulose. In addition, saccharide polyols are also found in the atmosphere [Wang et al., 2006a], including sorbitol, xylitol, mannitol, arabitol, and others. Saccharides and polyols in aerosols are primarily derived from biomass burning [Graham et al., 2002; Simoneit et al., 1999], pollen and fungal metabolism [Simoneit et al., 2004b, 2004c].

[5] Tropospheric chemistry of water-soluble organic compounds is a challenging issue because their temporal and spatial distributions are sometimes region-specific [Huang et al., 2005; Wang et al., 2006a]. Although they have been studied in many regions including the urban [Kawamura and Ikushima, 1993; Kawamura and Kaplan, 1987] and remote marine atmosphere [Kawamura and Sakaguchi, 1999], observations of tropospheric aerosols over high mountains were rarely conducted [Decesari et al., 2005; Limbeck and Puxbaum, 1999]. The mountain geography and meteorology may have a strong influence on tropospheric chemistry, which may be different from urban

areas on flat lowlands. In the high mountains, boundary layers become low at night and the mountaintop is often in the free troposphere. Further, ambient temperature significantly drops at night in high mountains and thus relative humidity increases, potentially causing unique behaviors of water-soluble organic compounds.

[6] In this study, we collected summertime tropospheric aerosols at the summit of Mount Tai in North China Plain (NCP), where agricultural activity such as wheat harvest is maximized in early summer and subsequent field burning of agricultural wastes is operated. We analyzed both $\text{PM}_{2.5}$ aerosols and eight-stage size-segregated particles for various water-soluble organic species using GC and GC/MS. Here we report molecular compositions, concentrations and size distributions of dicarboxylic acids, ketocarboxylic acids, α -dicarbonyls, sugars, polyols and polyacids in the Mount Tai aerosols. The detailed size distributions of saccharides are, for the first time, reported in the mountain aerosols.

2. Experiment

2.1. Sampling Site and Procedures

[7] Mount Tai is the highest peak (elevation, 1534 m) in central NCP, which faces to the East China Sea, Korean Peninsula and Japanese Islands (Figure 1). Mount Tai is surrounded by the economically most developed regions in China, where air quality is very poor [Akimoto, 2003; Gao et al., 2005; Richter et al., 2005]. The aerosol sampling was conducted at a meteorological observatory at the mountaintop, which is far away from historical temples and local restaurants on the mountain. Thus, emissions from local sources are minimal [Gao et al., 2005]. Using a high-

Table 1. Meteorological Data During the Mount Tai Sampling Period^a

	Daytime			Nighttime		
	0700–1700 Local Time			1900–0500 Local Time		
	Range	Mean	Std	Range	Mean	Std
Temperature, °C	21.0–35.0	27.7	3.4	19.2–31.5	23.6	2.7
Relative humidity, %	47–96	69	13	40–97	85	12
Pressure, hPa	986–994	991	2.1	987–994	991	1.9
Wind speed, m s ⁻¹	4–49	22	10	1–65	16	11
Wind direction, deg	72–291	162	63	20–323	184	82

^aSampling period is 22–28 June 2006. All the meteorological parameters were measured hourly. Std denotes standard deviation.

volume air sampler (Andersen GM2000, USA, flow rate of air was 1.13 m³ min⁻¹) and prebaked (450°C, 6 h) quartz fiber filters (20 × 25 cm²), PM_{2.5} aerosol samples were collected on a day/night basis (0700–1700 local time (LT) for daytime and 1900–0500 LT for nighttime) during a week (22–28 June 2006) on the square in front of the Meteorological Station.

[8] An eight-stage impactor sampler (Andersen 20–800, USA, flow rate: 28.3 L min⁻¹) was used with glass fiber filters (ϕ 90 mm, precombusted at 450°C for 6 h) for size-segregated aerosol sampling. Cutoff sizes of each stage were > 9.0, 9.0–5.8, 5.8–4.7, 4.7–3.3, 3.3–2.1, 2.1–1.1, 1.1–0.70, 0.7–0.40, and <0.40 μ m. Particles (>9.0 μ m) impacted into a pre-separation aluminum vessel were collected from the container using a brush after sampling. However, those particles were not analyzed owing to the potential collection loss by the brush. The size-segregated sampling was continued for 1 week to obtain enough materials.

[9] After the sample collection, filters and field blanks were taken in a prebaked (450°C, 6 h) glass jar with a Teflon-lined screw cap, transported to the laboratory in Sapporo and stored at –20°C prior to analysis. Meteorological parameters, hourly monitored during the campaign, are shown in Table 1.

2.2. Sample Extraction, Derivatization, and Determination

[10] In this study, we analyzed 14 PM_{2.5} samples and eight size-segregated samples for various organic compounds using two methodologies as follows.

[11] For dicarboxylic acids, an aliquot of the filter was extracted with pure water, and the extracts were filtered through a Pasteur pipette packed with quartz wool to remove filter debris and insoluble particles. The extracts were concentrated to nearly dryness using a rotary evaporator under vacuum, and then reacted with 14% BF₃/*n*-butanol at 100°C for 30 min to convert the COOH and CHO groups to butyl esters and dibutoxy acetals, respectively. The derivatives were extracted with *n*-hexane and concentrated to 100 μ L prior to GC injection. Water-extractable diacids, ketocarboxylic acids and dicarbonyls were quantified by GC/FID. More details of sample extraction, derivatization and GC and GC/MS measurements are described elsewhere [Kawamura and Yasui, 2005; Wang et al., 2006a].

[12] For saccharides, polyols and polyacids, another aliquot of the filter was extracted with a mixture of methanol and dichloromethane (v/v, 1/2) under ultra sonication. The extracts were concentrated to dryness using a

rotary evaporator and nitrogen blow-down system, and then reacted with *N,O*-bis-(trimethylsilyl)trifluoroacetamide (BSTFA) at 70°C for 3 h to convert the OH and COOH groups into trimethylsilyl (TMS) ethers and esters, respectively. Saccharides, polyols and polyacids were determined by GC/MS. More details are described elsewhere [Wang et al., 2006a].

[13] Most of the target compounds were undetectable in the field blanks, except for glycerol and oxalic acid. However, their amounts were less than 2% of the actual samples. Recoveries of all the compounds determined were better than 80%. Data reported here are corrected for the field blanks but not corrected for the recoveries. The mass of aerosol particles with a diameter less than 2.5 μ m in the size-segregated samples was calculated to be 80 μ g m⁻³ using a standard method described by the sampler manufacturer. It accounts for 80% of the mass collected by the high-volume PM_{2.5} sampler, suggesting that particle bounce-off from the upper stage to the lower stage was not serious. A strong linear correlation was found for the major species determined in the PM_{2.5} samples and in the summed size fractions (PM_{2.1}) of the size-segregated samples (Figure 2). The intercomparison suggests a good consistency between two data sets. But concentrations of the major species obtained in the size-segregated aerosols are 5–20% higher than those obtained by the PM_{2.5} samples. Unlike the size-segregated aerosol sampler, the PM_{2.5} sampler collect aerosols by filtration, thus compound losses by evaporation may occur owing to the pressure drop in the sampler. Such a negative artifact has been reported [Katz and Chan, 1980; McMurtry, 2000; Peters and Seifert, 1980].

3. Results and Discussion

3.1. Daytime and Nighttime Concentrations of Organic Compound Classes

[14] Five organic compound classes were determined in the PM_{2.5} samples (Table 2), among which DCAs were found




Figure 2. An intercomparison of concentrations of the major compounds determined in the PM_{2.5} samples and in the PM_{2.1} fraction of the eight-stage samples (the PM_{2.5} concentrations are the averages of all the day and night samples, while the PM_{2.1} concentrations are the sum of the compounds in the stages of < 0.4, 0.4–0.7, 0.7–1.1, and 1.1–2.1 μ m).

Table 2. Concentrations of Water-Soluble Organic Compounds in PM_{2.5} Over Mount Tai^a

	Daytime (n = 7)				Nighttime (n = 7)			
	Min	Max	Mean	Std	Min	Max	Mean	Std
<i>Dicarboxylic Acids</i>								
Oxalic (C ₂)	378	623	455	87	83	284	158	82
Malonic (C ₃)	64	103	77	13	13	43	29	11
Succinic (C ₄)	67	101	84	13	17	48	30	13
Glutaric (C ₅)	18	32	23	6.1	3.7	10	6.8	2.8
Adipic (C ₆)	7.6	13	9.7	1.7	1.5	5.1	3.3	1.3
Pimelic (C ₇)	5.7	14	8.5	2.8	BDL	4.6	2.4	2.1
Suberic (C ₈)	2.9	7.8	5.3	1.8	BDL	2.8	1.7	1.1
Azelaic (C ₉)	9.4	32	21	8.2	1.4	13	6.0	4.1
Decanedioic (C ₁₀)	0.5	1.8	1.0	0.5	BDL	0.7	0.3	0.3
Undecanedioic (C ₁₁)	2.2	6.7	4.0	1.4	0.5	4.1	1.9	1.2
Methylsuccinic (iC ₅)	4.9	8.3	7.0	1.2	0.8	3.6	2.1	1.4
Methylglutaric (iC ₆)	1.5	2.4	2.0	0.3	BDL	0.9	0.4	0.4
Maleic (M)	4.1	7.2	5.4	1.2	0.7	5.9	2.8	2.1
Fumaric (F)	2.8	5.0	3.7	0.7	BDL	2.1	1.1	0.9
Methylmaleic (mM)	7.4	18	12	3.3	0.4	8.1	4.1	3.8
Phthalic (Ph)	22	37	28	5.0	5.5	16	11	4.2
Iso-Phthalic (i-Ph)	1.5	2.7	2.0	0.4	0.2	1.3	0.7	0.4
Tere-phthalic (t-Ph)	0.6	1.2	0.9	0.2	BDL	0.3	0.1	0.2
4-Oxopimelic (kC ₇)	BDL	10.7	6.4	3.4	BDL	4.3	1.9	1.7
Subtotal	642	1025	756	139	140	434	263	123
<i>Ketocarboxylic Acids</i>								
Glyoxylic (ωC ₂)	47	84	61	14	6.9	28	17	11
3-Oxopropanoic (ωC ₃)	5.4	14	7.5	2.8	0.9	5.1	2.9	1.8
4-Oxobutanoic (ωC ₄)	4.0	19	9.2	4.8	2.0	6.8	4.0	1.9
9-Oxopelargonic (ωC ₉)	BDL	3.4	2.2	1.1	0.3	1.9	1.0	0.7
Pyruvic (Pyr)	18	43	27	8.3	2.9	14	8.2	5.4
Subtotal	82	163	107	29	13	54	33	21
<i>α-Dicarbonyls</i>								
Methylglyoxal (mGly)	13	47	24	11	0.4	14	7.1	6.4
Glyoxal (Gly)	5.5	18	9.5	4.0	0.7	6.0	3.0	2.4
Subtotal	19	65	33	15	1	20	10	9
<i>Sugars</i>								
Levogluconan	21	123	43	36	1.6	151	36	58
Arabitol	0.9	5.9	3.2	1.9	0.2	1.7	0.8	0.6
Fructose	2.8	6.2	4.7	1.1	0.6	4.6	1.5	1.5
Glucose	7.2	13	9.2	2.1	1.8	4.8	3.2	1.2
Mannitol	1.5	11	6.0	3.6	0.2	2.5	0.9	0.9
Inositol	0.5	2.2	1.5	0.6	0.1	1.8	0.5	0.6
Sucrose	0.9	2.9	1.7	0.7	0.4	5.5	2.2	2.1
Trehalose	1.0	4.4	2.4	1.2	0.3	1.1	0.7	0.3
Subtotal	40	149	72	36	7	160	46	58
<i>Polyols and Polyacids</i>								
Glycerol	8.0	23	17	6.2	8.5	21	14	4.9
Glyceric acid	9.6	17	12	2.4	1.7	6.8	3.7	2.4
Malic acid	3.7	45	28	13	3.5	25	11	9.3
Tartaric acid	1.5	37	7.4	13	BDL	4.9	2.3	2.1
Citric acid	BDL	1.8	0.3	0.7	BDL	1.1	0.4	0.5
Subtotal	54	75	64	9	20	55	32	17

^aSampling period is 22–28 June 2006. Unit is ng m⁻³. BDL, below detection limit; Min, minimum; Max, maximum; and Std, standard deviation.

as the most abundant compound class (756 ± 139 ng m⁻³ in daytime versus 263 ± 123 ng m⁻³ in nighttime), followed by ketocarboxylic acids (107 ± 29 versus 33 ± 21 ng m⁻³), saccharides (72 ± 36 versus 46 ± 58 ng m⁻³), polyols and polyacids (64 ± 9 versus 32 ± 17 ng m⁻³), and dicarbonyls (33 ± 15 versus 10 ± 9 ng m⁻³). Interestingly, daytime concentrations are 2–3 times higher than nighttime concentrations for most of the compound classes determined. In particular, photochemical reaction products such as diacids,

ketoacids and polyacids showed a strong diurnal pattern, suggesting that photochemical process in daytime is an important factor to cause the daytime maxima. This process should be shut down at night, which intensifies the diurnal patterns at the mountaintop.

[15] However, higher daytime concentrations were also reported for metals [Y. Wang *et al.*, 2007] and black carbon [Liu *et al.*, 2007] in the aerosols from Mount Tai. This may suggest that there is another important factor that also contribute to the diurnal distributions of water-soluble organic compounds. It was found that the heights of boundary layer in the Mount Tai frequently reduced to less than 1000 m at night during the campaign [Takigawa *et al.*, 2007], which keeps the mountaintop (1534 m) in the free troposphere at night. In contrast, the boundary layer came up far above the mountaintop in daytime. This local meteorology of Mount Tai should restrain the upward transport of pollutants from lowland to the summit at night and enhance the transport in daytime. This seems to be supported by the results of levoglucosan (biomass burning tracer and not produced by photochemistry). Concentrations of levoglucosan are similar between day (average 43 ng m⁻³, range 21–123 ng m⁻³) and night samples (average 36 ng m⁻³, range 1.6–151 ng m⁻³), but we can still recognize the lower average concentrations at night, suggesting that decrease in the height of boundary layer at night suppress the upward transport of biomass burning products from the lowland to the mountaintop.

3.2. Dicarboxylic and Ketocarboxylic Acids and α-Dicarbonyls

[16] Homologous dicarboxylic acids (DCAs, C₂–C₁₁), together with several ketocarboxylic acids and dicarbonyls, were detected in the PM_{2.5} samples with oxalic acid being the dominant species, followed by malonic, succinic, and phthalic acids (Table 2 and Figure 3a). Similar distribution has been reported in East Asian aerosols from the lower troposphere within the boundary layer [Wang *et al.*, 2006b] to the middle troposphere [Kawamura *et al.*, 2003]. Relative abundance of oxalic acid in total DCAs in the Mount Tai samples was 60 ± 1.3% in daytime and 59 ± 3.6% in nighttime. These values are higher than those (49 ± 7.4%) reported from urban Nanjing, China [Wang *et al.*, 2002], but much lower than those (70–75%) reported in the remote marine aerosols from the central Pacific [Kawamura and Sakaguchi, 1999]. Oxalic acid (C₂) is mostly produced by photooxidation of organic precursors [Kawamura *et al.*, 1996; Limbeck and Puxbaum, 1999], although it is also reported in gasoline and diesel car exhausts as a primary product [Kawamura and Kaplan, 1987]. The higher relative abundance of oxalic acid in the Mount Tai aerosols compared to the flat lowland samples suggests an enhanced photochemical oxidation of organic precursors during the transport from the lower parts of NCP to the mountaintop.

[17] Succinic acid (C₄) has been suggested as one of the precursors of oxalic (C₂) and malonic (C₃) acids [Kawamura *et al.*, 1996]. Adipic acid (C₆) is believed to be produced from oxidation of anthropogenic cyclic hexene [Grosjean *et al.*, 1978], whereas azelaic acid (C₉) is a photooxidation product of unsaturated fatty acids (e.g., oleic and linoleic acids) derived from biogenic emissions [Kawamura and Gagosian, 1987] and/or from anthropo-




Figure 3. Daytime and nighttime molecular compositions of water-soluble organic compound classes in the $PM_{2.5}$ aerosols during the campaign at Mount Tai: (a) dicarboxylic acids, (b) ketoacids and α -dicarbonyls, (c) sugars, and (d) polyols and polyacids. (Bars represent the standard deviations.)

genic cooking activities [Robinson *et al.*, 2006]. Maleic acid (M) can be produced by photooxidation of aromatic hydrocarbons such as benzene and toluene in the atmosphere, and be further isomerized into fumaric acid (F) [Kawamura and Ikushima, 1993]. We found that C_6/C_9 and M/F concentration ratios were slightly higher in nighttime than in daytime (Figure 4). This may suggest that oxidative production of C_9 and M to F isomerization are more significant in daytime. In contrast, *i*-phthalic to *t*-phthalic acid (*i*-Ph/*t*-Ph) ratios were lower in nighttime than in daytime. Such diurnal differences may suggest different origins of air masses over the mountaintop between day and night, although no significant differences were observed for C_2/C_4 , C_3/C_4 and *i*-phthalic to phthalic acid (*i*-Ph/Ph) ratios.

[18] Homologous ketocarboxylic acids (ωC_2 – ωC_4 , ωC_9 and Pyr) and α -dicarbonyls (mGly and Gly) were detected in the samples with total concentrations of 13–163 $ng\ m^{-3}$ (average 70 $ng\ m^{-3}$) and 1–65 $ng\ m^{-3}$ (average 22 $ng\ m^{-3}$), respectively (Table 2 and Figure 3b). These ketoacids and carbonyls are secondarily produced via atmospheric photooxidation of organic precursors and/or primarily formed by fossil fuel combustion and biomass burning, and further oxidized into saturated DCAs [Chebbi and Carlier, 1996; Kawamura *et al.*, 1996; Volkamer *et al.*, 2006]. Glyoxal, the smallest α -dicarbonyl, generally originates from the oxidation of anthropogenic and biogenic VOCs and serves as a novel indicator for fast VOC chemistry [Volkamer *et al.*,

2005; Wittrock *et al.*, 2006]. Glyoxal and methylglyoxal concentrations at the summit of Mount Tai are comparable to those in urban areas of coastal China [Wang *et al.*, 2002], but much higher than those reported in Jeju Island, the East China Sea (average 12 $ng\ m^{-3}$) [Kawamura *et al.*, 2004]. This contrast is consistent with the enhanced concentrations




Figure 4. Differences in the mass concentration ratios of selected dicarboxylic acids between the daytime and nighttime samples ($PM_{2.5}$) collected at Mount Tai. For abbreviations C_2 , C_3 , C_4 , C_6 , C_9 , M, F, Ph, *i*-Ph, and *t*-Ph, see Table 2. Bars represent the standard deviations. The *t* test for the paired samples showed $p \leq 0.05$.




Figure 5. Size distributions of water-soluble organic compound classes in the atmospheric aerosols over the summit of Mount Tai.

of glyoxal and formaldehyde over the NCP region as reported by satellite observation [Wittrock *et al.*, 2006]. Although Wittrock *et al.* [2006] reported gaseous carbonyls, our study demonstrates that particulate α -dicarbonyls are also abundant in the atmosphere over NCP.

3.3. Sugars, Polyols, and Polyacids

[19] Concentrations of saccharides including levoglucosan and several primary sugars in $PM_{2.5}$ samples were $7\text{--}160\text{ ng m}^{-3}$ (average 59 ng m^{-3}) (Table 2 and Figure 3c), accounting for 66% of those ($90 \pm 32\text{ ng m}^{-3}$) in total suspended particles (TSP) collected at the same site during the same period [Fu *et al.*, 2008]. Levoglucosan, a tracer for biomass burning, was found as a dominant sugar (43 ± 36

and $36 \pm 58\text{ ng m}^{-3}$ at day and night, respectively) (Table 2), followed by glucose, sucrose and/or mannitol. The predominance of levoglucosan in the aerosols is consistent with the ground surface samples from Chinese megacities [Wang *et al.*, 2006a]. Although the concentrations of the saccharides at Mount Tai are less than those ($9.4\text{--}735\text{ ng m}^{-3}$, average 132 ng m^{-3}) on the lowlands in China [Wang *et al.*, 2006a], they are higher than those ($31 \pm 25\text{ ng m}^{-3}$) measured in $PM_{2.5}$ aerosols collected by aircraft ($0.5\text{--}3.0\text{ km}$ altitude) over coastal China [G. H. Wang *et al.*, 2007]. Sucrose, a minor species in this group, is a metabolism product of bacteria in soil. Its higher concentrations in nighttime ($2.2 \pm 2.1\text{ ng m}^{-3}$) than in daytime ($1.7 \pm 0.7\text{ ng m}^{-3}$) (Table 2) were probably caused by an enhanced bacteria activity under higher humid conditions at night.

[20] Polyols and polyacids, including glycerol and glyceric, malic, tartaric and citric acids, were detected in the $PM_{2.5}$ aerosols (Table 2 and Figure 3d) with glycerol and glyceric and malic acids being the major species in this group. Concentrations of these compounds in the mountain aerosols ($9.6 \pm 10.7\text{ ng m}^{-3}$) are less than those ($38 \pm 35\text{ ng m}^{-3}$) reported in the lowland aerosols from Chinese megacities [Wang *et al.*, 2006a], but are very close to those ($7.0 \pm 8.1\text{ ng m}^{-3}$) in the summertime tropospheric aerosols collected by aircraft ($0.5\text{--}3.0\text{ km}$ altitude) over inland China [G. H. Wang *et al.*, 2007]. Glyceric and malic acids are secondary photooxidation products in the atmosphere [Kawamura and Ikushima, 1993; Simoneit *et al.*, 2004c]. The sum of glyceric and malic acids (average 27 ng m^{-3}) (Table 2) in the fine particles over Mount Tai is 3 times higher than that (average 7.5 ng m^{-3}) reported in the aircraft samples over the North Pacific [Simoneit *et al.*, 2004b]. This may suggest that the mountain aerosols are photochemically aged, to some extent, like marine aerosols, although the aircraft samples are more aged but may be more diluted and/or further subjected to photochemical decomposition during the transport over the Pacific.

[21] Glycerol is a reduced sugar primarily produced by fungal metabolism in soils and is emitted to the atmosphere as wind-blown dusts [Simoneit *et al.*, 2004c]. Its concentration (average 15 ng m^{-3} , Table 2) in the mountain aerosols is much less than that (average 38 ng m^{-3}) of the aircraft aerosols over the North Pacific [Simoneit *et al.*, 2004b]. Such a difference may be associated with the fact that Mount Tai samples are $PM_{2.5}$ whereas the North Pacific samples are total suspended particles (TSP), and thus soil fungus-derived glycerol that is enriched in coarse particles is less abundant in the $PM_{2.5}$ samples from Mount Tai. However, it is important to stress that the Mount Tai measurements represent a very limited time period (1 week) and may not be representative of typical concentrations measured at the site.

3.4. Size Distributions

[22] Various water-soluble organics were also determined in the eight-stage size-segregated aerosols from Mount Tai. As shown in Figures 5a, 5b, 5c, and 5e, bimodal distributions were found for diacids, ketoacids, α -dicarbonyls, and polyols and polyacids with two peaks at $0.70\text{--}1.1$ and $5.8\text{--}9.0\text{ }\mu\text{m}$ in diameter. Although major diacids (C_2 , C_3 and C_4) maximized at $0.70\text{--}1.1\text{ }\mu\text{m}$, some minor diacid species such as C_8 , C_9 and C_{10} showed a peak at larger size ($1.1\text{--}2.2\text{ }\mu\text{m}$)




Figure 6. Relative abundances and concentration ratios of water-soluble organic compounds in size-segregated aerosols collected at Mount Tai (total amount of each compound in the eight stages: C₂, 471 ng m⁻³; C₃, 76 ng m⁻³; C₄, 140 ng m⁻³; levoglucosan, 47 ng m⁻³; glucose, 13 ng m⁻³; glycerol, 30 ng m⁻³; glyceric acid, 16 ng m⁻³; and malic acid, 7.4 ng m⁻³).

(data are not shown here). In contrast, total sugars showed a different pattern with the highest concentration on the coarse mode (5.8–9.0 μm) (Figure 5d), mainly owing to the high concentration of sucrose. Such a bimodal distribution of dicarboxylic acids was not observed during the summer periods in urban Beijing, where diacids showed unimodal distribution with a peak in the fine mode [Yao *et al.*, 2003]. During the sampling period wind directions varied in a wide range of 20–323° (average 173° ± 74°), but there is no systematic difference in the wind direction between day and night (Table 1). Thus, influence of wind direction on the size distributions of diacids and related compounds may be insignificant. The coarse mode of diacids could in part be derived from biological formation in soil and/or adsorption of gaseous DCAs on alkali soil particles, but the bimodal patterns of the diacids on the mountaintop may be associated with other meteorological conditions of the high mountain regions.

[23] Over Mount Tai, fog or cloud events were often observed in nighttime during the campaign. Sizes of cloud/fog droplets are larger than the aerosol particles, ranging from a few micrometers to several hundreds of micrometers, because the droplets are grown by the condensation of water vapor from the particles that are activated as CCN [Seinfeld and Pandis, 1998]. In addition to the chemical components of aerosols, the cloud/fog droplets further uptake soluble gases via aqueous phase reactions [Blando and Turpin, 2000; Chen *et al.*, 2007]. Those droplets can be collected in the coarse fractions during the impactor sampling. These

situations may cause an enrichment of diacids on the coarse mode of size-segregated aerosols. Unfortunately, the largest fraction (>9 μm) was not analyzed for the chemical composition as described above. A formation of cloud/fog droplets that may enrich water-soluble compounds including diacids [Raja *et al.*, 2008] and/or hygroscopic growth of particles at night could lead to a decrease in the concentrations of water-soluble organics, to some extent, in PM_{2.5}.

[24] During the campaign, high-volume air sampling was conducted to collect bulk aerosol (TSP) samples without any particle cutoff. The TSP samples during the sampling period showed similar diurnal trend for diacids (C₂–C₁₁) with higher concentrations in daytime (Kawamura *et al.*, unpublished results, 2007). However, the difference in the TSP samples between day and night is much smaller (<40%) than those (200–300%) found in the PM_{2.5} samples. The similar but more distinct trend in PM_{2.5} can be explained by selective entrainment of diacids into cloud/fog droplets or hygroscopic growth of fine to coarse particles at night. These processes may cause a significant decrease in the concentrations of diacids in PM_{2.5} in nighttime because at least the sizes of fog/cloud particles are much larger (>7 μm) [Seinfeld and Pandis, 1998] than fine aerosols such as PM_{2.5}.

[25] In addition, the normalized abundances of C₂–C₄ diacids maximized in the fine mode (0.7–1.1 μm) and, to a lesser extent, in the coarse mode (5.8–9.0 μm), as shown in Figure 6a. Concentration ratios of diacids (C₂/C₄ and C₃/C₄) maximized in the finest mode (<0.4 μm) (Figure 6b),

suggesting that oxidative degradation of C_4 to C_2 and C_3 diacids occurred in fine aerosols and/or secondary production in gas phase followed by adsorption on preexisting fine aerosols. A strong linear correlation ($R^2 = 0.99$) was found between the C_2/C_4 and C_3/C_4 ratios among the different stages. This is consistent with the previously reported mechanism that C_2 and C_3 diacids are produced by the oxidation of C_4 in the atmosphere [Kawamura and Ikushima, 1993; Kawamura *et al.*, 1996, 2007].

[26] As shown in Figure 6c, levoglucosan was found to exist mostly in fine particles with a diameter less than $1.1 \mu\text{m}$. Levoglucosan is emitted from biomass burning and subsequently aggregates and/or condenses on preexisting particles owing to the higher surface area of fine particles [Simoneit *et al.*, 2004a; Wang *et al.*, 2006b]. Glucose is an important component of pollen and abundantly present in soil microorganisms. This sugar compound can be emitted into the air by wind motion from vegetation and soil systems. Therefore, it is reasonable that soil-derived glucose was found abundantly in coarse particles larger than $3.0 \mu\text{m}$, being in contrast to levoglucosan (Figure 6c), whose origin is associated with biomass burning. Recently, CCN efficiencies of saccharides have been reported to be lower than those of organic acids owing to their absence of dissociation in water, and their contributions to cloud formation are less than those of diacids [Ekstrom *et al.*, 2009; Rosenorn *et al.*, 2006; Varga *et al.*, 2007]. Thus, less abundance of levoglucosan on coarse mode (Figure 6c) may mean that cloud/fog formation over the mountaintop does not significantly affect its size distribution.

[27] Glycerol is also derived from soil suspension as discussed above. It was not detected in the particles smaller than $0.40 \mu\text{m}$ (Figure 6d). Glyceric and malic acids are secondary oxidation products and thus are enriched in the fine mode ($<2.1 \mu\text{m}$), although a significant fraction of both acids was also found on coarse mode ($5.8\text{--}9.0 \mu\text{m}$) (Figure 6d). The peak in the fine mode can be explained by the condensation via photooxidation of gaseous precursors [Kawamura *et al.*, 2007], whereas the second peak in the coarse mode may be attributable to the size shift of aerosols by hygroscopic growth of fine particles, which may occur at night under a high humidity condition. This hygroscopic growth of aerosols occasionally followed by fog/cloud formation may also be responsible for the second peak of diacids and related compounds on coarse mode (Figures 5a–5c). Because the size-segregated sampling was conducted for 7 days at the mountaintop, this process may have often happened in nighttime when ambient temperature decrease and relative humidity increase.

4. Summary and Conclusions

[28] On the basis of the analyses of size-segregated and day/night $\text{PM}_{2.5}$ aerosol samples, we found diurnal variations of water-soluble organic compounds with daytime maxima and bimodal size distributions for most of the compounds in the Mount Tai atmosphere over North China Plain (NCP). The results are interpreted by long-range atmospheric transport of pollutants and photochemical productions of water-soluble organics during the transport to the mountaintop under unique meteorological conditions of

high mountains (changes in the heights of boundary layer and relative humidity), as follows.

[29] Fine ($\text{PM}_{2.5}$) aerosol samples were collected for 7 days in 2006 summer on a day and night basis at the summit of Mount Tai (1534 m) in NCP and were analyzed for water-soluble organic compounds including dicarboxylic acids, ketocarboxylic acids, α -dicarbonyls, saccharides, polyols and polyacids using GC and GC/MS. A distinct difference was found in daytime and nighttime concentrations for most of the water-soluble organics. Their daytime concentrations are mostly 2–3 times higher than the nighttime concentrations, which were interpreted by photochemical production of the compounds in daytime and decrease in the heights of the boundary layer at night. Because the heights frequently reduced to less than 1000 m at night in the Mount Tai area, the mountaintop was often in the free troposphere, being isolated from the polluted ambient air of lowland origin at night. This is in contrast to the nighttime maxima generally observed in urban areas owing to the development of inversion layer. Compared with TSP samples, $\text{PM}_{2.5}$ samples showed more distinct diurnal difference of diacids and related compounds, suggesting the growth of fine particles to larger sizes ($>2.5 \mu\text{m}$) during the formation of cloud/fog droplets at night.

[30] Size-segregated aerosol samples (eight stages from $<0.4 \mu\text{m}$ to $9 \mu\text{m}$) collected during the campaign showed bimodal distributions of diacids, ketocarboxylic acids, and dicarbonyls with concentration peaks at $0.70\text{--}1.1 \mu\text{m}$ and $5.8\text{--}9.0 \mu\text{m}$. On the other hand, levoglucosan showed a unimodal distribution on a fine mode whereas glucose showed a peak on a coarse mode. The peaks of the water-soluble organics in smaller particles result from the condensations of organics from combustion process and photooxidation products from gaseous precursors. In contrast, the peaks in coarse particles are largely derived from the suspended soils and pollens. However, the peaks of diacids and ketoacids on the coarse mode may also be derived by a hygroscopic growth of fine particles to larger ones at night when relative humidity increases.

[31] **Acknowledgments.** We thank Xin Zhao of Nanjing University, China, for his help during the sampling. This work was financially supported by China Natural Science Foundation (40873083 and 40475050), Chinese Academy of Sciences (KZCX2-YW-148), and the Japanese Ministry of Education, Science and Culture through grant-in-aid 17340166. G.W. appreciates Japan Society for the Promotion of Science (JSPS) for the invitation fellowship (L08525). We also thank the staff at Meteorological Station of Mount Tai for providing the meteorological data and Phil Meyers of University of Michigan for his critical reading and comments on the early version of the manuscript.

References

- Akimoto, H. (2003), Global air quality and pollution, *Science*, *302*, 1716–1719, doi:10.1126/science.1092666.
- Blando, J. D., and B. J. Turpin (2000), Secondary organic aerosol formation in cloud and fog droplets: A literature evaluation of plausibility, *Atmos. Environ.*, *34*, 1623–1632, doi:10.1016/S1352-2310(99)00392-1.
- Chebbi, A., and P. Carlier (1996), Carboxylic acids in the troposphere, occurrence, sources, and sinks: A review, *Atmos. Environ.*, *30*, 4233–4249, doi:10.1016/1352-2310(96)00102-1.
- Chen, J., R. J. Griffin, A. Grini, and P. Tulet (2007), Modeling secondary organic aerosol formation through cloud processing of organic compounds, *Atmos. Chem. Phys.*, *7*, 5343–5355.
- Claeys, M., W. Wang, A. C. Ion, I. Kourtchev, A. Gelencser, and W. Maenhaut (2004), Formation of secondary organic aerosols from isoprene and its gas-phase oxidation products through reaction with

- hydrogen peroxide, *Atmos. Environ.*, **38**, 4093–4098, doi:10.1016/j.atmosenv.2004.06.001.
- Decesari, S., M. C. Facchini, S. Fuzzi, G. B. McFiggans, H. Coe, and K. N. Bower (2005), The water-soluble organic component of size-segregated aerosol, cloud water and wet depositions from Jeju Island during ACE-Asia, *Atmos. Environ.*, **39**, 211–222, doi:10.1016/j.atmosenv.2004.09.049.
- Ekstrom, S., B. Noziere, and H.-C. Hansson (2009), The cloud condensation nuclei (CCN) properties of 2-methyltetrols and C3–C6 polyols from osmolality and surface tension measurements, *Atmos. Chem. Phys.*, **9**, 973–980.
- Fu, P., K. Kawamura, K. Okuzawa, S. G. Aggarwal, G. Wang, Y. Kanaya, and Z. Wang (2008), Molecular characteristics, sources and temporal variations of summertime organic aerosols in the troposphere over Mt. Tai, North China Plain, *J. Geophys. Res.*, **113**, D19107, doi:10.1029/2008JD009900.
- Gao, J., T. Wang, A. J. Ding, and C. B. Liu (2005), Observational study of ozone and carbon monoxide at the summit of Mount Tai (1534 m a.s.l.) in central-eastern China, *Atmos. Environ.*, **39**, 4779–4791, doi:10.1016/j.atmosenv.2005.04.030.
- Graham, B., O. L. Mayol-Bracero, P. Guyon, G. C. Roberts, S. Decesari, M. C. Facchini, P. Artaxo, W. Maenhaut, P. Koll, and M. O. Andreae (2002), Water-soluble organic compounds in biomass burning aerosols over Amazonia: 1. Characterization by NMR and GC-MS, *J. Geophys. Res.*, **107**(D20), 8047, doi:10.1029/2001JD000336.
- Grosjean, D., K. V. Cauwenbergh, J. P. Schmid, P. E. Kelly, and J. N. Pitts Jr. (1978), Identification of C3–C10 aliphatic dicarboxylic acids in airborne particulate matter, *Environ. Sci. Technol.*, **12**, 313–317, doi:10.1021/es60139a005.
- Hemming, B. L., and J. H. Seinfeld (2001), On the hygroscopic behavior of atmospheric organic aerosols, *Ind. Eng. Chem. Res.*, **40**, 4162–4171, doi:10.1021/ie000790l.
- Huang, X. F., M. Hu, L. Y. He, and X. Y. Tang (2005), Chemical characterization of water-soluble organic acids in PM_{2.5} in Beijing, China, *Atmos. Environ.*, **39**, 2819–2827, doi:10.1016/j.atmosenv.2004.08.038.
- Hyvarinen, A.-P., H. Lihavainen, A. Gaman, L. Vairila, H. Ojala, M. Kulmala, and Y. Viisanen (2006), Surface tensions and densities of oxalic, malonic, succinic, maleic, malic, and cis-pinonic acids, *J. Chem. Eng. Data*, **51**, 255–260, doi:10.1021/je050366x.
- Katz, M., and C. Chan (1980), Comparative distribution of eight polycyclic aromatic hydrocarbons in airborne particles collected by conventional high-volume sampling and by size fractionation, *Environ. Sci. Technol.*, **14**, 838–842, doi:10.1021/es60167a010.
- Kawamura, K., and R. B. Gagosian (1987), Implication of ω -oxocarboxylic acids in the remote marine atmosphere for photo-oxidation of unsaturated fatty acids, *Nature*, **325**, 330–332, doi:10.1038/325330a0.
- Kawamura, K., and K. Ikushima (1993), Seasonal changes in the distribution of dicarboxylic acids in the urban atmosphere, *Environ. Sci. Technol.*, **27**, 2227–2235, doi:10.1021/es00047a033.
- Kawamura, K., and I. R. Kaplan (1987), Motor exhaust emissions as a primary source for dicarboxylic acids in Los Angeles ambient air, *Environ. Sci. Technol.*, **21**, 105–110, doi:10.1021/es00155a014.
- Kawamura, K., and F. Sakaguchi (1999), Molecular distribution of water soluble dicarboxylic acids in marine aerosols over the Pacific Ocean including tropics, *J. Geophys. Res.*, **104**(D3), 3501–3509, doi:10.1029/1998JD100041.
- Kawamura, K., and O. Yasui (2005), Diurnal changes in the distribution of dicarboxylic acids, ketocarboxylic acids and dicarbonyls in the urban Tokyo atmosphere, *Atmos. Environ.*, **39**, 1945–1960, doi:10.1016/j.atmosenv.2004.12.014.
- Kawamura, K., H. Kasukabe, and L. A. Barrie (1996), Source and reaction pathways of dicarboxylic acids, ketoacids and dicarbonyls in Arctic aerosols: One year of observations, *Atmos. Environ.*, **30**, 1709–1722, doi:10.1016/1352-2310(95)00395-9.
- Kawamura, K., N. Umemoto, M. Mochida, T. Bertram, S. Howell, and B. J. Huebert (2003), Water-soluble dicarboxylic acids in the tropospheric aerosols collected over east Asia and western North Pacific by ACE-Asia C-130 aircraft, *J. Geophys. Res.*, **108**(D23), 8639, doi:10.1029/2002JD003256.
- Kawamura, K., M. Kobayashi, N. Tsubonuma, M. Mochida, T. Watanabe, and M. Lee (2004), Organic and inorganic compositions of marine aerosols from East Asia: Seasonal variations of water-soluble dicarboxylic acids, major ions, total carbon and nitrogen, and stable C and N isotopic composition, in *Geochemical Investigation in Earth and Space Science: A Tribute to Issac Kaplan, R.*, edited by J. Hill et al., pp. 243–265, Elsevier, New York.
- Kawamura, K., Y. Imai, and L. A. Barrie (2005), Photochemical production and loss of organic acids in high Arctic aerosols during long-range transport and polar sunrise ozone depletion events, *Atmos. Environ.*, **39**, 599–614, doi:10.1016/j.atmosenv.2004.10.020.
- Kawamura, K., M. Narukawa, S. M. Li, and L. A. Barrie (2007), Size distributions of dicarboxylic acids and inorganic ions in atmospheric aerosols collected during polar sunrise in the Canadian high Arctic, *J. Geophys. Res.*, **112**, D10307, doi:10.1029/2006JD008244.
- Latif, M. T., and P. Brimblecombe (2004), Surfactants in atmospheric aerosols, *Environ. Sci. Technol.*, **38**, 6501–6506, doi:10.1021/es049109n.
- Limbeck, A., and H. Puxbaum (1999), Organic acids in continental background aerosols, *Atmos. Environ.*, **33**, 1847–1852, doi:10.1016/S1352-2310(98)00347-1.
- Liu, Y., P. Pochanart, Y. Kanaya, Z. Wang, J. Li, and H. Akimoto (2007), Synoptic meteorology study on observation ozone, carbon monoxide and black carbon at East Asia mountain areas in summer and winter, paper presented at 2nd Data Workshop of Taishan Campaign in Yokohama, Japan, Japan Agency for Marine-Earth Sci. and Technol., Yokohama, Japan.
- McMurry, P. H. (2000), A review of atmospheric aerosol measurements, *Atmos. Environ.*, **34**, 1959–1999, doi:10.1016/S1352-2310(99)00455-0.
- Mochida, M., K. Kawamura, N. Umemoto, M. Kobayashi, S. Matsunaga, H. J. Lim, B. J. Turpin, T. S. Bates, and B. R. T. Simoneit (2003), Spatial distributions of oxygenated organic compounds (dicarboxylic acids, fatty acids, and levoglucosan) in marine aerosols over the western Pacific and off the coast of East Asia: Continental outflow of organic aerosols during the ACE-Asia campaign, *J. Geophys. Res.*, **108**(D23), 8638, doi:10.1029/2002JD003249.
- Nepotchaykh, O. V., and P. A. Ariya (2002), Degradation of dicarboxylic acids (C₂/C₉) upon liquid-phase reactions with O₃ and its atmospheric implications, *Environ. Sci. Technol.*, **36**(15), 3265–3269, doi:10.1021/es0114310.
- Peters, J., and J. H. Seifert (1980), Losses of benzo (a)pyrene under the conditions of high-volume sampling, *Atmos. Environ.*, **14**, 117–120, doi:10.1016/0004-6981(80)90114-6.
- Raja, S., et al. (2008), Fog chemistry in the Texas-Louisiana Gulf Coast corridor, *Atmos. Environ.*, **42**, 2048–2061, doi:10.1016/j.atmosenv.2007.12.004.
- Richter, A., J. Burrows, H. Nub, C. Granier, and U. Niemeier (2005), Increase in tropospheric nitrogen dioxide over China observed from space, *Nature*, **437**, 129–132, doi:10.1038/nature04092.
- Robinson, A. L., R. Subramanian, N. M. Donahue, A. Bernadop-Bricker, and W. Rogge (2006), Source apportionment of molecular markers and organic aerosol. 3. Food cooking emissions, *Environ. Sci. Technol.*, **40**, 7820–7827, doi:10.1021/es060781p.
- Rosenorn, T., G. Kiss, and M. Bilde (2006), Cloud droplet activation of saccharides and levoglucosan particles, *Atmos. Environ.*, **40**, 1794–1802, doi:10.1016/j.atmosenv.2005.11.024.
- Seinfeld, J. H., and S. N. Pandis (1998), *Atmospheric Chemistry and Physics*, John Wiley, Hoboken, N.J.
- Simoneit, B. R. T., W. F. Rogge, M. A. Mazurek, L. J. Standley, L. M. Hildemann, and G. R. Cass (1993), Lignin pyrolysis products, lignans and resin acids as specific tracers of plant classes in emissions from biomass combustion, *Environ. Sci. Technol.*, **27**, 2533–2541, doi:10.1021/es00048a034.
- Simoneit, B. R. T., J. J. Schauer, C. G. Nolte, D. R. Oros, V. O. Elias, M. P. Fraser, W. F. Rogge, and G. R. Cass (1999), Levoglucosan: A tracer for cellulose in biomass burning and atmospheric particles, *Atmos. Environ.*, **33**, 173–182, doi:10.1016/S1352-2310(98)00145-9.
- Simoneit, B. R. T., V. O. Elias, M. Kobayashi, K. Kawamura, A. I. Rushdi, P. M. Medeiros, W. F. Rogge, and B. M. Dydik (2004a), Sugars: Dominant water-soluble organic compounds in soils and characterization as tracers in atmospheric particulate matter, *Environ. Sci. Technol.*, **38**, 5939–5949, doi:10.1021/es0403099.
- Simoneit, B. R. T., M. Kobayashi, M. Mochida, K. Kawamura, and B. J. Huebert (2004b), Aerosol particles collected on aircraft flights over the northwestern Pacific region during the ACE-Asia campaign: Composition and major sources of the organic compounds, *J. Geophys. Res.*, **109**, D19S09, doi:10.1029/2004JD004565.
- Simoneit, B. R. T., M. Kobayashi, M. Mochida, K. Kawamura, M. Lee, H. J. Lim, B. J. Turpin, and Y. Komazaki (2004c), Composition and major sources of organic compounds of aerosol particulate matter sampled during the ACE-Asia campaign, *J. Geophys. Res.*, **109**, D19S10, doi:10.1029/2004JD004598.
- Takigawa, M., K. Yamaji, and J. Li (2007), Model intercomparison during intensive field campaign at Taishan in 2006, paper presented at 2nd Data Workshop of Taishan Campaign in Yokohama, Japan, Japan Agency for Marine-Earth Sci. and Technol., Yokohama, Japan.
- Varga, Z., G. Kiss, and H.-C. Hansson (2007), Modelling the cloud condensation nucleus activity of organic acids on the basis of surface tension and osmolality measurements, *Atmos. Chem. Phys.*, **7**, 4601–4611.
- Volkamer, R., L. T. Molina, M. J. Molina, T. Shirley, and W. H. Brune (2005), DOAS measurement of glyoxal as an indicator for fast VOC

- chemistry in urban air, *Geophys. Res. Lett.*, *32*, L08806, doi:10.1029/2005GL022616.
- Volkamer, R., J. L. Jimenez, F. S. Martini, K. Dzepina, Q. Zhang, D. Salcedo, L. T. Molina, D. R. Worsnop, and M. J. Molina (2006), Secondary organic aerosol formation from anthropogenic air pollution: Rapid and higher than expected, *Geophys. Res. Lett.*, *33*, L17811, doi:10.1029/2006GL026899.
- Wang, G. H., S. L. Niu, C. Liu, and L. S. Wang (2002), Identification of dicarboxylic acids and aldehydes of PM10 and PM2.5 aerosols in Nanjing, China, *Atmos. Environ.*, *36*, 1941–1950, doi:10.1016/S1352-2310(02)00180-2.
- Wang, G. H., K. Kawamura, S. C. Lee, K. F. Ho, and J. J. Cao (2006a), Molecular, seasonal and spatial distributions of organic aerosols from fourteen Chinese cities, *Environ. Sci. Technol.*, *40*, 4619–4625, doi:10.1021/es060291x.
- Wang, G. H., K. Kawamura, T. Watanabe, S. C. Lee, K. F. Ho, and J. J. Cao (2006b), Heavy loadings and source strengths of organic aerosols in China, *Geophys. Res. Lett.*, *33*, L22801, doi:10.1029/2006GL027624.
- Wang, G. H., K. Kawamura, S. Hatakeyama, A. Takami, H. Li, and W. Wang (2007), Aircraft measurement of organic aerosols over China, *Environ. Sci. Technol.*, *41*, 3115–3120, doi:10.1021/es062601h.
- Wang, Y., H. Xu, and Y. Yang (2007), Chemical characteristics of aerosols at the summit of Mt. Tai, paper presented at 2nd Data Workshop of Taishan Campaign in Yokohama, Japan, Japan Agency for Marine-Earth Sci. and Technol., Yokohama, Japan.
- Wilkering, K. E., L. A. Barrie, and M. Engle (2000), Trans-Pacific air pollution, *Science*, *290*, 65–67, doi:10.1126/science.290.5489.65.
- Wittrock, F., A. Richter, H. Oetjen, J. P. Burrows, M. Kanakidou, S. Myriokefalitakis, R. Volkamer, S. Beirle, U. Platt, and T. Wagner (2006), Simultaneous global observations of glyoxal and formaldehyde from space, *Geophys. Res. Lett.*, *33*, L16804, doi:10.1029/2006GL026310.
- Yao, X., A. P. S. Lau, M. Fang, C. K. Chan, and M. Hu (2003), Size distributions and formation of ionic species in atmospheric particulate pollutants in Beijing, China: 2-dicarboxylic acids, *Atmos. Environ.*, *37*, 3001–3007, doi:10.1016/S1352-2310(03)00256-5.
- Yao, X., M. Fang, C. K. Chan, K. F. Ho, and S. C. Lee (2004), Characterization of dicarboxylic acids in PM2.5 in Hong Kong, *Atmos. Environ.*, *38*, 963–970, doi:10.1016/j.atmosenv.2003.10.048.
- Zhang, R. Y., I. Suh, J. Zhao, D. Zhang, E. C. Fortner, X. X. Tie, L. T. Molina, and M. J. Molina (2004), Atmospheric new particle formation enhanced by organic acids, *Science*, *304*, 1487–1490, doi:10.1126/science.1095139.
-
- S. Hu and M. Xie, School of the Environment, State Key Laboratory of Pollution Control and Resources Reuse, Nanjing University, Nanjing 210093, China.
- K. Kawamura and N. Umemoto, Institute of Low Temperature Science, Hokkaido University, Sapporo 060-0819, Japan.
- G. Wang, State Key Laboratory of Loess and Quaternary Geology, Institute of Earth Environment, Chinese Academy of Sciences, Xi'an 710075, China. (wanggh@ieecas.cn)
- Z. Wang, Institute of Atmospheric Physics, Chinese Academy of Sciences, Beijing 100029, China.