



Title	Retinol binding protein 4 in dairy cows : its presence in colostrum and alteration in plasma during fasting, inflammation, and the peripartum period
Author(s)	Eldaim, Mabrouk A. Abd; Kamikawa, Akihiro; Soliman, Mohamed M.; Ahmed, Mohamed M.; Okamatsu-Ogura, Yuko; Terao, Akira; Miyamoto, Toru; Kimura, Kazuhiro
Citation	Journal of Dairy Research, 77(1), 27-32 <a href="https://doi.org/10.1017/S0022029909990276">https://doi.org/10.1017/S0022029909990276</a>
Issue Date	2010-02
Doc URL	<a href="http://hdl.handle.net/2115/44774">http://hdl.handle.net/2115/44774</a>
Rights	Copyright © Proprietors of Journal of Dairy Research 2009
Type	article
File Information	JDR77-1_27-32.pdf



[Instructions for use](#)

## Retinol binding protein 4 in dairy cows: its presence in colostrum and alteration in plasma during fasting, inflammation, and the peripartum period

Mabrouk A Abd Eldaim<sup>1</sup>, Akihiro Kamikawa<sup>1</sup>, Mohamed M Soliman<sup>1</sup>, Mohamed M Ahmed<sup>1</sup>, Yuko Okamoto-Ogura<sup>1</sup>, Akira Terao<sup>1</sup>, Toru Miyamoto<sup>2</sup> and Kazuhiro Kimura<sup>1\*</sup>

<sup>1</sup> Laboratory of Biochemistry, Department of Biomedical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Japan

<sup>2</sup> Production Disease Team, National Institute of Animal Health, Japan

Received 26 May 2009; accepted for publication 4 August 2009; first published online 29 September 2009

Retinol-binding protein 4 (RBP4) is a plasma protein involved in retinol transportation, and recent evidence in rodents suggests that RBP4 is also a metabolic regulator that modifies insulin sensitivity. To assess how RBP4 levels are regulated in ruminants, we determined the RBP4 concentrations in bovine plasma and milk using Western blot analysis. Plasma RBP4 levels in non-pregnant non-lactating (control) cows were around 45 µg/ml, which were sustained during 60-h fasting, but decreased significantly 4 h after lipopolysaccharide (LPS) administration. Basal plasma retinol concentration was around 30 µg/dl, but this decreased to approximately one-third and one-half of these values during fasting and 8 h after LPS challenge, respectively. Plasma RBP4 and retinol levels in cows 3–6 d before parturition were comparable to those of the controls. However, on the day of parturition both were significantly decreased and had returned to basal levels by two weeks after calving. Interestingly, RBP4 was clearly detected in colostrum (16.4±5.6 µg/ml) but was only faintly detected in milk from cows at 7 d and 15 d after calving. Retinol concentrations in colostrum were almost 10-fold higher than those in plasma, while those in milk were comparable to those in plasma. These results suggest that RBP4 and retinol levels are independently regulated under physiological and pathophysiological conditions and that RBP4, like retinol, is transferred from maternal stores to calves through colostrum.

**Keywords:** Colostrum, cattle, RBP4, retinol, vitamin A.

Vitamin A (retinol) is essential for a variety of physiological processes, including vision, immune functions, reproduction, embryonic development as well as cellular growth and differentiation (Stephensen, 2001; Blomhoff & Blomhoff, 2006).

Retinol represents the most abundant retinoid in bovine blood (Van Merris et al. 2004) and its levels in steers are maintained for at least 145 d even if the animals are fed a diet with no supplemental vitamin A (Gorocica-Buenfil et al. 2008). However, plasma retinol and β-carotene levels in heifers and cows decrease at the end of gestation, reach their lowest values around parturition and increase again during the first week of lactation (Johnston & Chew, 1984; Goff & Stabel, 1990; Lindberg et al. 1999; Kumagai et al. 2000; Kumagai et al. 2001; Nonnecke et al. 2001; Goff et al. 2002; Van Merris et al. 2004; Debier et al.

2005; Rezamand et al. 2007). The decrease in plasma retinol concentration is probably due to the transfer of large amounts of retinol and its derivatives into colostrum (Goff et al. 2002), resulting in the accumulation of retinoids in colostrum rather than in milk (Debier et al. 2005). In addition, plasma retinol concentrations decrease immediately after intramammary injection of *Escherichia coli* (Van Merris et al. 2004) suggesting that vitamin A metabolism is also affected by the acute phase reactions that occur during coliform mastitis. Indeed, the plasma retinol levels in post-partum cows with mastitis are significantly lower than those of cows without mastitis (Johnston & Chew, 1984), and there is a significant delay in the recovery of the plasma β-carotene concentration in post-partum cows with a subclinical intramammary infection (Rezamand et al. 2007).

In newborn calves, plasma retinol levels at birth are low and increase markedly after colostrum intake (Blum et al. 1997; Kumagai et al. 2001; Debier et al. 2005; Puvogel

\*For correspondence; e-mail: k-kimura@vetmed.hokudai.ac.jp

et al. 2008). As colostrum provokes drastic morphological and functional changes in the gastrointestinal tract (GIT) in neonatal calves (Blum, 2006) and vitamin A plays a role in maintaining gut integrity (Thurnham et al. 2000; Quadro et al. 2000; Carroll & Forsberg, 2007) the retinol in colostrum is likely to contribute to the maturation of the GIT together with non-nutrient factors such as IGF-1 (Blum, 2006).

Retinoids are small hydrophobic compounds and therefore associate *in vivo* with soluble proteins. Retinol-binding protein 4 (RBP4) is the protein responsible for plasma retinol transport and binds with retinol in a 1:1 ratio (Noy, 2000). Recently, it has been reported that RBP4 induces insulin resistance in mice (Yang et al. 2005) and that the serum RBP4 levels in humans correlate with the magnitude of insulin resistance in subjects with impaired glucose tolerance or type 2 diabetes (Graham et al. 2006). More recently, STRA6 was identified as a cell surface receptor for RBP4 (Kawaguchi et al. 2007), indicating that RBP4 is a novel biologically active factor.

It was reported that the mean serum RBP4 and retinol concentrations in cows 4 weeks before calving were 42 µg/ml (~2 µM) and 53 µg/dl (~1.8 µM), respectively. RBP4 levels decreased one week before parturition and recovered to control levels at one week after calving (Lindberg et al. 1999) suggesting the possibility that RBP4, like retinol, is transferred from maternal stores to calves through colostrum and milk. However, a similar decrease in plasma RBP4 levels was found only in cows with sub-clinical intramammary infections, but not in uninfected cows (Rezamand et al. 2007), suggesting the decrease might be due to suppression of hepatic RBP4 synthesis during acute-phase reactions, as demonstrated in rats (Rosales et al. 1996).

In the present study, we measured the RBP4 and retinol concentrations in the plasma, colostrum and milk of cows to assess how plasma RBP4 and retinol levels are regulated in ruminants under physiological and pathophysiological conditions. Moreover, owing to the lack of such reports in any mammalian species, the study was undertaken to describe RBP4 levels in colostrum and/or milk.

## Materials and Methods

### *Blood sampling and collection of colostrum and milk*

All experimental procedures were performed in accordance with the guidelines outlined by the Animal Care and Use Committee of Hokkaido University.

In Experiment I, five non-pregnant non-lactating Holstein cows weighing 450–550 kg were used to examine the effects of fasting on plasma RBP4 and retinol levels. They were housed in individual stalls with free access to water and trace mineral blocks and fed a mixture of forage (orchard-grass hay, alfalfa hay cubes and corn silage) and concentrates. At the start of the experiment, feed was completely withdrawn, and food deprivation was maintained


for 60 h, when the metabolic effects of starvation in cows were clearly observed (Baird et al. 1972). Water was provided throughout the study. Blood was collected from the jugular vein at 0, 12, 24, 36, 48 and 60 h after the onset of food deprivation. Plasma was separated by centrifugation and stored at –30 °C until use.

In Experiment II, six non-pregnant non-lactating Holstein cows were housed as described above and used to examine the effects of lipopolysaccharide (LPS) on plasma RBP4 and retinol levels. On the day of the experiment, the cows were divided into two groups (groups A and B, three per group). Group A received an intravenous injection of LPS (*Escherichia coli* 055: B5, Difco, Detroit MI, USA) solution dissolved in saline at a dose 500 ng/kg, while group B was injected with saline as a control. Blood was collected at 0, 2, 4, 6 and 8 h after the injection, and the plasma was stored at –30 °C.

In Experiment III, six pregnant Holstein cows were housed as described above. Blood and milk (colostrum) samples were collected 3–6 d before calving, on the day of parturition and 1 and 2 weeks after calving. These samples were stored at –30 °C.

### *Western blot analysis of RBP4 in plasma, colostrum and milk*

Bovine RBP4 was purified as previously described (Miyamoto et al. 1989) and anti-sera were obtained by immunizing purified RBP4 to rabbits. Plasma was diluted 20-times with distilled water, and the whole milk and colostrum were undiluted and diluted 10-times with distilled water, respectively. These samples (7.5 µl for plasma and 24 µl for milk) and increasing concentrations of purified RBP4 were separated simultaneously by SDS-PAGE (13% gel) under reducing conditions. Thereafter, the proteins were electroblotted onto a PVDF membrane (Immobilon™, Millipore, Bedford MA, USA). The membrane was blocked for 1 h at room temperature in 5% (w/v) skimmed milk in a buffer [20 mM-Tris-HCl (pH 7.5), 0.15 M-NaCl and 0.1% Tween 20] and then incubated with the anti-RBP4 sera (1:2000) overnight at 4 °C. The membrane was washed five times with the wash buffer and incubated with horseradish peroxidase-conjugated goat anti-rabbit IgG antibody (1:4000, Zymed Laboratories Inc., South San Francisco CA, USA) for 1 h at room temperature. Visualization was performed using an enhanced chemiluminescence detection system (Millipore) according to the manufacturer's instructions. The intensities of the immunoreactive bands were analysed densitometrically using the NIH Image program. RBP4 concentration was measured by comparing the intensity of the bands with those of purified RBP4. As shown in Fig. 1, the band intensity of purified bovine RBP4 was closely correlated with the protein amount, at least from 10 ng to 80 ng of RBP4. Purified RBP4, added to bovine plasma and milk, was recovered at 96.2±0.4% and 95.4±0.8%, respectively. The intra- and inter-assay variations in



**Fig. 1.** Quantitative analysis of RBP4 by Western blot analysis. Representative blot of the increasing amounts of purified RBP4 and a standard curve obtained by densitometric analysis of the bands are shown.

plasma RBP4 determination for five independent experiments were less than 5.1% and 2.4%, respectively.

#### Spectrophotometric measurement of retinol in plasma, colostrum and milk

Retinol concentrations in plasma, colostrum and milk were determined by modifying the method described by Suzuki & Katoh (1990). In brief, 50  $\mu$ l of ethanol and 150  $\mu$ l of hexane were added to 50  $\mu$ l of plasma, colostrum, or milk, and the hexane phase was recovered after 40-min mixing and 10-min centrifugation at 6500 *g*. Retinol concentrations were calculated based on the absorbance of hexane extracts at 325 nm and 453 nm using the equations described (Suzuki & Katoh, 1990). Recovery of retinol added to milk was  $97.8 \pm 0.4\%$ .


#### Statistical analysis

Results are expressed as means  $\pm$  SEM. Statistical analysis was performed using one-way repeated measures ANOVA for Experiments I and III and two-way repeated measures ANOVA for Experiment II, and Fisher's post hoc test, with  $P < 0.05$  being considered statistically significant.


## Results

#### Changes in plasma RBP4 levels during fasting and after LPS challenge

Basal plasma RBP4 levels in the two experimental groups were almost the same ( $41.8 \pm 1.9$   $\mu$ g/ml in Experiment I and  $46.1 \pm 4.6$   $\mu$ g/ml in Experiment II) and were sustained during the 60-h fasting (Fig. 2A) and after saline administration (Fig. 3A). However, RBP4 levels were significantly reduced 4 h after LPS administration and decreased for at




**Fig. 2.** Effects of fasting on plasma RBP4 and retinol levels in cows. Cows ( $n=5$ ) were housed without feeding for 60 h. Blood was collected at 0, 12, 24, 36, 48 and 60 h after the onset of food deprivation, and isolated plasma samples were stored. Plasma concentrations of RBP4 (A) and retinol (B) were quantified by Western blot analysis and a spectrophotometric method, respectively. (C) The molar ratio of retinol to RBP4 was calculated. Values are expressed as means  $\pm$  SEM. \*:  $P < 0.05$  v. 0 h.



**Fig. 3.** Effect of LPS administration on plasma RBP4 and retinol levels in cows. Cows ( $n=6$ ) were injected intravenously with either LPS (500 ng/kg,  $n=3$ ) or saline ( $n=3$ ) as a control. Blood samples were collected at 0, 2, 4, 6 and 8 h after the injection. Plasma concentrations of RBP4 (A) and retinol (B) and their molar ratio (C) are shown. Values are expressed as means  $\pm$  SEM. \*:  $P < 0.05$  v. control.

least 8 h after the injection (Fig. 3A). Basal plasma retinol concentrations were  $36.2 \pm 3.2$   $\mu$ g/dl (Experiment I) and  $25.5 \pm 1.8$   $\mu$ g/dl (Experiment II). During fasting, the concentration progressively decreased to  $13.1 \pm 2.7$   $\mu$ g/dl at 60 h (Fig. 2B), and the retinol/RBP4 molar ratio fell from  $0.65 \pm 0.05$  to  $0.23 \pm 0.05$  (Fig. 2C). Plasma retinol concentration was also significantly decreased at 2 h and had decreased by almost 50% by 8 h after the LPS injection



**Fig. 4.** Changes in plasma RBP4 and retinol levels in cows before and after parturition. Blood was collected from 6 cows before and after parturition. Plasma concentrations of RBP4 (A) and retinol (B) and their molar ratio (C) are shown. Values are expressed as means  $\pm$  SEM. \*:  $P < 0.05$  v. day 0.


(Fig. 3B). In contrast to the fasting experiment, the retinol/RBP4 ratio was not changed significantly after LPS injection, although there was a tendency for it to decrease in the LPS-treated cows 2–4 h after the injection (Fig. 3C).

#### Changes in plasma RBP4 and retinol levels in cows before and after parturition

Plasma RBP4 levels at 3–6 d before parturition were  $50.7 \pm 5.3$   $\mu\text{g/ml}$ , which were comparable to those of non-pregnant, non-lactating cows in Experiments I and II. However, on the day of parturition, RBP4 levels were significantly decreased ( $29.8 \pm 5.4$   $\mu\text{g/ml}$ ) but they had returned to basal levels by 2 weeks after parturition ( $52.1 \pm 2.1$   $\mu\text{g/ml}$ ) (Fig. 4A). Plasma retinol levels before parturition were  $26.6 \pm 3.3$   $\mu\text{g/dl}$ , which were comparable to those of non-lactating cows. Similarly to RBP4, the retinol levels were significantly decreased on the day of parturition ( $18.5 \pm 2.7$   $\mu\text{g/dl}$ ) and had returned to basal levels by 15 d after parturition ( $33.9 \pm 4.3$   $\mu\text{g/dl}$ ) (Fig. 4B). There was no significant difference among the retinol/RBP4 ratios obtained before and after parturition.

#### Changes in RBP4 and retinol levels in colostrum and milk from cows before and after parturition

In the two milk samples obtained 3–6 d before parturition, RBP4 was hardly detected. However, in colostrum, RBP4 was clearly present at a concentration of  $16.4 \pm 5.6$   $\mu\text{g/ml}$  ( $n=6$ , Fig. 5A). RBP4 had almost disappeared from the milk of the cows at 7 d and 15 d after parturition. Retinol concentrations in milk ( $43.5 \pm 6.9$   $\mu\text{g/dl}$  at 7 d after



**Fig. 5.** Changes in RBP4 and retinol levels in colostrum and milk. Milk was collected from cows before ( $n=2$ ) and after ( $n=6$ ) parturition. Concentrations of RBP4 (A) and retinol (B) in colostrum and milk were measured by Western blot analysis and a spectrophotometric method, respectively. (C) The molar ratio of retinol to RBP4 is also shown. Values are expressed as means  $\pm$  SEM. \*:  $P < 0.05$  v. day 7 and day 15.

delivery) were comparable to plasma retinol concentrations, but those in colostrum were markedly higher ( $295.3 \pm 5.9$   $\mu\text{g/dl}$ ). The molar ratio of retinol to RBP4 in colostrum was almost 20 : 1.

#### Discussion

In the present study, we have demonstrated for the first time that considerable amounts of RBP4 are present in colostrum, but that it is scarce in milk. The increase in RBP4 levels in colostrum was accompanied by a comparable decrease in plasma RBP4 on the day of calving. The lack of alteration in plasma RBP4 levels during the 60-h fasting experiment suggests that its decrease on the day of calving may not be due to food availability or negative energy status during and shortly after parturition. The mild inflammation induced during the process of parturition may acutely decrease in plasma RBP4 as evidenced by the experiment with LPS. However, in bovine colostrum, a variety of non-nutrient bioactive compounds such as immunoglobulin G (IgG), growth hormone, prolactin, insulin and IGF-I are present, many of which are derived from blood (Barrington et al. 2001; Van de Perre, 2003; Taylor et al. 2004; Blum, 2006). In addition, since RBP4 was substantially lacking in milk, it is unlikely that RBP4 is synthesized in the mammary gland. Therefore, it is most likely that RBP4 is transferred from the blood to colostrum, resulting in a decrease its plasma levels.

In milk,  $\beta$ -lactoglobulin acts as a retinol carrier and thereby enhances intestinal retinol uptake in calves (Said

et al. 1989; Perez & Calvo, 1995). As one molecule of RBP4 binds to one molecule of retinol (Noy, 2000) the presence of almost a 20-fold excess of retinol over RBP4 in colostrum suggests that  $\beta$ -lactoglobulin is the primary retinol carrier in colostrum as well as in milk. Thus, the physiological relevance of colostrum RBP4 as a retinol carrier is currently unknown, but it is plausible that RBP4 contributes to calf development as a metabolic regulator, as described in rodents (Yang et al. 2005). It is, however, unlikely that maternal RBP4 contributes to the circulating RBP4 level in calves, although some macromolecules in colostrum such as IgG are transported across the intestinal epithelium into the neonatal circulation (Barrington et al. 2001; Van de Perre, 2003; Blum, 2006). This is because RBP4 is detected in the plasma of calves shortly after birth (before being fed colostrum) and its levels are unchanged one day after colostrum feeding, although its levels are approximately 50% of those in dairy heifers and cows (Nonnecke et al. 2001).

Nonnecke et al. (2001) also showed a positive correlation between plasma retinol and RBP4 levels in pre-ruminant calves (from birth to 27 d of age) fed a milk replacement with different amounts of supplemental vitamin A. A similar relationship between circulating retinol and RBP4 levels is observed in children and infants (Shenai et al. 1990; Craft, 2001; Aeberli et al. 2007) suggesting that RBP4 is a surrogate marker for retinol. However, the apparent dissociation of the plasma retinol levels from the plasma RBP4 levels seen during the 60-h fasting indicates that their levels in the plasma of cows are regulated independently of each other. Moreover, there were rapid changes in both plasma retinol and RBP4 levels after LPS challenge, and both changes seemed to be independent. The decrease of plasma retinol caused by LPS challenge was confirmed by a similar finding showing a decrease in plasma retinol concentrations immediately after intramammary injection of *Esch. coli* (Van Merris et al. 2004) and may be attributed to an increase of vitamin A consumption in various tissues such as the lung during inflammation (Kanda et al. 1990). On the other hand, the decrease of plasma RBP4 induced by LPS injection was probably due to the inhibition of hepatic RBP4 synthesis, as observed in rats (Rosales et al. 1996). It is therefore suggested that plasma RBP4 levels in cows are regulated independently of retinol concentrations under these conditions.

In summary, RBP4 was found in abundance in cow colostrum, accompanied by a comparable decrease in plasma RBP4, suggesting that RBP4 is transported from the blood to colostrum. In addition, RBP4 and retinol levels are independently regulated under physiological and pathophysiological conditions.

We are grateful to Dr Masato Fukui (Nippon Zenyaku Kogyo Co., Ltd.) for providing the blood, colostrum and milk for Experiment III. This study was supported by grants from the Japan Society for the Promotion of Science (JSPS) to KK and YO-O.

The study was also supported by a JSPS Research Fellowship for Young Scientists awarded to AK.

## References

- Aeberli I, Biebinger R, Lehmann R, l'Allemand D, Spinaz GA & Zimmermann MB 2007 Serum retinol-binding protein 4 concentration and its ratio to serum retinol are associated with obesity and metabolic syndrome components in children. *Journal of Clinical Endocrinology and Metabolism* **92** 4359–4365
- Baird GD, Heitzman RJ & Hibbitt KG 1972 Effects of starvation on intermediary metabolism in the lactating cow. *Biochemical Journal* **128** 1311–1318
- Barrington GM, McFadden TB, Huyler MT & Besser TE 2001 Regulation of colostrum synthesis in cattle. *Livestock Production Science* **70** 95–104
- Blomhoff R & Blomhoff HK 2006 Overview of retinoid metabolism and function. *Journal of Neurobiology* **66** 606–630
- Blum JW, Hadorn U, Sallmann H-P & Schuep W 1997 Delaying colostrum intake by one day impairs the plasma lipid, essential fatty acid, carotene, retinol and  $\alpha$ -tocopherol status in neonatal calves. *Journal of Nutrition* **127** 2024–2029
- Blum JW 2006 Nutritional physiology of neonatal calves. *Journal of Animal Physiology and Animal Nutrition* **90** 1–11
- Carroll JA & Forsberg NE 2007 Influence of stress and nutrition on cattle immunity. *Veterinary Clinics of North America—Food Animal Practice* **23** 105–149
- Craft NE 2001 Innovative approaches to vitamin A assessment. *Journal of Nutrition* **131** 1626S–1630S
- Debier C, Potter J, Goffe C & Larondell Y 2005 Present knowledge and unexpected behaviors of vitamin A and E in colostrum and milk. *Livestock Production Science* **98** 135–147
- Goff JP & Stabel JR 1990 Decreased plasma retinol,  $\alpha$ -tocopherol, and zinc concentration during the periparturient period: Effect of milk fever. *Journal of Dairy Science* **73** 3195–3199
- Goff JP, Kimura K & Horst RL 2002 Effect of mastectomy on milk fever, energy, and vitamins A, E, and  $\beta$ -carotene status at parturition. *Journal of Dairy Science* **85** 1427–1436
- Gorocica-Buenfil MA, Fluharty FL & Loerch SC 2008 Effect of vitamin A restriction on carcass characteristics and immune status of beef steers. *Journal of Animal Science* **86** 1609–1616
- Graham E, Yang Q, Blüher M, Hammarstedt A, Ciaraldi TP, Henry RR, Wason CJ, Oberbach A, Jansson P-A, Smith U & Kahn BB 2006 Retinol-binding protein 4 and insulin resistance in lean, obese, and diabetic subjects. *New England Journal of Medicine* **354** 2552–2563
- Johnston LA & Chew BP 1984 Peripartum changes of plasma and milk vitamin A and  $\beta$ -carotene among dairy cows with or without mastitis. *Journal of Dairy Science* **67** 1832–1840
- Kanda Y, Yamamoto N & Yoshino Y 1990 Utilization of vitamin A in rats with inflammation. *Biochimica et Biophysica Acta* **1034** 337–341
- Kawaguchi R, Yu J, Honda J, Hu J, Whitelegge J, Ping P, Wiita P, Bok D & Sun H 2007 A membrane receptor for retinol binding protein mediates cellular uptake of vitamin A. *Science* **315** 820–825
- Kumagai H, Chaipan Y & Mitani K 2000 Effect of supplementary vitamin A on plasma retinol concentrations, fertility and milk yield of dairy cows. *Animal Science Journal* **71** 143–149
- Kumagai H, Chaipan Y & Mitani K 2001 Effects of periparturient vitamin A supplementation on vitamin A concentrations in colostrum and milk from dairy cows, and plasma retinol concentrations, feed intake and growth of their calves. *Animal Science Journal* **72** 126–133
- Lindberg L-A, Sinkkonen H, Poso AR, Tesfä AT & Schroder J 1999 Production of monoclonal antibodies and enzyme immunoassay to bovine retinol-binding protein and determination of retinol-binding protein serum levels and retinol concentrations in serum and liver in dairy cows before and after parturition. *Research in Veterinary Science* **66** 259–263

- Miyamoto T, Katoh N, Motoi Y, Ohashi T, Nagasawa S & Shimbayashi K 1989 Retinol transport system in cattle and purification of retinol-binding protein from bovine serum. *Japanese Journal of Veterinary Science* **51** 408–415
- Nonnecke BJ, Roberts MP, Godkin JD, Horst RL, Hammell DC & Franklin ST 2001 Influence of supplemental, dietary vitamin A on retinol-binding protein concentrations in the plasma of preruminant calves. *Journal of Dairy Science* **84** 641–648
- Noy N 2000 Retinoid-binding proteins: mediators of retinoid action. *Biochemical Journal* **348** 481–495
- Perez MD & Calvo M 1995 Interaction of  $\beta$ -lactoglobulin with retinol and fatty acids and its role as a possible biological function for this protein: a review. *Journal of Dairy Science* **78** 978–988
- Puvogel G, Baumrucker C & Blum JW 2008 Plasma vitamin A status in calves fed colostrum from cows that were fed vitamin A during late pregnancy. *Journal of Animal Physiology and Animal Nutrition* **92** 614–620
- Quadro L, Gamble MV, Vogel S, Lima AAM, Piantedosi V, Moore SR, Colantuoni V, Gottesman ME, Guerrant RL & Blaner WS 2000 Retinol and retinol-binding protein: gut integrity and circulating immunoglobulins. *Journal of Infectious Diseases* **182** S97–S102
- Rezamand P, Hoagland TA, Moyes KM, Silbart LK & Andrew SM 2007 Energy status, lipid-soluble vitamins, and acute phase proteins in periparturient Holstein and Jersey dairy cows with or without sub-clinical mastitis. *Journal of Dairy Science* **90** 5097–5107
- Rosales FJ, Ritter SJ, Zolfaghari R, Smith JE & Ross AC 1996 Effects of acute inflammation on plasma retinol, retinol-binding protein, and its mRNA in the liver and kidneys of vitamin A-sufficient rats. *Journal of Lipid Research* **37** 962–971
- Said HM, Ong DE & Shingleton JL 1989 Intestinal uptake of retinol: enhancement by bovine milk  $\beta$ -lactoglobulin. *American Journal of Clinical Nutrition* **49** 690–694
- Shenai JP, Rush MG, Stahlman MT & Chytil F 1990 Plasma retinol-binding protein response to vitamin A administration in infants susceptible to bronchopulmonary dysplasia. *Journal of Pediatrics* **116** 607–614
- Stephensen CB 2001 Vitamin A, infection, and immune function. *Annual Review of Nutrition* **21** 167–192
- Suzuki J & Katoh N 1990 A simple and cheap method for measuring serum vitamin A in cattle using only a spectrophotometer. *Japanese Journal of Veterinary Science* **52** 1281–1283
- Taylor VJ, Cheng Z, Pushpakumara PGA, Beever DE & Wathes DC 2004 Relationships between the plasma concentrations of insulin-like growth factor-I in dairy cows and their fertility and milk yield. *Veterinary Record* **155** 583–588
- Thurnham DI, Northrop-Clews CA, McCullough FSW, Das BS & Lunn PG 2000 Innate immunity, gut integrity, and vitamin A in Gambian and Indian infants. *Journal of Infectious Diseases* **182** S23–S28
- Van de Perre P 2003 Transfer of antibody via mother's milk. *Vaccine* **21** 3374–3376
- Van Merris V, Meyer E, Duchateau L, Blum J & Burvenich C 2004 All-trans retinoic acid is increased in the acute phase-related hyporetinemia during *Escherichia coli* mastitis. *Journal of Dairy Science* **87** 980–987
- Yang Q, Graham TE, Mody N, Preitner F, Peroni OD, Zabolotny JM, Kotani K, Quadro L & Kahn BB 2005 Serum retinol binding protein 4 contributes to insulin resistance in obesity and type 2 diabetes. *Nature* **436** 356–362