Title	Effect of chemical fertilizer and manure application on N ₂ O emission from reed canary grassland in Hokkaido, Japan
Author(s)	Jin, Tao; Shimizu, Mariko; Marutani, Satoru; Desyatkin, Alexey R.; Iizuka, Natsumi; Hata, Hiroshi; Hatano, Ryusuke
Citation	Soil Science & Plant Nutrition, 56(1), 53-65 https://doi.org/10.1111/j.1747-0765.2010.00447.x
Issue Date	2010-02
Doc URL	http://hdl.handle.net/2115/44778
Rights	The definitive version is available at www.blackwell-synergy.com
Type	article (author version)
File Information	SSPN56-1_53-65.pdf

[Instructions for use](#)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Title

Effect of chemical fertilizer and manure application on N₂O emission from reed canary
grassland in Hokkaido, Japan

Names of authors

Tao Jin¹, Mariko Shimizu¹, Satoru Marutani¹, Alexey R. Desyatkin¹, Natsumi Iizuka¹,
Hiroshi Hata², Ryusuke Hatano¹

Affiliations

¹Soil Science Laboratory, Graduate School of Agriculture, Hokkaido University, Kita 9
Nishi 9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan
²Field Science Center for Northern Biosphere, Agro-ecosystem Research Station,
Shizunai Livestock Farm, Hokkaido University, Kita 11 Nishi 10, Kita-ku, Sapporo,
Hokkaido 060-0811, Japan

Corresponding author

Tao Jin
Tel.: +81 11 706 2503
Fax.: +81 11 706 2494

19 E-mail address: jintao@chem.agr.hokudai.ac.jp

20 Soil Science Laboratory, Graduate School of Agriculture, Hokkaido University, Kita 9

21 Nishi 9, Kita-ku, Sapporo, Hokkaido 060-8589, Japan

For review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

Abstract

We evaluated the effect of chemical fertilizer and manure applications on N₂O emission from a managed grassland by establishing three treatment plots of chemical N fertilizer (chemical fertilizer), manure combined chemical N fertilizer (manure), and no N fertilizer (control) at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. N₂O fluxes from soils were measured by a closed-chamber method from May 2005 to April 2008. Soil denitrifying enzyme activity (DEA) in root-mat layer (0-2.5cm) and mineral soil layer (2.5-5cm) of each treatment plot was measured by an acetylene inhibition method after treatment with NO₃⁻-N addition, glucose addition, both NO₃⁻-N and glucose addition and neither NO₃⁻-N nor glucose addition, respectively. Annual N₂O emission ranged from 0.6 4.9 kg N₂O-N ha⁻¹yr⁻¹, with the highest observed in manure plot and lowest in control plot. Chemical fertilizer-induced emission factor (EF) (range: 0.85 - 1.32%) was significantly higher than manure-induced EF (range 0.35 - 0.85%). Denitrification potential of soil horizons was measured with addition of both NO₃⁻-N and glucose, which was significantly higher in root-mat soil than that in mineral soil. Soil DEA in root-mat in NO₃⁻-N addition with and without addition of glucose had a significantly positive correlation with soil pH (P < 0.05). Soil pH was significantly

influenced by N source, which was significantly lower in chemical fertilizer plot than that in control and manure plot. For a fixed quantity of available N, application of manure could result in higher N₂O emission compared to chemical fertilizer owing to high soil pH values under manure application than under chemical fertilizer application.

Key words: chemical fertilizer, grassland, manure, N₂O emission, soil DEA.

INTRODUCTION

N₂O is one of the most important radiatively active trace gases in the atmosphere ~~that~~ contributes at least 5% to the observed global warming at present (Myhre *et al.* 1998).

The atmospheric concentration of N₂O has increased from a pre-industrial value of about 270 ppb to 319 ppb in 2005 and continues to increase as a result of human activities (IPCC 2007). Agriculture as a whole (i.e. animal excreta, denitrification of leached nitrate, etc.) contributes about 80% of the anthropogenic N₂O emissions (Brown *et al.* 2001). Direct and indirect emissions from agricultural systems are now thought to contribute approximately 6.2 Tg N₂O-N yr⁻¹ to the total global source strength of 17.7 Tg N₂O-N yr⁻¹ (Kroeze *et al.* 1999). About 57% of the global atmospheric sources of N₂O are estimated to be related to emissions from soils (Mosier *et al.* 1998).

N₂O is produced in soils primarily by microbial processes of nitrification and

Deleted: We evaluated the effect of chemical fertilizer and manure applications on N₂O emission from a managed grassland by establishing three treatment plots of chemical fertilizer, manure, and control at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. Seasonal N₂O and NO fluxes were measured by a closed-chamber method at 4 to 6 replications in each treatment plot from May 2005 to April 2008. Soil samples were collected from a 0-5 cm top soil layer at three replications on each gas sampling date for measuring pH, NO₃⁻-N, NH₄⁺-N, and DOC. Soil samples were collected from the root-mat layer (0-2.5cm) and the mineral soil layer (2.5-5cm) of each treatment plot in April, June, and August 2007, which was followed by measuring soil denitrifying enzyme activity (DEA). The soil DEA was measured by an acetylene inhibition method under the four treatments with and without the addition of NO₃⁻-N and glucose. The cumulative N₂O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0, and 2.1 to 4.9 kg N₂O-N ha⁻¹yr⁻¹, respectively. The application of both

Formatted: Font: Times New Roman

Deleted: contributing

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

denitrification (Tiedje 1988; Conrad 1996). Nitrification is the biological oxidation of ammonium to nitrite or nitrate under aerobic conditions, while denitrification is the reduction of nitrate to N₂O and N₂ when the supply of oxygen is limited. Increasing soil N availability as a result of increased N inputs by the application of chemical fertilizer and manure and atmospheric deposition have greatly enhanced N₂O emissions from soils (Kroeze *et al.* 1999) by influencing nitrifying and denitrifying enzyme activity. Chemical fertilizer and animal wastes are the two most important sources of direct N₂O emissions from agricultural soils (Mosier *et al.* 1998). The default IPCC emission factor, i.e. the percentage of applied N emitted as N₂O, is 1.0%, regardless of the fertilizer type (IPCC 2006). However, the type of N inputs to the fields may affect the N₂O emission rate in different ways, leading to different patterns of N₂O emissions from inorganic and organic N fertilizers. Addition of inorganic N increases N₂O emission through affecting the process of nitrification and denitrification by increasing the available NH₄⁺-N and NO₃⁻-N substrates. Organic fertilizers with a high and easily mineralizable organic C content stimulate microbial activity and thus N₂O emissions (Chadwick *et al.* 2000). The application of chemical N fertilizer resulted in short-lived N₂O peaks (Dobbie and Smith 2003). However, N₂O losses from manure plots extended over a longer period of time and were greater in magnitude than from chemical N fertilization (Jones *et al.* 2007). Higher

Deleted: agricultural

Deleted: through affecting

Deleted: activities

Deleted: type of fertilizer

Deleted: the

Deleted: is reported to

Deleted: the

Deleted: by

Deleted: providing

Deleted: additional

Deleted: .

1
2
3
4
5 76 N₂O fluxes from manure and sewage applications compared to that from chemical N
6
7
8 77 fertilizers were also observed in other studies (Scott *et al.* 2000). High N₂O fluxes from
9
10 78 manure treatments can be partly explained by the higher total N input than chemical
11
12 fertilizer treatments, providing more available N that was mineralized over a longer
13
14 79
15
16 80 period of time. Another reason for the increased N₂O losses could be an addition of
17
18 81 organic C by the manures, which is known to stimulate denitrification. McTaggart *et al.*
19
20 82 (1997) reported that C supply from slurry spread onto grasslands stimulated N₂O
21
22 83 production, resulting in a four times larger N₂O loss compared to the application of
23
24 84 NH₄NO₃ although the total rate of N application was similar in both treatments.
25
26
27
28 85 Grassland is an important ecosystem to support the production of herbivorous livestock
29
30 86 (Soussana *et al.* 2007). Application of chemical fertilizer and animal manure to
31
32 87 grasslands has been conducted to increase grass production, especially in developed
33
34 88 countries where grassland-based livestock production is important (Bouwman *et al.*
35
36 89 2002). However, N application to grasslands also poses a risk of N loss to the
37
38 90 environment in the form of N₂O emission. The objective of this study is to clarify the
39
40 91 effect of fertilizer and manure application on N₂O emission and to identify the factors
41
42
43
44
45
46
47 92 controlling N₂O emission from the grassland.
48
49
50 93

Deleted: partly

Deleted: additional

Deleted: clarify

1
2
3
4
5 94 MATERIALS AND METHODS
6

7
8 95 Study site
9

10 96 This study was conducted on a managed grassland located at the Shizunai
11
12
13 97 Experimental Livestock Farm, Field Science Center for Northern Biosphere of Hokkaido
14
15
16 98 University in Southern Hokkaido, Japan (42°26'N, 142°29'E). The site is characterized
17
18
19 99 by a humid continental climate with cold winters and cool summers. During 1979 to 2000,
20
21 100 the mean annual precipitation and air temperature for this region were 1365 mm and
22
23 101 7.9°C, respectively. The soil is derived from Tarumae (b) volcanic ash, and is classified as
24
25
26 102 Thaptic Melanudands (Soil Survey Staff, 2006; Mollic Andosol (IUSS Working Group,
27
28
29 103 WRB, 2006)). A layer of three cm thick root-mat was found on the top, and a 21 cm thick
30
31 104 Ap-layer was found under the root-mat in a survey conducted in August 2004 (Shimizu *et*
32
33
34 105 *al.* 2009). The C and N contents in the Ap-layer were 3.7% and 0.33%, respectively, and
35
36 106 the C:N ratio was 11.1. Dominant grass species at this site were reed canary grass
37
38
39 107 (*Phalaris arundinacea* L.) and foxtail grass (*Alopecurus pratensis* L.). The harvesting of
40
41
42 108 grass was carried out twice a year (21st June and 11th August in 2005, 27th June and 23rd
43
44
45 109 August in 2006 and 18th June and 18th August in 2007) in accordance with the local
46
47
48 110 practice.
49
50 111

Formatted: Justified

Deleted: in

Formatted: Superscript

Formatted: Superscript

Experimental setup

Three experimental plots were set up on the study site; one for treatment with chemical fertilizer (chemical fertilizer plot), another with beef cattle manure and chemical fertilizer (manure plot), and the other with no N fertilizer or manure (control plot). Setting of the treatments was initiated in the spring of 2005. Eighteen subplots (5 m × 4 m) were established for the chemical fertilizer, manure, and control plots with six and four replications from May 2005 to April 2007 and from May 2007 to April 2008, respectively.

Table 1 shows the information on the date of application and the application rates of chemical fertilizer and manure. The N application rates in the chemical fertilizer plot were at the recommended level for this site on the basis of soil tests, and were 164 kg N ha⁻¹ year⁻¹ in 2005 and 184 kg N ha⁻¹ year⁻¹ in 2006 as ammonium sulfate and ammonium phosphate (Table 1). For learning the N₂O emission from grassland soil under the local N fertilization level, N fertilization decreased to 74 kg N ha⁻¹ year⁻¹ in chemical fertilizer plot according to the recommends of Shizunai Experimental Livestock Farm staffs. The pH value of manure that used in 2005, 2006 and 2007 were 8.3, 8.8 and 9.1, respectively.

The manure application rates were the optimum rates used by farmers in the region, and were based on adequate amounts of potassium (K) application to the fields. Beef cattle

Deleted: ,

Deleted: in

Formatted: Font color: Auto

Formatted: Subscript

Formatted: Font color: Auto

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

manure with bedding litter (bark) was applied to the manure plot, and the application rates were 44 Mg FM ha⁻¹ (236 kg N ha⁻¹ and 5.8 Mg C ha⁻¹) in May 2005, 43 Mg FM ha⁻¹ (310 kg N ha⁻¹ and 6.0 Mg C ha⁻¹) in May 2006, and 43 Mg FM ha⁻¹ (331 kg N ha⁻¹ and 7.7 Mg C ha⁻¹) in May 2007 (Table 1). In the manure plot, the N supply rates from manure were estimated by multiplying the application rates by the N mineralization rate, and the differences between the supply rates in manure and the application rates in the fertilizer plot were supplied by chemical fertilizer. The N mineralization rates were estimated based on Uchida's model (Shiga *et al.* 1985) which was developed in Japan and were 13.2%, 7.0% and 5.5%, respectively in the first, second and third years after application. The mineralization rates of P and K from the manure were estimated based on the handbook of animal waste management and utilization in Hokkaido 2004 (Anon. 2004). The P mineralization rate was 20%, 10%, and 0% and the K mineralization rate was 70%, 10%, and 0% in the first year, second year, and third year after application, respectively.

N₂O and NO fluxes

We defined the crop growing season as a 7-day moving average of daily air temperature above 5 °C and the non-growing season as the rest (Shimizu *et al.* 2009). The growing season was 215 days in 2005 (From 10th April 2005 to 10th November 2005), 218 days in

Deleted: nutrient

1
2
3
4
5 148 2006 (From 15th April 2006 to 18th November 2006) and 220 days in 2007 (From 13th
6
7
8 149 April 2007 to 18th November 2007. N₂O and NO fluxes from the soil to the atmosphere
9
10 150 were measured by the static closed chamber method on the control, fertilizer, and manure
11
12
13 151 plots (Shimizu *et al.* 2009). The flux measurements were conducted in 2 - 28 day intervals
14
15
16 152 during the crop growing season and 10 - 30 day intervals during the non growing season
17
18 153 and between 8:00 and 11:00 h in each measuring day to minimize the effect of diurnal
19
20
21 154 temperature variation. The stainless steel chambers were 40 cm in diameter and 30 cm
22
23
24 155 high in the chemical fertilizer and manure plots, and 20 cm in diameter and 25 cm high in
25
26 156 the control plots. The chambers were placed directly into the soil to a depth of about 3 cm,
27
28
29 157 12 hours before the measurement of each subplot, and contained no aboveground biomass
30
31
32 158 in the chemical fertilizer, manure, and control plots. Before closing the chamber, a 250 ml
33
34 159 gas sample from the headspace of each chamber was extracted into a Tedlar bag for NO
35
36 160 analysis, and a 20 ml gas sample was injected into an evacuated vial (10 ml) for N₂O
37
38
39 161 analysis. This measurement was regarded as time 0 min. After 20 min or 30 min under a
40
41
42 162 closed-chamber condition, 250 ml of the headspace gas sample was extracted from each
43
44
45 163 chamber into a bag, and 20 ml was injected into a vial. From these bag samples, NO gas
46
47 164 concentrations were determined in a laboratory within 16 hours using a
48
49
50 165 Chemiluminescence N Oxide Analyzer (Model 265P, Kimoto Electric, Osaka, Japan).
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

N₂O gas concentrations were determined in a laboratory within 1 month using an ECD (Electron capture detector) gas chromatograph (model GC-14B, Shimadzu, Kyoto, Japan) from the samples of vials.

Gas fluxes were calculated from the change in gas concentration in the chamber against closure time:

$$F = \rho \times h \times (\Delta c / \Delta t) \times [273 / (273 + T)]$$

where F is the gas flux ($\mu\text{g N m}^{-2} \text{ h}^{-1}$ for N₂O), ρ is the gas density (N₂O-N = $1.26 \times 10^9 \mu\text{g m}^{-3}$), h is the height of the chamber from the soil surface (m), $\Delta c / \Delta t$ is the change in gas concentration inside the chamber during the sampling period ($\text{m}^3 \text{ m}^{-3} \text{ h}^{-1}$), and T is the average air temperature during the sampling period ($^{\circ}\text{C}$). A positive flux denotes the emission from the soil, whereas a negative flux denotes the uptake from the atmosphere.

The cumulative gas flux was calculated as follow:

$$\text{Cumulative gas flux} = \sum_{i=1}^n (R_i \times 24 \times D_i)$$

where R_i is the mean gas flux ($\text{mg m}^{-2} \text{ hr}^{-1}$) of the two successive sampling dates, D_i is the number of days in the sampling interval, and n is the number of sampling times. The cumulative period of 2005, 2006 and 2007 were calculated from 10th April 2005 to 14th April, from 15th April 2006 to 18th April 2007 and from 19th April 2007 to 4th April 2008, respectively.

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Emission factor

N₂O Emission factor (EF) for chemical fertilizer and manure (kg N₂O–N (kg N input)⁻¹)

was calculated as follows:

Chemical fertilizer-induced EF = {[N₂O emission (chemical fertilizer plot)] – [N₂O emission (control plot)]} / [chemical fertilizer N application rate (chemical fertilizer plot)]

Manure-induced EF = {[N₂O emission (manure plot)] – [chemical fertilizer N application rate (manure plot)] × chemical fertilizer-induced EF – [N₂O emission (control plot)]} / [manure N application rate (manure plot)]

Environmental variables

Daily precipitation was obtained at the Sasayama AMeDAS (Automated Meteorological Data Acquisition System) station by the Japan Meteorological Agency. Air temperature and soil temperature at a 5 cm depth were measured at the same time with the flux measurements using a thermistor thermometer (CT220, CUSTOM, Tokyo, Japan), and soil moisture content at a 0 - 6 cm depth was measured using the Frequency Domain Reflectometry (FDR) method (DIK-311A, Daiki, Saitama, Japan). Soil core samples (14 cm diameter, 13 cm height) were collected in April 2007, and calibration curves were made to calculate water-filled pore space (WFPS) from the FDR device reading (m³ m⁻³)

Deleted: and daily maximum depths of snow were

Deleted:

Deleted: closest

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

and percent total porosity (Linn and Doran 1984). The percent total porosity was measured using a 100 ml soil core collected in April 2007 and was regarded as constant throughout the study period because of no tillage.

Soil chemical analyses

Soil samples at a depth of 0 to 5 cm from ground surface were collected during a period from April to November at three replicates in all treatment plots. Within 48 hours of soil sampling, soil samples were sieved through the 2 mm sieve and stones and roots were removed. Soil samples were then immediately extracted in deionized water (1:5) and in 2 M KCl (1:10), and the extracts were stored at 4 °C until analysis for dissolved nutrients after filtered through 0.2-µm membrane filters. Water-soluble organic carbon (DOC) content in the deionized-water-extract solution was analyzed using a TOC analyzer (TOC 5000A, Shimadzu). The concentration of NO₂-N and NO₃-N in the deionized-water-extract solution was analyzed by ion chromatography (Dionex QIC Analyzer, Dionex Japan, Osaka, Japan). The concentration of NH₄⁺-N in the 2 M KCl extracted solution was determined by the indophenol-blue method (UV mini 1240, Shimadzu, Kyoto, Japan). Soil pH was measured in the deionized-water-extract solution with a combined electrode pH meter (F-8 pH meter, Horiba, Japan).

Measurement of soil denitrifying enzyme activity (DEA)

Deleted: -----Page Break-----

Deleted: under the root-mat

For measuring soil denitrifying enzyme activity (DEA), soil samples were taken from all treatment plots in the root-mat layer (0 - 2.5cm depth) and mineral soils layer (2.5 - 5cm depth) with 3 replications in April, June, and August 2007. The root-mat soil samples were cut into small pieces with 1cm diameter, and stones or roots were removed from the mineral soil samples by passing through the 2mm sieve within 48 h after sampling. Then we mixed three replicates of soil samples and kept in refrigerator at 4 °C until analysis. The DEA was determined by an acetylene block technique, which inhibits the final conversion of N_2O to N_2 gas (Tiedje, 1994). Soil samples were incubated under anaerobic condition at 25°C with a solution treated with 1) chloramphenicol (1g L^{-1}) (Chl), 2) with chloramphenicol (1g L^{-1}) and $\text{NO}_3^- \text{-N}$ (200 mg N L^{-1} as KNO_3) (Chl+N), 3) with chloramphenicol (1g L^{-1}) and organic-C (2 g C L^{-1} as glucose) (Chl+C), and 4) with chloramphenicol (1g L^{-1}), $\text{NO}_3^- \text{-N}$ (200 mg N L^{-1} as KNO_3) and organic-C (2 g C L^{-1} as glucose) (Chl+N+C). Fresh soil of 15g was placed into a 100 conical flask, and 15 ml treated solution was added to the flask. The flasks were evacuated and flushed four times with N_2 to ensure anaerobic conditions, and acetylene (C_2H_2) gas was added to a final concentration of 10% (10 kPa) in the headspace. The headspace gas was sampled by a syringe at 2 and 4 h and denitrification rates were calculated from the linear increment of N_2O production against time. Denitrification potential of the soil horizons was measured

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

with the addition of both NO_3^- -N and soluble C source as proposed by D'Haene *et al.* (2003).

Statistical analyses

Analysis of variance (ANOVA) and Pearson correlation analysis were performed using SPSS 13.0. Linear regression and other statistical analyses were carried out by using Excel 2003. Two-way ANOVA and Tukey test were used to compare the mean difference ($P < 0.05$) of a given variable between treatment plots and years. Three-way ANOVA and Tukey test were used to compare the mean difference ($P < 0.05$) in soil N_2O fluxes among the seasons, treatment plots of field experiment, and years, and in soil DEA among the treatment plots of field experiment, soil layers, and incubation treatments.

Deleted: ,

Deleted: plots

RESULTS

Soil temperature and moisture

Daily precipitation is shown in Fig. 1a. Annual precipitations were 1176 mm from the mid April 2005 to the mid April 2006, 1047 mm from the mid April 2006 to the mid April 2007, and 879 mm from the mid April 2007 to the beginning of April 2008. These values are smaller than that the mean annual precipitation (1365 ± 215 mm) from 1989 to 2000.

Soil moisture content expressed as WFPS in a 0 to 6 cm depth is shown in Fig. 1b. The

Deleted: These values are smaller than the long-term average (1365 ± 215 mm).¶

WFPS from April to November was influenced by precipitation and the low soil moisture was observed with low precipitation in August 2006 and June 2007. From December to March, soil moisture was not observed because of soil freezing. In the winter of 2006/2007, soil freezing began from beginning of December 2006, reaching the maximum depth of 17.75 cm in 9th March 2007, and then thawed in early April in 2007. In the winter of 2007/2008, soil freezing also began from beginning of December 2007, but reaching the maximum 29 cm in 12th March 2008, and then thawed in early April in 2008. The soil freezing depth was not observed in the winter of 2005/2006.

Deleted: ¶

Soil temperature at a 5 cm depth is shown in Fig. 1c. The soil temperature increased from April, reaching its maximum from July through August, and then decreased gradually. The soil temperature was around 0 °C from December to March. There was no difference in soil temperature between the chemical fertilizer and manure plots, but soil temperature was higher in the control plot than in the chemical fertilizer and manure plots ($P < 0.05$).

N₂O fluxes

The seasonal patterns of N₂O fluxes were mainly driven by a seasonal variation as varieties of air and soil temperature which were higher in summer and lower in winter, and influenced by fertilization (Fig. 2a). N₂O fluxes in the chemical fertilizer and manure

Deleted: pattern of N₂O fluxes was mainly driven by a seasonal variation

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

plots increased after application of manure or chemical fertilizer. These remained at a higher level than in the control plot until the beginning of September (Fig. 2a). Three-way ANOVA showed that there was a significant difference in N₂O fluxes between the non-growing and growing seasons (P < 0.001) and between each treatment (P < 0.05). In the meantime, an interaction between the season and treatment in the N₂O fluxes was observed (P < 0.05) (Table 2). In the growing season, the mean N₂O fluxes in 2005, 2006, and 2007 were 12.3, 12.9, and 16.7 ug N₂O-N m⁻²h⁻¹ for control plot, 85.3, 83.9, and 36.5 ug N₂O-N m⁻²h⁻¹ for chemical fertilizer plot, and 101.9, 187.6, and 50.6 ug N₂O-N m⁻²h⁻¹ for manure plot, respectively (Table 3). The N₂O fluxes in the growing season were significantly higher in the chemical fertilizer and manure plots than that in the control plot (P < 0.01), but there was no significant difference between the chemical fertilizer and manure plots. In the non-growing season, the N₂O fluxes were lower and stable with the mean values of 0.3, 2.6 and 4.7 ug N₂O-N m⁻²h⁻¹ in the control plot, 2.7, 6.0 and 7.4 ug N₂O-N m⁻²h⁻¹ in the chemical fertilizer plot, and 2.7, 3.6 and 3.5 ug N₂O-N m⁻²h⁻¹ in the manure plot in 2005, 2006, and 2007, respectively (Table 3). There was no significant difference in the mean of N₂O fluxes between each treatment plot.

The annual N₂O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0 and 2.1 to 4.9 kg N₂O-N ha⁻¹yr⁻¹ during 2005 to 2007, respectively

Deleted: N₂O fluxes in the chemical fertilizer and manure plots increased after the application of N fertilizer.

Deleted: 8

Deleted: 4

Deleted: cumulative

Deleted: in

Deleted: , 2006, and

(Table 4). Application of both chemical fertilizer and manure stimulated the annual cumulative N_2O emissions, and the significantly highest annual cumulative N_2O emission was observed in the manure plot, followed by the chemical fertilizer plot. Application of chemical fertilizer contributed to 76.9, 79.2 and 47.2% of the total N_2O emission from the chemical fertilizer plot in 2005, 2006, and 2007, respectively. In the manure plot, N_2O emission from the applied chemical fertilizer and manure contributed to 81.8, 87.4, and 67.6% of the total N_2O emission in 2005, 2006, and 2007, respectively. Chemical fertilizer-induced EF was 1.32, 1.30, and 0.85 % in 2005, 2006, and 2007, respectively. Manure-induced EF was significantly lower than the fertilizer-induced EF ($P < 0.001$), and was 0.51, 0.85, and 0.35 % in 2005, 2006, and 2007, respectively (Table 5).

NO fluxes showed a seasonal variation that similar with the seasonal pattern of N_2O fluxes, which were higher in summer and lower in winter and also influenced by fertilization (Fig. 2b). The NO fluxes ranged from -1.2 to 91.3 $\mu\text{g NO-N m}^{-2}\text{h}^{-1}$ which was smaller than the N_2O fluxes (-3.6 to 1290.7 $\mu\text{g N}_2\text{O-N m}^{-2}\text{h}^{-1}$). Large NO fluxes were observed mainly after the manure and chemical fertilizer application (Fig. 2b). Most of the values of $\text{N}_2\text{O}/\text{NO}$ ratio were distributed from 1 to 100, and a significant positive correlation was found ($P < 0.01$) between the $\text{N}_2\text{O}/\text{NO}$ ratio and the N_2O fluxes (Fig. 3).

Deleted: variability in a

Deleted:

Deleted: seasonal variation and

Deleted: 249.0

Deleted: of fertilizer

Deleted:

Deleted: ly

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Soil chemical properties

During 2005 to 2007, soil pH in chemical fertilizer plot was obviously lower than that in control and manure plot (Fig.4a). Chemical fertilizer application could lead to the soil pH decreasing not only in chemical fertilizer plot but also in manure plot (Fig.4a). The mean soil pH during 2005 to 2007, in control, chemical fertilizer and manure plots were 5.2, 4.6, and 5.1, respectively. The soil pH in the chemical fertilizer plot was significantly lower than that in the manure and control plots ($P < 0.001$). There was no significant difference in soil pH between the manure and control plots.

The mean $\text{NH}_4^+\text{-N}$ concentrations, during 2005 to 2007, in control, chemical fertilizer, and manure plots were 4.4, 23.7, and 17.5 mg kg^{-1} , respectively. Soil $\text{NH}_4^+\text{-N}$ concentration in chemical fertilizer and manure plots exhibited a wide range of 0.4 - 245 mg kg^{-1} (Fig. 4b). In contrast, soil $\text{NH}_4^+\text{-N}$ concentration in the control plot was stable, and was always below 12 mg kg^{-1} . The pattern of soil $\text{NH}_4^+\text{-N}$ concentration was not influenced by manure application only; however, it was influenced by the chemical fertilizer application (Fig. 4b). Soil $\text{NH}_4^+\text{-N}$ concentration in chemical fertilizer and manure plots increased rapidly right after chemical fertilizer application, but then decreased within a few days (Fig. 4b). In 2005 and 2006, peak concentrations of soil $\text{NH}_4^+\text{-N}$ were always observed in both chemical fertilizer and Manure plot after chemical

Deleted: From

Deleted: soil pH from the 0 - 5 cm top soil layer in control, chemical fertilizer and manure plots showed a similar seasonal pattern (Fig. 4a).

Formatted: Font color: Aqua

Deleted: However

Deleted: , t

Deleted: in 2005, 2006, and 2007,

Deleted:

Deleted: in 2005, 2006, and 2007

Deleted: T

Deleted: he pattern of soil $\text{NH}_4^+\text{-N}$ concentration

Deleted: N

Deleted: the N

Deleted: days

Deleted: twice

328 fertilizer application. But in 2007, only one small peak was observed in the chemical
 329 fertilizer plot following the base fertilizer application in May. After chemical fertilizer
 330 application in July 2007, soil NH_4^+ -N concentration peak was not observed may caused
 331 by low precipitation and low soil moisture that limited the additional NH_4^+ -N by chemical
 332 fertilizer application go into more deep soil layer by water dynamics before it was
 333 assumed on the soil surface and in root mat through NH_4^+ -N volatilization or microbial,
 334 chemical and physical reaction.

Deleted: and manure plots

335 The mean NO_3^- -N concentrations, during 2005 to 2007, were 1.4, 2.7, and 2.2 mg kg^{-1} in
 336 the control, chemical fertilizer, and manure plot, respectively. The pattern of soil NO_3^- -N
 337 concentration was also influenced by N application, and the peaks were observed slightly
 338 later than those of soil NH_4^+ -N concentration (Fig. 4c).

Deleted:

Deleted: in 2005, 2006, and 2007

339 During the study period, the soil DOC concentration ranged from 48 to 121 mg kg^{-1} in
 340 the control plot, from 23 to 116 mg kg^{-1} in the chemical fertilizer plot, and from 43 to 199
 341 mg kg^{-1} in the manure plot (Fig. 4d). The mean soil DOC concentrations in control,
 342 chemical fertilizer, and manure plots were 73.3, 59.4, and 97.8 mg kg^{-1} , respectively. The
 343 soil DOC concentration in the manure plot was significantly higher than that in the
 344 control and chemical fertilizer plots ($P < 0.01$), but application of chemical fertilizer had
 345 no significant influence on the soil DOC concentration compared with that in the control

Deleted: 156

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

plot. Continuous 3 years manure application significantly increased the soil DOC concentration, which was significantly higher in 2007 than in 2005 and 2006 (P<0.01).

Pearson correlation analysis showed that instantaneous N₂O flux had a strong positive correlation with soil temperature (P < 0.01), soil NO₃⁻-N concentration (P < 0.01), and soil NH₄⁺-N concentration (P < 0.01) (Table 6).

Soil denitrifying enzyme activity (DEA)

Table 7 shows the result of DEA. Result of a 3-way ANOVA shows that there was a significant difference in soil DEA among soil layers (root-mat and mineral) (P < 0.001) and treatments (with and without NO₃ and glucose) (P < 0.001), but there was no significant difference in soil DEA among plots (control, chemical fertilizer, and manure) (P = 0.058) (Table 8). However, there was a significant interaction between the soil layer and treatments (P < 0.001). In the root-mat layer, soil DEA was significantly increased by the addition of NO₃-N with (P < 0.001) or without (P < 0.001) the addition of glucose. But there was no significant effect on soil DEA by the addition of only glucose. There was no significant difference in soil DEA between the treatments Chl+N and Chl+N+C. On the other hand, there was no significant effect of single addition of NO₃-N (Chl+N) or glucose (Chl+C) on soil DEA in the mineral soil. However, a combination of NO₃-N and glucose addition (Chl+N+C) increased the soil DEA significantly (P < 0.05). The soil

DEA with the addition of both $\text{NO}_3\text{-N}$ and glucose in the root-mat soil was significantly higher than that in the mineral soil ($P < 0.001$). The soil DEA in the root-mat soil with the addition of $\text{NO}_3\text{-N}$ and both $\text{NO}_3\text{-N}$ and glucose had a significantly positive correlation with soil pH ($P < 0.05$) (Fig. 5).

368

DISCUSSION

Seasonal pattern of N_2O emission

Soil N_2O fluxes were significantly higher in the growing season than that in the non-growing season. This is attributed to the high soil temperature (Table 6) in the growing season. Granli and Bøckman (1994) found an increased rate of N_2O production with an increase in soil temperature up to 20–40 °C. High peaks of N_2O fluxes were usually observed in both chemical fertilizer and manure plots within a few weeks after the application of manure or chemical fertilizer in our study (Fig. 2a). This can be attributed to the result of rapid increase in soil $\text{NH}_4^+\text{-N}$ and $\text{NO}_3^-\text{-N}$ concentrations immediately after the application of fertilizer, which decreased within a few days (Fig 4). It is well established that the rate of N_2O emission usually increases with an increase in soil available N (Skiba and Smith 2000; Sehy *et al.* 2003). Several studies reported that N_2O fluxes significantly increased after the application of N fertilizers. Mu *et al.* (2008)

Deleted: fertilizer

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

382 reported that N₂O fluxes increased rapidly to higher emission levels in soils cultivated
383 with wheat (from 242 to 433 µg N m⁻² h⁻¹) and onion (from 47.2 to 157 µg N m⁻² h⁻¹) after
384 N fertilization and that the fluxes lasted for about three weeks. Schils *et al.* (2008) also
385 reported high N₂O fluxes occurred in the first week after the application of chemical
386 fertilizer or cattle slurry.

387 The bacterial processes of nitrification and denitrification are the most important
388 sources of N₂O in soil (Granli and Bøckmann 1994). According to Davidson (1992) and
389 Skiba *et al.* (1993), nitrification produces more NO than N₂O; conversely, denitrification
390 produces more N₂O than NO. The ratio of N₂O-N/NO-N is the index of N₂O production
391 from nitrification or denitrification (Lipschultz *et al.* 1981). Lipschultz *et al.* (1981)
392 reported that the ratio of production of N₂O-N / NO-N ranged from 0.2 to 1.0 in
393 nitrification and 100 in denitrification. A significant positive correlation between the N₂O
394 fluxes and the ratio of N₂O-N / NO-N and was found (Fig. 3) in our study (P < 0.01),
395 indicating that the high N₂O emissions are primarily due to denitrification.

396 **Cumulative N₂O emission**

397 Chemical fertilizer and animal wastes are [the](#) two most important sources of direct N₂O
398 emissions from agricultural soils (Mosier *et al.* 1998). Increasing soil N availability
399 associated with application of N by chemical fertilizer and manure has greatly enhanced

N₂O emissions from agricultural soils (Kroeze *et al.* 1999). Meng *et al.* (2005) found that chemical fertilizer and manure contributed to 74–82% of the total N₂O emissions. Mori *et al.* (2008) also reported the N₂O emission predominantly derived from the manure and the chemical fertilizer N application on a volcanic grassland soil in Nasu, Japan. Generally, emissions of N₂O increase with an increase in N application rates (Granli and Bøckman 1994; MacKenzie *et al.* 1997). In our study, the N₂O emission from applied chemical fertilizer and manure contributed to 77–85% of the total N₂O emission in 2005 and 2006. The contribution of chemical fertilizer and manure to N₂O emission in 2007 decreased to 47–65% due to the lower application rates than that in 2005 and 2006. The chemical fertilizer-induced EF ranged from 0.85 to 1.32%, which was comparable to the IPCC default value 1% (IPCC 2006), but was higher than that reported by Akiyama and Tsuruta (2003) from the Japanese Andisols amended with chemical fertilizer (ranging from 0.06% to 0.29%). The manure-induced EF of our study ranged from 0.35 to 0.85%, which was significantly lower than the chemical fertilizer-induced EF and the IPCC default value, but close to that reported by Akiyama and Tsuruta (2003), which was 0.55%.

Soil DEA

The soil DEA with an addition of NO₃⁻-N and glucose in the root-mat soil was

Deleted: average

Deleted: was 1.17%

Deleted: average

Deleted: was

Deleted: 57

Deleted: Denitrification potential of the soil horizons was measured with the addition of both NO₃⁻-N and soluble C source as proposed by D'Haene *et al.* (2003).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

significantly higher than that in the mineral soil, indicating that the soil denitrification potential in the root-mat soil was significantly higher than that in the mineral soil. Microbial activities in the surface soil are reported to be higher than in the deeper soil (Speir *et al.* 1984; Higashida and Takao, 1985). Parkin and Meisinger (1989) reported that total viable bacteria and numbers of denitrifying bacteria were found to decrease exponentially with an increase in soil depth on a well-drained silt loam soil.

Soil DEA in the root-mat soil significantly increased by the addition of NO₃-N with (P < 0.001) or without (P < 0.001) the addition of glucose, indicating that the availability of soil NO₃⁻-N could be the major limiting factor for soil DEA in our study grassland. In the mineral soil, only addition NO₃-N or glucose could not increase the soil DEA, but addition NO₃-N and glucose together increased the soil DEA that means not only the NO₃-N but also the carbon is the limited factor for soil DEA in mineral soil. The soil DEA in the root-mat soil with NO₃-N addition and both NO₃-N and glucose addition had a significantly positive correlation with soil pH (P < 0.05, Fig. 5). The soil pH is supposed to be a major variable of soil, controlling the microbial community in general and the community of denitrifiers in particular (Simek and Hopkins 1999). Simek and Hopkins (1999) detected an optimum pH value for denitrification in soils as a range from 7 to 8. Simek and Cooper (2002) reported that both the overall rates of denitrification under field

Formatted: Subscript

Formatted: Subscript

Formatted: Subscript

conditions (i.e. the formation of N_2O , N_2 and NO and their subsequent emission) and DEA were influenced by soil pH, and they were less in acidic soils than in neutral or slightly alkaline soils. Ellis *et al.* (1998) observed that the production of N_2O decreased with decreasing pH under anaerobic conditions through an incubation experiment. These results suggest that the highest N_2O emission in the manure plot in our study was resulted from the soil DEA that could have been controlled by the soil pH.

The application of chemical fertilizer significantly decreased the soil pH in the chemical fertilizer plot than in the control plot. However, the soil pH in manure plot was not significantly different with control plot. That maybe because higher pH of manure (8.3-9.1) input decrease the effect of chemical fertilizer application on the soil pH. Soil acidity is controlled by the amount of H^+ and Al^{3+} which is either contained in or generated by the soil and soil components. According to Kirikae *et al.* (2001), nitrification is a source of H^+ through two nitrification pathways of NH_4^+ origin and organic N origin. They reported that the ratio of H^+ to NO_3^- was 2 in the pathway of NH_4^+ origin and 1 in the pathway of organic N origin. In the meanwhile, NO_3^- uptake by vegetation was the sink of H^+ . Therefore, the organic N has less effect on H^+ production than $\text{NH}_4^+\text{-N}$. On the other hand, application of manure increased the value of cation exchange capacity (CEC) compared to that of chemical fertilizer (Bulluck *et al.* 2002).

Deleted: manure application had no significant effect on t

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

The soils with a high CEC have a greater capacity to contain or generate sources of acidity.

The soil pH is higher in soils with manure than that with chemical fertilizer as reported by several studies (Bulluck *et al.* 2002 and Gil *et al.* 2008).

Conclusions

The application of both chemical fertilizer and manure to grassland stimulated the annual N₂O emission. The chemical fertilizer-induced EF (range: 0.85 - 1.32%) was significantly higher than the manure-induced EF (range 0.35 - 0.85%). However, annual N₂O emission was significantly higher in the manure plot than that in the chemical fertilizer plot. The soil DEA in the NO₃-abundant root-mat layer significantly decreased with a decrease in soil pH. Moreover, application of chemical fertilizer could significantly decrease soil pH, but the manure application had no significant effect on soil pH. Therefore, for the a fixed quantity of available N, application of manure could result in higher N₂O emission compared to chemical fertilizer owing to high soil pH values under manure application than under chemical fertilizer application.

ACKNOWLEDGMENTS

We would like to thank the technical staff of Shizunai Livestock Farm for their help in the field measurements. This study was partly supported by a research grant provided by the

Deleted: cumulative

Deleted: 89

Deleted: 33

Deleted: similar

Deleted: application of manure could result in higher N₂O emission compared to chemical fertilizer owing to high pH values in manures.¶

Project entitled 'Establishment of good practices to mitigate Greenhouse Gas emissions from Japanese grasslands' funded by Racing and Livestock Association.

474

475 REFERENCES

476 Akiyama H, Tsuruta H 2003: Effect of organic matter application on N₂O, NO, and NO₂
477 fluxes from an Andisol field. *Global Biogeochem. Cycles*, **17**, 1-16.

478 Anon., 2004: Handbook of animal waste management and utilization in Hokkaido.
479 Hokkaido Prefectural Experiment Stations and Hokkaido Animal Research Center,
480 Sapporo, 64-67 (in Japanese).

481 Bouwman AF, Boumans LJM, Batjes NH 2002: Modeling global annual N₂O and NO
482 emissions from fertilized fields. *Global Biogeochem. Cycles*, **16**, 1080.

483 Brown CD, Fryer CJ, Walker A 2001: Influence of topsoil tilth and soil moisture status on
484 losses of pesticide to drains from a heavy clay soil. *Pest Management Science*, **57**,
485 1127-1134.

486 Bulluck III LR, Brosius M, Evanylo GK, Ristaino JB 2002: Organic and synthetic
487 fertility amendments influence soil microbial, physical and chemical properties on
488 organic and conventional farms. *Applied Soil Ecology*, **19**, 147-160.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

489 Chadwick DR, Pain BF, Brookman SKE 2000: Nitrous oxide and methane emissions
490 following application of animal manures to grassland. *J. Environ. Qual.*, **29**, 277-287.

491 Conrad R 1996: Soil microorganisms as controllers of atmospheric trace gases (H₂, CO,
492 CH₄, OCS, N₂O, and NO). *Microbiol. Rev.*, **60**, 609 - 640.

493 D’Haene K, Moreels E, Neve S, Daguiar BC, Boeckx P, Hofman G, Van Cleemput O
494 2003: Soil properties influencing the denitrification potential of Flemish agricultural
495 soils. *Biol. Fertil. Soils.*, **38**, 358–366.

496 Davidson EA 1992: Source of nitric oxide and nitrous oxide following wetting of dry soil.
497 *Soil Sci. Soc. Am. J.*, **56**, 95-102.

498 Dobbie, KE, Smith KA 2003: Nitrous oxide emission factors for agricultural soil in Great
499 Brita: the impact of soil water-filled pore space and other controlling variables.
500 *Global Change Biol.*, **9**, 204-218.

501 Ellis S, Howe MT, Goulding WT, Muggleston M.A, Dendooven, L 1998: Carbon and
502 nitrogen dynamics in a grassland soil with varying pH: effect of pH on the
503 denitrification potential and dynamics of the reduction enzymes. *Soil Biol. Biochem.*,
504 **30**, 359-367.

- 1
2
3
4
5 505 Gil MV, Carballo MT and Calvo LF 2008: Fertilization of maize with compost from
6
7
8 506 cattle manure supplemented with additional mineral nutrients. *Waste Management*,
9
10 507 **28**, 1432-1440.
11
12
13 508 Granli T and Bøckman OC 1994: Nitrous oxide from agriculture. *Nor. J. Agric. Sci.*,
14
15 509 *Suppl.*, **12**, 7-128.
16
17
18 510 Higashida S, and Takao K 1985: Seasonal fluctuation patterns of microbial numbers in
19
20
21 511 the surface of a grassland. *Soil Sci. plant Nutr.*, **31**, 113-121.
22
23
24 512 IPCC 2006: Guidelines for National Greenhouse Gas Inventories, prepared by the
25
26 513 National Greenhouse Gas Inventories Programme. Intergovernmental Panel on
27
28 514 Climate Change.
29
30
31 515 IPCC 2007: Climate changes 2007: The Physical Science Basis. Cambridge University
32
33
34 516 Press, Cambridge.
35
36 517 IUSS Working Group WRB 2006: World Reference Base for Soil Resources 2006.
37
38
39 518 World Soil Resources Reports 103, FAO, Rome.
40
41
42 519 Jones SK, Rees RM, Skiba UM, Ball BC 2007: Influence of organic and mineral N
43
44 520 fertiliser on N₂O fluxes from a temperate grassland. *Agr. Ecosyst. Environ.*, **121**,
45
46
47 521 74–83.
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

522 Kirikae M, Hatano R, Shibata H, Tanaka Y 2001: Analysis of proton generation and
523 consumption of forest surface soils in Hokkaido, northern Japan. *Water, Air and Soil
524 Pollution*, **130**, 697-702.

525 Kroeze C, Mosier AR, Bouwman AF 1999: Closing the global N₂O budget: A
526 retrospective analysis 1500 - 1994. *Global Biogeochem. Cycles*, **13**, 1-8.

527 Linn DM, Doran JW 1984: Effect of water-filled pore-space on carbon-dioxide and
528 nitrous-oxide production in tilled and nontilled soils. *Soil Sci. Soc. Am. J.*, **48**,
529 1267-1272.

530 Lipschultz F, Zafirou OC, Wofsy SC, McElroy MB, Valois FW, Watson SW 1981:
531 Production of NO and N₂O by soil nitrifying bacteria. *Nature*, **294**, 641-643.

532 MacKenzie AF, Fan MX, Cadrin F 1997: Nitrous oxide emission as affected by tillage,
533 corn-soybean-alfalfa rotations and nitrogen fertilization. *Can. J. Soil Sci.*, **77**,
534 145-152.

535 McTaggart IP, Douglas JT, Clayton H, Smith KA 1997: Nitrous oxide emissions from
536 slurry and mineral nitrogen applied to grassland. In: Jarvis, S.C., Pain, B.F. (Eds.),
537 Gaseous Nitrogen Emission from Grasslands. CAB International, Wallingford, UK,
538 209-210.

- 539 Meng L, Ding W, Cai Z 2005: Long-term application of organic manure and nitrogen
540 fertilizer on N₂O emissions, soil quality and crop production in a sandy loam soil. *Soil
541 Biol. Biochem.*, **37**, 2037-2045
- 542 Mogge B, Kaiser EA, Munch JC 1999: Nitrous oxide emissions and denitrification
543 N-losses from agricultural soils in the Bornhoved Lake region: influence of organic
544 fertilizers and land-use. *Soil Biol. Biochem.*, **31**, 1245-1252.
- 545 Mori A, Hojito M, Shimizu M, Matsuura S, Miyaji T, Hatano R 2008: N₂O and CH₄
546 fluxes from a volcanic grassland soil in Nasu, Japan: Comparison between manure plus
547 fertilizer plot and fertilizer-only plot. *Soil Sci. Plant Nutr.*, **54**, 606-617.
- 548 Mosier A, Kroeze C, Nevison C, Oenema O, Seitsinger S, van Cleemput O 1998: Closing
549 the global N₂O budget: nitrous oxide emissions through the agricultural nitrogen
550 cycle. *Nutr. Cycling Agroecosyst.*, **52**, 225-248.
- 551 Mu Z, Kimura SD, Toma Y, Hatano R 2008: Nitrous oxide fluxes from upland soils in
552 central Hokkaido, Japan. *J. Environ. Qual.*, **20**, 1312-1322.
- 553 Myhre G, Highwood EK, Shine KP, Stordal F 1998: New estimates of radiative forcing
554 due to well mixed greenhouse gases. *Geophys. Res. Lett.*, **25**, 2715-2718.
- 555 Parkin TB, Meisinger JJ 1989: Denitrification below the crop rooting zone as influenced
556 by surface tillage. *J. Environ. Qual.*, **18**, 12-16.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

557 Schils RLM, van Groenigen JW, Velthof GL, Kuikman PJ 2008: Nitrous oxide emissions
558 from multiple combined applications of fertiliser and cattle slurry to grassland. *Plant*
559 *and Soil*, **310**, 89-101.

560 Scott A, Ball BC, Crichton IJ, Aitken MN 2000: Nitrous oxide and carbon dioxide
561 emissions from grassland amended with sewage sludge. *Soil Use Manage.*, **16**,
562 36-41.

563 Sehy U, Ruser R, Munch JC 2003: Nitrous oxide fluxes from maize fields: relationship to
564 yield, site-specific fertilization, and soil conditions. *Agric. Ecosyst. Environ.*, **99**,
565 97-111.

566 Shiga H, Ohyama N, Maeda K, Suzuki M 1985: An evaluation of different organic
567 materials based on their decomposition pattern in paddy soils. *Res. Bull. Natl. Agric.*
568 *Res. Cetr.*, **5**, 1-19 (in Japanese with English summary).

569 Shimizu M, Marutani S, Desyatkin AR, Jin T, Hata H, Hatano R 2009: The effect of
570 manure application on carbon dynamics and budgets in a managed grassland of
571 Southern Hokkaido, Japan. *Agric. Ecosyst. Environ.*, **130**, 31-40.

572 Simek M, Cooper JE 2002: The influence of soil pH on denitrification-progress towards
573 the understanding of this interaction over the last 50 years. *European Journal of Soil*
574 *Science*, **53**, 345-354.

Formatted: Font: Italic

Deleted: Title:

Formatted: Font: Italic

- 575 Simek M, Hopkins DW 1999: Regulation of potential denitrification by soil pH in
576 long-term fertilized arable soils. *Biol. Fertil. Soils*, **30**, 41-47.
- 577 Skiba U, Smith K.A 2000: The control of nitrous oxide emissions from agricultural and
578 natural soils. *Chemosphere Global Change Sci.*, **2**, 379-386.
- 579 Skiba U, Smith KA, Fowler D 1993: Nitrification and denitrification as source of nitric
580 oxide and nitrous oxide in a sandy loam soils. *Soil Biol. Biochem.*, **25**, 1527-1536.
- 581 Soil Survey staff 2006: Keys to Soil Taxonomy, 10th ed. USDA-Natural Resources
582 Conservation Service, Washington DC.
- 583 Soussana JF, Allard V, Pilegaard K 2007: Full accounting of the greenhouse gas (CO₂,
584 N₂O, CH₄) budget of nine European grassland sites. *Agric. Ecosyst. Environ.*, **121**,
585 121-134.
- 586 Speir T, Ross DJ, Orchard VA 1984: Spatial Variability of biochemical properties in a
587 taxonomically-uniform soil under grazed pasture. *Soil Biol Biochem.*, **16**, 153-160.
- 588 Tiedje JM 1988: Ecology of denitrification and dissimilatory nitrate reduction to
589 ammonium. In: Zehnder, A.J.B., (Ed.), *Biology of Anaerobic Microorganisms*,
590 Wiley, New York, 179-244.
- 591 Tiedje JM, 1994: Denitrifiers. *Methods of Soil Analysis, Part 2. Microbiological and*
592 *Biochemical Properties*. Soil Sci. Soc. Amer., Madison, Wisconsin, 245-267.

Deleted: ,

Deleted: .

Deleted: .

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

593

For review

594

595 **Figure captions**

596 **Figure 1** Seasonal patterns of precipitation (a), WFPS at 6 cm depth (b), and soil
 597 temperature at 5 cm depth (c). Data of WFPS and soil temperature represent means $\pm$
 598 SD (n = 4 to 6).

599 **Figure 2** Seasonal patterns of soil N₂O fluxes (a) and soil NO fluxes (b). Data represent
 600 means $\pm$ SD (n = 4 to 6). Full arrow indicates date of the chemical fertilizer
 601 application, and dotted arrow indicates date of manure application. The growing
 602 season was 215 days in 2005 (From 10th April 2005 to 10th November 2005), 218
 603 days in 2006 (From 15th April 2006 to 18th November 2006) and 220 days in 2007
 604 (From 13th April 2007 to 18th November 2007).

Deleted: The arrows indicate the
date of fertilizer application.

605 **Figure 3** Relationships between N₂O fluxes and the ratio of N₂O to NO.

606 **Figure 4** Seasonal patterns of Soil pH (a), soil NH₄⁺-N (b), soil NO₃⁻-N (c), and soil DOC
 607 (d) at a depth of 0 - 5 cm. Data represent means $\pm$ SD (n = 3). Full arrow indicates
 608 date of the chemical fertilizer application, and dotted arrow indicates date of manure
 609 application. The arrows indicate the date of fertilizer application.

610 **Figure 5** Relationship between the soil DEA and the soil pH in root-mat soil.

We evaluated the effect of chemical fertilizer and manure applications on N₂O emission from a managed grassland by establishing three treatment plots of chemical fertilizer, manure, and control at the Shizunai Experimental Livestock Farm in southern Hokkaido, Japan. Seasonal N₂O and NO fluxes were measured by a closed-chamber method at 4 to 6 replications in each treatment plot from May 2005 to April 2008. Soil samples were collected from a 0-5 cm top soil layer at three replications on each gas sampling date for measuring pH, NO₃⁻-N, NH₄⁺-N, and DOC. Soil samples were collected from the root-mat layer (0-2.5cm) and the mineral soil layer (2.5-5cm) of each treatment plot in April, June, and August 2007, which was followed by measuring soil denitrifying enzyme activity (DEA). The soil DEA was measured by an acetylene inhibition method under the four treatments with and without the addition of NO₃⁻-N and glucose. The cumulative N₂O emission in control, chemical fertilizer, and manure plots ranged from 0.6 to 0.7, 1.4 to 3.0, and 2.1 to 4.9 kg N₂O-N ha⁻¹yr⁻¹, respectively. The application of both chemical fertilizer and manure significantly increased the cumulative N₂O emission, and the highest cumulative N₂O emission was observed in the manure plot. N₂O fluxes were positively correlated with the N₂O:NO ratio (P < 0.01), indicating high N₂O fluxes resulting from the increased denitrification activities. The denitrification potential of the soil horizons was significantly higher in the root-mat soil than that in the mineral soil. The soil DEA in the root-mat soil in NO₃⁻-N addition with and without the addition of glucose had a significantly positive correlation with soil pH (P < 0.05). The soil pH was significantly influenced by N fertilization, which was significantly lower in the chemical fertilizer plot than that in the control and manure plots. For the similar quantity of available N,

1
2
3 application of manure could result in higher N₂O emission compared to chemical fertilizer
4
5
6 owing to high pH values in manures.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1 The applied date and the application rates of chemical fertilizer and manure during the study period

Treatment	Date	Fertilizer type	Application rates (kg ha ⁻¹)			
			C	N	P ₂ O ₅	K ₂ O
Control	2005/5/11	Chemical fertilizer ^a	0	0	0	0
	2005/7/4	Chemical fertilizer ^a	0	0	0	0
	2006/5/9	Chemical fertilizer ^a	0	0	0	0
	2006/7/10	Chemical fertilizer ^a	0	0	0	0
	2007/5/12	Chemical fertilizer ^a	0	0	14	73
	2007/7/5	Chemical fertilizer ^a	0	0	7	37
Chemical fertilizer	2005/5/11	Chemical fertilizer ^a	0	103	23	168
	2005/7/4	Chemical fertilizer ^a	0	61	23	97
	2006/5/9	Chemical fertilizer ^a	0	124	50	177
	2006/7/10	Chemical fertilizer ^a	0	59	18	97
	2007/5/12	Chemical fertilizer ^a	0	49	14	73
	2007/7/5	Chemical fertilizer ^a	0	25	7	37
Manure	2005/5/11	Manure ^b	5833	236	191	266
	2005/7/4	Chemical fertilizer ^a	0	133	7	70
	2006/5/9	Manure ^b	5958	310	212	167
	2006/5/9	Chemical fertilizer ^a	0	71	0	33
	2006/7/10	Chemical fertilizer ^a	0	59	6	97
	2007/5/12	Manure ^b	7714	331	342	336
	2007/7/5	Chemical fertilizer ^a	0	21	0	0

^a Chemical fertilizer is comprised of ammonium sulfate, ammonium phosphate, potassium sulfate, and potassium magnesium sulfate.

^b Beef cattle manure with bedding litter was applied in the manure plot.

Table 2 ANOVA results for mean N₂O fluxes

Source	df	Mean Square	F	P value
Season	1	179,974.04	17.521	0
Year	2	13,990.75	1.362	0.258
Treatment	2	39,922.53	3.887	0.022
Season * Year	2	15,307.66	1.49	0.228
Season * Treatment	2	38,549.47	3.753	0.025
Year * Treatment	4	7,659.25	0.746	0.562
Season * Year * Treatment	4	7,454.57	0.726	0.575
Error	210	10,271.71		

Deleted: Sig

For review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 3 Mean N₂O fluxes from the control, chemical fertilizer, and manure plots.

Season	Treatment	Mean N ₂ O fluxes (ug N ₂ O-N m ⁻² h ⁻¹)		
		2005	2006	2007
Growing season	Control	12.3 (12.7)	12.9 (15.23)	16.7 (27.6)
	Chemical fertilizer	85.3 (94.1)	83.9 (66.3)	36.5 (53.2)
	Manure	101.9 (102.1)	187.6 (301.6)	50.6 (48.9)
Non-growing season	Control	0.3 (0.6)	2.6 (6.0)	4.7 (3.1)
	Chemical fertilizer	2.7 (2.3)	6.0 (11.4)	7.4 (13.7)
	Manure	2.7 (3.9)	3.6 (5.1)	3.5 (8.0)

Data represent means (SD), n=4 to 6. We defined the crop growing season as a 7-day moving average of daily air temperature above 5 °C and the non-growing season as the rest (Shimizu *et al.* 2009). The growing season was 215 days in 2005 (From 10th April 2005 to 10th November 2005), 218 days in 2006 (From 15th April 2006 to 18th November 2006) and 220 days in 2007 (From 13th April 2007 to 18th November 2007).

Formatted: Font: 12 pt

Deleted: .

Formatted: Font: 12 pt

Table 4 Cumulative N₂O emissions from the control, chemical fertilizer, and manure plots.

Treatment	Cumulative N ₂ O emissions (kg N ₂ O-N ha ⁻¹ yr ⁻¹)		
	2005	2006	2007
Control ^a	0.7 (0.4)	0.6 (0.3)	0.7 (0.5)
Chemical fertilizer ^b	2.8 (0.7)	3.0 (0.8)	1.4 (0.5)
Manure ^c	3.6 (1.2)	4.9 (2.8)	2.1 (0.6)

Data represent means (SD), n=4 to 6, and different small letters denote the significant difference at the 0.05 level between each treatment.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 5 N₂O Emission factor for chemical fertilizer and manure (kg N₂O–N (kg N input)⁻¹)

	N ₂ O emission factor (%)		
	2005	2006	2007
Chemical fertilizer	1.32 (0.43)	1.30 (0.44)	0.85 (0.97)
Manure	0.51 (0.42)	0.85 (0.89)	0.35 (0.23)

Data represent means (SD), n=4 to 6.

For review

Table 6 Relationships (Pearson correlation coefficient, r) between instantaneous N_2O fluxes and environmental factors using the whole data

	N_2O	Soil T.	WFPS	pH	NO_3^-	NH_4^+	DOC
N_2O	1						
Soil T.	0.330 **	1					
WFPS	0.020	-0.288 **	1				
pH	-0.117	-0.275 **	0.378 **	1			
NO_3^-	0.307 **	0.378 **	-0.061	-0.201 *	1		
NH_4^+	0.329 **	0.156 *	-0.048	-0.273 **	0.380 **	1	
DOC	-0.104	-0.173 *	-0.204 *	0.341 **	-0.282 **	-0.040	1

* $p < 0.05$; ** $p < 0.01$

Deleted: .

Deleted: 1

... [1]

Deleted: 1

... [2]

Deleted: 1

... [3]

Deleted: 1

... [4]

Deleted: 1

... [5]

Deleted: 1

... [6]

Deleted: 1

For review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 7 Denitrification enzyme activity (DEA) of soil samples from three treatment plots of control, chemical fertilizer, and manure

Date	Soil layer	Treatment	DEA (mgN ₂ O-N kg ⁻¹ h ⁻¹)					
			Control plot		Chemical fertilizer plot		Manure plot	
2007/4/29	Root-mat	Chl	0.04	(0.04)	0.04	(0.01)	0.04	(0.02)
		Chl+N	9.99	(1.07)	3.66	(0.50)	16.30	(2.98)
		Chl+C	0.06	(0.01)	0.07	(0.02)	0.14	(0.07)
		Chl+N+C	10.98	(6.71)	3.08	(0.49)	13.56	(3.16)
	Mineral	Chl	2.68	(3.84)	0.16	(0.11)	0.58	(0.34)
		Chl+N	0.52	(0.64)	0.16	(0.03)	1.56	(0.25)
		Chl+C	0.02	(0.12)	0.22	(0.01)	0.07	(0.14)
		Chl+N+C	1.00	(0.20)	2.21	(1.50)	1.97	(0.36)
2007/6/11	Root-mat	Chl	0.15	(0.02)	0.01	(0.00)	0.10	(0.05)
		Chl+N	14.05	(5.03)	2.63	(0.68)	15.05	(2.28)
		Chl+C	0.08	(0.02)	0.02	(0.01)	0.13	(0.03)
		Chl+N+C	9.47	(0.33)	4.66	(0.40)	17.35	(0.81)
	Mineral	Chl	0.09	(0.03)	0.07	(0.02)	0.71	(0.09)
		Chl+N	1.43	(0.00)	0.35	(0.10)	1.24	(0.55)
		Chl+C	0.03	(0.00)	0.04	(0.01)	0.05	(0.14)
		Chl+N+C	2.14	(0.21)	0.54	(0.15)	1.71	(2.15)
2007/8/20	Root-mat	Chl	0.12	(0.05)	0.20	(0.06)	0.04	(0.01)
		Chl+N	27.84	(3.87)	9.80	(0.18)	13.58	(1.83)
		Chl+C	0.13	(0.06)	0.14	(0.05)	0.03	(0.02)
		Chl+N+C	21.33	(0.23)	35.19	(37.29)	19.91	(1.73)
	Mineral	Chl	1.14	(0.31)	0.76	(0.37)	3.01	(0.20)
		Chl+N	1.34	(0.17)	0.55	(0.09)	2.49	(1.83)
		Chl+C	0.37	(0.03)	0.64	(0.33)	1.03	(1.22)
		Chl+N+C	2.92	(0.23)	0.83	(0.33)	7.65	(0.87)

Data represent means (SD), n=3.

Table 8 ANOVA results for soil denitrification enzyme activity (DEA).

Source	df	Mean Square	F	P value
Plot	2	94,796,698	3.351	0.058
Soil layer	1	991,237,940	35.040	0.000
Treatment	3	832,401,859	29.426	0.000
Plot * Soil layer	2	9,030,352	0.319	0.390
Plot * Treatment	6	53,775,775	1.901	0.174
Soil layer * Treatment	3	451,061,076	15.945	0.000
Plot * Soil layer * Treatment	5	25,358,833	0.896	0.258
Error	37	28,288,420		

Deleted: .

Deleted: Sig.

For review

Page 6: [1] Deleted	Jin	11/24/2009 7:25:00 AM
1		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
0.330 **		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
0.020		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
-0.117		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
0.307 **		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
0.329 **		
Page 6: [1] Deleted	Jin	11/24/2009 7:24:00 AM
-0.104		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
1		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
-0.288 **		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
-0.275		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
0.378 **		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
0.156 *		
Page 6: [2] Deleted	Jin	11/24/2009 7:24:00 AM
-0.173 *		
Page 6: [3] Deleted	Jin	11/24/2009 7:25:00 AM
1		
Page 6: [3] Deleted	Jin	11/24/2009 7:24:00 AM
0.378 **		
Page 6: [3] Deleted	Jin	11/24/2009 7:24:00 AM
-0.061		
Page 6: [3] Deleted	Jin	11/24/2009 7:24:00 AM
-0.048		
Page 6: [3] Deleted	Jin	11/24/2009 7:24:00 AM
-0.204 *		
Page 6: [4] Deleted	Jin	11/24/2009 7:25:00 AM
1		
Page 6: [4] Deleted	Jin	11/24/2009 7:25:00 AM
-0.201 *		
Page 6: [4] Deleted	Jin	11/24/2009 7:25:00 AM

-0.273 **

Page 6: [4] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

0.341 **

Page 6: [5] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

1

Page 6: [5] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

0.380 **

Page 6: [5] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

-0.282 **

Page 6: [6] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

1

Page 6: [6] Deleted	Jin	11/24/2009 7:25:00 AM
---------------------	-----	-----------------------

-0.040

For review

Follow the 'Figure captions' in the manuscript.