Title	Phylogenetic history of mustelid fauna in Taiwan inferred from mitochondrial genetic loci
Author(s)	Hosoda, T.; Sato, J. J.; Lin, L. -K.; Chen, Y. -J.; Harada, M.; Suzuki, H.
Citation	Canadian Journal of Zoology, 89(6), 559-569 https://doi.org/10.1139/Z11-029
Issue Date	2011-06
Doc URL	http://hdl.handle.net/2115/46203
Type	article (author version)
File Information	CJZ89-6_559-569.pdf

[Instructions for use](#)

Phylogenetic history of mustelid fauna in Taiwan inferred from mitochondrial genetic loci.

T. Hosoda¹, J. J. Sato^{2,*}, L.-K. Lin³, Y.-J. Chen⁴, M. Harada⁵, and H. Suzuki⁶

¹ *Taikyu High School, 1985 Yuasa-cho, Arida-gun, Wakayama 643-0004, Japan (E-mail: tehosoda@cameo.plala.or.jp)*

² *Laboratory of Animal Cell Technology, Faculty of Life Science and Technology, Fukuyama University, Higashimura-cho, Aza, Sanzo, 985, Fukuyama 729-0292, Japan (E-mail: jsato@bt.fybt.fukuyama-u.ac.jp)*

³ *Laboratory of Wildlife Ecology, Department of Biology, Tunghai University, R. O. C., Taichung 407, Taiwan (E-mail: lklin@thu.edu.tw)*

⁴ *Department of Zoology, National Museum of Natural Science, 1, Kuan Chien RD., Taichung 404, Taiwan (E-mail: cyj@mail.nmns.edu.tw)*

⁵ *Laboratory Animal Center, Graduate School of Medicine, Osaka City University, Osaka 545-8585, Japan (E-mail: haradam@med.osaka-cu.ac.jp)*

⁶ *Laboratory of Ecology and Genetics, Graduate School of Environmental Earth Science, Hokkaido University, Kita-ku, Sapporo 060-0810, Japan (E-mail: htsuzuki@ees.hokudai.ac.jp)*

Running title: Phylogenetic history of the Taiwanese mustelid fauna

*Corresponding author.

Tel: +81-84-936-2111

Fax: +81-84-936-2459

E-mail address: jsato@bt.fybt.fukuyama-u.ac.jp (JJ Sato)

Authors: T. Hosoda, J. J. Sato, L.-K. Lin, Y.-J. Chen, M. Harada, and H. Suzuki

Title: Phylogenetic history of mustelid fauna in Taiwan inferred from mitochondrial genetic loci.

Abstract: Phylogenetic relationships among species of the family Mustelidae were examined using the combined nucleotide sequences of the three mitochondrial genetic loci, cytochrome *b* (MT-CYB; 1140 bp), NADH dehydrogenase subunit 2 (MT-ND2; 1044 bp), and displacement loop (MT-DLOOP; 540 bp), with special emphasis on the phylogenetic history of four Taiwanese mustelid species: *Martes flavigula* (Boddaert, 1785), *Melogale moschata* (Gray, 1831), *Mustela nivalis* L., 1766, and *Mustela sibirica* Pallas, 1773. Maximum likelihood phylogenetic analysis of the combined sequences of the mitochondrial genetic loci produced a topology largely congruent with that of previous studies at species level. Analyses of intraspecific genetic variations revealed two *Melogale moschata* individuals from Taiwan and Vietnam that showed genetic distances comparable to interspecific variations within the mustelid lineages. Furthermore, *Mustela nivalis*, recently discovered in Taiwan, was not as genetically differentiated from other continental conspecific individuals as a previous morphological survey suggested. Divergence time estimations for the mustelid lineages of Taiwan and the Eurasian continent by the Bayesian relaxed-molecular clock approach suggested multiple colonization of Taiwan by mustelids from the continent during the Pleistocene, creating a hierarchical pattern of endemism based on the differential isolation history of the mustelid species in Taiwan.

Introduction

Islands are hotspots of biodiversity (Whittaker and Fernandez-Palacios 2007), and the eastern peripheral islands of the Eurasian continent have played an important role in creating novel evolutionary lineages and harboring historically diverse organisms (Suzuki 2009). The unique characteristics of island organisms are thought to be the result of repeated isolations from and connections to the continental mainland (Ota 1998; Otsuka and Takahashi 2000). Thus, the endemism of island biota is achieved through a process of iterative receptions of continental organisms and subsequent geographic isolations, culminating in global centers of endemic richness (Kier et al. 2009). Among these centers of richness are the East Asian marginal islands (Whittaker and Fernandez-Palacios 2007).

The islands of Taiwan are located at the southeastern edge of the Eurasian continent. The main island of Taiwan is thought to have been formed by the collision of the Philippine Sea and continental Eurasian plates (Page and Suppe 1981; Ho 1986; Teng 1990; Huang et al. 1997). Since the emergence of the island in the Pliocene (Teng 1990; Huang et al. 1997), intermittent connections to the Asian mainland are thought to have occurred (Kizaki and Oshiro 1980; Kimura 2000), possibly due to sea-level lowering in glacial periods (Gascoyne et al. 1979; Voris 2000; Wang 2004). Therefore, the island likely has received multiple invasions of organisms from the Asian continent, as suggested in the palaeontological record (e.g., Otsuka and Takahashi 2000; Takahashi et al. 2001). The complicated biota of Taiwan is an intermingling of various organisms with different histories. The zoogeographic (faunal) region of Taiwan is near the boundary between the northern Palaearctic subregion of the Holarctic region and the southern Oriental subregion of the Paleotropical region. Thus, various northern and southern elements of the Asian continent are found in the Taiwan islands (Kano 1940; Tzeng 1986; Lin 2000), and no chronologically and biogeographically congruent evolutionary scenario explains the current patterns of species distributions in Taiwan.

Recent phylogeographic studies based on molecular genetic approaches have sought to unravel the complex genealogies of Taiwanese fauna. Two major hypotheses have been proposed to explain genetic variations observed on the island. The first proposes *in situ* differentiation, in which variations have been generated due to interruption of within-island gene-flow by physical barriers, e.g., a central mountain range for Pallas's squirrel *Callosciurus erythraeus* (Pallas, 1779) (Oshida et al. 2006) and the Indian rice frog *Rana limnocharis* Gravenhorst, 1829 (Toda et al. 1997, 1998). Some refugia during the Pleistocene may also have functioned as a mechanism of *in situ* differentiation (Wang et al. 1999; Oshida et al. 2006; Yuan et al. 2006). The alternative hypothesis is immigration from the mainland, in which multiple colonization events occurred via land bridges formed sporadically during the Pleistocene at the current Taiwan Strait (Yu 1995; Yu et al., 1996; Tu et al. 2000; Creer et al. 2001, 2004; Lin et al. 2002; Jang-Liaw et al. 2008; Zhong et al. 2008). Determining which hypothesis better explains the genetic variations of specific organisms in Taiwan is difficult (Oshida et al. 2006; Watanabe et al. 2007), owing to limited information about the geology of Taiwan-continent interactions and the lack of comparisons between Taiwanese and continental organisms. Knowledge of the influence of island history on the modern fauna in Taiwan is thus fragmentary. A comparison of Taiwanese lineages of diverse organisms to corresponding continental counterparts with different origins and histories would help to disentangle the complex histories of the Taiwanese fauna. In particular, comparisons of the possible colonization ages of different organisms would provide insight into the importance of the landbridge hypothesis.

We evaluated the genetic status of the lineages of the four Taiwanese species in the family Mustelidae (Carnivora, Mammalia): *Martes flavigula* (Yellow-throated marten), *Melogale moschata* (Chinese ferret badger), *Mustela nivalis* (Least weasel), and *Mustela sibirica* (Siberian weasel). The former two species mostly inhabit the southern Oriental subregion,

Mustela nivalis has expanded to the northern Holarctic (both Palaearctic and Nearctic) region, and *Mustela sibirica* has both Oriental and Palaearctic distributions (Wozencraft 2005). Thus, the mustelid fauna in Taiwan provides an excellent opportunity to test the role of different processes involved in the assembly of an island fauna. We examined nucleotide sequences of three mitochondrial genetic loci, cytochrome *b*, NADH dehydrogenase subunit 2, and displacement loop of mustelids, with special emphasis on Taiwanese and continental individuals of the four species present in Taiwan. *Lutra lutra* (L., 1758) and *Aonix cinerea* (Illiger, 1815), which inhabit Taiwan, were not examined in this study. On the basis of the genetic diversity of the Taiwanese mustelids, we considered their taxonomy and evolutionary history.

Materials and Methods

Sample collection and gene nomenclature

In total, we examined 24 species of the family Mustelidae (Carnivora, Mammalia). Four species (*Martes flavigula*, *Melogale moschata*, *Mustela nivalis*, and *Mustela sibirica*) included individuals from both Taiwan and the continental mainland. One to two representatives of the other species were analyzed. We determined the nucleotide sequences of one to three mitochondrial genetic loci for 45 individuals from 24 mustelid species and downloaded 32 sequences from the DDBJ/EMBL/GenBank international DNA database (Table 1). Gene locus symbols and names follow the nomenclature of the homologous human locus, as approved by the Human Genome Organisation (HUGO) Gene Nomenclature Committee, and are listed in the Human Gene Nomenclature Database (<http://www.genenames.org/>). Cytochrome *b* and NADH dehydrogenase subunit 2 were designated MT-CYB and MT-ND2, respectively. For the displacement loop region, which has been called the D loop or control region, sub-regions are named differently in the database.

For brevity, however, we designated it as the MT-DLOOP, following the naming system for MT-CYB and MT-ND2. Novel sequences were deposited in the international DNA databases as accession numbers AB564113–AB564168 and AB601552–AB601597 (Table 1).

PCR amplification and sequencing strategy

Total genomic DNA was extracted from tissues preserved in ethanol using the conventional phenol–chloroform method (Sambrook and Russell 2001). The DNA was amplified with nested polymerase chain reactions (PCRs) using an automated thermal cycler (model PC 808, Astec). For the PCR of MT-CYB and MT-ND2, we used Ex *Taq* Hot Start Version (Takara) according to the manufacturer's instructions. Each PCR mix was composed of Ex *Taq* Buffer containing 2.0 mM MgCl₂, 0.2 mM of each dNTP, 0.8 μM of each primer, 1.25 units of TAKARA Ex *Taq*™ HS polymerase, and 0.1–0.2 μg of template total genomic DNA in a total volume of 50 μl. The thermal cycling parameters for the first PCR consisted of 94°C for 3 min, 30 cycles of 94°C for 30 s, 50°C for 30 s, and 72°C for 1.5 min, and a final 72°C for 10 min. A 1-μl aliquot of each reaction mixture from the first PCR product was used as template for the second nested PCR in a 50-μl reaction mixture. The second nested PCR mixture included the same reagents as the first PCR, except for the primer pairs. The thermal cycling for the second nested PCR was performed using essentially the same conditions as the first PCR, except that the extension was decreased from 1.5 min to 40 s based on the expected length of the amplified fragments. For PCR of the MT-DLOOP, we used AmpliTaq Gold PCR Master Mix (Life Technologies) according to the manufacturer's instructions. Each PCR mix contained the appropriate amount of AmpliTaq DNA polymerase as well as the correct buffer, 2.5 mM MgCl₂, 0.2 mM of dNTP, 0.5 μM of each primer, and 0.1–0.2 μg of template total genomic DNA in a total volume of 20 μl. The thermal cycling parameters of the PCR were the same as in the second PCR for the MT-CYB and MT-ND2 analyses. The primers

used in this study are the same as those reported in Sato *et al.* (2009b). As a negative control, PCR without DNA template was performed, and no amplification was seen in the negative control. The final PCR products were sequenced according to the manufacturer's instructions using the BigDye Terminator Cycle Sequencing kit v3.1 (ABI), followed by automated sequencing on an ABI3130 genetic analyzer.

Multiple alignments and applied data matrix

The MT-CYB and MT-ND2 sequences generated in this study and collected from the DNA database were aligned manually, without using an alignment program, because the absence of sequence-length variation, which likely stemmed from functional constraints on these protein-coding regions, made alignment straightforward. Conversely, we used the program MUSCLE version 3.8 (Edgar 2004) for the multiple alignment of the MT-DLOOP due to the non-trivial nature of the alignment caused by sequence length polymorphisms. Sequence fragments that could not be aligned unambiguously were excluded from the analyses to avoid reconstruction of a phylogeny based on erroneous sequence homology. From the 574-bp alignment length of the MT-DLOOP sites after the MUSCLE analysis, we excluded 34 sites (sites 21–30 and 77–100) because of alignment ambiguities. Part of the MT-CYB sequence could not be determined for one *Martes flavigula* (HS844, the initial 593 bp were missing), two *Melogale moschata* (AK703 and HS2372, the initial 528 bp were missing), and one *Mustela erminea* L., 1758 (HS678, the final 6 bp were missing) because of unknowns in the PCR and sequencing processes. Despite some missing characters in our data, the large set of existing characters could lead to a correct phylogenetic reconstruction (Wolsan and Sato 2010). In our data, the sequences were determined for all taxa for many regions. Therefore, we treated the above-mentioned undetermined data as missing characters with the assumption that a small proportion of missing characters does not significantly affect the

phylogenetic reconstruction (*e.g.*, Wiens 2006). The assembled data matrix is available from TreeBASE (<http://www.treebase.org/treebase-web/home.html>) as study accession number S10586.

Phylogenetic analyses

Phylogenetic analyses were conducted on the combined mitochondrial genetic data consisting of 2,724 -bp nucleotide sequences of MT-CYB, MT-ND2, and MT-DLOOP using the maximum likelihood (ML; Felsenstein 1981) approach. Trees were rooted using species of the subfamily Guloninae (martens, wolverine, and allies), based on the phylogenetic hypothesis that the subfamilies Helictidinae (ferret badgers), Lutrinae (otters and allies), and Mustelinae (minks, polecats, weasels, and allies) are more closely related to each other, than to Guloninae (Sato et al. 2004, 2006, 2009*a*; Fulton and Strobeck 2006; Koepfli et al. 2008; Wolsan and Sato 2010). See Sato et al. (2009*a*) for the rationale for using the subfamily name Guloninae, instead of Martinae, as proposed by Fulton and Strobeck (2006) and followed by Koepfli et al. (2008).

The ML analysis was conducted using the program GARLI, version 0.96 (Zwickl 2006), and the nucleotide substitution model that best fit the data was applied. The model was determined using Akaike's information criterion (AIC) from the program Modeltest, version 3.7 (Posada and Crandall 1998; see also Posada and Buckley 2004). The GTR+I+ Γ model was selected as the best model for the combined dataset. The ML tree was inferred from heuristic searches involving three runs of the genetic algorithm with 20,000 generations of a mutation-selection-reproduction cycle. The starting tree was generated from the ML stepwise-addition-sequence option. All the other parameters were set by default. Clade support was assessed using nonparametric bootstrap analyses (BP; 200 replicates; Felsenstein 1985).

Divergence-time estimations

To estimate the divergence times for the separation of the Taiwanese and Eurasian mustelids, we adopted the Bayesian relaxed molecular-clock dating approach as implemented in the program BEAST, version 1.6.1 (Drummond and Rambaut 2007). Before the BEAST analyses, we tested the global molecular clock by comparing the log likelihood of the unconstrained ML topology with the molecular-clock enforced ML topology in the program PAUP, version 4.0b10 (Swofford 2002). The likelihood-ratio test rejected the strict molecular clock ($-\ln L_{\text{unconstrained}} = 19645.29052$ and $-\ln L_{\text{molecular clock}} = 19690.24655$; $\chi^2 = 90$, $df = 42$, $P < 0.001$). Therefore, we used the approach that considers rate variation among different branches, in which the rates for each branch were drawn independently from a lognormal distribution and uncorrelated (Drummond et al. 2006). First, we created the input file for the BEAST using the program BEAUTi, version 1.6.1 (provided in the BEAST package). In BEAUTi, we set the substitution model for the sequence evolution, priors, and conditions of Markov Chain Monte Carlo (MCMC) for estimating posterior distributions of the time to the most recent common ancestor (MRCA) for the divergence between mustelids in Taiwan and those on the continent. We set the GTR+I+ Γ for the substitution model as selected above, and the parameters in the model were unlinked among the three partitions (MT-CYB, MT-ND2, and MT-DLOOP). We selected the Yule speciation process as a prior on phylogeny. Two independent MCMC analyses were run for 10 million generations with trees sampled every 1,000 generations. Each log file was checked to confirm the convergence to the stationary posterior distribution and the sufficient effective sample size (ESS) of each parameter in the program Tracer, version 1.5 (Rambaut and Drummond 2009). We obtained sufficient ESS values exceeding 200 for most of the parameters, and over 2,000 for the divergence-time parameters of each node representing the MRCA of relevant Taiwanese and continental

individuals. The log files of both runs were then combined in LogCombiner version 1.6.1 (provided in the BEAST package), with the first 25% of the sampled parameters discarded as burn-in. We provided parameter estimates based on the combined samples from the two independent runs.

Multiple calibration points were set using the first appearance of fossils of some mustelids. The fossil information was adopted in the form of the lognormal prior distribution in line with the recommendation of Ho (2007) and Ho and Phillips (2009). The values of the time constraints based on fossil evidence were set to the zero-offset of the lognormal distribution as the minimum bound. The mean of the lognormal distribution was unavoidably set to somewhat subjective values (Ho 2007). The standard deviation was assigned so that upper limit of the 95% credible interval (CI) of the lognormal distribution took a value of times for divergences one step older in the phylogenetic topologies (the time for the first emergence of the stem lineage of the clade of interest) based on the estimations from Koepfli et al. (2008) and Sato et al. (2009a). When the age of the fossil record directly ancestral to the extant species could be used, the younger age for this ancestor (possible extinction age) was used for the upper limit of the 95% CI of the lognormal distribution. We used three time constraints for calibrations as minimum bounds. First, 3.4 Ma was adopted for the MRCA of *Mustela* and *Neovison* based on the earliest fossil findings of previously circumscribed *Mustela* (including *Neovison*) in the lower Pliocene strata (European Neogene Land Mammal Zone MN15), following Morlo and Kunderát (2001; also see Sato et al. 2003). The mean value was set to 2.0. The standard deviation was set so that the upper limit of the 95% CI took the value of 9.6, as the divergence time for *Mustela-Neovison* and the closest clade (consisting of the Lutrinae species) was estimated to be 8.7–9.0 Ma by Koepfli et al. (2008) and 9.6 Ma by Sato et al. (2009a). Second, 3.3 Ma was applied for the MRCA of the subgenus *Martes*, *M. americana* (Turton, 1806), *M. foina* (Erxleben, 1777), *M. martes* (L., 1758), *M. melampus*

(Wagner, 1840), and *M. zibellina* (L., 1758), based on the earliest fossil remains of the subgenus *Martes* represented by *Martes wenzensis* Stach, 1959 (see Sato et al. 2003 for more detail). The mean value was set to 1.0. The standard deviation was set so that the upper limit of the 95% CI took the value of 5.4, as the stem lineage of the subgenus *Martes* was estimated to have occurred at 4.7–5.1 Ma (Koepfli et al. 2008) and 5.4 Ma (Sato et al. 2009a). Third, 0.13 Ma was used for the MRCA of the two individuals of *Mustela erminea* based on the first fossil record of this species in the Illinoian age in North America (King 1983). The mean value was set to 0.5. The standard deviation was set so that the upper limit of the 95% CI took the value of 0.7, because *Mustela palerminea* L., 1758, a direct ancestor of *M. erminea*, was common until 0.7 Ma (King, 1983). Koepfli et al. (2008) and Sato et al. (2009a) dated the divergence of the *Gulo-Martes* and *Melogale-Lutrinae-Mustela-Neovison* lineages to be 11.6–11.9 Ma and 12.0 Ma, respectively. Therefore, we applied 12.0 Ma to the root height prior as a mean of the normal distribution with a standard deviation of 1.0.

Results

Phylogenetic relationships and genetic distances

The phylogenetic tree inferred from the combined sequences of MT-CYB, MT-ND2, and MT-DLOOP loci using the ML criterion supported the monophyly of the subfamilies Guloninae, Mustelinae, and Lutrinae, albeit with low support for the subfamily Lutrinae (Fig. 1). The interrelationships among these subfamilies were very weakly supported (BP = 44%). By contrast, basal intrasubfamilial relationships within the clade of Mustelinae comprising *Mustela* and *Neovison* were resolved with high support values. *Mustela kathiah* Hodgson, 1835 represented the monotypic lineage sister to the clade that included other musteline species except *M. nudipes* Desmarest, 1822, *M. strigidorsa* Gray, 1853, and *Neovison vison* (Schreber, 1777) (BP = 100%). *Mustela nudipes* and *M. strigidorsa* formed a well-supported

clade (BP = 100%), which was the second-oldest lineage within the clade of Mustelinae (BP = 100%). *Neovison vison* was the first offshoot of the musteline clade (BP = 97%).

Here, we described the relationships of the four mustelids that occur on Taiwan in more detail. The monophyly of *Martes flavigula*, *Melogale moschata*, *Mustela nivalis*, and *Mustela sibirica* was highly supported (BP = 100%). *Martes flavigula* was supported as a sister lineage to the clade that included species in the subgenus *Martes* (*M. americana*, *M. foina*, *M. martes*, *M. melampus*, and *M. zibellina*), albeit with low reliability (BP = 42%). Each *M. flavigula* individual from Kunming, Primorye, and Taiwan had different haplotypes. Additionally, an individual from Taiwan was closely related to one from Primorye, Russia, albeit with low reliability (BP = 53%). The average genetic distance (ML genetic distances inferred with parameters for the GTR+I+ Γ model obtained in the Modeltest analysis) between the individuals from Taiwan and Primorye was 0.003. *Melogale moschata* was very weakly identified as the most basal lineage within the ingroup mustelids (BP = 44%). Each individual *M. moschata* from Taiwan and Vietnam had a different haplotype, with a genetic distance of 0.043. *Mustela nivalis* was the closest relative of *Mustela altaica* Pallas, 1811 (BP = 100%). *Mustela nivalis* individuals from Germany, Korea, and Taiwan formed distinct clades (BP > 95%). The clade composed of two Taiwanese individuals was closely related to that consisting of individuals from Germany, Korea, Russia (Primorye), and Japan (Hokkaido) with relatively high support (BP = 84%). The average genetic distance between individuals from Taiwan and the closest clade individuals was 0.009. *Mustela sibirica* was strongly supported as sister to the clade of *Mustela furo* L., 1758, *Mustela eversmanii* Lesson, 1827, *Mustela lutreola* (L., 1761), and *Mustela putorius* L., 1758 (BP = 100%). The Taiwanese lineage appeared in the basal position of the lineage of this species, albeit with low reliability (BP = 52%). The close relationship among individuals from Korea, Russia (Primorye), and

the Tsushima Islands was strongly supported (BP = 100%). The average genetic distance between the Taiwanese and other individuals was 0.016.

Among all of the individuals examined, the minimum and maximum genetic distances observed within species were 0 (two Taiwanese *Mustela nivalis* individuals and two Kunming *Martes flavigula* individuals) and 0.043 (Taiwanese and Vietnamese *Melogale moschata* individuals), respectively. Genetic distances less than the largest value of the intraspecific variation (0.043) were observed within the interspecific variation: 0.013 (*Mustela eversmanii* vs. *Mustela putorius*), 0.014 (*Mustela eversmanii* vs. *Mustela lutreola*), 0.014 (*Mustela lutreola* vs. *Mustela putorius*), 0.035 (*Martes martes* vs. *Martes zibellina*), and 0.041 (*Martes martes* vs. *Martes melampus*). The 0.048 value for *Martes melampus* vs. *Martes zibellina* was close to the maximum intraspecific difference observed within *Melogale moschata*.

Divergence-time estimation

The divergence of the mustelid lineages from Taiwan and from the continent was dated using the Bayesian relaxed-molecular clock approach with the program BEAST. The 95% highest posterior density (HPD) of the covariance of parent and child branch rates varied from -0.20 to 0.21, with a mean near zero, showing no strong evidence of autocorrelation of the rates among adjacent branches (see the BEAST manual). This validated our use of the uncorrelated lognormal distribution as a rate-variation model. The times to the most recent common ancestor for the Taiwanese individual and the closest clade or individual with upper and lower HPD intervals are indicated in Table 2. The mean divergence times obtained in the analyses (0.11, 1.46, 0.36, and 0.63 Ma for *Martes flavigula*, *Melogale moschata*, *Mustela nivalis*, and *Mustela sibirica*, respectively) gave four different ages for the divergence between the Taiwanese and closest lineages.

Discussion

Interspecific relationships within the mustelid phylogeny

Our combined phylogenetic analyses based on the ML criterion produced a topology largely congruent with previous hypotheses. As a whole, shallower interspecific relationships were well supported, with the opposite true for deeper nodes, a phenomenon consistent with the previous notion that the mitochondrial genetic loci suffer from the multiple-substitution problem in the phylogenetic reconstruction of the older past (*e.g.*, Sato et al. 2003; Sato and Suzuki 2004). Nonetheless, the relationships we obtained among the four subfamilies studied (Guloninae, (Helictidinae, (Lutrinae, Mustelinae))) are consistent with the findings of recent molecular phylogenetic studies (Koepfli and Wayne 2003; Sato et al. 2004, 2006, 2009a; Fulton and Strobeck 2006; Koepfli et al. 2008). Interspecific relationships within the subfamilies Mustelinae and Guloninae mostly agreed with previous reports (*e.g.*, Sato et al. 2003, 2004, 2006, 2009a; Koepfli et al. 2008; Wolsan and Sato 2010), although our data showed unprecedented strong support for the phylogenetic position of *Mustela kathiah*.

Intraspecific variations and two taxonomic implications

The intraspecific variation has implications for taxonomic issues concerning the Taiwanese mustelids. *Mustela nivalis* was first found in Taiwan in 1998 (Lin 2000; Lin et al. 2010). Its presence in Taiwan was unexpected, as this is a Holarctic species, and it is mainly distributed in the northern parts of Eurasia and North America (Wozencraft 2005). The considerable divergence of morphological characteristics of this newly discovered weasel from other described subspecies led Lin and Harada (1998) to initially regard it as a new species (but see Lin et al. 2010). However, our study demonstrated that Taiwanese individuals are not appreciably genetically differentiated from those of continental regions. This result is congruent with our previous study examining a 402-bp MT-CYB sequence of the Taiwanese

individual (Hosoda et al. 2000). Saarma and Tumanov (2006) used part of our MT-CYB sequence (312 bp) from the DNA database and compared it to the other *Mustela nivalis* subspecies. Their results agreed with ours and indicated that the genetic divergence of the Taiwanese individual was within intraspecific variation. Therefore, the newly discovered species was classified as *Mustela nivalis*, as noted in the most recent classification book (Wozencraft 2005). Previous observations of the morphological differentiation of *Mustela nivalis* in Taiwan are in accordance with the notion of the island effect as accelerating morphological evolution (Millien 2006).

Another taxonomic implication of this study is the evidence of significant differentiation between two *Melogale moschata* individuals from Taiwan and Vietnam, which showed genetic divergence comparable to that of interspecific genetic variation. To date, six subspecies of this species have been recognised (Storz and Wozencraft 1999; Wozencraft 2005). The Taiwanese and Vietnamese individuals that we examined belong to *M. moschata subaurantiaca* and *M. moschata taxilla*, respectively. The other subspecies are distributed from southeast China to northeast India, between the distributions of the two subspecies examined in this study. Therefore, the large genetic divergence we observed may not be due exclusively to the geographical separation of the Taiwan islands from the continent. Namely, the large genetic divergence might be established within the continent. Although it was impossible to specify the mechanism producing the large genetic diversity in this study by examining only two individuals, two possible explanations exist for the genetic divergence of the ferret badger in Taiwan. First, it might have anciently dispersed into Taiwan and have been isolated for a long time by the Taiwan–continent separation. Second, the genetic diversity might have arisen on the continent, and a diverged lineage might have recently dispersed into Taiwan. If the latter is correct, there should be a closely related lineage on the continent. Clearly, more samples, including those of other continental subspecies, are required

to understand the mechanism leading to such considerable divergence and to revise the taxonomy within this species. Regarding the other two mustelids on Taiwan (*Martes flavigula* and *Mustela sibirica*), no novel implications were gleaned from a taxonomical perspective.

Phylogenetic history of mustelid fauna in Taiwan

The divergence times of the Taiwanese and continental mustelids estimated from combined mitochondrial loci were 0.11, 1.46, 0.36, and 0.63 million years ago for *Martes flavigula*, *Melogale moschata*, *Mustela nivalis*, and *Mustela sibirica*, respectively (Table 2). These ages are all within the Pleistocene, suggesting that these divergences were influenced by the glacial–interglacial cycles (*e.g.*, Luthi et al. 2008). The continental fauna likely moved dynamically between northern and southern regions in response to these glacial cycles (Riddle 1996; Hewitt 2000; Sommer and Zachos 2009). Additionally, the sea between Taiwan and the continent was shallow enough during glacial periods to enable various organisms to cross the Taiwan Strait (Vorisi 2000). Consequently, Taiwan has probably received historically and geographically diverse lineages. We found no consistency in the migration scenarios of the four Taiwanese mustelids. *Martes flavigula* and *Melogale moschata*, from the southern Orient, are the youngest and oldest residents of Taiwan, respectively. Northern Holarctic *Mustela nivalis* and *Mustela sibirica*, which expanded both to the north and south, appear to have established themselves in Taiwan in periods between the migrations of the two Oriental species. The ability of four species of the family Mustelidae to coexist in Taiwan may be attributed to ecological niche differences, with arboreal martens (*Martes flavigula*), fossorial badgers (*Melogale moschata*), and open habitat weasels (*Mustela* spp.). These differences presumably permit coexistence with little competition. This idea, although implied from a smaller island system, is consistent with the suggestion from a study on a larger continental scale (Koepfli et al. 2008). They stressed that a phylogenetically and ecologically

heterogeneous set of species could coexist within the same mustelid community. The existence of two congeneric *Mustela* species, *M. nivalis* and *M. sibirica*, which might be expected to compete owing to similar ecological preferences, may be possible because of differences in foraging and breeding strategies, micro-habitats, and social dominance (King and Moors 1979; Powell and Zielinski 1983; Erlinge and Sandell 1988). Grant (1970) suggested a trend in island colonisation from ecological generalists to specialists, ultimately allowing closely related species to coexist on an island. In terms of diet, *M. nivalis*, which feeds almost entirely on small rodents, has characteristics of a specialist, whereas *M. sibirica* is a generalist that feeds more diversely (King and Moors 1979; Nowak 1991; Masuda 2009; Sasaki 2009). Therefore, the observation of Grant (1970) concurs with our estimated divergence times of *M. sibirica* (0.63 Ma) and *M. nivalis* (0.36 Ma) in Table 2. As an alternative explanation of the order of migration, *M. sibirica* may simply have had more opportunity, as its distribution in the continent is in closer proximity to the Taiwan islands than that of *M. nivalis*. Clearly, additional phylogenetic and ecological evidence, with larger samples, is necessary to elucidate the coexistence history of the Taiwanese mustelid fauna.

To conclude, our study found that the Taiwanese mustelids colonised the island at four different times during the Pleistocene, likely across land bridges that emerged owing to the lowering of sea levels during glacial periods. The results support the hypothesis of multiple episodes of immigration from the mainland, in agreement with other studies of the evolution of the Taiwanese fauna (Yu et al. 1996; Tu et al. 2000; Creer et al. 2001, 2004; Lin et al. 2002; Jang-Liaw et al. 2008; Zhong et al. 2008). The Taiwanese fauna (and flora) provide a good model for understanding the assembly and evolution of island communities that contain both old and new lineages.

ACKNOWLEDGEMENTS

We are so grateful to two anonymous reviewers for constructive comments for our previous manuscript. We thank Hisashi Abe, Kevin Campbell, Kay Fuhrman, Alexei Kryukov, Mitsuru Mukohyama, Takashi Saitoh, Sang-hoon Han, Hiroshi Sasaki, Masaya Tatara, and Kimiyuki Tsuchiya for providing tissue samples for molecular analyses. We also thank Masahiro A. Iwasa and Shumpei P. Yasuda for valuable and constructive discussions. This work was supported in part by the Ministry of Education, Culture, Sports, Science, and Technology, Japan (grants-in-aid for scientific research to Hitoshi Suzuki) and the Heiwa Nakajima Foundation (grant to Jun J. Sato).

REFERENCES

- Creer, S., Malhotra, A., Thorpe, R.S., and Chou, W.-H. 2001. Multiple causation of phylogeographical pattern as revealed by nested clade analysis of the bamboo viper (*Trimeresurus stejnegeri*) within Taiwan. *Mol. Ecol.* **10**: 1967–1981.
- Creer, S., Thorpe, R.S., Malhotra, A., Chou, W.-H., and Stenson, A.G. 2004. The utility of AFLPs for supporting mitochondrial DNA phylogeographical analyses in the Taiwanese bamboo viper, *Trimeresurus stejnegeri*. *J. Evol. Biol.* **17**: 100-107.
- Davis, C.S., Delisle, I., Stirling, I., Siniff, D.B., and Strobeck, C. 2004. A phylogeny of the extant Phocidae inferred from complete mitochondrial DNA coding regions. *Mol. Phylogenet. Evol.* **33**: 363-377.
- Delisle, I., and Strobeck, C. 2005. A phylogeny of the Caniformia (order Carnivora) based on 12 complete protein-coding mitochondrial genes. *Mol. Phylogenet. Evol.* **37**: 192-201.
- Drummond, A.J., Ho, S.Y.W., Phillips, M.J., Rambaut, A. 2006. Relaxed phylogenetics and dating with confidence. *Plos Biol.* **4**: e88.
- Drummond, A.J., and Rambaut A. 2007. BEAST: Bayesian evolutionary analysis by sampling trees. *BMC Evol. Biol.* **7**: 214.

- Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res.* **32**: 1792-1797.
- Erlinge, S., and Sandell, M. 1988. Coexistence of stoat, *Mustela erminea*, and weasel, *M. nivalis*: social dominance, scent communication, and reciprocal distribution. *Oikos*, **53**: 242-246.
- Felsenstein, J. 1981. Evolutionary trees from DNA sequences: A maximum likelihood approach. *J. Mol. Evol.* **17**: 368-376.
- Felsenstein, J. 1985. Confidence limits on phylogenies: An approach using the bootstrap. *Evolution*, **39**: 783-791.
- Flynn, J.J., Finarelli, J.A., Zehr, S., Hsu, J., and Nedbal, M.A. 2005. Molecular phylogeny of the Carnivora (Mammalia): assessing the impact of increased sampling on resolving enigmatic relationships. *Syst. Biol.* **54**: 317–337.
- Fulton, T.L., and Strobeck, C. 2006. Molecular phylogeny of the Arctoidea (Carnivora): effect of missing data on supertree and supermatrix analyses of multiple gene data sets. *Mol. Phylogenet. Evol.* **41**: 165-181.
- Gascoyne, M., Benjamin, G.J., Schwarcz, H.P., and Ford, D.C. 1979. Sea-level lowering during the Illinoian glaciation: evidence from a Bahama “Blue Hole”. *Science (Washington, D.C.)*, **205**: 806-808.
- Grant, P.R. 1970. Colonization of islands by ecologically dissimilar species of mammals. *Can. J. Zool.* **48**: 545-553.
- Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. *Nature (London)*, **405**: 907-913.
- Ho, C.C. 1986. A synthesis of the geologic evolution of Taiwan. *Tectonophysics* **125**:1-16.
- Ho, S.Y.W. 2007. Calibrating molecular estimates of substitution rates and divergence times in birds. *J. Avian Biol.* **38**: 409-414.

- Ho, S.Y.W., and Phillips, M.J. 2009. Accounting for calibration uncertainty in phylogenetic estimation of evolutionary divergence time. *Syst. Biol.* **58**: 367-380.
- Hosoda, T., Suzuki, H., Harada, M., Tsuchiya, K., Han, S.-H., Zhang, Y.-P., Kryukov, A.P., and Lin, L.-K. 2000. Evolutionary trends of the mitochondrial lineage differentiation in species of genera *Martes* and *Mustela*. *Genes. Genet. Syst.* **75**: 233-243.
- Huang, C.-Y., Wu, W.-Y., Chang, C.-P., Tsao, S., Yuan, P.B., Lin, C.-W., and Xia, K.-Y. 1997. Tectonic evolution of accretionary prism in the arc-continent collision terrane of Taiwan. *Tectonophysics*, **281**: 31-51.
- Jang-Liaw, N.-H., Lee, T.-H., and Chou, W.-H. 2008. Phylogeography of *Sylvirana latouchii* (Anura, Ranidae) in Taiwan. *Zool. Sci. (Tokyo)*, **25**: 68-79.
- Kano, T. 1940. Zoogeographic studies of the Tsugitaka mountains of Formosa (in Japanese). A report for Shibusawa Institute of ethonology (also see Kawamura, T. *Acta phytotaxonomica et geobotanica*, **10**: 150-151, 1941-06-30 in Japanese)
- Kier, G., Kreft, H., Lee, T.-M., Jetz, W., Ibisch, P.L., Nowicki, C., Mutke, J., and Barthlott, W. 2009. A global assessment of endemism and species richness across island and mainland regions. *Proc. Natl. Acad. Sci. U.S.A.* **106**: 9322-9327.
- Kimura, M. 2000. Paleogeography of the Ryukyu Islands. *Tropics*, **10**: 5-24.
- King, C.M., and Moors, P.J. 1979. On Co-existence, foraging strategy and the biogeography of weasels and stoats (*Mustela nivalis* and *M. erminea*) in Britain. *Oecologia*, **39**: 129-150.
- King, C.M. 1983. *Mustela erminea*. *Mammalian Species Account*, **195**: 1-8.
- Kizaki, K., and Oshiro, I. 1980. The origins of the Ryukyu Islands. *In* Natural History of Ryukyu. *Edited by* K. Kizaki. Tsukiji-Shokan, Tokyo. pp. 8-37 (in Japanese).
- Koepfli, K.-P., and Wayne, R.K. 1998. Phylogenetic relationships of otters (Carnivora: Mustelidae) based on mitochondrial cytochrome *b* sequences. *J. Zool. (Lond.)*

246:401-416.

- Koepfli, K.-P., and Wayne, R.K. 2003. Type I STS markers are more informative than cytochrome *b* in phylogenetic reconstruction of the Mustelidae (Mammalia: Carnivora). *Syst. Biol.* **52**: 571–593.
- Koepfli, K.-P., Deere, K.A., Slater, G.J., Begg, C., Begg, K., Grassman, L., Lucherini, M., Veron, G., and Wayne, R.K. 2008. Multigene phylogeny of the Mustelidae: resolving relationships, tempo and biogeographic history of a mammalian adaptive radiation. *BMC Biol.* **6**: 1-22.
- Kurose, N., Abramov, A.V., and Masuda, R. 2000. Intrageneric diversity of the cytochrome *b* gene and phylogeny of Eurasian species of the genus *Mustela*. *Zool. Sci. (Tokyo)*, **17**: 673-679.
- Lin, L.-K. 2000. The status and conservation of Taiwan's mustelids. *In* “Mustelids in a modern world: management and conservation aspects of small carnivore: human interactions.” *Edited by* H.I. Griffiths. Backhuys Publishers, Leiden. pp. 331–335.
- Lin, L.-K., and Harada, M. 1998 A new species of *Mustela* from Taiwan. *Proceedings of the Euro-American Mammal Congress, Santiago de Compostela, Spain*: **765**.
- Lin, L.-K., Motokawa, M., and Harada, M. 2010. A new subspecies of the least weasel *Mustela nivalis* (Mammalia: Carnivora) from Taiwan. *Mammal Study*, **35**: 191-200.
- Lin, S.-M., Chen, C.A., and Lue, K.-Y. 2002. Molecular phylogeny and biogeography of the grass lizards genus *Takidromus* (Reptilia: Lacertidae) of East Asia. *Mol. Phylogenet. Evol.* **22**: 276-288.
- Luthi, D., Floch, M.L., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T.F. 2008. High-resolution carbon dioxide concentration record 650,000-800,000 years before present. *Nature (London)*, **453**: 379-382.

- Masuda, R. 2009. *Mustela erminea* Linnaeus, 1766. In *The Wild Mammals of Japan Edited by* S.D. Ohdachi., Y. Ishibashi., M.A. Iwasa., T. Saitoh. Shokadoh, Kyoto. pp 242-243.
- Millien, V. 2006. Morphological evolution is accelerated among island mammals. *Plos Biol.* **4**: 1863-1868.
- Morlo, M., and Kunderát, M. 2001. The first carnivoran fauna from the Ruscium (Early Pliocene, MN 15) of Germany. *Paläontol. Z.* **75**: 163–187.
- Nowak, R.M. 1991. *Walker's Mammals of the World Vol2.* 5th edn, The Johns Hopkins University Press, Baltimore.
- Oshida, T., Lee, J.-K., Lin, L.-K., and Chen, Y.-J. 2006. Phylogeography of Pallas's squirrel in Taiwan: geographical isolation in an arboreal small mammal. *J. Mammal.* **87**: 247-254.
- Ota, H. 1998. Geographic patterns of endemism and speciation in amphibians and reptiles of the Ryukyu archipelago, Japan, with special reference to their paleogeographical implications. *Res. Popul. Ecol.* **40**: 189-204.
- Otsuka, H., and Takahashi, A. 2000. Pleistocene vertebrate faunas in the Ryukyu Islands: their migration and extinction. *Tropics*, **10**: 25-40.
- Page, B.M., and Suppe, J. 1981. The Pliocene lichi melange of Taiwan: Its plate tectonic and olistostromal origin. *Am. J. Sci.* **281**: 193-227.
- Posada, D., and Crandall, K.A. 1998. MODELTEST: Testing the model of DNA substitution. *Bioinformatics*, **14**: 817-818.
- Posada, D., and Buckley, T.R. 2004. Model selection and model averaging in phylogenetics: Advantages of Akaike Information Criterion and Bayesian approaches over likelihood ratio tests. *Syst. Biol.* **53**: 793-808.
- Powell, R.A., and Zielinski, W.J. 1983. Competition and coexistence in mustelid communities. *Acta Zool. Fenn.* **174**: 223-227.
- Rambaut, A., and Drummond, A.J. 2009. Tracer v1.5, Available from

<http://beast.bio.ed.ac.uk/Tracer>

- Riddle, B.R. 1996. The molecular phylogeographic bridge between deep and shallow history in continental biotas. *Trends Ecol. Evol.* **11**: 207-211.
- Saarma, U., and Tumanov, I.L. 2006. Phylogenetic Evaluation of Three Subspecies from the *Mustela nivalis* Group. *Zool. Stud.* **45**: 435-442.
- Sambrook, J., and Russell, D.W. 2001. *Molecular Cloning: A Laboratory Manual*, 3rd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York.
- Sasaki, H. 2009. *Mustela sibirica* Pallas, 1773. In *The Wild Mammals of Japan Edited by S.D. Ohdachi., Y. Ishibashi., M.A. Iwasa., T. Saitoh. Shokadoh, Kyoto.* pp 242-243.
- Sato, J.J., and Suzuki, H. 2004. Phylogenetic relationships and divergence times of the genus *Tokudaia* within Murinae (Muridae; Rodentia) inferred from the nucleotide sequences encoding the mitochondrial cytochrome *b* and nuclear recombination-activating gene 1 and interphotoreceptor retinoid-binding protein. *Can. J. Zool.* **82**: 1343-1351.
- Sato, J.J., Hosoda, T., Wolsan, M., Tsuchiya, K., Yamamoto, Y., and Suzuki, H. 2003. Phylogenetic relationships and divergence times among mustelids (Mammalia: Carnivora) based on nucleotide sequences of the nuclear interphotoreceptor retinoid binding protein and mitochondrial cytochrome *b* genes. *Zool. Sci. (Tokyo)*, **20**: 243-264.
- Sato, J.J., Hosoda, T., Wolsan, M., and Suzuki, H. 2004. Molecular phylogeny of arctoids (Mammalia: Carnivora) with emphasis on phylogenetic and taxonomic positions of the ferret-badgers and skunks. *Zool. Sci. (Tokyo)*, **21**: 111-118.
- Sato, J.J., Wolsan, M., Suzuki, H., Hosoda, T., Yamaguchi, Y., Hiyama, K., Kobayashi, M., and Minami, S. 2006. Evidence from nuclear DNA sequences sheds light on the phylogenetic relationships of Pinnipedia: single origin with affinity to Musteloidea. *Zool. Sci. (Tokyo)*, **23**: 125-146.
- Sato, J.J., Wolsan, M., Minami, S., Hosoda, T., Sinaga, M.H., Hiyama, K., Yamaguchi, Y., and

- Suzuki, H. 2009a. Deciphering and dating the red panda's ancestry and early adaptive radiation of Musteloidea. *Mol. Phylogenet. Evol.* **53**: 907-922.
- Sato, J.J., Yasuda, S.P., and Hosoda, T. 2009b. Genetic diversity of the Japanese marten (*Martes melampus*) and its implications for the conservation unit. *Zool. Sci. (Tokyo)*, **26**: 457-466.
- Sommer, R.S., and Zachos, F.E. 2009. Fossil evidence and phylogeography of temperate species: 'glacial refugia' and post-glacial recolonization. *J. Biogeogr.* **36**: 2013–2020.
- Storz, J.F., and Wozencraft, W.C. 1999. *Melogale moschata*. *Mamm Species*, No. **631**: 1-4.
- Suzuki, H. 2009. A molecular phylogenetic view of mammals in the “three-story museum” of Hokkaido, Honshu, and Ryukyu Islands, Japan. *In The Wild Mammals of Japan Edited by S.D. Ohdachi., Y. Ishibashi., M.A. Iwasa., T. Saitoh. Shokadoh, Kyoto.* pp 261-263.
- Swofford, D.L. 2002. PAUP*. Phylogenetic Analyses Using Parsimony (* and Other Methods). Version 4. Sinauer Associates, Sunderland, MA.
- Takahashi, K., Chang, C.-H., and Cheng, Y.-N. 2001. Proboscidean fossils from the Japanese Archipelago and Taiwan Islands and their relationship with the Chinese mainland. The world of elephants: Proceedings of the 1st international congress, Rome, pp.148-151.
- Teng, L.S. 1990. Geotectonic evolution of late Cenozoic arc-continent collision in Taiwan. *Tectonophysics*, **183**: 57-76.
- Toda, M., Nishida, M., Matsui, M., Wu, G.-F., and Ota, H. 1997. Allozyme variation among east asian populations of the Indian Rice Frog, *Rana limnocharis* (Amphibia: Anura). *Biochem. Syst. Ecol.* **25**: 143–159.
- Toda, M., Nishida, M., Matsui, M., Lue, K.-Y., and Ota, H. 1998. Genetic variation in the Indian rice frog, *Rana limnocharis* (Amphibia: Anura), in Taiwan, as revealed by allozyme data. *Herpetologica*, **54**: 73–82.
- Tu, M.-C., Wang, H.-Y., Tsai, M.-P., Toda, M., Lee, W.-J., Zhang, F.-J., and Ota, H. 2000.

- Phylogeny, Taxonomy, and Biogeography of the Oriental Pitvipers of the Genus *Trimeresurus* (Reptilia: Viperidae: Crotalinae): A Molecular Perspective. *Zool. Sci.* (Tokyo), **17**: 1147–1157.
- Tzeng, C.S. 1986. Distribution of the freshwater fishes of Taiwan. *J. Taiwan Mus.* **39**: 127–146.
- Voris, H.K. 2000. Maps of Pleistocene sea levels in Southeast Asia: shorelines, river systems and time durations. *J. Biogeogr.* **27**: 1153-1167.
- Wang, H.-Y., Tsai, M.-P., Yu, M.-J., and Lee, S.-C. 1999. Influence of glaciation on divergence patterns of the endemic minnow, *Zacco pachycephalus*, in Taiwan. *Mol. Ecol.* **8**: 1879-1888.
- Wang, P.X. 2004. Cenozoic deformation and the history of sea-land interactions in Asia. *In* “Continent-Ocean interactions in the East Asian Marginal Seas” *Edited by* P.Clift et al. Geophysical Monograph **149**, American Geophysical Union, 1-22.
- Watanabe, K., Jang-Liaw, N.-H., Zhang, C.-G., Jeon, S.-R., and Nishida, M. 2007. Comparative phylogeography of bagrid catfishes in Taiwan. *Ichthyol. Res.* **54**: 253-261.
- Whittaker, R.J., and Fernandez-Palacios, J.M. 2007. *Island Biogeography* second edition. Oxford University Press, New York.
- Wiens, J.J. 2006. Missing data and the design of phylogenetic analyses. *J. Biomed. Inform.* **39**: 34-42.
- Wolsan, M., and Sato, J.J. 2010. Effects of data incompleteness on the relative performance of parsimony and Bayesian approaches in a supermatrix phylogenetic reconstruction of Mustelidae and Procyonidae (Carnivora). *Cladistics*, **26**: 168-194.
- Wozencraft, W.C. 2005. Order Carnivora. *In* *Mammal Species of the World: A Taxonomic and Geographic Reference*, Third Edition, Volume 1. *Edited by* D.E. Wilson., D.M. Reeder. Johns Hopkins Univ. Press, Baltimore, Maryland. pp. 532–628.

- Yuan, S.-L., Lin, L.-K., and Oshida, T. 2006. Phylogeography of the mole-shrew (*Anourosorex yamashinai*) in Taiwan: implications of interglacial refugia in a high-elevation small mammal. *Mol. Ecol.* **15**: 2119-2130.
- Yu, H.-T. 1995. Patterns of diversification and genetic population structure of small mammals in Taiwan. *Biol. J. Linn. Soc.* **55**: 69-89.
- Yu, H.-T., Fang, Y.-P., Chou, C.-W., Huang, S.-W., and Yew, F.-H. 1996. Chromosomal evolution in three species of murid rodents of Taiwan. *Zool. Stud.* **35**: 195-199.
- Yu, L., and Zhang, Y.P. 2006. Phylogeny of the caniform carnivora: evidence from multiple genes. *Genetica*, **127**: 65-79.
- Zhong, J., Liu, Z.-Q., and Wang, Y.-Q. 2008. Phylogeography of the Rice Frog, *Fejervarya multistriata* (Anura: Ranidae), from China Based on mtDNA D-loop Sequences. *Zool. Sci. (Tokyo)*, **25**: 811–820.
- Zwickl, D.J. 2006. Genetic Algorithm approaches for the phylogenetic analysis of large biological sequence datasets under the maximum likelihood criterion. Ph.D. dissertation, The University of Texas at Austin.

The English in this document has been checked by at least two professional editors, both native speakers of English. For a certificate, please see:

<http://www.textcheck.com/certificate/v2GWU7>

FIGURE CAPTIONS

Figure 1. Phylogenetic relationships of the mustelids inferred from maximum likelihood analyses of the combined MT-CYB, MT-ND2, and MT-DLOOP sequence (2,724 bp). The negative log likelihood scores for optimal ML topology ($-\ln L$) was -19649.848271. Numbers attached near nodes are bootstrap proportions calculated from the 200 pseudo-replicate topologies. The vertical thin lines indicated mustelid species including multiple individuals from different localities. Species names in bold faces indicate Taiwanese mustelid species. *Ma*, *Mu*, and *Me* indicate the genus names *Martes*, *Mustela*, and *Melogale*, respectively. The vertical thick lines indicate the subfamily names. For the rationale of those subfamily's names, see Sato et al. (2009a).

Fig. 1

Table 1. Species and DNA sequences used in this study

Taxon	Voucher number ^a	Locality ^a	DDBJ/EMBL/GenBank accession numbers		
			MT-CYB	MT-ND2	MT-DLOOP
<i>Enhydra lutris</i> , sea otter	TH257	Alaska, USA	AF057120 ^b	AY750618 ^c	AB601556 ^g
<i>Gulo gulo</i> , wolverine	HS1603	Sakhalin, Russia	AB051245 ^d	AY377284 ^e	AB601557 ^g
<i>Lontra canadensis</i> , North American river otter	TH332	Unknown, Canada	AF057121 ^b	AY598557 ^f	AB601558 ^g
<i>Martes americana</i> , American marten	TH330	Manitoba, Canada	AB051234 ^d	AY598546 ^f	AB601559 ^g
<i>Martes flavigula</i> , yellow-throated marten	HS844	Kunming, China	AB564113 ^g	AB601552 ^g	AB601560 ^g
	HS858	Kunming, China	AB564114 ^g	AB564137 ^g	AB601561 ^g
	NMNS707	Nantou county, Taiwan	AB564115 ^g	AB564138 ^g	AB601562 ^g
	TH444	Primorye, Russia	AB564116 ^g	AB564139 ^g	AB601563 ^g
<i>Martes foina</i> , beech marten	HS1753	Gera, Thuringia, Germany	AB051236 ^d	AB564140 ^g	AB601564 ^g
<i>Martes martes</i> , European pine marten	AK702	Moscow, Russia	AB051237 ^d	AB564141 ^g	AB601565 ^g
<i>Martes melampus</i> , Japanese marten			AB455595 ^h	AB455695 ^h	AB455645 ^h
<i>Martes pennanti</i> , fisher	TH331	Manitoba, Canada	AF057131 ⁱ	AY750624 ^c	AB601566 ^g
<i>Martes zibellina</i> , sable			AB455641 ^h	AB455741 ^h	AB455691 ^h
<i>Melogale moschata</i> , Chinese ferret-badger	AK703	Unknown, Vietnam	AB564117 ^g	AB564142 ^g	AB601567 ^g
	HS2372	Taichung County, Taiwan	AB564118 ^g	AB564143 ^g	AB601568 ^g
<i>Mustela altaica</i> , mountain weasel	AK805	Altai region, Russia	AB051239 ^d	AB564144 ^g	AB601569 ^g
<i>Mustela erminea</i> , ermine	HS678	Novosibirsk, Russia	AB564119 ^g	AB564145 ^g	AB601570 ^g
	HS1759	Iwate, Japan	AB564120 ^g	AB564146 ^g	AB601571 ^g
	HS2169	Chita region, Russia	AB026102 ⁱ	AB564148 ^g	AB601572 ^g
<i>Mustela eversmanii</i> , steppe polecat			AB026103 ⁱ	AB564149 ^g	AB601573 ^g
<i>Mustela furo</i> , ferret	TH27	Experimental Animal (The Jikei University)			
<i>Mustela itatsi</i> , Japanese weasel	TH50	Ishikari, Hokkaido, Japan	-	AB564150 ^g	AB601574 ^g
	TH89	Aomori, Japan	AB564121 ^g	-	-
<i>Mustela kathiah</i> , yellow-bellied weasel	TH321	Kunming, China	AB285331 ^k	AY882063 ^l	AB601575 ^g
<i>Mustela lutreola</i> , European weasel	AK13	Novosibirsk, Russia	AB026105 ⁱ	AY750628 ^c	AB601576 ^g
<i>Mustela nivalis</i> , least weasel	HS597	Eniwa, Hokkaido, Japan	AB564122 ^g	AB564152 ^g	AB601577 ^g
	HS686	Aomori, Japan	AB564123 ^g	AB564153 ^g	AB601578 ^g
	HS928	Yeongyang, South Korea	AB564124 ^g	AB601553 ^g	AB601579 ^g
	HS1119	Kedrovaya, Primorye, Russia	AB564125 ^g	AB564154 ^g	AB601580 ^g
	HS1745	Chizchik River, Uzbekistan	AB564126 ^g	AB564155 ^g	AB601581 ^g
	HS1758	Thuringia, Germany	AB564127 ^g	AB564156 ^g	AB601582 ^g
	HS1913	Mt. Odaesan, South Korea	AB564128 ^g	AB564157 ^g	AB601583 ^g

	NMNS3302	Mt. Hohuanshan, Nantou County, Taiwan	AB046612 ^g	AB564158 ^g	AB601584 ^g
	NMNS5244	Mt. Hohuanshan, Nantou County, Taiwan	AB564129 ^g	AB601554 ^g	AB601585 ^g
	TH229	Thuringia, Germany	AB564130 ^g	AB564160 ^g	AB601586 ^g
<i>Mustela nudipes</i> , Malayan weasel	BM2002-227	Tasek Merimbun Heritage Park, Tutong, Brunei Darussalam	AB285332 ^k	AB564161 ^g	AB601587 ^g
<i>Mustela putorius</i> , European polecat	AK710	Moscow, Russia	AB026107 ⁱ	AY750630 ^c	AB601588 ^g
<i>Mustela sibirica</i> , Siberian weasel	AK#9	Primorye, Russia	AB564131 ^g	AB564162 ^g	AB601589 ^g
	HS1121	Primorye, Russia	AB564132 ^g	AB564163 ^g	AB601590 ^g
	HS1145	Mt. Hohuanshan, Nantou County, Taiwan	AB564133 ^g	AB564164 ^g	AB601591 ^g
	HS1146	Mt. Hohuanshan, Nantou County, Taiwan	AB051243 ^d	AB601555 ^g	AB601592 ^g
	TH51	Tsushima, Japan	AB564134 ^g	AB564165 ^g	AB601593 ^g
	TH56	Mt. Chiri, South Korea	AB564135 ^g	AB564166 ^g	AB601594 ^g
	TH98	Chikushi, Fukuoka, Japan	AB564136 ^g	AB564167 ^g	AB601595 ^g
<i>Mustela strigidorsa</i> , back-striped weasel	ANWC M32057	Oudomsouk, Nakai, Khammouan, Laos	AB305635 ^k	AB564168 ^g	AB601596 ^g
<i>Neovison vison</i> , American mink	TH49	Nayoro, Hokkaido, Japan (Introduced)	AF057129 ^b	AY377285 ^e	AB601597 ^g

^a Voucher number and Locality refer only to sequences generated in this study.

^b Koepfli and Wayne (1998); ^c Flynn et al. (2005); ^d Hosoda et al. (2000); ^e Davis et al. (2004); ^f Delistle and Strobeck (2005); ^g Sequences generated in this study; ^h Sato et al. (2009b); ⁱ Kurose et al. (2000); ^j Koepfli and Wayne (2003); ^k Sato et al. (unpublished); ^l Yu and Zhang (2006).

Footnote

Sato et al. (unpublished): Sato, J.J., Wolsan, M., Prevosti, F.J., D'Elía, G., Begg, C., Begg, K., Hosoda, T, Campbell, K.L., and Suzuki, H. Evolutionary and biogeographical history of weasel-like carnivorans (Musteloidea)

Table 2. Divergence times of mustelid lineages from Taiwan and the continent estimated with the Bayesian relaxed molecular-clock approach.

	Posterior mean (Ma)	95% HPD [*]
<i>Martes flavigula</i>	0.11	0.04 - 0.19
<i>Melogale moschata</i>	1.46	1.00 - 1.97
<i>Mustela nivalis</i>	0.36	0.26 - 0.48
<i>Mustela sibirica</i>	0.63	0.43 - 0.82

^{*}: HPD, Highest Posterior Densities