Title	Volume-rendering and endocardial views of partially unroofed coronary sinus with 64-slice multidetector CT
Author(s)	Oyama, Noriko; Ooka, Tomonori; Sasaki, Tsukasa; Kubota, Suguru; Onodera, Yuya; Matsui, Yoshiro; Terae, Satoshi; Shirato, Hiroki
Citation	Journal of Cardiovascular Computed Tomography, 3(5), 346-347 https://doi.org/10.1016/j.jcct.2009.05.017
Issue Date	2009-09
Doc URL	http://hdl.handle.net/2115/46805
Type	article (author version)
File Information	JCCT3-5_346-347.pdf

[Instructions for use](#)

Title: Volume rendering and Endocaridal views of Partially Unroofed Coronary Sinus with
64-slice MDCT

Noriko Oyama, MD, PhD,^{a)} Tomonori Ooka, MD, PhD,^{b)} Tsukasa Sasaki, RT,^{a)} Suguru Kubota,
MD, PhD,^{b)}, Yuya Onodera, MD, PhD,^{a)} Yoshiro Matsui, MD, PhD,^{a)} Satoshi Terae MD, PhD,^{a)},
and Hiroki Shirato MD, PhD,^{a)}

^{a)} Department of Radiology,

^{b)} Department of Cardiovascular Medicine,

Hokkaido University Hospital, Sapporo, Japan

There is no conflict of interests for all.

There is no financial disclosure.

Address for correspondence:

Noriko Oyama, MD, PhD

Department of Radiology

Hokkaido University Hospital

Kita 15, Nishi 7, Kita-ku

Sapporo 060-8638, Japan

TEL 81-11-706-5977

FAX 81-11-706-7876

E-mail: norikooyama@med.hokudai.ac.jp

A 71-year-old woman presented with one year of moderate-effort dyspnea and a past history of brain infarction. Image quality on trans-thoracic echocardiography was suboptimal due to obesity. An ECG-gated 64-slice MDCT image showed a dilated proximal coronary sinus (CS); the distal CS was normal in caliber. The 3D volume rendering (VR) (Figure 1a) and multi-planner reconstruction (Figure 1b) images revealed a partial defect of the CS. The relationship between the mitral valve and the defect was clearly visualized with the endocardial view (Figure 1c). The diameter of the CS defect was 2.3 cm. The pulmonary-to-systemic blood flow ratio ($Q_p:Q_s$) was more than 3 by cardiac catheterization. The patient had a direct surgical closure of the defect.

Preoperative diagnosis of unroofed coronary sinus has been reported by catheterization, angiography (1), echocardiography, CT (2) and MRI (3, 4). We successfully diagnosed unroofed coronary sinus preoperatively by the 3D VR and endocardial views with 64-slice MDCT, and those clearly showed the defect of the roof of the CS and the relationship with the surrounding structure. The 3D VR view was helpful for visualizing the relationship between the right coronary artery and dilated CS. The endocardial view was helpful for surgical planning as it was similar to the surgeon's view and delineated the relationship between the CS defect and the mitral valve. In conclusion, MDCT is a good non-invasive tool for the evaluation of unroofed coronary sinus, and the reconstructed endocardial view is useful for preoperative

surgical planning.

References

1. Freedom RM, Culham JA, Rowe RD. Left atrial to coronary sinus fenestration (partially unroofed coronary sinus). Morphological and angiocardiographic observations. *Br Heart J* 1981; 46:63-68.
2. Brancaccio G, Miraldi F, Ventriglia F, Michielon G, Di Donato RM, De Santis M. Multidetector-row helical computed tomography imaging of unroofed coronary sinus. *Int J Cardiol* 2003; 91:251-253.
3. Iga K, Hori K, Matsumura T, Gen H, Yamashita M, Takahashi S. Unroofed coronary sinus syndrome: diagnostic consideration by contrast echocardiography and usefulness of transesophageal echocardiography and magnetic resonance imaging. *Heart Vessels* 1992; 7:91-94.
4. Hamada Y, Ebihara H, Tanimoto Y, Kobayashi Y, Matsuda Y. Unroofed coronary sinus demonstrated by two-dimensional echocardiography. *Am Heart J* 1984; 108:1558-1560.

Figure Legends:

Figure 1a: MDCT 3D volume rendering (VR) image (the point-of-view is posterior to the patient)

The 3D VR image shows the dilated, partially unroofed CS and its relationship to the LA. The asterisk (*) indicates the unroofed portion of the CS. Note that the distal CS (white arrowheads) is normal in size.

Figure 1b: Multi-planar reconstructed image

It shows a virtual viewpoint (arrow) from the LA toward the coronary sinus defect (*) and left ventricle, from which the images in Figure 1c and 1d were observed.

Figure 1c: 3D endocardial view image in the left atrium

Note the relationship between the MV and the defect of the CS roof (asterisk*).

Figure 1d: 3D endocardial cine image

The cine shows views from right upper side of the LA towards the defect of the CS.

CS: coronary sinus, LA: left atrium, LUPV: left upper pulmonary vein, LLPV: left lower pulmonary vein, MV: mitral valve, RLPV: right lower pulmonary vein

1a	1b
1c	