



Title	OsJAR1 and OsJAR2 are jasmonyl-L-isoleucine synthases involved in wound- and pathogen-induced jasmonic acid signalling
Author(s)	Wakuta, Shinji; Suzuki, Erika; Saburi, Wataru; Matsuura, Hideyuki; Nabeta, Kensuke; Imai, Ryozi; Matsui, Hirokazu
Citation	Biochemical and Biophysical Research Communications, 409(4), 634-639 https://doi.org/10.1016/j.bbrc.2011.05.055
Issue Date	2011-06-17
Doc URL	http://hdl.handle.net/2115/46840
Type	article (author version)
File Information	BBRC409-4_634-639.pdf



[Instructions for use](#)

OsJAR1 and OsJAR2 are jasmonyl-L-isooleucine synthases involved in wound- and pathogen-induced jasmonic acid signalling

Shinji Wakuta^a, Erika Suzuki^a, Wataru Saburi^{a,*}, Hideyuki Matsuura^a, Kensuke Nabeta^a,
Ryozo Imai^b and Hirokazu Matsui^a

^a *Division of Applied Bioscience, Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan*

^b *National Agricultural Research Center of Hokkaido region, National Agriculture and Food Research Organization, Sapporo 062-8555, Japan*

*Corresponding author. Tel./Fax: +81-11-706-2508

E-mail address: saburiw@chem.agr.hokudai.ac.jp

Division of Applied Bioscience, Research Faculty of Agriculture

Hokkaido University, N-9, W-9, Kita-Ku

Sapporo 060-8589, Japan

Key words: jasmonyl-L-isooleucine synthase; jasmonic acid; jasmonyl-L-isooleucine; wound response; response to blast infection

Abbreviations: IAA, indole-3-acetic acid; JA, jasmonic acid; JA-Ile, jasmonyl-L-isooleucine; MS/MS, tandem mass spectrometry; UPLC, ultra-performance liquid chromatography.

Abstract

The synthesis of JA-Ile was catalysed by JA-Ile synthase, which is a member of the group I GH3 family of proteins. Here, we showed evidence that OsGH3.5 (OsJAR1) and OsGH3.3 (OsJAR2) are the functional JA-Ile synthases in rice, using recombinant proteins. The expression levels of *OsJAR1* and *OsJAR2* were induced in response to wounding with the concomitant accumulation of JA-Ile. In contrast, only the expression of *OsJAR1* was associated with the accumulation of JA-Ile after blast infection. Our data suggest that these two JA-Ile synthases are differentially involved in the activation of JA signalling in response to wounding and pathogen challenge in rice.

Highlights

- 1) Two genes from rice plant, OsGH3.5 (*OsJAR1*) and OsGH3.3 (*OsJAR2*), had been confirmed to encode functional JA-Ile synthase.
- 2) The expression of *OsJAR1* and *OsJAR2* were induced in response to wounding with the concomitant accumulation of JA-Ile.
- 3) Only the expression of *OsJAR1* was associated with the accumulation of JA-Ile after blast infection.

OsJAR1 and OsJAR2 are jasmonyl-L-isoleucine synthases involved in wound- and pathogen-induced jasmonic acid signalling

Shinji Wakuta^a, Erika Suzuki^a, Wataru Saburi^{a,*}, Hideyuki Matsuura^a, Kensuke Nabeta^a, Ryozo Imai^b and Hirokazu Matsui^a

^a *Division of Applied Bioscience, Research Faculty of Agriculture, Hokkaido University, Sapporo 060-8589, Japan*

^b *National Agricultural Research Center of Hokkaido region, National Agriculture and Food Research Organization, Sapporo 062-8555, Japan*

*Corresponding author. Tel./Fax: +81-11-706-2508

E-mail address: saburiw@chem.agr.hokudai.ac.jp

Division of Applied Bioscience, Research Faculty of Agriculture

Hokkaido University, N-9, W-9, Kita-Ku

Sapporo 060-8589, Japan

Key words: jasmonyl-L-isoleucine synthase; jasmonic acid; jasmonyl-L-isoleucine; wound response; response to blast infection

Abbreviations: IAA, indole-3-acetic acid; JA, jasmonic acid; JA-Ile, jasmonyl-L-isoleucine; MS/MS, tandem mass spectrometry; UPLC, ultra-performance liquid chromatography.

Abstract

The synthesis of JA-Ile was catalysed by JA-Ile synthase, which is a member of the group I GH3 family of proteins. Here, we showed evidence that OsGH3.5 (OsJAR1) and OsGH3.3 (OsJAR2) are the functional JA-Ile synthases in rice, using recombinant proteins. The expression levels of *OsJAR1* and *OsJAR2* were induced in response to wounding with the concomitant accumulation of JA-Ile. In contrast, only the expression of *OsJAR1* was associated with the accumulation of JA-Ile after blast infection. Our data suggest that these two JA-Ile synthases are differentially involved in the activation of JA signalling in response to wounding and pathogen challenge in rice.

1. Introduction

Jasmonic acid (JA) is an oxylipin phytohormone that is essential for regulating stress responses against insect herbivory and pathogenic attack in plants [1,2]. JA also regulates plant growth and development, including root elongation, fruit ripening, and pollen maturation [3-6]. JA undergoes various modifications that are implicated in activation, inactivation, storage, transport, and volatilisation. Among these modifications, amino acid conjugation controls the physiological activity of JA. JA in conjugation with Ile, Leu, Val, Phe, and ACC have been detected in a wide range of plant species [7,8]. Among them, JA-Ile is a major active compound in JA signalling [9,10].

Similar to auxin, the amino acid conjugation of JA is catalysed by GH3 proteins. The GH3 protein was first isolated from *Glycine max* as an early auxin-responsive gene [11]. GH3 is encoded by a multigene family in various plant species, and the protein sequences are classified into three groups [12,13]. One of the group I GH3 proteins in *Arabidopsis*, AtJAR1, has been shown to synthesise JA-amino acid conjugates, such as JA-Ile, JA-Leu, and JA-Val [14], while group II proteins produce indole-3-acetic acid (IAA) conjugates, such as IAA-Asp, IAA-Glu, and IAA-Ala [15]. Group III proteins are involved in the conjugation of amino acids to 4-substituted benzoate [16].

The physiological roles of group I GH3 proteins have been elucidated in the *Arabidopsis jar1* mutant, which shows decreased sensitivity to exogenous JA [3,17] and decreased levels of JA-Ile [14]. JAR1 is involved in resistance to the soil fungus *Pythium irregular* [18], induced systemic resistance to rhizobacteria [19,20], and protection against ozone damage [21,22]. Group I GH3 proteins with JA-Ile synthase activity have been identified in *Nicotiana attenuate* (JAR4 and JAR6) and tomato (SIJAR1), and they are involved in the activation of defence systems after wounding [23-25].

Three group I GH3 proteins, GH3.3, GH3.5, and GH3.12, are encoded by the rice genome. Although GH3.5 (OsJAR1) has been implicated in phytochrome and jasmonic acid signalling [26], no active JA-Ile synthase has been identified thus far in rice. In this study, we report that OsJAR1 and OsJAR2 (GH3.3) are the functional JA-Ile synthases, and their expression correlates with JA-Ile accumulation after wounding and pathogen

attack.

2. Materials and methods

2.1. Plant materials, growth conditions, and stress treatments

Seeds of *O. sativa* L. cv. Nipponbare were surface-sterilised in 70% ethanol for 3 min and further sterilised in 50% commercial bleach for 30 min. The sterilised seeds were then soaked in distilled water for 4 days and set to germinate in the dark at 20°C. Germinated seeds were uniformly distributed into a seed bed. Seeds were then grown under a 10.5 h light period in a growth chamber at 25°C, 60% humidity, and 530 $\mu\text{mol}/\text{m}^2/\text{s}$ illumination.

At the fourth leaf stage, seedlings were subjected to wounding or pathogenic stress treatments. Seedlings were nipped tightly with a pair of tweezers. Shoots of the treated seedlings were collected at 0 (control), 0.5, 1, 2, and 4 h after the wounding treatment, immediately frozen in liquid nitrogen, and stored at -80°C until further analysis.

For the blast infection, *Magnaporthe grisea* (race, 007) was grown on an oatmeal medium (Difco Laboratories, Detroit, MI, USA) for 1 week at 25°C in the dark, and then spore formation was induced under a 15 W BLB light (FL15S, BLB15W; Toshiba, Tokyo, Japan) for 4 days at 25°C. A spore suspension (2.5×10^7 spores/ml) in 0.05% Tween-20 was sprayed on the seedlings at the fifth leaf stage. The sprayed plants were incubated at 25°C with high humidity in the dark for 20 h and then moved back to the original conditions. Shoots of the treated seedlings were collected at 0 (control), 24, 48, and 72 h after the blast treatment and stored as mentioned above.

2.2. cDNA cloning of three cDNAs from the group I GH3

Based on the amino acid sequences of OsJAR1, OsJAR2, and OsJAR3 obtained from the MSU-RGAP database [27], primers were designed and are shown in Supplementary Table 1. Total RNA was extracted from the rice leaf sheath and roots by RNAqueous (Applied Biosystems, Foster City, CA), and cDNA was synthesised by reverse transcription, using the BcaBest RNA PCR Kit Ver. 1.1 (Takara Bio, Kyoto, Japan). Prime Star Polymerase

(Takara Bio) was used for the cDNA amplification. The reaction temperatures were set to 98°C for 10 s and 68°C for 1 min 30 s with 30 cycles. The DNA fragments were inserted into a pBluescript II SK (+) plasmid (Stratagene, La Jolla, CA, USA), and sequenced to verify that no errors had occurred.

2.3. Construction of expression vectors

Expression plasmids were constructed to produce recombinant OsJAR1 and OsJAR2 as GST and 6x His tagged proteins. The ORFs of *OsJAR1* and *OsJAR2* were amplified with *PciI* and *HindIII* linker sequences from the cDNA clones, respectively. These PCR products were inserted into pET42a (Novagen, Madison, WI, USA) via *NcoI* and *HindIII*.

2.4. Induction of expression of *OsJAR1* and *OsJAR2* and analysis of JA-Ile formation in recombinant *E. coli* cells

E. coli cells carrying each expression vector were cultured overnight in 2 ml of Luria-Bertani (LB) medium, containing 50 µg/ml of kanamycin at 37°C. The culture was transferred into fresh LB medium (2 ml) and incubated until the absorbance at 600 nm reached 0.1. Then, isopropyl thio-β-D-galactoside was added at a final concentration of 1 mM, and the culture was incubated at 37°C for 2 h to induce protein production. Subsequently, JA was added to the culture at a final concentration of 0.1 mM and incubation was conducted at 37°C for 3 h with shaking. After extraction with an equal volume of MeOH and centrifugation at 6,000 x g for 10 min, the supernatant was collected. As an internal standard, JA-Ile *d-6* (5 ng) was added to the extract, and the mixture was analysed, using the ultra-performance liquid chromatography tandem mass spectrometry (UPLC-MS/MS) system. The UPLC-MS/MS instrumentation conditions for the analysis and synthesis of JA and JA-amino acid conjugates were set as previously described [28,29].

2.5. Analyses of the endogenous JA and JA-amino acid conjugates after wounding

The frozen sample was crushed and soaked in 15 ml of 80% aqueous MeOH with 0.5% acetic acid for 12 h. The mixture was filtered to give a crude extract, and JA *d-6* (400 ng),

JA-Ile *d*-6 (100 ng), JA-Leu *d*-6 (100 ng), JA-Trp *d*-6 (100 ng), JA-Gly *d*-6 (100 ng), JA-Val *d*-6 (100 ng), JA-Phe *d*-6 (100 ng), and JA-ACC *d*-6 (100 ng) were added to the extract as internal standards, and analysed as previously described[28,29].

2.6. Real-time PCR analysis

Total RNA from shoots treated with wounding and blast infection were prepared with RNAqueous as described above, and cDNAs were synthesised from the RNA, using random hexamer primers and Superscript III (Invitrogen, Carlsbad, CA, USA). The real-time PCR reaction was performed, using SYBR Premix Ex Taq (Takara Bio), and the signals were detected on a Smart Cycler II System (Takara Bio). The sequences of each specific primer are described in Supplementary Table 2. The real-time PCR reaction was normalised, using the rice *18S rRNA* gene as an internal control. The results shown are the mean values of at least three independent RNA samples.

3. Results

3.1. Phylogenetic relationship of the group I GH3 proteins in plants

To identify the sequence diversity of group I GH3 proteins in plant genomes, sequences were collected from public genome databases for monocots (*Oryza sativa*, *Zea mays*, *Brachypodium distachyon*, and *Sorghum bicolor*) and dicots (*Arabidopsis thaliana*, *Vitis vinifera*, *Glycine max*, and *Populus trichocarpa*). In total, 35 protein sequences were selected for the multiple alignment and phylogenetic analyses, using the Clustal W software [30]. The phylogenetic tree identified four subgroups within the group I GH3 proteins (Fig. 1). Five enzymatically active JA-Ile synthases that had been previously characterised fell into the dicot subgroup I. The three group I GH3 proteins from rice, OsJAR1/OsGH3.5, OsJAR2/OsGH3.3, and OsJAR3/OsGH3.12 were classified into different subgroups. OsJAR1 was classified into the monocot subgroup I, while OsJAR2 and OsJAR3 were classified into the monocot subgroup II.

3.2. Isolation of three cDNAs encoding *OsJAR1*, *OsJAR2*, and *OsJAR3*

To identify the functional JA-Ile synthase genes in rice, the three *OsJAR* cDNAs were isolated. RT-PCR was performed with specific primer sets that were designed using genome annotation information available at MSU-RGAP (Release 6.1). Because two mRNA variants were predicted for *OsJAR2* by the MSU-RGAP database, both *OsJAR2.1* and *OsJAR2.2* cDNAs were isolated. The *OsJAR3* cDNA could not be amplified from the leaf sheath. Among the several tissues tested, the *OsJAR3* cDNA could only be amplified from a callus containing two introns. Therefore, we assumed that *OsJAR3* is not functional in most tissues. Both *OsJAR1* and *OsJAR2.1* have three motifs involved in ATP/AMP binding [31] (Supplementary Fig. 1): motif 1, SSGTT(H/Q)QKRK; motif 2, YGASEGW; and motif 3, YRLGDVVK. However, *OsJAR2.2* does not contain motif 3, suggesting the possibility that *OsJAR2.2* may lack enzymatic activity. The amino acid sequences of *OsJAR1* and *OsJAR2.1* share 60% and 59% identity to *AtJAR1*, respectively.

3.3. Enzymatic production of JA-Ile by *OsJAR1* and *OsJAR2.1*

To examine the JA-Ile-producing activities of *OsJAR1*, *OsJAR2.1*, and *OsJAR2.2*, *E. coli* cells expressing these genes were incubated with JA, and the production of JA-Ile was determined by UPLC-MS/MS (Fig. 2). The analysis of the reaction products in the culture supernatant clearly demonstrated that *OsJAR1* and *OsJAR2.1* catalyse the conjugation of JA to Ile, using endogenous ATP and Ile from *E. coli*. In contrast, no JA-Ile production was detected in the cells producing *OsJAR2.2* (data not shown). Therefore, we concluded that *OsJAR2.1* is a functional *OsJAR2* transcript.

3.4. Induction of mRNAs encoding *OsJAR1* and *OsJAR2* by wounding

The up-regulation of JA-Ile synthase mRNA and accumulation of JA-Ile were concomitantly observed, following leaf wounding in Arabidopsis and tomato [25,29]. Therefore, the expression levels of *OsJAR1* and *OsJAR2* and accumulation of JA-Ile in response to wounding stress were assessed. A real-time PCR analysis showed that the transcripts of *OsJAR1* and *OsJAR2* were induced transiently, reaching peaks at 1 h and 0.5

h, respectively (Fig. 3A). Analyses of endogenous JA, JA-Ile, and other JA-conjugates in wounded rice plants showed that the endogenous levels of JA and JA-Ile were increased after wounding. Maximum levels of accumulation of JA (3.29 nmol/g FW) and JA-Ile (1.59 nmol/g FW) were observed at 1 h after wounding (Fig. 3B). Among the analysed JA-conjugates, JA-Val (0.12 nmol/g FW) and JA-Phe (<0.01 nmol/g FW) were detected, but they were induced at much lower levels than JA-Ile (Fig. 3C). These data indicate that OsJAR1 and OsJAR2 are both involved in the synthesis of JA-Ile in response to wounding stress.

3.5. Determination of mRNA levels of *OsJAR1* and *OsJAR2* after blast infection

To elucidate the influence of pathogenic attack on jasmonic acid signalling, blast infections were performed by spraying intact plants with a spore suspension of rice blasts. The real-time PCR analysis revealed that *OsJAR1* mRNA levels increased gradually from 48 to 72 h after the blast spray, while *OsJAR2* expression was not significantly altered (Fig. 4A). Endogenous JA-Ile levels were also induced from 48 to 72 h after infection and reached maximum levels (0.5 nmol/g FW) at 72 h (Fig. 4B). In infected leaves, 0.069 nmol/g FW of JA-Val was also detected at 72 h (Fig. 4C), while JA-Phe was not detected under these conditions. These data showed that OsJAR1 mainly produces JA-Ile in response to pathogen attack. JA levels gradually increased from 0 to 72 h after inoculation, while a dramatic increase in JA-Ile was observed after 24 h (Fig. 4).

4. Discussion

OsJAR1 (OsGH3.5) has been implicated in phytochrome and JA signalling [26]. However, enzymes that synthesise JA-amino acid conjugates have not been identified in rice. In this study, we demonstrated that OsJAR1 (OsGH3.5) and OsJAR2 (OsGH3.3), belonging to the group I GH3 family proteins, exert their activities to form JA-Ile from JA and Ile in response to wounding and pathogenic stress in rice.

A phylogenetic analysis of group I GH3 proteins from eight plant species whose genome sequences are known revealed that they are divided into four subgroups. Five previously

characterised group I GH3s, AtJAR1, SIJAR1, NgJAR1, NaJAR4, and NaJAR6, are all clustered into the dicot subgroup I (Fig. 2). It was not known if the other subgroup members within the group I GH3 proteins also displayed JA-Ile synthase activities. Because OsJAR1 and OsJAR2 are positioned in distinct monocot subgroups (Fig. 1), our data suggest the possibility that a wider cluster within the group I GH3 proteins displays JA-Ile synthase activity.

In contrast to *OsJAR1* and *OsJAR2*, the existence of the *OsJAR3* transcript was questionable. No corresponding full-length cDNA or EST for OsJAR3 could be identified in the databases [13]. The *OsJAR3* mRNA was not detectable in tissues, including those from the shoot and roots, where *OsJAR1* and *OsJAR2* mRNA are present [13]. However, another group reported the detection of *OsJAR3* mRNA in the roots [12]. In this study, RT-PCR was performed, using mRNA from the roots, shoot, leaf sheath, leaf blade, panicles, immature seeds and callus cells, but only the unspliced form of *OsJAR3* transcripts were detected in the callus. The resulting transcript encodes a truncated protein due to a nonsense mutation. The putative spliced form of the *OsJAR3* mRNA, encoding a protein with three conserved ATP/AMP binding motifs, may exist in tissues under specific stress conditions or in certain developmental stages. However, it is also possible that OsJAR3 is a pseudogene because there is one group I GH3 protein in each subgroup in *Brachypodium*.

The expressions of *OsJAR1* and *OsJAR2* were both rapidly and transiently induced in response to wounding (Fig. 3). The accumulation pattern of JA-Ile in rice leaves after wounding was in accordance with the expression patterns of *OsJAR1* and *OsJAR2* (Fig. 3). Similarly, the accumulation of JA-Ile in response to wounding stress is rapid and transient in *Arabidopsis*, tobacco, and tomato leaves [23,25,32]. These observations suggest that the transcriptional activation of *OsJAR1* and *OsJAR2* is required for the production of JA-Ile, which turns on jasmonic acid signalling in rice as was reported for *Arabidopsis*. The formation of JA-Leu, JA-Trp, JA-Gly, JA-Val, JA-Phe, and JA-ACC were monitored after wounding, and JA-Val and JA-Phe could be detected at low levels compared to JA-Ile.

The accumulation of JA-Ile was also observed in blast-infected leaves. However, the

accumulation pattern was quite different from that caused by wounding. Compared with wound stress, JA-Ile accumulated slowly and continuously and reached higher levels than JA. Relatively slow accumulations of JA and JA-Ile may be due to the fact that pathogen infection occurs progressively at the site of inoculation. A dramatic increase in JA-Ile levels was observed after 24 h (Fig. 4). The correlation of *OsJAR1* expression and JA-Ile accumulation suggests that OsJAR1 is involved in JA-Ile synthesis during blast infection.

In summary, we have demonstrated that two functionally active JA-Ile synthases are present in rice. They are differentially regulated in response to multiple stressors. Further enzymatic and physiological characterisations of OsJAR1 and OsJAR2 are required to understand the role of JA signalling in rice.

5. References

- [1] M. McConn, R.A. Creelman, E. Bell, J.E. Mullet, J. Browse, Jasmonate is essential for insect defense in *Arabidopsis*, Proc. Natl. Acad. Sci. USA 94 (1997) 5473-5477.
- [2] P. Vijayan, J. Shockey, C.A. Lévesque, R.J. Cook, J. Browse, A role for jasmonate in pathogen defense of *Arabidopsis*, Proc. Natl. Acad. Sci. USA 95 (1998) 7209-7214.
- [3] P.E. Staswick, W. Su, S.H. Howell, Methyl jasmonate inhibition of root growth and induction of a leaf protein are decreased in an *Arabidopsis thaliana* mutant, Proc. Natl. Acad. Sci. USA 89 (1992) 6837-6840.
- [4] X. Fan, J.P. Mattheis, J.K. Fellman, A role for jasmonates in climacteric fruit ripening, Planta 204 (1998) 444-449.
- [5] A. Stintzi, J. Browse, The *Arabidopsis* male-sterile mutant, *opr3*, lacks the 12-oxophytodienoic acid reductase required for jasmonate synthesis, Proc. Natl. Acad. Sci. USA 97 (2000) 10625-10630.
- [6] C. Wasternack, Jasmonates: An update on biosynthesis, signal transduction and action in plant stress response, growth and development, Annals of Botany 100 (2007) 681-697.
- [7] R. Kramell, R. Atzorn, G. Schneider, O. Miersch, C. Bruckner, G. Schmidt, B. Parthier, Occurrence and identification of jasmonic acid and its amino acid conjugates induced by osmotic stress in barley leaf tissue, J. Plant Growth Regul. 14, (1995) 29-36.

- [8] B. Hause, W. Maier, O. Miersch, R. Kramell, D. Strack, Induction of jasmonate biosynthesis in arbuscular mycorrhizal barley roots, *Plant Physiol.* 130 (2002) 1213-1220.
- [9] A. Chini, S. Fonseca, G. Fernandez, B. Adie, J.M. Chico, O. Lorenzo, G. Gracia-Casado, I. Lopez-Vidriero, F.M. Lozano, M.R. Ponce, J.L. Micol, R. Solano, The JAZ family of repressors is the missing link in jasmonate signalling, *Nature* 448 (2007) 666-671.
- [10] B. Thines, L. Katsir, M. Melotto, Y. Niu, A. Mandaokar, G. Liu, K. Nomura, S. Y. He, G.A. Howe, J. Browse, JAZ repressor proteins are targets of the SCFCO11 complex during jasmonate signalling, *Nature* 448 (2007) 661-665.
- [11] G. Hagen, T.J. Guilfoyle, Rapid induction of selective transcription by auxins, *Mol. Cell Biol.* 5 (1985) 1197-1203.
- [12] M. Jain, N. Kaur, A.K. Tyagi, J.P. Khurana, J.P. The auxin-responsive GH3 gene family in rice (*Oryza sativa*), *Funct. Integr Genomics* 6 (2006) 36-46.
- [13] J. Terol, C. Domingo, M. Talon, The GH3 family in plants: Genome wide analysis in rice and evolutionary history based on EST analysis, *Gene* 371 (2006) 279-290.
- [14] P.E. Staswick, I. Tiryaki, The oxylipin signal jasmonic acid is activated by an enzyme that conjugates it to isoleucine in Arabidopsis, *Plant Cell* 16 (2004) 2117-2127.
- [15] P.E. Staswick, B. Serban, M. Rowe, I. Tiryaki, M.T. Maldonado, M.C. Maldonado, W.P. Suza, Characterization of an Arabidopsis enzyme family that conjugates amino acids to indole-3-acetic acid, *Plant Cell* 17 (2005) 616-627.
- [16] R.A. Okrent, M.D. Brooks, M.C. Wildermuth, Arabidopsis GH3.12 (PBS3) conjugates amino acids to 4-substituted benzoates and is inhibited by salicylate, *J. Biol. Chem.* 284 (2009) 9742-9754.
- [17] P.E. Staswick, I. Tiryaki, M. Rowe, M. Jasmonate response locus JAR1 and several related Arabidopsis genes encode enzymes of the firefly luciferase superfamily that show activity on jasmonic, salicylic, and indole-3-acetic acids in an assay for adenylation, *Plant Cell* 14 (2002) 1405-1415.
- [18] P.E. Staswick, G.Y. Yuen, C.C. Lehman, Jasmonate signaling mutants of *Arabidopsis* are susceptible to the soil fungus *Pythium irregular*, *Plant J.* 15 (1998) 747-754.

- [19] C.M.J. Pieterse, S.C.M. van Wees, J.A. van Pelt, M. Knoester, R. Laan, H. Gerrits, P.J. Weisbeek, L.C. van Loon, A novel signaling pathway controlling induced systemic resistance in Arabidopsis, *Plant Cell* 10 (1998) 1571-1580.
- [20] J.D. Clarke, S.M. Volko, H. Ledford, F.M. Ausubel, X. Dong, Roles of salicylic acid, jasmonic acid, and ethylene in cpr-induced resistance in Arabidopsis, *Plant Cell* 12 (2000) 2175-2190.
- [21] K. Overmyer, H. Tuominen, R. Kettunen, C. Betz, C. Langebartels, H. Sandermann, J. Kangasjarvi, Ozone-sensitive Arabidopsis *rcd1* mutant reveals opposite roles for ethylene and jasmonate signaling pathways in regulating superoxide-dependent cell death, *Plant Cell* 12 (2000) 1849-1862.
- [22] M.V. Rao, H. Lee, R.A. Creelman, J.E. Mullet, K.R. Davis, Jasmonic acid signaling modulates ozone-induced hypersensitive cell death, *Plant Cell* 12 (2000) 1633-1646.
- [23] J.H. Kang, L. Wang, A. Giri, I.T. Baldwin, Silencing threonine deaminase and JAR4 in *Nicotiana attenuata* impairs jasmonic acid-isoleucine-mediated defences against *Manduca sexta*, *Plant Cell* 18 (2006) 3303-3320.
- [24] L. Wang, S. Allmann, J. Wu, I.T. Baldwin, Comparisons of LIPOXYGENASE3- and JASMONATE-RESISTANT4/6-silenced plants reveal that jasmonic acid and jasmonic acid-amino acid conjugates play different roles in herbivore resistance of *Nicotiana attenuata*, *Plant Physiol.* 146 (2007) 904-915.
- [25] W.P. Suza, M.L. Rowe, M. Hamberg, P.E. Staswick, A tomato enzyme synthesizes (+)-7-*iso*-jasmonoyl-L-isoleucine in wounded leaves, *Planta* 231 (2010) 717-728.
- [26] M. Riemann, M. Riemann, M. Takano, Rice JASMONATE RESISTANT 1 is involved in phytochrome and jasmonate signalling, *Plant Cell Environ.* 31 (2008) 783-792.
- [27] S. Ouyang, W. Zhu, J. Hamilton, H. Lin, M. Campbell, K. Childs, T. Thibaud-Nissen, R.L. Malek, Y. Lee, L. Zheng, J. Orvis, B. Haas, J. Wortman, C.R. Buell, The TIGR rice genome annotation resource: improvements and new features, *Nucleic Acids Res.* 35 (2007) 883-887.
- [28] H. Matsuura, A. Aoi, C. Satou, M. Nakaya, C. Masuta, K. Nabeta. Simultaneous UPLC MS/MS analysis of endogenous jasmonic acid, salicylic acid, and their related

- compounds, *Plant Growth Regul.* 57 (2009) 293-301.
- [29] C. Sato, Y. Seto, K. Nabeta, H. Matsuura, Kinetics of the accumulation of jasmonic acid and its derivatives in systemic leaves of tobacco (*Nicotiana tabacum* cv. Xanthi nc) and translocation of deuterium-labeled jasmonic acid from the wounding site to the systemic site, *Biosci. Biotechnol. Biochem.* 73 (2009) 1962-1970.
- [30] J.D. Thompson, D.G. Higgins, T.J. Gibson, CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice, *Nucleic Acids Res.* 22 (1994) 4673-4680.
- [31] K.H. Chang, H. Xiang, D. Dunaway, Mariano, Acyl-adenylate motif of the acyl-adenylate/thioester-forming enzyme superfamily: a site-directed mutagenesis study with the *Pseudomonas* sp. strain CBS3 4-chlorobenzoate: coenzyme A ligase, *Biochemistry* 36 (1997) 15650-15659.
- [32] W.P. Suza, P.E. Staswick, The role of JAR1 in jasmonoyl-L-isoleucine production during Arabidopsis wound response, *Planta* 227 (2008) 1221-1232.
- [33] N. Saitou, M. Nei, The neighbor-joining method: a new method for reconstructing phylogenetic trees, *Mol. Biol. Evol.* 4 (1987) 406-425.
- [34] Arabidopsis Genome Initiative, Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*, *Nature* 408 (2000) 796-815.
- [35] G.A. Tuskan, et al. The genome of black cottonwood, *Populus trichocarpa* (Torr. & Gray), *Science* 313 (2006) 1596-1604.
- [36] Grapevine Genome Characterization, The grapevine genome sequence suggests ancestral hexaploidization in major angiosperm phyla, *Nature* 449 (2007) 463-467.
- [37] A.H. Paterson, et al, The *Sorghum bicolor* genome and the diversification of grasses, *Nature* 457 (2009) 551-556.
- [38] P.S. Schnable, et al, The B73 maize genome: complexity, diversity, and dynamics, *Science* 326 (2009) 1112-1115.
- [39] J. Schmutz, et al, Genome sequence of the palaeopolyploid soybean, *Nature* 463 (2010) 178-183.
- [40] International Brachypodium Initiative, Genome sequencing and analysis of the model

grass *Brachypodium distachyon*, Nature 463 (2010) 763-768.

[41] M. Jain, A. Nijhawan, A.K. Tyagi, J.P. Khurana, Validation of housekeeping genes as internal control for studying gene expression in rice by quantitative real-time PCR, Biochem. Biophys. Res. Commun. 345 (2006) 646-651.

6. Figure captions

Fig. 1. A phylogenetic tree of the amino acid sequences from the group I GH3 family proteins from various plant species.

The tree was constructed, using ClustalX (version 2.0.11) (neighbour-joining method; [33]), and visualised with NJ plot, using amino acid sequences gathered from various databases [32,34-40]. Numbers indicate bootstrap values. The positions of proteins targeted in this study are indicated with open triangles: OsJAR1 (LOC_Os05g50890), OsJAR2 (LOC_Os01g12160), and OsJAR3 (LOC_Os11g08340). Known JA-Ile synthase proteins are indicated by closed triangles: AtJAR1 (*Arabidopsis thaliana*); NaJAR4 and NaJAR6 (*Nicotiana attenuata*); NgJAR1 (*Nicotiana glutinosa*); and SIJAR1 (*Solanum lycopersicum*). Protein sequences were obtained from genome databases of *Glycine max* (Glyma16g03010, Glyma07g06370, Glyma19g44310, Glyma03g41700, Glyma13g37550, Glyma12g32910, Glyma12g11890, and Glyma06g45120), *Populus trichocarpa* (POPTR_0002s16960, POPTR_0014s09120, POPTR_13s14050, and POPTR_0019s13450), *Vitis vinifera* (GSVIVT01027057001 and GSVIVT01030558001), *Brachypodium distachyon* (Bradi2g14830 and Bradi2g07230), *Zea mays* (GRMZM2G091276_P02, GRMZM2G162413_P01, GRMZM2G001421_P01, GRMZM2G060991_P01, and GRMZM2G061005_P01), *Sorghum bicolor* (Sb03g028240, Sb09g030050, Sb03g001490, and Sb03g001500), and *Arabidopsis thaliana* (AtGH3.10: At4g03400).

Fig. 2. Metabolism of JA in *E. coli* cells carrying the *OsJAR1* and *OsJAR2* genes.

The trace chromatogram in the selected reaction monitoring mode shows the ion peaks m/z 328.03/129.68 for JA-Ile *d*-6 and m/z 322.03/129.68 for JA-Ile. JA-Ile *d*-6 was used

as an internal standard. The metabolism in culture supernatant of *E. coli* cells with pET42a as a control (A), pET42a:OsJAR1 (B), and pET42a:OsJAR2 (C) were analysed.

Fig. 3. The expression levels of *OsJAR1* and *OsJAR2* mRNA and the accumulation of JA and JA-amino acid conjugates in response to wounding.

(A) The accumulation of *OsJAR1* and *OsJAR2* transcripts after mechanical wounding was quantified by quantitative real-time RT-PCR with the specific primer sets. Data were normalised with respect to *18s rRNA* levels [41]. The relative amount of each mRNA was calculated, using intact (unwounded) plants as standards. Each value is the mean \pm S.D. of three independent experiments. (B),(C) Endogenous amounts of JA (triangles), JA-Ile (circles), and JA-Val (squares) were assayed in plants after wound treatments by the UPLC-MS/MS system. JA-Val was not detected before wounding. Data are mean \pm S.D. for three independent experiments.

Fig. 4. The expression levels of *OsJAR1* and *OsJAR2* mRNA and the accumulation of JA and JA-amino acid conjugates in response to blast infection.

(A) The accumulation of *OsJAR1* and *OsJAR2* transcripts after blast infection was quantified by quantitative real-time RT-PCR with the specific primer sets. Data were normalised with respect to *18s rRNA* levels [41]. The relative amount of each mRNA was calculated against the amount in intact (uninfected) rice. Each value is the mean \pm S.D. of three independent experiments. (B), (C) Endogenous levels of JA (triangles), JA-Ile (circles), and JA-Val (squares) were assayed in plants after infection, using the UPLC-MS/MS system. Data are the mean \pm S.D. of three independent experiments.




Fig. 1.




Fig. 2.




Fig. 3.




Fig. 4.

OsJAR1 1 : -----MTICSCEETINEFEMLTRDAARVQKDTLKKILEINASAEYLQNFGLGG--RTDAESYKSCIPICVHNDI
OsJAR2 1 : MLEKKATRSTRVDGVSGEAVIEEFERVTRDAANVQRETLRRILAENGGVEYLRGLGLAG--ATDPATFRARVPLATHADL
AtJAR1 1 : -----MLEKVETFDMNRVIDEEDEMTRNAHQVQKQTLKEILLKNQSAIYLQNCGLNGNATDPEEAFKSMVEIVTDVEL

Motif 1

OsJAR1 68 : EPYIQRIVDGDITSPVVTGEPITNLSLSSGTHCKPKFIPFNDELLETLOIYRISYAFRNREYPIGO-GKALQFVYGSKQ
OsJAR2 79 : EPYIDRIADGDASPVLTAKPATSISLSSGTOGKRKYLLENEELVKSIMQIYRISYAFRNREFPVEN-GKALQFIYSSRE
AtJAR1 74 : EPYIKRMVDGDITSPILTGHVPVAISLSSGTSQGRPKFIPFTDELMENTLQLFRTAFAFRNRDFPIDDNGKALQEIFSSKQ

OsJAR1 147 : VITKGGILATATTNLYRRORYKEGMDIOSQCSPDEVIFGPDFHOSLYCHLLCGLIYSEEVHSVFSTFAHSLVHAFOT
OsJAR2 158 : TRTKGGLTATTATTNVYRSEEFKATMRDIOSQCCSPDEVIFGPDFAOSLYCHLLAGLLAAGDVQIVSATFAHSLVLAFO
AtJAR1 154 : YISTGCVPGVTATTNVYRNPNFKAGMKSITSPSCSPDEVIFSPDVHQAALYCHLLSGILFRDOVQYVFAVFAHGLVHAERT

OsJAR1 227 : FEEVWEDLCTDIRDG-VLSKKVITAPSIREAVSKILKP-NPELADSIYKCIGLSNWYGVIPALWPNAKVYVGIMTGSMEP
OsJAR2 238 : FERAWEDLCADIRRCVSPSRVTSFAVRRAMAALLAAPNPLGLADEVARKCAALSNWYGVIPALWPNARVYVGIMTGSMEH
AtJAR1 234 : FEQVWEEIVTDIKDG-VLSNRIITVSVRTAMSKLLTP-NPELAETIRTKMSLSNWYGLIPALFPNAKVVYVGIMTGSMEP

Motif 2

OsJAR1 305 : YLKCLRHYAGNPLISADYGASEGWVGSNIDFTVPPEQVTYAVLPQVGYFEEIPLKPIGEETENSASIHYESDPVGLT
OsJAR2 318 : YVKCLRHYAGGLPLVAAEYGASEGWVGANVEFGTPPERATFTVLPDIAYFEETPLKPVAGDGGY-----AEAEPVGLT
AtJAR1 312 : YVPKLRHYAGDLPLVSHDYGSSEGWIAANVTPELSPPEEATFAVIPNLGYFEELEVSETGEGEK-----PVGLT

Motif 3

OsJAR1 385 : EVEVGKIYEVVITNFAGLYRYRLGDVVKIARFHNSTPELOFICRRSLVLSINIDKNTEKDLQLAVEEASKFLEG-EKLEV
OsJAR2 391 : EVAAGELYEVVMTTFAGLYRYRLGDVVKVAGFYNATPKLKFVCCRNLMLSNINIDKNSEQDLQLAVDAAARAVLAGEKLEV
AtJAR1 381 : QVKIGEEYEVVITNYAGLYRYRLGDVVKVIGFYNTPOLKFCRRNLILSNINIDKNTERDLQLSVESAKRLSE-EKIEV

OsJAR1 464 : MDFTSFVERSSDPGRYVIFWELSG----DASDEVLSSCANALDLAFIDAGYTGSRKIKTIGPLELRILRKGTKEILDHF
OsJAR2 471 : VDYTSHADVSSDPGHYVVFLELNAADPAAVDGDVMQACCDELDRAFADAGYVGSRKSGAIAPLELRVLQRGTQKVLRYH
AtJAR1 460 : IDFSYIDVSTDPGHYAFWEISG----ETNEDVLQDCCNCLDRAFIDAGYVSSRKCKTIGALELRVVAKGTFRKIQEHF

OsJAR1 540 : LSLCGAVSQFKTPREFVNPSNSKVLQILSRNVTQSYFSTAYGF
OsJAR2 551 : LSLGAPVSQFKSPRCVSRNSGVLQILAGCTVNVFFSSAYD-
AtJAR1 536 : LGLGSSAGQFKMPCVKPSNAKVLQILCENVVSSYFSTAF--

Supplemental Fig. 1.

Supplementary Table 1. Primer sequences used for the cloning of OsJAR1, OsJAR2.1, OsJAR2.2, and OsJAR3

Primer name	Primer sequence
OsJAR1-F	5'-GCCAGGATTCTACCATTACAATGACG-3'
OsJAR1-R	5'-TGTATCAAATCCATAGGCAGTACTG-3'
OsJAR2.1-F	5'-ATGTTGGTCGATCGTGCAGG-3'
OsJAR2.1-R	5'-GCGTCTTTGGTTTGCAATTCG-3'
OsJAR2.2-F	5'-GGCGAGATGTTGGAGAAGAAGGC-3'
OsJAR2.2-R	5'-TTATTTTTGGGCAAACATTATATGGCC-3'
OsJAR3-F	5'-GATCCTCTTGACATTGTTGCAG-3'
OsJAR3-R	5'-CACACATTCAAACACCGGC-3'

Supplementary Table 2. Primer sequences used for the real-time PCR analysis

Gene name	Primer sequence
<i>OsJAR1</i>	5'-AAGGTTTGTGAACCCATCAAACAGC-3' 5'-ATAAATACTTTGCAGCACTTGTTACG-3'
<i>OsJAR2</i>	5'-CGCCACTACCTCTCCCTCGGCG-3' 5'-GCGTCTTTGGTTTGCAATTCG-3'
<i>18S rRNA</i> ^a	5'-CTACGTCCCTGCCCTTTGTACA-3' 5'-ACACTTCACCGGACCATTCAA-3'

^a [41]

Supplemental Fig. 1. Multiple alignment of amino acid sequences of OsJAR1, OsJAR2, and AtJAR1.

Amino acid sequences were aligned using the ClustalW program. Database accession numbers are as follows: OsJAR1, LOC_Os05g50890.1; OsJAR2, LOC_Os01g12160.1; and AtJAR1, NP_850453. Conserved residues are highlighted in black. The bars above the sequences show the three conserved motifs responsible for ATP/AMP binding.