Title	Synthesis of Cyclic ADP-Carbocyclic-Xylose and its 3'-OO-Methyl Analogue as Stable and Potent Ca ²⁺ -Mobilizing Agents
Author(s)	Kudoh, Takashi; Matsuda, Akira; Shuto, Satoshi; Murayama, Takashi; Ogawa, Yasuo
Citation	Nucleosides, Nucleotides and Nucleic Acids, 25(4-6), 583-599 https://doi.org/10.1080/15257770600685867
Issue Date	2006-04
Doc URL	http://hdl.handle.net/2115/46970
Rights	This is an electronic version of an article published in Nucleosides, Nucleotides & Nucleic Acids, 25(4/6), 583-599, 2006. Nucleosides, Nucleotides & Nucleic Acids is available online at: http://www.informaworld.com/openurl?genre=article&issn=15257770&volume=25&issue=4-6&spage=583
Type	article (author version)
File Information	366_NsNt Kudo.pdf

[Instructions for use](#)

SYNTHESIS OF CYCLIC ADP-CARBOCYCLIC-XYLOSE AND ITS 3''-O-METHYL ANALOGUE AS STABLE AND POTENT Ca^{2+} -MOBILIZING AGENTS

Takashi Kudoh,^a Takashi Murayama,^b Yasuo Ogawa,^b Akira Matsuda^a and Satoshi^a Shuto*

^a*Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan*

^b*Department of Pharmacology, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113-8421, Japan*

Corresponding Author: Satoshi Shuto

Phone & Fax: +81-11-706-3769.

E-mail: shu@pharm.hokudai.ac.jp

This paper is dedicated to Professor Eiko Ohtsuka on the occasion of her 70th birthday.

ABSTRACT

We previously showed that 3''-deoxy-cyclic ADP-carbocyclic-ribose (3''-deoxy-cADPcR, **2**) is a stable and highly potent analogue of cyclic ADP-ribose (cADPR, **1**), a Ca^{2+} -mobilizing second messenger. From these results, we newly designed another 3''-modified analogues of cADPcR and identified the N1-“xylo”-type carbocyclic analogue, i.e., cADPcX (**4**), as one of the most potent cADPR-related compounds reported so far.

INTRODUCTION

Much attention has been focused on cyclic ADP-ribose (cADPR, **1**, **FIGURE 1**), a naturally occurring metabolite of NAD^+ ,² due to the biological interest.³ cADPR has been shown to mobilize intracellular Ca^{2+} in various cells, and is now recognized as a general mediator involved in Ca^{2+} signaling.³ Under neutral conditions, cADPR is in a zwitterionic form with a positive charge around the N(1)-C(6)-N⁶ moiety ($\text{pK}_a = 8.3$),

making the molecule unstable. The charged adenine moiety attached to the anomeric carbon of the N1-ribose can be an efficient leaving group. Accordingly, cADPR is readily hydrolyzed at the unstable *N*-1-ribosyl linkage of its adenine moiety to produce ADP-ribose (ADPR), even in neutral aqueous solution.⁴ Under physiological conditions, cADPR is also hydrolyzed at the *N*-1-ribosyl linkage by cADPR hydrolase to give the inactive ADPR.⁴

cADPR analogues can be used in proving the mechanism of cADPR-mediated Ca^{2+} signaling pathways and are also expected to be lead structures for the development of drugs, since cADPR has been shown to play important physiological roles.³ Therefore, the synthesis of cADPR analogues has been extensively investigated by enzymatic and chemo-enzymatic methods using ADP-ribosyl cyclase-catalyzed cyclization. However, the analogues obtained by these methods are limited due to the substrate-specificity of the ADP-ribosyl cyclase.³

On the other hand, in the chemical synthesis of cADPR and its analogues, construction of the large 18-membered ring structure is the key step, and we recently developed an efficient method for forming the 18-membered ring employing phenylthiophosphate-type substrates.⁵ When these substrates were activated by AgNO_3 or I_2 in the presence of molecular sieves in pyridine, the corresponding 18-membered ring products were obtained in high yields.^{5b,c} Using this method, we successfully synthesized cyclic ADP-carbocyclic-ribose (cADPcR, **2**),^{5c} designed as a stable mimic of cADPR, in which the oxygen atom in the *N*-1-ribose ring of cADPR is replaced by a methylene group. Biological evaluation of cADPcR showed that it actually act as biologically and chemically stable equivalent of cADPR.^{5c}

Based on these results, we have investigated further synthetic and biological studies on N1-carbocyclic derivatives of cADPR.⁶ In the course of these studies, we describe here the synthesis and biological evaluation of newly designed analogues of cADPcR, which are cyclic ADP-carbocyclic-xylose (cADPcX, **4**) and the corresponding 3''-*O*-methyl derivative (3''-OMe-cADPcX, **5**).

RESULTS AND DISCUSSION

Design and Synthetic Plan. We previously showed 1) that cADPcR (**2**) is actually resistant to both enzymatic and chemical hydrolysis, since it has a chemically and biologically stable *N*-alkyl linkage instead of the unstable N1-glycosidic linkage of cADPR,^{5b} 2) that cADPcR has a conformation similar to that of cADPR,^{6c} and 3) that cADPcR, like cADPR, effectively mobilizes intracellular Ca^{2+} in sea urchin eggs and neuronal cells.^{5c,6b,c} Furthermore, we have investigated SRA of the N1-ribose moiety of cADPcR and clarified that modification at the N1-ribose moiety changes the biological potency.^{6b,c} Throughout these studies, we also found that although deletion of the 2''-hydroxy group resulted in a marked reduction of potency, deletion of the adjacent 3''-hydroxy group (3''-deoxy-cADPcR, **3**) greatly potentiated the Ca^{2+} -mobilizing ability in sea urchin eggs.^{6c} These results suggest that modification at the 3''-position may

improve the biological potency of cADPR and its analogues. Thus, we newly designed another 3''-modified analogues of cADPr, which were the N1-''xylo''-type carbocyclic analogue, i.e., cADPrX (**4**) and the corresponding *O*-methyl analogue **5**.

As described above, we have developed an efficient total synthetic method for cADPR analogues,⁵ which we⁶ and other groups⁷ have effectively used in the synthesis of a variety of cADPR analogues. Thus, we planned to synthesize the target compounds based on the previous total synthetic method.

The synthetic plan is shown in **SCHEME 1** as a retrosynthetic analysis. The chiral carbocyclic-xylosyl amines **III**, composing the N1-substituted moiety in the targets **4** and **5**, could be prepared from commercially available (1*R*)-(-)-2-azabicyclo[2.2.1]hept-5-en-3-one (**8**). From these carbocyclic amines **III** and the known imidazole nucleoside derivative **6**,^{5b} the 5'-phenylthiophosphate-type substrates **II** for the key intramolecular condensation could be prepared. Treatment of **II** with AgNO₃/MS 3A as a promoter^{5a,b} was expected to form the cyclized products **I**, and subsequent acidic treatment for deprotection would furnish the desired cADPr analogues **4** and **5**.

Synthesis. The carbocyclic-xylosyl amine units **14a** and **14b** were synthesized from the optically active bicyclic lactam **8**, as summarized in Scheme 2. After protection of the primary hydroxyl with a triisopropylsilyl (TIPS) group of carbocyclic-ribose derivative **9**, prepared from **8** according to the previously reported method,^{6c} treatment of the resulting **10** under Mitsunobu reaction conditions with ClCH₂CO₂H/DIAD/Ph₃P gave the corresponding 3-position-inverted product **11**, and subsequent removal the 3-*O*-chloroacetyl group with NaOMe/MeOH to afford the carbocyclic-xylose derivative **12**. The 3-hydroxy of **12** was protected with a MOM group under usual conditions or methylated with MeOTf and 2,6-di-*tert*-butyl-4-methylpyridine (DTBMP), followed by removal of the TIPS group, to give the desired 3-*O*-MOM-protected unit **14a** or the 3-*O*-methyl unit **14b**, respectively.

The target 3''-modified cADPr analogues **4** and **5** were successfully synthesized from the carbocyclic amines **14a** or **14b** and the imidazole nucleoside **6**, as shown in **SCHEME 3**.

The *N*-1-substituted adenosine derivatives **15a** and **15b** were obtained in high yield by the treatment of a mixture of **6** and either amine **14a** or amine **14b** with K₂CO₃ in MeOH at room temperature. The 5''-hydroxy group of **15a** or **15b** was protected with a dimethoxytrityl (DMTr) group, and the 5'-*O*-TBS group of the product was subsequently removed with TBAF to give **16a** or **16b**. Treatment of **16a** or **16b** under the conditions reported by Hata and co-workers with an *S,S'*-diphenylphosphorodithioate (PSS)/2,4,6-triisopropylbenzenesulfonyl chloride (TPSCl)/pyridine system⁸ gave the 5'-bis(phenylthio)phosphate **17a** or **17b**, respectively. The 5''-*O*-DMTr group of **17a** or **17b** was removed to give **18a** or **18b**, respectively. A phosphoryl group was introduced at the resulting 5''-primary hydroxyl of **18a** or **18b** by Yoshikawa's method with POCl₃/(EtO)₃PO,⁹ followed by treatment of the product with

H₃PO₂ and Et₃N¹⁰ in the presence of *N*-methylmaleimide (NMM)^{5c} in pyridine, to afford the corresponding 5'-phenylthiophosphate **19a** or **19b**, respectively, which was the substrate of the key intramolecular condensation reaction. When a solution of **19a** in pyridine was added slowly to a mixture of a large excess of AgNO₃ and Et₃N in the presence of MS 3A in pyridine at room temperature,^{5b,c} the intramolecular pyrophosphate linkage was successfully formed as the previously reported cases to give the desired cyclization product **20a** in 46% yield. The other substrate **19b** was similarly condensed and the cyclization product **19b** was obtained. Finally, the protecting groups of **19a** and **19b** were simultaneously removed by acidic treatment with aqueous HCO₂H furnished the target cADPcX (**4**) and 3''-OMe-cADPcX (**5**).

Ca²⁺-mobilizing Activity in Sea Urchin Egg Homogenate. The Ca²⁺-mobilizing ability of the newly synthesized compounds **4** and **5** was evaluated by the fluorometrically Ca²⁺-monitoring method with *H. pulcherrimus* sea urchin egg homogenate,^{6b,11} and the results were compared those of the natural second messenger cADPR (**1**) and the related carbocyclic analogues cADPcR (**2**) and 3''-deoxy-cADPcR (**3**). Both of the two newly synthesized compounds **4** and **5** released Ca²⁺ from the homogenate in a dose-dependent manner, as shown in **FIGURE 2**, where the maximal Ca²⁺-mobilizing activity was almost equal to that of cADPR. Thus, **4** and **5** were shown to be full agonists as cADPcR (**2**) and 3''-deoxy-cADPcR (**3**). cADPcX (**4**), the 3''-epimer of cADPcR, showed marked Ca²⁺-mobilizing activity with an EC₅₀ value of 69 nM, which was similar to that of cADPcR with an EC₅₀ value of 79 nM and was 3 times more potent than the natural ligand cADPR (EC₅₀ = 220 nM). The 3''-*O*-methyl cADPcX (**5**) showed an EC₅₀ value of 740 nM, which was about 10 times weaker than cADPcX. Similar to the case of cADPcR, 3''-methylation of the 3''-hydroxy group of cADPcX markedly reduced the activity.

CONCLUSION

We successfully synthesized cADPcX (**4**), the 3''-epimer of cADPcR (**2**), and the corresponding 3''-*O*-methyl analogue (3''-OMe-cADPcX, **5**) and identified cADPcX as one of the most potent cADPR-related compounds reported so far. Therefore, irrespective of the 3''-configuration, both cADPcX and cADPcR have strong Ca²⁺-mobilizing activity to suggest that the 3''-hydroxy group seems to be unnecessary in their binding with the target biomolecule, which is in accord with the previous results on the 3''-deoxy-cADPcR with remarkable activity.^{6c}

EXPERIMENTAL

General Methods. Chemical shifts are reported in ppm downfield from Me₄Si (¹H), MeCN (¹³C) or H₃PO₄ (³¹P). All of the ¹H NMR assignments described were in agreement with COSY spectra. Thin-layer chromatography was done on Merck coated

plate 60F₂₅₄. Silica gel chromatography was done on Merck silica gel 5715. Reactions were carried out under an argon atmosphere.

(1R,2S,3R,4R)-1-tert-Butoxycarbonylamino-2-(methoxymethoxy)-3-hydroxy-4-(triisopropylsilyloxymethyl)cyclopentane (10). A mixture of **9** (1.20 g, 4.12 mmol), TIPSCl (2.12 mL, 9.88 mmol), imidazole (1.01 g, 14.8 mmol) and DMAP (0.604 g, 4.94 mmol) in DMF (40 mL) was stirred at room temperature for 4 h. After addition of MeOH (10 mL), the resulting mixture was evaporated. The residue was partitioned between EtOAc and H₂O, and the organic layer was washed with brine, dried (Na₂SO₄), and evaporated. The residue was purified by column chromatography (SiO₂, 25% EtOAc in hexane) to give **10** (1.68 g, 91%) as a colorless oil: ¹H-NMR (CDCl₃, 500 MHz) δ 4.98 (brs, 1H, NH), 4.78 (d, 1H, MOM-CH₂, *J* = 6.4 Hz), 4.72 (d, 1H, MOM-CH₂, *J* = 6.4 Hz), 4.08 (m, 1H, H-3), 3.99 (m, 1H, H-1), 3.88 (dd, 1H, H-6a, *J*_{6a,6b} = 9.6 Hz, *J*_{6a,4} = 2.8 Hz), 3.80 (m, 1H, H-2), 3.69 (dd, 1H, H-6b, *J*_{6b,6a} = 9.6 Hz, *J*_{6b,4} = 3.2 Hz), 3.41 (s, 3H, MOM-CH₃), 2.66 (brs, 1H, OH), 2.37 (m, 1H, H-5a), 2.04 (m, 1H, H-4), 1.42 (s, 9H, *tert*-Bu), 1.33 (m, 1H, H-5b), 1.07-1.18 (m, 21H, TIPS); ¹³C-NMR (CDCl₃, 125 MHz) δ 155.10, 96.53, 84.21, 79.13, 72.64, 63.66, 55.73, 53.62, 45.40, 30.33, 28.36, 18.00, 12.05; HRMS (FAB, positive) calcd for C₂₂H₄₆NO₆Si 448.3094 (MH⁺), found 448.3109.

(1R,2S,3S,4R)-1-tert-Butoxycarbonylamino-2-(methoxymethoxy)-3-(chloroacethoxy)-4-(triisopropylsilyloxymethyl)cyclopentane (11). To a mixture of **10** (0.160 g, 0.357 mmol), chloroacetic acid (0.135 g, 1.43 mmol), and Ph₃P (0.374 g, 1.43 mmol) in toluene (2 mL), a solution of DIAD (281 μL, 1.43 mmol) in toluene (1.5 mL) was added slowly at 0 °C, and the mixture was stirred at room temperature for 6 h and then evaporated. The residue was purified by column chromatography (SiO₂, 10% EtOAc in hexane) to give **11** (0.139 g, 74%) as a colorless oil: ¹H-NMR (CDCl₃, 500 MHz) δ 5.21 (dd, 1H, H-3, *J*_{3,2} = 3.9 Hz, *J*_{3,4} = 6.0 Hz), 4.86 (m, 1H, NH), 4.72 (d, 1H, MOM-CH₂, *J* = 6.6 Hz), 4.69 (d, 1H, MOM-CH₂, *J* = 6.6 Hz), 4.05 (s, 2H, Cl-Ac), 3.97 (m, 1H, H-1), 3.92 (m, 1H, H-2), 3.70 (m, 2H, H-6 x 2), 3.36 (s, 3H, MOM-CH₃), 2.48 (m, 1H, H-4), 2.38 (m, 1H, H-5a), 1.43 (m, 10H, H-5b, *tert*-Bu), 1.09 (m, 21H, TIPS); ¹³C-NMR (CDCl₃, 125 MHz) δ 166.33, 155.16, 95.70, 85.43, 79.78, 73.06, 70.82, 61.80, 55.48, 54.77, 41.55, 40.78, 32.43, 28.35, 17.95, 11.89; HRMS (FAB, positive) calcd for C₂₄H₄₇ClNO₇Si 524.2810 (MH⁺), found 524.2816.

(1R,2S,3S,4R)-1-tert-Butoxycarbonylamino-2-(methoxymethoxy)-3-hydroxy-

4-(triisopropylsilyloxymethyl)cyclopentane (12). A mixture of **11** (0.173 g, 0.330 mmol) and NaOMe (1 M in MeOH, 50 μ L, 50 μ mol) in MeOH (3 mL) was stirred at room temperature for 30 min. After addition of aqueous saturated NH_4Cl (1 mL) at 0 $^\circ\text{C}$, the resulting mixture was extracted with EtOAc, and the organic layer was washed with H_2O and brine, dried (Na_2SO_4), and evaporated. The residue was purified by column chromatography (SiO_2 , 25% EtOAc in hexane) to give **12** (0.130 g, 88%) as a colorless oil: ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz) δ 6.69 (d, 1H, NH, J = 7.3 Hz), 4.70 (d, 1H, OH, J = 4.6 Hz), 4.63 (d, 1H, MOM- CH_2 , J = 6.4 Hz), 4.60 (d, 1H, MOM- CH_2 , J = 6.4 Hz), 3.84 (m, 2H, H-6a, H-3), 3.62 (m, 1H, H-1), 3.58 (m, 2H, H-2, H-6b), 3.22 (s, 3H, MOM- CH_3), 2.02 (m, 1H, H-4), 1.96 (m, 1H, H-5a), 1.36 (s, 10H, H-5b, *tert*-Bu), 1.03 (m, 21H, TIPS); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz) δ 156.04, 95.36, 88.30, 78.17, 74.92, 63.31, 55.46, 55.04, 43.62, 32.28, 28.56, 18.16, 11.67; HRMS (FAB, positive) calcd for $\text{C}_{22}\text{H}_{46}\text{NO}_6\text{Si}$ 448.3094 (MH^+), found 448.3092.

(1R,2S,3S,4R)-1-tert-Butoxycarbonylamino-2,3-bis(methoxymethyloxy)-4-hydroxy methylcyclopentane (13a). A mixture of **12** (0.871 g, 1.95 mmol), MOMCl (739 μ L, 9.73 mmol) and *i*-Pr₂NEt (3.39 mL, 19.5 mmol) in CH_2Cl_2 (20 mL) was stirred at room temperature for 75 h. After addition of MeOH (10 mL), the resulting mixture was evaporated. The residue was partitioned between EtOAc and 0.1 M HCl, and the organic layer was washed H_2O and brine, dried (Na_2SO_4), and evaporated. A mixture of the residue and TBAF (1.0 M in THF, 3.0 mL, 3.0 mmol) in THF (20 mL) was stirred at room temperature for 1 h and then evaporated. The residue was purified by column chromatography (SiO_2 , EtOAc) to give **13a** (0.634 g, 97%) as a yellow oil: ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz) δ 6.85 (d, 1H, NH, J = 7.9 Hz), 4.64 (d, 1H, MOM- CH_2 , J = 6.5 Hz), 4.59 (m, 3H, MOM- CH_2 x 3), 4.34 (m, 1H, OH), 3.84 (m, 1H, H-3), 3.80 (m, 1H, H-2), 3.66 (m, 1H, H-1), 3.50 (m, 1H, H-6a), 3.38 (m, 1H, H-6b), 3.26 (s, 3H, MOM- CH_3), 3.23 (s, 3H, MOM- CH_3), 2.06 (m, 1H, H-4), 1.93 (m, 1H, H-5a), 1.33 (s, 9H, *tert*-Bu), 1.32 (m, 1H, H-5b); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz) δ 155.15, 95.14, 94.64, 85.42, 80.11, 77.69, 60.05, 55.65, 54.98, 54.72, 42.69, 32.25, 28.39; FAB-MS m/z 336 (MH^+); Anal. Calcd for $\text{C}_{15}\text{H}_{29}\text{NO}_7$: C, 53.72; H, 8.72; N, 4.18. Found; C, 53.49; H, 8.57; N, 4.00.

(1R,2S,3S,4R)-1-tert-Butoxycarbonylamino-2-(methoxymethyloxy)-3-methoxy-4-hydroxymethylcyclopentane (13b). A mixture of **12** (0.513 g, 1.15 mmol), MeOTf (648 μ L, 5.73 mmol) and DTBMP (1.29 g, 6.30 mmol) in CH_2Cl_2 (11 mL) was stirred at

room temperature for 12 h. After addition of aqueous saturated NaHCO_3 (10 mL) at 0 °C, the mixture was extracted with EtOAc, and the organic layer was washed with H_2O and brine, dried (Na_2SO_4), and evaporated. The residue was purified by column chromatography (SiO_2 , 10% EtOAc in hexane) to give the 3-*O*-methylated product (0.259 g, 49%) as a colorless oil. A mixture of the oil and TBAF (1.0 M in THF, 0.84 mL, 0.84 mmol) in THF (5 mL) was stirred at room temperature for 2 h and then evaporated. The residue was purified by column chromatography (SiO_2 , 75% EtOAc in hexane) to give **13b** (0.154 g, 90%) as a yellow oil: ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz) δ 6.84 (d, 1H, NH, $J = 8.0$ Hz), 4.65 (d, 1H, MOM- CH_2 , $J = 6.5$ Hz), 4.58 (d, 1H, MOM- CH_2 , $J = 6.5$ Hz), 4.29 (m, 1H, OH), 3.77 (dd, 1H, H-2, $J_{2,1} = 5.1$ Hz, $J_{2,3} = 3.0$ Hz), 3.65 (m, 1H, H-1), 3.55 (m, 1H, H-6a), 3.47 (dd, 1H, H-3, $J_{3,2} = 3.0$ Hz, $J_{3,4} = 5.8$ Hz), 3.32 (m, 1H, H-6b), 3.24 (s, 6H, OMe, MOM- CH_3), 2.04 (m, 1H, H-4), 1.91 (m, 1H, H-5a), 1.36 (s, 9H, *tert*-Bu), 1.30 (m, 1H, H-5b); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz) δ 155.16, 94.70, 84.84, 84.45, 77.68, 59.93, 56.82, 55.69, 54.74, 42.65, 32.30, 28.39; FAB-MS m/z 306 (MH^+); Anal. Calcd for $\text{C}_{14}\text{H}_{27}\text{NO}_6$: C, 55.07; H, 8.91; N, 4.59. Found: C, 55.09; H, 8.76; N, 4.51.

(1*R*,2*S*,3*S*,4*R*)-1-Amino-2,3-bis(methoxymethoxy)-4-hydroxymethylcyclopentane (14a). A solution of **13a** (0.614 g, 1.83 mmol) in H_2O (18 mL) was stirred at 100 °C for 12 h and then evaporated. The residue was azeotroped with toluene (10 mL x 3) to give **14a** (0.430 g, quant.) as a brown oil: ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz) δ 4.67 (d, 1H, MOM- CH_2 , $J = 6.5$ Hz), 4.59 (m, 3H, MOM- CH_2 x 3), 3.85 (dd, 1H, H-3, $J_{3,2} = 3.5$ Hz, $J_{3,4} = 6.0$ Hz), 3.57 (m, 1H, H-2), 3.50 (dd, 1H, H-6a, $J_{6a,6b} = 10.3$ Hz, $J_{6a,4} = 7.0$ Hz), 3.37 (dd, 1H, H-6b, $J_{6b,6a} = 10.3$ Hz, $J_{6b,4} = 6.5$ Hz), 3.26 (s, 6H, MOM- CH_3 x 2), 2.96 (m, 1H, H-1), 2.06 (m, 1H, H-4), 1.92 (m, 1H, H-5a), 1.15 (s, 1H, H-5b); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz) δ 95.24, 95.18, 89.75, 80.81, 60.41, 56.84, 54.98, 54.86, 42.57, 35.50; HRMS (FAB, positive) calcd for $\text{C}_{10}\text{H}_{22}\text{NO}_5$ 236.1498 (MH^+), found 236.1508.

(1*R*,2*S*,3*S*,4*R*)-1-Amino-2-(methoxymethoxy)-3-methoxy-4-hydroxymethylcyclopentane (14b). Compound **14b** (0.239 g, quant.) was obtained from **13b** (0.351 g, 1.15 mmol) as described for the synthesis of **14a**: ^1H -NMR ($\text{DMSO}-d_6$, 500 MHz) δ 4.68 (d, 1H, MOM- CH_2 , $J = 6.6$ Hz), 4.62 (d, 1H, MOM- CH_2 , $J = 6.6$ Hz), 3.53 (m, 1H, H-2), 3.49 (m, 2H, H-3, H-6a), 3.33 (m, 1H, H-6b), 3.27 (s, 3H, MOM- CH_3 or OMe), 3.25 (s, 3H, MOM- CH_3 or OMe), 2.95 (m, 1H, H-1), 2.06 (m, 1H, H-4), 1.91 (m, 1H, H-5a), 1.14 (s, 1H, H-5b); ^{13}C -NMR ($\text{DMSO}-d_6$, 125 MHz) δ 95.30, 89.03, 85.62, 60.28,

56.96, 56.91, 54.87, 42.50, 35.44; HRMS (FAB, positive) calcd for C₉H₂₀NO₄ 206.1392 (MH⁺), found 206.1390.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2,3-Bis(methoxymethoxy)-4-(hydroxymethyl)cyclopentyl]-5'-*O*-(*tert*-butyldimethylsilyl)-2',3'-*O*-isopropylideneadenosine (**15a**).** A mixture of **6** (0.765 g, 1.75 mmol), **14a** (0.417 g, 1.77 mmol), and K₂CO₃ (12 mg, 88 μmol) in MeOH (18 mL) was stirred at room temperature for 12 h and then evaporated. The residue was partitioned between EtOAc and H₂O, and the organic layer was washed with brine, dried (Na₂SO₄), and evaporated. The residue was purified by column chromatography (SiO₂, 25% MeOH in EtOAc) to give **15a** (0.930 g, 83%) as a white foam: ¹H-NMR (CDCl₃, 500 MHz) δ 7.91 (s, 1H, H-2 or H-8), 7.82 (s, 1H, H-2 or H-8), 6.02 (d, 1H, H-1', *J*_{1',2'} = 2.7 Hz), 5.37 (m, 1H, H-1''), 5.09 (dd, 1H, H-2', *J*_{2',1'} = 2.7 Hz, *J*_{2',3'} = 6.1 Hz), 4.89 (dd, 1H, H-3', *J*_{3',2'} = 6.1 Hz, *J*_{3',4'} = 2.4 Hz), 4.81 (d, 1H, MOM-CH₂, *J* = 6.6 Hz), 4.76 (d, 1H, MOM-CH₂, *J* = 6.6 Hz), 4.69 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.65 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.48 (m, 1H, H-2''), 4.39 (ddd, 1H, H-4', *J*_{4',3'} = 2.4 Hz, *J*_{4',5'a} = 6.4 Hz, *J*_{4',5'b} = 3.7 Hz), 4.19 (dd, 1H, H-3'', *J*_{3'',2''} = 3.5 Hz, *J*_{3'',4''} = 5.8 Hz), 3.83 (m, 2H, H-5'a, H-5''a), 3.77 (m, 2H, H-5'b, H-5''b), 3.45 (s, 3H, MOM-CH₃), 3.21 (s, 3H, MOM-CH₃), 2.47 (m, 1H, H-4''), 2.39 (m, 1H, H-6'a), 2.02 (m, 1H, H-6'b), 1.61, 1.39 (each s, each 3H, isopropylidene), 0.86 (s, 9H, *tert*-Bu), 0.046, 0.035 (each s, each 3H, TBS-Me x 2); ¹³C-NMR (CDCl₃, 125 MHz) δ 154.45, 146.66, 140.62, 136.78, 123.40, 114.16, 96.39, 95.63, 91.26, 87.03, 85.32, 84.38, 81.76, 81.33, 63.46, 61.56, 60.08, 56.02, 55.49, 42.10, 31.22, 27.23, 25.86, 25.38, 18.32, -5.43, -5.54; FAB-MS *m/z* 640 (MH⁺); UV (MeOH) λ_{max} = 261, 295 (sh) nm; Anal. Calcd for C₂₉H₄₉N₅O₉Si: C, 54.44; H, 7.72; N, 10.95. Found; C, 54.24; H, 7.66; N, 10.84.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2,3-Bis(methoxymethoxy)-4-(dimethoxytrityloxymethyl)cyclopentyl]-2',3'-*O*-isopropylideneadenosine (**16a**).** A mixture of **15a** (0.884 g, 1.38 mmol) and DMTrCl (1.40 g, 4.14 mmol) in pyridine (15 mL) was stirred at room temperature for 10 min. After addition of MeOH (10 mL), the resulting mixture was evaporated. The residue was partitioned between EtOAc and H₂O, and the organic layer was washed with brine, dried (Na₂SO₄), and evaporated. A mixture of the residue, TBAF (1.0 M in THF, 2.76 mL, 2.76 mmol) and AcOH (79 μL, 1.38 mmol) in THF (10 mL) was stirred at room temperature for 3 h and then evaporated. The residue was purified

by column chromatography (SiO₂, 25% MeOH in EtOAc) to give **16a** (1.19 g, quant.) as a white foam: ¹H-NMR (CDCl₃, 500 MHz) δ 8.00 (s, 1H, H-2 or H-8), 7.62 (s, 1H, H-2 or H-8), 6.82-7.43 (m, 13H, DMTr), 5.79 (m, 2H, H-1', H-1''), 5.02 (m, 2H, H-2', H-3'), 4.81 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.67 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.61 (d, 1H, MOM-CH₂, *J* = 6.7 Hz), 4.54 (d, 1H, MOM-CH₂, *J* = 6.7 Hz), 4.47 (m, 1H, H-4'), 4.20 (m, 1H, H-3''), 4.12 (m, 1H, H-2''), 3.91 (m, 1H, H-5'a), 3.80 (s, 6H, DMTr-OMe x 2), 3.72 (m, 1H, H-5'b), 3.33 (s, 3H, MOM-CH₃), 3.29 (m, 1H, H-5''a), 3.25 (s, 3H, MOM-CH₃), 3.13 (m, 1H, H-5''b), 2.68 (m, 1H, H-4''), 2.56 (m, 1H, H-6''a), 1.62, 1.35 (each s, each 3H, isopropylidene), 1.49 (m, 1H, H-6''b); ¹³C-NMR (CDCl₃, 125 MHz) δ 158.45, 146.70, 144.96, 137.95, 136.31, 136.11, 129.97, 129.93, 128.12, 127.79, 126.76, 114.20, 113.09, 113.06, 96.41, 94.93, 93.90, 86.00, 85.82, 83.61, 81.37, 63.19, 61.56, 60.37, 58.81, 55.98, 55.60, 55.20, 42.94, 35.23, 27.55, 25.18, 21.02, 14.19; HRMS (FAB, positive) calcd for C₄₄H₅₄N₅O₁₁ 828.3820 (MH⁺), found 828.3818; UV (MeOH) λ_{max} = 260, 295 (sh) nm.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2,3-Bis(methoxymethoxy)-4-(dimethoxytrityloxymethyl)cyclopentyl]-5'-*O*-[bis(phenylthio)phospholyl]-2',3'-*O*-isopropylideneadenosine (**17a**). After stirring a mixture of PSS (1.56 g, 4.09 mmol) and TPSCl (1.12 g, 3.69 mmol) in pyridine (14 mL) at room temperature for 1 h, **16a** (1.13 g, 1.36 mmol) was added, and the resulting mixture was stirred at room temperature for further 2 h and then evaporated. The residue was partitioned between EtOAc and H₂O, and the organic layer was washed with brine, dried (Na₂SO₄), and evaporated. The residue was purified by column chromatography (SiO₂, 2% MeOH in CHCl₃) to give **17a** (0.890 g, 65%) as a white foam: ¹H-NMR (CDCl₃, 500 MHz) δ 7.95 (s, 1H, H-2 or H-8), 7.64 (s, 1H, H-2 or H-8), 6.81-7.51 (m, 23H, DMTr, SPh x 2), 5.96 (d, 1H, H-1', *J*_{1,2'} = 2.6 Hz), 5.76 (m, 1H, H-1''), 5.07 (dd, 1H, H-2', *J*_{2',1'} = 2.6 Hz, *J*_{2',3'} = 6.3 Hz), 4.85 (dd, 1H, H-3', *J*_{3',2'} = 6.3 Hz, *J*_{3',4'} = 2.7 Hz), 4.76 (d, 1H, MOM-CH₂, *J* = 6.8 Hz), 4.64 (d, 1H, MOM-CH₂, *J* = 6.8 Hz), 4.57 (d, 1H, MOM-CH₂, *J* = 6.7 Hz), 4.53 (d, 1H, MOM-CH₂, *J* = 6.7 Hz), 4.41 (m, 1H, H-4'), 4.38 (m, 2H, H-5' x 2), 4.20 (d, 1H, H-3'', *J*_{3'',4''} = 4.0 Hz), 4.14 (d, 1H, H-2'', *J*_{2'',1''} = 2.6 Hz), 3.79 (s, 6H, DMTr-OMe x 2), 3.30 (m, 1H, H-5''a), 3.27 (s, 3H, MOM-CH₃), 3.19 (s, 3H, MOM-CH₃), 3.16 (m, 1H, H-5''b), 2.61 (m, 1H, H-4''), 2.54 (m, 1H, H-6''a), 1.59, 1.34 (each s, each 3H, isopropylidene), 1.54 (m, 1H, H-6''b); ¹³C-NMR (CDCl₃, 125 MHz)**

δ 158.43, 154.52, 146.61, 144.99, 140.49, 136.94, 136.35, 136.17, 135.34, 135.29, 135.21, 135.17, 129.98, 129.96, 129.66, 129.63, 129.46, 129.44, 128.13, 127.76, 126.72, 125.93, 125.87, 125.79, 125.74, 123.52, 114.69, 113.05, 113.02, 95.88, 94.99, 90.61, 86.00, 85.87, 84.67, 84.61, 84.45, 81.06, 80.78, 66.42, 66.36, 61.80, 58.76, 55.86, 55.51, 55.19, 42.79, 35.17, 27.13, 25.30, 21.02, 14.19; ^{31}P -NMR (CDCl_3 , 202 MHz) δ 50.74 (s); HRMS (FAB, positive) calcd for $\text{C}_{56}\text{H}_{63}\text{N}_5\text{O}_{12}\text{PS}_2$ 1092.3652 (MH^+), found 1092.3660; UV (MeOH) $\lambda_{\text{max}} = 295$ (sh) nm.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2,3-Bis(methoxymethyloxy)-4-(hydroxymethyl)cyclopentyl]-5'-*O*-[bis(phenylthio)phospholyl]-2',3'-*O*-isopropylideneadenosine (18a).** A solution of **17a** (0.916 g, 0.839 mmol) in aqueous 60% AcOH (8 mL) was stirred at room temperature for 4 h. After addition of aqueous saturated NaHCO_3 (60 mL) at 0 °C, the mixture was extracted with EtOAc, and the organic layer was washed with aqueous saturated NaHCO_3 , H_2O and brine, dried (Na_2SO_4), and evaporated. The residue was purified by column chromatography (SiO_2 , 10% MeOH in CHCl_3) to give **18a** (0.551 g, 83%) as a white foam: ^1H -NMR (CDCl_3 , 500 MHz) δ 7.93 (s, 1H, H-2 or H-8), 7.69 (s, 1H, H-2 or H-8), 7.30-7.52 (m, 10H, SPh x 2), 5.99 (m, 1H, H-1'), 5.32 (m, 1H, H-1''), 5.11 (m, 1H, H-2'), 4.90 (dd, 1H, H-3', $J_{3',2'} = 6.2$ Hz, $J_{3',4'} = 2.5$ Hz), 4.76 (d, 1H, MOM- CH_2 , $J = 6.5$ Hz), 4.72 (d, 1H, MOM- CH_2 , $J = 6.5$ Hz), 4.66 (d, 1H, MOM- CH_2 , $J = 6.7$ Hz), 4.62 (d, 1H, MOM- CH_2 , $J = 6.7$ Hz), 4.45 (m, 1H, H-2''), 4.43 (m, 2H, H-4', H-5'a), 4.38 (m, 1H, H-5'b), 4.18 (m, 1H, H-3''), 3.84 (dd, 1H, H-5''a, $J_{5''a,5''b} = 11.2$ Hz, $J_{5''a,4''} = 3.8$ Hz), 3.76 (dd, 1H, H-5''b, $J_{5''b,5''a} = 11.2$ Hz, $J_{5''b,4''} = 5.9$ Hz), 3.41 (s, 3H, MOM- CH_3), 3.19 (s, 3H, MOM- CH_3), 2.47 (m, 1H, H-4''), 2.40 (m, 1H, H-6''a), 2.04 (m, 1H, H-6''b), 1.60, 1.36 (each s, each 3H, isopropylidene); ^{13}C -NMR (CDCl_3 , 125 MHz) δ 154.27, 146.89, 137.38, 135.31, 135.27, 135.17, 135.13, 129.69, 129.66, 129.64, 129.62, 129.44, 129.42, 125.87, 125.82, 125.77, 125.72, 114.66, 96.35, 95.77, 90.74, 84.74, 84.68, 84.43, 81.64, 81.12, 66.39, 66.32, 61.49, 55.95, 55.50, 42.02, 31.10, 27.11, 25.29; ^{31}P -NMR (CDCl_3 , 202 MHz) δ 51.10 (s); FAB-MS m/z 790 (MH^+); UV (MeOH) $\lambda_{\text{max}} = 259, 295$ (sh) nm; Anal. Calcd for $\text{C}_{35}\text{H}_{44}\text{N}_5\text{O}_{10}\text{PS}_2$: C, 53.22; H, 5.61; N, 8.87. Found: C, 53.20; H, 5.66; N, 8.68.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2,3-Bis(methoxymethyloxy)-4-(phosphonoxymethyl)cyclopentyl]-5'-*O*-[(phenylthio)phospholyl]-2',3'-*O*-isopropylideneadenosine (19a).** A mixture of POCl_3 (93 μL , 1.0 mmol) and **18a** (79 mg, 0.10 mmol) in $\text{PO}(\text{OEt})_3$ (1.0 mL)

was stirred at 0 °C for 1 h. After addition of aqueous saturated NaHCO₃ (3.0 mL), the resulting mixture was stirred at 0 °C for 10 min. To the mixture was added triethylammonium acetate (TEAA, 2.0 M, pH 7.0, 1.0 mL) buffer and H₂O (5.0 mL), and the resulting solution was applied to a C₁₈ reversed phase column (1.1 × 14 cm). The column was developed using a linear gradient of 0–65% MeCN in TEAA buffer (0.1 M, pH 7.0, 400 mL). Appropriate fractions were evaporated, and excess TEAA was removed by C₁₈ reversed phase column chromatography (1.1 × 17 cm, eluted with 70% aqueous MeCN). Appropriate fractions were evaporated, and the residue was co-evaporated with pyridine (2.0 mL × 3). A mixture of the residue, NMM (77 mg, 0.69 mmol), H₃PO₂ (70 µL, 1.39 mmol), and Et₃N (97 µL, 0.69 mmol) was stirred at 0 °C for 4 h under shading. After addition of TEAA buffer (1.0 M, pH 7.0, 2.0 mL), the resulting mixture was evaporated. The residue was partitioned between EtOAc and H₂O, and the aqueous layer was evaporated. A solution of the residue in H₂O (5.0 mL) was applied to a C₁₈ reversed phase column (1.1 × 16 cm), and the column was developed using a linear gradient of 0–40% MeCN in TEAA buffer (0.1 M, pH 7.0, 400 mL). Appropriate fractions were evaporated, and excess TEAA was removed by C₁₈ reversed phase column chromatography (1.1 × 17 cm, eluted with 50% aqueous MeCN). Appropriate fractions were evaporated, and the residue was lyophilized to give **19a** (41 mg, 46%) as a triethylammonium salt: ¹H-NMR (D₂O, 500 MHz) δ 8.71 (s, 1H, H-2 or H-8), 8.40 (s, 1H, H-2 or H-8), 7.01–7.30 (m, 5H, SPh), 6.31 (d, 1H, H-1', *J*_{1',2'} = 2.1 Hz), 5.33 (dd, 1H, H-2', *J*_{2',1'} = 2.1 Hz, *J*_{2',3'} = 6.0 Hz), 5.00 (m, 1H, H-1''), 4.94 (dd, 1H, H-3', *J*_{3',2'} = 6.0 Hz, *J*_{3',4'} = 1.2 Hz), 4.86 (s, 2H, MOM-CH₂), 4.72 (d, 1H, MOM-CH₂, *J* = 7.3 Hz), 4.70 (m, 1H, H-4'), 4.68 (d, 1H, MOM-CH₂, *J* = 7.3 Hz), 4.51 (m, 1H, H-2''), 4.33 (m, 1H, H-3''), 4.15 (m, 2H, H-5' × 2), 4.07 (m, 2H, H-5'' × 2), 3.48 (s, 3H, MOM-CH₃), 3.27 (s, 3H, MOM-CH₃), 3.18 (q, 6H, Et₃NH-CH₂ × 3, *J* = 7.3 Hz), 2.74 (m, 1H, H-4''), 2.68 (m, 1H, H-6''a), 2.08 (m, 1H, H-6''b), 1.62, 1.39 (each s, each 3H, isopropylidene), 1.26 (t, 9H, Et₃NH-CH₃ × 3, *J* = 7.3 Hz); ¹³C-NMR (D₂O, 125 MHz) δ 151.02, 146.31, 146.10, 143.91, 133.21, 133.17, 129.90, 129.85, 129.44, 128.28, 119.61, 115.17, 97.10, 96.75, 91.87, 87.05, 86.67, 86.59, 84.64, 81.91, 80.44, 66.27, 63.81, 56.40, 56.14, 47.12, 40.75, 40.68, 26.38, 24.73, 8.69; ³¹P-NMR (D₂O, 202 MHz) δ 17.10 (s), 0.84 (s); HRMS (FAB, positive) calcd for C₂₉H₄₂N₅O₁₄P₂S 778.1919 (MH⁺), found 778.1934; UV (H₂O) λ_{max} 260 nm.

2'',3''-Bis-*O*-Methoxymethyl-cyclic ADP-carbocyclic-xylose 2',3'-Acetonide

(20a). To a mixture of AgNO₃ (310 mg, 1.83 mmol), Et₃N (255 µL, 1.83 mmol), and MS 3A (400 mg) in pyridine (70 mL), a solution of **19a** (76 mg, 0.087 mmol) in pyridine (70 mL) was added slowly over 15 h, using a syringe-pump, at room temperature under shading. The MS 3A was filtered off with Celite and washed with H₂O. To the combined filtrate and washings was added TEAA buffer (2.0 M, pH 7.0, 2 mL), and the resulting solution was evaporated. The residue was partitioned between EtOAc and H₂O, and the aqueous layer was evaporated. A solution of the residue in H₂O (5.0 mL) was applied to a C₁₈ reverse phase column (1.1 × 17 cm), and the column was developed using a linear gradient of 0–25% MeCN in TEAA buffer (0.1 M, pH 7.0, 400 mL). Appropriate fractions were evaporated, and excess TEAA was removed by C₁₈ reverse phase column chromatography (1.1 × 17 cm, eluted with 40% aqueous MeCN). Appropriate fractions were evaporated, and the residue was lyophilized to give **20a** (40 mg, 46%) as a triethylammonium salt: ¹H-NMR (D₂O, 500 MHz) δ 9.20 (s, 1H, H-2 or H-8), 8.44 (s, 1H, H-2 or H-8), 6.41 (d, 1H, H-1', *J*_{1',2'} = 1.6 Hz), 5.53 (dd, 1H, H-2', *J*_{2',1'} = 1.6 Hz, *J*_{2',3'} = 6.1 Hz), 5.46 (dd, 1H, H-3', *J*_{3',2'} = 6.1 Hz, *J*_{3',4'} = 2.5 Hz), 5.04 (m, 1H, H-1''), 4.80 (s, 2H, MOM-CH₂), 4.76 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.67 (d, 1H, MOM-CH₂, *J* = 6.9 Hz), 4.63 (m, 1H, H-4'), 4.62 (m, 1H, H-2''), 4.37 (m, 1H, H-3''), 4.23 (m, 1H, H-5''a), 4.16 (m, 1H, H-5'a), 4.09 (m, 2H, H-5'b, H-5''b), 3.42 (s, 3H, MOM-CH₃), 3.11 (s, 3H, MOM-CH₃), 3.19 (q, 6H, Et₃NH-CH₂ × 3, *J* = 7.3 Hz), 2.99 (m, 1H, H-6''a), 2.76 (m, 1H, H-4''), 2.36 (m, 1H, H-6''b), 1.64, 1.44 (each s, each 3H, isopropylidene), 1.27 (t, 9H, Et₃NH-CH₃ × 3, *J* = 7.3 Hz); ¹³C-NMR (D₂O, 125 MHz) δ 151.93, 146.22, 146.28, 144.80, 119.64, 115.13, 97.69, 96.82, 92.47, 88.26, 86.92, 86.82, 85.13, 81.87, 81.29, 64.82, 64.61, 60.93, 56.25, 47.10, 37.29, 37.21, 28.70, 26.43, 24.74, 8.67; ³¹P-NMR (D₂O, 202 MHz) δ -9.58 (d, *J* = 11.2 Hz), -10.56 (d, *J* = 11.2 Hz); HRMS (FAB, positive) calcd for C₂₃H₃₆N₅O₁₄P₂ 668.1729 (MH⁺), found 668.1730; UV (H₂O) λ_{max} 259 nm.

Cyclic ADP-carbocyclic-xylose (4). A solution of **20a** (31 mg, 0.040 mmol) in aqueous 80% HCO₂H (1 mL) was stirred at room temperature for 48 h and then evaporated. After evaporation of the residue in H₂O and *i*-PrOH, aqueous 28% NH₃ (1 mL) was added to the residue, and the mixture was stirred at room temperature for 90 min. After evaporation of the residue in H₂O and *i*-PrOH, the resulting residue was dissolved in TEAB buffer (0.1 M, pH 7.0, 600 µL), and the solution was lyophilized to give **4** (21 mg, 90%) as a triethylammonium salt: ¹H-NMR (D₂O, 500 MHz, K⁺ salt) δ

9.11 (s, 1H, H-2 or H-8), 8.42 (s, 1H, H-2 or H-8), 6.11 (d, 1H, H-1', $J_{1',2'} = 6.0$ Hz), 5.20 (dd, 1H, H-2', $J_{2',1'} = 6.0$ Hz, $J_{2',3'} = 4.8$ Hz), 4.95 (m, 1H, H-1''), 4.64 (dd, 1H, H-3', $J_{3',2'} = 4.8$ Hz, $J_{3',4'} = 2.6$ Hz), 4.62 (m, 1H, H-5'a), 4.51 (m, 1H, H-2''), 4.45 (m, 1H, H-4'), 4.32 (m, 1H, H-3''), 4.23 (m, 1H, H-5''a), 4.14 (m, 2H, H-5'b, H-5''b), 2.95 (m, 1H, H-6''a), 2.65 (m, 1H, H-4''), 2.46 (m, 1H, H-6''b); ^{13}C -NMR (D_2O , 125 MHz) δ 152.20, 146.70, 145.43, 144.83, 120.40, 91.07, 85.29, 82.52, 76.53, 73.89, 71.16, 65.13, 64.20, 63.12, 39.52, 28.02; ^{31}P -NMR (D_2O , 202 MHz) δ -9.41 (d, $J = 11.4$ Hz), -10.35 (d, $J = 11.4$ Hz); HRMS (FAB, positive) calcd for $\text{C}_{16}\text{H}_{24}\text{N}_5\text{O}_{12}\text{P}_2$ 540.0891 (MH^+), found 540.0875; UV (H_2O) λ_{max} 259 nm.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2-(Methoxymethoxy)-3-methoxy-4-(hydroxymethyl)cyclopentyl]-5'-*O*-(*tert*-butyldimethylsilyl)-2',3'-*O*-isopropylideneadenosine (15b).** Compound **15b** (0.565 g, 84%) was obtained from **6** (0.480 g, 1.10 mmol) and **14b** (0.227 g, 1.11 mmol) as described for the synthesis of **15a**: ^1H -NMR (CDCl_3 , 500 MHz) δ 7.91 (s, 1H, H-2 or H-8), 7.83 (s, 1H, H-2 or H-8), 6.02 (d, 1H, H-1', $J_{1',2'} = 2.6$ Hz), 5.54 (m, 1H, H-1''), 5.11 (dd, 1H, H-2', $J_{2',1'} = 2.6$ Hz, $J_{2',3'} = 6.1$ Hz), 4.89 (dd, 1H, H-3', $J_{3',2'} = 6.1$ Hz, $J_{3',4'} = 2.5$ Hz), 4.74 (d, 1H, MOM-CH₂, $J = 6.8$ Hz), 4.65 (d, 1H, MOM-CH₂, $J = 6.8$ Hz), 4.39 (ddd, 1H, H-4', $J_{4',3'} = 2.5$ Hz, $J_{4',5'a} = 3.7$ Hz, $J_{4',5'b} = 6.3$ Hz), 4.35 (m, 1H, H-2''), 3.90 (dd, 1H, H-5''a, $J_{5''a,4'} = 3.0$ Hz, $J_{5''a,5''b} = 11.3$ Hz), 3.84 (dd, 1H, H-5'a, $J_{5'a,4'} = 3.8$ Hz, $J_{5'a,5''b} = 11.2$ Hz), 3.79 (m, 2H, H-3'', H-5'b), 3.76 (m, 1H, H-5''b), 3.50 (s, 3H, MOM-CH₃ or OMe), 3.25 (s, 3H, MOM-CH₃ or OMe), 2.42 (m, 2H, H-4'', H-6''a), 2.08 (m, 1H, H-6''b), 1.61, 1.39 (each s, each 3H, isopropylidene), 0.86 (s, 9H, *tert*-Bu), 0.048, 0.037 (each s, each 3H, TBS-Me x 2); ^{13}C -NMR (CDCl_3 , 125 MHz) δ 154.48, 146.35, 140.70, 136.89, 123.26, 114.17, 95.33, 91.27, 87.04, 86.99, 85.28, 83.40, 81.34, 63.45, 61.54, 59.37, 57.70, 55.51, 42.02, 31.65, 27.22, 25.86, 25.37, 18.32, -5.43, -5.54; FAB-MS m/z 610 (MH^+); Anal. Calcd for $\text{C}_{28}\text{H}_{47}\text{N}_5\text{O}_8\text{Si}$: C, 55.15; H, 7.77; N, 11.48. Found; C, 54.95; H, 7.59; N, 11.26.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2-(Methoxymethoxy)-3-methoxy-4-(dimethoxytrityloxy)methyl)cyclopentyl]-2',3'-*O*-isopropylideneadenosine (16b).** Compound **16b** (0.697 g, quant.) was obtained from **15b** (0.527 g, 0.864 mmol) as described for the synthesis of **16a**: ^1H -NMR (CDCl_3 , 500 MHz) δ 7.92 (brs, 1H, H-2 or H-8), 7.62 (brs, 1H, H-2 or H-8), 6.82-7.45 (m, 13H, DMTr), 5.78 (m, 2H, H-1', H-1''), 5.02 (m, 2H, H-2', H-3'), 4.82 (d, 1H, MOM-CH₂, $J = 6.7$ Hz), 4.68 (d, 1H, MOM-CH₂, $J = 6.7$ Hz), 4.49 (m, 1H,

H-4'), 4.11 (m, 1H, H-2''), 3.91 (m, 1H, H-3''), 3.80 (s, 7H, H-5'a, DMTr-OMe x 2), 3.75 (m, 1H, H-5'b), 3.37 (s, 3H, MOM-CH₃ or OMe), 3.35 (s, 3H, MOM-CH₃ or OMe), 3.25 (m, 1H, H-5''a), 3.19 (m, 1H, H-5''b), 2.56 (m, 1H, H-4''), 2.48 (m, 1H, H-6''a), 1.62, 1.36 (each s, each 3H, isopropylidene), 1.46 (m, 1H, H-6''b); ¹³C-NMR (CDCl₃, 125 MHz) δ 180.59, 165.74, 163.74, 152.56, 152.49, 144.47, 142.10, 141.56, 140.21, 136.11, 124.30, 122.87, 122.84, 99.99, 98.40, 88.24, 87.29, 86.38, 85.52, 82.67, 59.34, 56.69, 54.38, 51.96, 49.88, 49.24, 33.87, 23.65, 18.31, 14.03, 11.04, 9.88, 4.19, -3.60, -21.04; HRMS (FAB, positive) calcd for C₄₃H₅₂N₅O₁₀ 798.3714 (MH⁺), found 798.3721.

N-1-[(1R,2S,3S,4R)-2-(Methoxymethoxy)-3-methoxy-4-(dimethoxytrityloxy methyl)cyclopentyl]-5'-O-[bis(phenylthio)phospholyl]-2',3'-O-isopropylideneadenosine (17b). Compound **17b** (0.539 g, 62%) was obtained from **16b** (0.659 g, 0.826 mmol) as described for the synthesis of **17a**: ¹H-NMR (CDCl₃, 500 MHz) δ 7.91 (s, 1H, H-2 or H-8), 7.64 (s, 1H, H-2 or H-8), 6.81-7.51 (m, 23H, DMTr, SPh x 2), 5.96 (d, 1H, H-1', *J*_{1',2'} = 2.6 Hz), 5.76 (m, 1H, H-1''), 5.08 (dd, 1H, H-2', *J*_{2',1'} = 2.6 Hz, *J*_{2',3'} = 6.3 Hz), 4.86 (dd, 1H, H-3', *J*_{3',2'} = 6.3 Hz, *J*_{3',4'} = 2.7 Hz), 4.79 (d, 1H, MOM-CH₂, *J* = 6.8 Hz), 4.63 (d, 1H, MOM-CH₂, *J* = 6.8 Hz), 4.41 (m, 3H, H-4', H-5' x 2), 4.11 (m, 1H, H-2''), 3.79 (s, 6H, DMTr-OMe x 2), 3.78 (m, 1H, H-3''), 3.33 (s, 3H, MOM-CH₃ or OMe), 3.30 (s, 3H, MOM-CH₃ or OMe), 3.25 (m, 1H, H-5''a), 3.18 (m, 1H, H-5''b), 2.54 (m, 1H, H-4''), 2.47 (m, 1H, H-6''a), 1.59, 1.35 (each s, each 3H, isopropylidene), 1.48 (m, 1H, H-6''b); ¹³C-NMR (CDCl₃, 125 MHz) δ 158.38, 154.52, 146.79, 145.20, 140.52, 136.97, 136.42, 136.38, 135.34, 135.29, 135.22, 135.18, 130.04, 130.02, 129.65, 129.63, 129.43, 128.15, 127.69, 126.63, 125.94, 125.89, 125.82, 125.77, 123.51, 114.69, 113.00, 112.98, 95.09, 90.65, 85.79, 85.23, 84.71, 84.64, 84.45, 81.04, 66.45, 66.38, 61.23, 58.69, 57.39, 55.56, 55.19, 43.12, 35.07, 27.13, 25.30, 11.44; ³¹P-NMR (CDCl₃, 202 MHz) δ 48.14 (s); FAB-MS *m/z* 1062 (MH⁺); UV (MeOH) λ_{max} = 295 (sh) nm; Anal. Calcd for C₅₅H₆₀N₅O₁₁PS₂: C, 62.19; H, 5.69; N, 6.59. Found; C, 62.34; H, 5.83; N, 6.34.

N-1-[(1R,2S,3S,4R)-2-(Methoxymethoxy)-3-methoxy-4-(hydroxymethyl)cyclopentyl]-5'-O-[bis(phenylthio)phospholyl]-2',3'-O-isopropylideneadenosine (18b). Compound **18b** (0.312 g, 88%) was obtained from **17b** (0.496 g, 0.467 mmol) as described for the synthesis of **18a**: ¹H-NMR (CDCl₃, 500 MHz) δ 7.92 (s, 1H, H-2 or H-8), 7.69 (s, 1H, H-2 or H-8), 7.31-7.52 (m, 10H, SPh x 2), 5.99 (d, 1H, H-1', *J*_{1',2'} =

2.3 Hz), 5.48 (m, 1H, H-1''), 5.12 (dd, 1H, H-2', $J_{2',1'} = 2.3$ Hz, $J_{2',3'} = 6.2$ Hz), 4.91 (dd, 1H, H-3', $J_{3',2'} = 6.2$ Hz, $J_{3',4'} = 2.4$ Hz), 4.71 (d, 1H, MOM-CH₂, $J = 6.7$ Hz), 4.62 (d, 1H, MOM-CH₂, $J = 6.5$ Hz), 4.43 (m, 2H, H-4', H-5'a), 4.39 (m, 2H, H-2'', H-5'b), 3.90 (dd, 1H, H-5''a, $J_{5''a,5''b} = 11.4$ Hz, $J_{5''a,4''} = 2.8$ Hz), 3.79 (m, 1H, H-3''), 3.76 (dd, 1H, H-5''b, $J_{5''b,5''a} = 11.4$ Hz, $J_{5''b,4''} = 4.7$ Hz), 3.46 (s, 3H, MOM-CH₃ or OMe), 3.21 (s, 3H, MOM-CH₃ or OMe), 2.41 (m, 2H, H-4'', H-6''a), 2.06 (m, 1H, H-6''b), 1.60, 1.36 (each s, each 3H, isopropylidene); ¹³C-NMR (CDCl₃, 125 MHz) δ 154.37, 146.65, 140.61, 137.38, 135.34, 135.30, 135.21, 135.17, 129.70, 129.68, 129.66, 129.63, 129.44, 125.92, 125.86, 125.81, 125.76, 123.57, 114.70, 95.54, 90.76, 86.98, 84.79, 84.73, 84.47, 83.46, 81.14, 66.44, 66.38, 61.55, 60.36, 59.69, 57.74, 55.55, 41.90, 31.44, 27.13, 25.30, 21.01, 14.18; ³¹P-NMR (CDCl₃, 202 MHz) δ 51.05 (s); FAB-MS m/z 760 (MH⁺); UV (MeOH) $\lambda_{\max} = 259, 295$ (sh) nm; Anal. Calcd for C₃₄H₄₂N₅O₉PS₂: C, 53.75; H, 5.57; N, 9.22. Found; C, 53.59; H, 5.60; N, 8.97.

***N*-1-[(1*R*,2*S*,3*S*,4*R*)-2-(Methoxymethyloxy)-3-methoxy-4-(phosphonoxymethyl)cyclopentyl]-5'-*O*-[(phenylthio)phosphoryl]-2',3'-*O*-isopropylideneadenosine (19b).** Compound **19b** (90 mg, 58%) was obtained from **18b** (140 mg, 0.184 mmol) as described for the synthesis of **19a**: ¹H-NMR (D₂O, 500 MHz) δ 8.65 (s, 1H, H-2 or H-8), 8.39 (s, 1H, H-2 or H-8), 7.12-7.30 (m, 5H, SPh), 6.30 (d, 1H, H-1', $J_{1',2'} = 2.2$ Hz), 5.32 (dd, 1H, H-2', $J_{2',1'} = 2.2$ Hz, $J_{2',3'} = 5.9$ Hz), 5.01 (m, 1H, H-1''), 4.94 (dd, 1H, H-3', $J_{3',2'} = 5.9$ Hz, $J_{3',4'} = 1.6$ Hz), 4.74 (d, 1H, MOM-CH₂, $J = 7.1$ Hz), 4.69 (m, 2H, H-4', MOM-CH₂), 4.50 (m, 1H, H-2''), 4.15 (m, 1H, H-5'a), 4.11 (m, 1H, H-5'b), 4.03 (m, 2H, H-5'' x 2), 3.99 (m, 1H, H-3''), 3.54 (s, 3H, MOM-CH₃ or OMe), 3.26 (s, 3H, MOM-CH₃ or OMe), 3.18 (q, 6H, Et₃NH-CH₂ x 3, $J = 7.3$ Hz), 2.71 (m, 1H, H-4''), 2.60 (m, 1H, H-6''a), 2.06 (m, 1H, H-6''b), 1.62, 1.38 (each s, each 3H, isopropylidene), 1.26 (t, 9H, Et₃NH-CH₃ x 3, $J = 7.3$ Hz); ¹³C-NMR (D₂O, 125 MHz) δ 150.82, 146.42, 146.30, 143.88, 133.18, 133.14, 129.92, 129.87, 129.42, 128.25, 119.82, 119.61, 115.15, 97.02, 91.86, 86.61, 86.53, 84.62, 84.33, 81.89, 66.24, 63.67, 58.06, 56.12, 47.10, 40.69, 40.63, 26.39, 24.73, 8.68; ³¹P-NMR (D₂O, 202 MHz) δ 16.70 (s), 0.55 (s); HRMS (FAB, positive) calcd for C₂₈H₄₀N₅O₁₃P₂S 748.1813 (MH⁺), found 748.1804; UV (H₂O) $\lambda_{\max}$ 260 nm.

2''-*O*-Methoxymethyl-3''-*O*-methyl-cyclic ADP-carbocyclic-xylose 2',3'-Acetonide (20b). Compound **20b** (37 mg, 47%) was obtained from **19b** (90 mg, 0.11 mmol) as described for the synthesis of **20a**: ¹H-NMR (D₂O, 500 MHz) δ 9.10 (s,

1H, H-2 or H-8), 8.45 (s, 1H, H-2 or H-8), 6.41 (s, 1H, H-1'), 5.55 (d, 1H, H-2', $J_{2',3'} = 6.1$ Hz), 5.44 (dd, 1H, H-3', $J_{3',2'} = 6.1$ Hz, $J_{3',4'} = 2.4$ Hz), 5.02 (m, 1H, H-1''), 4.75 (d, 1H, MOM-CH₂, $J = 6.8$ Hz), 4.67 (d, 1H, MOM-CH₂, $J = 6.8$ Hz), 4.64 (m, 1H, H-4'), 4.57 (m, 1H, H-2''), 4.15 (m, 2H, H-5'a, H-5''a), 4.09 (m, 3H, H-5'b, H-5''b, H-3''), 3.48 (s, 3H, MOM-CH₃ or OMe), 3.16 (s, 3H, MOM-CH₃ or OMe), 3.19 (q, 6H, Et₃NH-CH₂ x 3, $J = 7.3$ Hz), 2.94 (m, 1H, H-6'a), 2.88 (m, 1H, H-4''), 2.37 (m, 1H, H-6'b), 1.64, 1.44 (each s, each 3H, isopropylidene), 1.27 (t, 9H, Et₃NH-CH₃ x 3, $J = 7.3$ Hz); ¹³C-NMR (D₂O, 125 MHz) δ 152.18, 145.29, 144.63, 119.65, 115.10, 97.24, 92.46, 87.79, 86.93, 86.83, 85.04, 84.82, 81.87, 64.60, 64.25, 58.47, 56.24, 47.10, 36.34, 36.26, 28.41, 26.42, 24.73, 8.67; ³¹P-NMR (D₂O, 202 MHz) δ -9.56 (d, $J = 11.4$ Hz), -10.46 (d, $J = 11.4$ Hz); HRMS (FAB, positive) calcd for C₂₂H₃₄N₅O₁₃P₂ 638.1623 (MH⁺), found 638.1627; UV (H₂O) λ_{max} 259 nm.

3''-O-Methyl-cyclic ADP-carbocyclic-xylose (5). Compound **5** (30 mg, quant.) was obtained from **20b** (37 mg, 0.050 mmol) as described for the synthesis of **4**: ¹H-NMR (D₂O, 500 MHz, K⁺ salt) δ 9.04 (s, 1H, H-2 or H-8), 8.41 (s, 1H, H-2 or H-8), 6.10 (d, 1H, H-1', $J_{1',2'} = 6.1$ Hz), 5.21 (m, 1H, H-2'), 4.95 (m, 1H, H-1''), 4.62 (m, 1H, H-5'a), 4.60 (m, 1H, H-3'), 4.56 (m, 1H, H-2''), 4.44 (m, 1H, H-4'), 4.11 (m, 3H, H-5'b, H-5'' x 2), 4.03 (m, 1H, H-3''), 3.52 (s, 3H, OMe), 2.94 (m, 1H, H-6'a), 2.90 (m, 1H, H-4''), 2.52 (m, 1H, H-6'b); ¹³C-NMR (D₂O, 125 MHz) δ 152.19, 146.69, 145.43, 144.52, 120.42, 91.00, 85.77, 85.34, 81.35, 73.82, 71.18, 65.14, 63.94, 63.46, 58.59, 36.81, 27.72; ³¹P-NMR (D₂O, 202 MHz) δ -9.70 (d, $J = 11.4$ Hz), -10.32 (d, $J = 11.4$ Hz); HRMS (FAB, positive) calcd for C₁₇H₂₆N₅O₁₂P₂ 554.1048 (MH⁺), found 554.1035; UV (H₂O) λ_{max} 259 nm.

REFERENCES

1. This report constitutes Part 239 of Nucleosides and Nucleotides. Part 238: Kodama, T.; Shuto, S., Matsuda, A. Synthesis of 1'-fluoronucleosides. Submitted for publication.
2. Clapper, D. L.; Walseth, T. F.; Dargie, P. J.; Lee, H. C. Pyridine nucleotide metabolite stimulate calcium release from sea urchin egg microsomes desensitized to inositol triphosphate. *J. Biol. Chem.* **1987**, 262, 9561-9568.
3. Shuto, S.; Matsuda, A. Chemistry of cyclic ADP-ribose and its analogs. *Curr.*

Med. Chem. **2004**, *11*, 827-845, and references therein.

4. Lee, H. C.; Aarhus, R. Wide distribution of an enzyme that catalyzes the hydrolysis of cyclic ADP-ribose. *Biochem. Biophys. Acta* **1993**, *1164*, 68-74.
5. (a) Shuto, S.; Shirato, M.; Sumita, Y.; Ueno, Y.; Matsuda, A. Synthesis of cyclic IDP-carbocyclic-ribose, a stable mimic of cyclic ADP-ribose. Significant facilitation of the intramolecular condensation reaction of *N*-1-(carbocyclic-ribosyl)inosine 5',6"-Diphosphate derivatives by an 8-bromo-substitution at the hypoxanthine moiety. *J. Org. Chem.* **1998**, *63*, 1986-1994. (b) Fukuoka, M.; Shuto, S.; Minakawa, N.; Ueno, Y.; Matsuda, A. An efficient synthesis of cyclic IDP- and cyclic 8-bromo-IDP-carbocyclic-riboses using a modified Hata condensation method to form an intramolecular pyrophosphate linkage as a key step. An entry to a general method for the chemical synthesis of cyclic ADP-ribose analogues. *J. Org. Chem.* **2000**, *65*, 5238-5248. (c) Shuto, S.; Fukuoka, M.; Manikowsky, M.; Ueno, T.; Nakano, T.; Kuroda, R.; Kuroda, H.; Matsuda, A. Total synthesis of cyclic ADP-carbocyclic-ribose, a stable mimic of Ca^{2+} -mobilizing second messenger cyclic ADP-ribose. *J. Am. Chem. Soc.* **2001**, *123*, 8750-8759.
6. (a) Shuto, S.; Fukuoka, M.; Kudoh, T.; Garnham, C.; Galione, A.; Potter, B. V. L.; Matsuda, A. Convergent synthesis and unexpected Ca^{2+} -mobilizing activity of 8-substituted analogues of cyclic ADP-carbocyclic-ribose, a stable mimic of the Ca^{2+} -mobilizing second messenger cyclic ADP-ribose. *J. Med. Chem.* **2003**, *46*, 4741-4749. (b) Hashii, M.; Shuto, S.; Fukuoka, M.; Kudoh, T.; Matsuda, A.; Higashida, H. Amplification of depolarization-induced and ryanodine-sensitive cytosolic Ca^{2+} elevation by synthetic carbocyclic analogues of cyclic ADP-ribose and their antagonistic effects in NG108-15 neuronal cells. *J. Neurochem.* **2005**, *94*, 316-323. (c) Kudoh, T.; Fukuoka, M.; Ichikawa, S.; Murayama, T.; Ogawa, Y.; Hashii, M.; Higashida, H.; Kunerth, S.; Weber, K.; Guse, A. H.; Potter, B. V. L.; Matsuda, A.; Shuto, S. Synthesis of stable and cell-type selective analogues of cyclic ADP-ribose, a Ca^{2+} -mobilizing second messenger. Structure-activity relationship of the N1-ribose moiety. *J. Am. Chem. Soc.* **2005**, *127*, 8846-8855.
7. For examples: (a) Galeone, A.; Mayol, L.; Oliviero, G.; Piccialli, G.; Varra, M. Synthesis of a novel N-1 carbocyclic, N-9 butyl analogue of cyclic ADP ribose (cADPR). *Tetrahedron* **2002**, *58*, 363-368. (b) Huang, L.-J.; Zhao, Y.-Y.; Yuan, L.; Min J.-M.; Zhang L.-H. Syntheses and calcium-mobilizing evaluations of N1-glycosyl-substituted stable mimics of cyclic ADP-ribose. *J. Med. Chem.* **2002**, *45*, 5340-5352.
8. Sekine, M.; Nishiyama, S.; Kamimura, T.; Osaki, Y.; Hata, T. Chemical synthesis of capped oligoribonucleotides, $\text{m}^7\text{G}^{5'}$ ppp AUGACC. *Bull. Chem. Soc. Jpn.* **1985**, *58*, 850-860.

9. Yoshikawa, M.; Kato, T.; Takenishi, T. Studies of Phosphorylation. III. Selective phosphorylation of unprotected nucleosides. *Bull, Chem. Soc. Jpn.* **1969**, 42, 3505-3508.
10. Hata, T.; Kamimura, T.; Urakami, K.; Kohno, K.; Sekine, M.; Kumagai, I.; Shinozaki, K.; Miura, K. A new method for the synthesis of oligodeoxyribonucleotides bearing a 5'-terminal phosphate group. *Chem. Lett.* **1987**, 117-120.
11. Shiwa, S.; Murayama, T.; Ogawa, Y. Molecular cloning and characterization of ryanodine receptor from unfertilized sea urchin eggs. *Am. J. Physiol., Regul. Integr. Comp. Physiol.* **2002**, 282, R727-R737.

FIGURE 1 cADPR (1), cADPcR (2), and the 3''-modified cADPcR analogues 3–5.

FIGURE 2 Dose-dependent Ca^{2+} -mobilizing activity of compounds in sea urchin egg homogenate. The Ca^{2+} -mobilizing activity of each compound was expressed as a percent change in ratio of fura-2 fluorescence (F340/F380) relative to that of 10 μM cADPR. The compounds examined are cADPR (**1**, filled circles), cADPcR (**2**, open circles), 3''-deoxy-cADPcR (**3**, open triangles), cADPcX (**4**, open squares), and 3''-O-methyl-cADPcX (**5**, open diamonds). Data are mean $\pm$ SEM of 3-6 experiments.

SCHEME 1

SCHEME 2

Reagents and conditions: a) TIPSCl, imidazole, DMAP, DMF, rt, 91%; b) $\text{ClCH}_2\text{CO}_2\text{H}$, DIAD, Ph_3P , toluene, 0 °C, 74%; c) NaOMe, MeOH, rt, 88%; d) 1) MOMCl, $i\text{-Pr}_2\text{NEt}$, CH_2Cl_2 , rt, 2) TBAF, THF, rt, 97% (**13a**) ; e) 1) MeOTf, DTBMP, CH_2Cl_2 , rt, 2) TBAF, THF, rt, 45% (**13b**); f) H_2O , reflux, quant (**14a**, **14b**).

SCHEME 3

Reagents and conditions: a) K_2CO_3 , MeOH, rt, 83% (**15a**), 84% (**15b**); b) 1) DMTrCl, pyridine, rt, 2) TBAF, AcOH, THF, rt, quant. (**16a**), quant. (**16b**); c) PSS, TPSCl, py, rt, 65% (**17a**), 62% (**17b**); d) aq. 60% AcOH, rt, 83% (**18a**), 88% (**18b**); e) 1) $POCl_3$, $(EtO)_3PO$, 0 °C, 2) H_3PO_2 , Et_3N , NMM, pyridine, 0 °C, 46% (**19a**), 58% (**19b**); f) $AgNO_3$, MS 3A, Et_3N , py, rt, 46% (**20a**), 47% (**20b**); h) aq. 80% HCO_2H , then aq. NH_3 , rt, 90% (**4**), quant. (**5**).