



Title	INCREASE OF Na ⁺ GRADIENT-DEPENDENT L-GLUTAMATE AND L-ASPARTATE TRANSPORT IN HIGH K ⁺ DOG ERYTHROCYTES ASSOCIATED WITH HIGH ACTIVITY OF (Na ⁺ , K ⁺)-ATPase
Author(s)	INABA, Mutsumi
Citation	Japanese Journal of Veterinary Research, 32(2), 96-96
Issue Date	1984-04-28
Doc URL	http://hdl.handle.net/2115/4703
Type	bulletin (article)
File Information	KJ00002374219.pdf



[Instructions for use](#)

INCREASE OF Na^+ GRADIENT-DEPENDENT L-GLUTAMATE AND L-ASPARTATE
TRANSPORT IN HIGH K^+ DOG ERYTHROCYTES ASSOCIATED WITH
HIGH ACTIVITY OF $(\text{Na}^+, \text{K}^+)\text{-ATPase}$

Mutsumi INABA

*Department of Veterinary Internal Medicine
Faculty of Veterinary Medicine
Hokkaido University, Sapporo, 060, Japan*

As reported previously (MAEDE, Y., INABA, M. & TANIGUCHI, N. (1983): *Blood*, **61**, 493-499), certain dogs possess red cells characterized by low Na^+ , high K^+ concentrations and high activity of $(\text{Na}^+, \text{K}^+)\text{-ATPase}$, although normal dog red cells contain low K^+ , high Na^+ , and lack $(\text{Na}^+, \text{K}^+)\text{-ATPase}$. Furthermore, these red cells show increased activities of L-glutamate and L-aspartate transport, resulting in high accumulations of such amino acids in their cells. The present study demonstrated: (i) Na^+ gradient-dependent L-glutamate and L-aspartate transport in the high K^+ and low K^+ red cells were dominated by a saturable component obeying Michaelis-Menten kinetics. Although no difference of the K_m values was observed between the high K^+ and low K^+ cells, the V_{max} values for both amino acids transport in the high K^+ cells were about 3 times those of low ones. (ii) L- and D-Aspartate but not D-glutamate competitively inhibited L-glutamate transport in both types of the cells. (iii) Ouabain decreased the uptake of the amino acids in the high K^+ dog red cells, whereas it was not effective on those in the low K^+ cells. (iv) The ATP-treated high K^+ cells ($[\text{K}^+]_{\text{in}} \cong [\text{K}^+]_{\text{out}}$, $[\text{Na}^+]_{\text{in}} > [\text{Na}^+]_{\text{out}}$) showed a marked decrease of uptake rate of both amino acids, which was almost the same as that of the low K^+ cells. (v) Valinomycin stimulated the transport of the amino acids in both of the high K^+ cells ($[\text{K}^+]_{\text{in}} > [\text{K}^+]_{\text{out}}$, $[\text{Na}^+]_{\text{in}} < [\text{Na}^+]_{\text{out}}$), suggesting that the transport system of L-glutamate and L-aspartate in both types of cells might be electrogenic. These results indicate that the increased transport activity in the high K^+ dog red cells was a secondary consequence of the Na^+ concentration-gradient created by $(\text{Na}^+, \text{K}^+)\text{-ATPase}$.