Title	Purification, characterization and amino acid sequence of a novel enzyme, D-threo-3-hydroxyaspartate dehydratase, from <i>Delftia</i> sp. HT23
Author(s)	Maeda, Takayuki; Takeda, Yuki; Murakami, Tomoko; Yokota, Atsushi; Wada, Masaru
Citation	The Journal of Biochemistry, 148(6), 705-712 https://doi.org/10.1093/jb/mvq106
Issue Date	2010-12
Doc URL	http://hdl.handle.net/2115/47109
Rights	This is a pre-copy-editing, author-produced PDF of an article accepted for publication in The Journal of Biochemistry following peer review. The definitive publisher-authenticated version, J Biochem (2010) 148 (6): 705-712, is available online at: http://jb.oxfordjournals.org/content/148/6/705
Type	article (author version)
File Information	JoB148-6_705-712.pdf

[Instructions for use](#)

Regular Paper

Field; Biochemistry (Topic; Enzymology)

Purification, characterization, and amino acid sequence of a novel enzyme, D-threo-3-hydroxyaspartate dehydratase, from *Delftia* sp. HT23

Takayuki Maeda, Yuki Takeda, Tomoko Murakami, Atsushi Yokota, and Masaru Wada*

Division of Applied Bioscience, Research Faculty of Agriculture, Hokkaido University, Kita-9, Nishi-9, Kita-ku, Sapporo, 060-8589, Japan

Running title: D-threo-3-hydroxyaspartate dehydratase from *Delftia*

*Correspondence to Masaru Wada, Division of Applied Bioscience, Research Faculty of Agriculture, Hokkaido University, Kita-9, Nishi-9, Kita-ku, Sapporo, 060-8589, Japan
Tel: +81-11-706-4185; Fax: +81-11-706-4961; E-mail: wada@chem.agr.hokudai.ac.jp

Abbreviations:

DH, dehydratase; D-EHA, D-erythro-3-hydroxyaspartate; D-THA, D-threo-3-hydroxyaspartate; EDTA, ethylenediaminetetraacetic acid; IPTG, isopropyl- β -D-thiogalactopyranoside; LB, Luria–Bertani; L-EHA, L-erythro-3-hydroxyaspartate; L-THA, L-threo-3-hydroxyaspartate; MALDI-TOF-MS, matrix-assisted laser desorption/ionization time-of-flight mass spectrometry; NADH, nicotinamide adenine dinucleotide; PAGE, polyacrylamide gel electrophoresis; PCR, polymerase chain reaction; PLP, pyridoxal 5'-phosphate; SDS, sodium dodecyl sulfate; TLC, thin layer chromatography.

Summary

D-*threo*-3-hydroxyaspartate dehydratase (D-THA DH) was purified from the cell-free extract of the soil-isolated bacterium *Delftia* sp. HT23. The enzyme exhibited dehydratase activity toward D-*threo*-3-hydroxyaspartate, L-*threo*-3-hydroxyaspartate, L-*erythro*-3-hydroxyaspartate, and D-serine.

Absorption of the purified enzyme at 412 nm suggests that it contains pyridoxal 5'-phosphate (PLP) as a cofactor. The NH₂-terminal and internal amino acid sequences showed significant similarity to hypothetical alanine racemase of genome-sequenced *Delftia acidovorans* SPH-1; however, the purified enzyme showed no alanine racemase activity. Using the sequence information of *Delftia acidovorans* SPH-1, the gene encoding D-THA DH was cloned. The deduced amino acid sequence, which belongs to the alanine racemase family, shows significant (26-36%) similarity to D-serine dehydratase of both *Saccharomyces cerevisiae* and chicken. In order to obtain purified D-THA DH efficiently, the gene was expressed in *Escherichia coli*. The recombinant enzyme was highly activated by divalent cations, such as Mn²⁺, Co²⁺, and Ni²⁺. Site-directed mutagenesis experiment revealed that lysine 43 is an important residue involved in PLP binding and catalysis. This is the first reported enzyme that acts on D-THA. In addition, this enzyme is the first example of a prokaryotic dehydratase belonging to the fold-type III PLP-dependent enzyme family.

Key words: D-*threo*-3-hydroxyaspartate dehydratase, alanine racemase, pyridoxal 5'-phosphate,

Delftia sp. HT23

3-Hydroxyaspartate and its derivatives have attracted the attention of biochemists because they are competitive blockers of the excitatory glutamate/aspartate transporters of the mammalian nervous system (1, 2). However, 3-hydroxyaspartate has two chiral centers, and their four stereoisomers, *i.e.*, *D-threo*-3-hydroxyaspartate (D-THA), *L-threo*-3-hydroxyaspartate (L-THA), *D-erythro*-3-hydroxyaspartate (D-EHA), and *L-erythro*-3-hydroxyaspartate (L-EHA), have been difficult to synthesize (3).

The biochemical activity of 3-hydroxyaspartate has been investigated in considerable detail (4). Nevertheless, little is known about the enzymes that act on 3-hydroxyaspartate isomers, although two microbial enzymes, *erythro*-3-hydroxyaspartate aldolase (EC 4.1.3.14) (5, 6) and *erythro*-3-hydroxyaspartate dehydratase (EC 4.3.1.20) (7), were identified many years ago. Recently, we reported that *L-threo*-3-hydroxyaspartate dehydratase (EC 4.3.1.16) isolated from the soil-isolated bacterium *Pseudomonas* sp. T62 exhibits dehydratase activity specifically toward L-THA and not toward other 3-hydroxyaspartate isomers (8, 9). The amino acid sequence and detailed biochemical features, including side reactions, of this enzyme have also been reported (9). However, no enzyme acting on the *D-threo* form of 3-hydroxyaspartate has been reported. The enzyme degrading D-THA might be useful for enzymatic optical resolution of DL-THA to produce optically pure L-THA. Because there has been no report about enzyme acting on D-THA so far, in order to obtain the amino acid sequence information of D-THA converting enzyme, enzyme purification from microorganism that produces D-THA converting enzyme was necessary. Thus, we screened microorganisms that can utilize D-THA as a sole carbon source and found that a newly isolated bacterium, *Delftia* sp. HT23, which produces an enzyme that catalyzes the dehydratase reaction of D-THA to oxaloacetate. We designated this enzyme as *D-threo*-3-hydroxyaspartate dehydratase (D-THA DH; *D-threo*-3-hydroxyaspartate ammonia-lyase).

We report here the purification, partial characterization, and amino acid sequence of this novel enzyme, D-THA DH, from *Delftia* sp. HT23. The comparison of amino acid sequence of this enzyme with known eukaryotic D-serine dehydratases is also discussed. This information may provide useful clues for understanding mechanisms of these PLP-dependent enzymes.

Materials and Methods

Materials.

The L-THA was purchased from Tocris Cookson, Ltd. (Bristol, UK); L-EHA, from Wako Pure Chemicals (Osaka, Japan); and DL-THA, from Tokyo Kasei Kogyo (Tokyo, Japan). D-THA was prepared from DL-THA by enzymatic resolution using L-THA dehydratase from *Saccharomyces cerevisiae* (10). Restriction endonucleases were obtained from Nippon Gene (Toyama, Japan). All other chemicals were of analytical grade and commercially available.

Screening of D-THA DH-producing microorganisms from soil.

Basal agar medium contained 3 g of D-THA, 1 g of $(\text{NH}_4)_2\text{SO}_4$, 1 g of KH_2PO_4 , 1 g of K_2HPO_4 , 0.5 g of yeast extract, 0.2 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, and 2% agar in 1 L of tap water, pH 7.0. Each of the soil samples collected from Sapporo city area was suspended in 0.85% NaCl solution, streaked onto the basal agar medium, and incubated at 30°C for 48 h. Strains forming colonies were isolated and transferred to the same agar medium, and cultured at 30°C until growth of the microorganisms was apparent. The isolated colonies were then transferred to liquid medium containing 10 g of glucose, 3 g of D-THA, 1 g of KH_2PO_4 , 1 g of K_2HPO_4 , 0.5 g of yeast extract, 0.2 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ in 1 L of tap water (pH 7.0), and cultivated 30°C for 16 h with shaking. The cells harvested by centrifugation were used for the reaction. Each reaction mixture contained 50 mM D-THA, 0.5 mM $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$,

and 50 mM potassium phosphate buffer (pH 7.0). The reaction was carried out at 30°C for 16 h with shaking. Degradation of D-THA in the reaction mixture was monitored by thin layer chromatography (TLC) using the developing solvent ethanol/28% ammonia solution/water=7/1/2 (v/v/v), and was visualized by ninhydrin. D-THA DH activity of the cell-free extract prepared from the strains showing high D-THA degrading activity on TLC was measured as described below. Determination of 16S rDNA was done using a MicroSeq 500 16S rDNA Bacterial Identification Sequencing kit (Applied Biosystems, CA, USA).

Microorganism and cultivation.

Delftia sp. HT23, isolated from soil and identified in our laboratory, was used. Bacteria were grown aerobically in medium containing 10 g of DL-THA, 10 g of glucose, 1 g of KH_2PO_4 , 1 g of K_2HPO_4 , 0.5 g of yeast extract, and 0.2 g of $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ in 1 L of tap water, pH 7.0. A loopful of *Delftia* sp. HT23 cells was inoculated into a test tube (16.5 × 165 mm) containing 5 mL of medium and was cultivated for 24 h at 30°C. The culture was transferred to a 2-L flask containing 400 mL of medium and grown for 48 h at 30°C, with shaking. *Escherichia coli* JM109 was used as the host cell for the cloning and expression of the D-THA DH gene (*dthadh*). The *E. coli* cells were grown at 37°C in Luria-Bertani (LB) medium containing 1% polypeptone, 0.5% yeast extract, and 1% NaCl (pH 7.0). When necessary, 100 $\mu\text{g mL}^{-1}$ ampicillin were added to the medium.

Enzyme assays.

The activity of 3-hydroxyaspartate dehydratase was determined as described previously (8). One unit of the enzyme was defined as the amount capable of catalyzing the oxidation of 1 μmol of NADH per min. Serine dehydratase activity (11) and alanine, serine, and aspartate racemase activities (12-14)

were measured as described previously. Protein concentrations were determined by the dye-binding method of Bradford (15) with a Bio-Rad protein assay kit, using bovine serum albumin as the standard.

Purification of the enzyme.

All purification procedures were carried out at 4°C in 10 mM Tris-HCl buffer (pH 8.0) containing 0.01 mM PLP, 0.1 mM MnCl₂, and 0.1 mM dithiothreitol, unless otherwise stated. *Delftia* sp. HT23 cells (30 g wet weight) obtained from a 2.4-L culture were disrupted with an ultrasonic oscillator. After centrifugation (8,000 × *g* for 40 min), the supernatant was fractionated with solid ammonium sulfate. The precipitate obtained at 40–60% saturation was collected, dialyzed against 10 L of the buffer for 18 h, and applied to a HiPrep Q FF 16/10 column (1.6 × 10 cm; GE healthcare, UK) equilibrated in the buffer. The enzyme was eluted with a linear gradient of 0 to 1.0 M NaCl in 160 mL of the buffer at a flow rate of 2.0 mL/min. The enzyme eluted at approximately 0.55 M NaCl.

The concentration of (NH₄)₂SO₄ was adjusted to 1 M by the addition of solid (NH₄)₂SO₄, and the enzyme solution was loaded onto a HiTrap Phenyl FF column (high sub, 1.0 × 10 cm; GE Healthcare), previously equilibrated with buffer containing 1 M (NH₄)₂SO₄, connected to a FPLC system (Pharmacia Biotech, Sweden). The enzyme was eluted with a linear gradient of 1 to 0 M (NH₄)₂SO₄ in 230 mL of buffer at a flow rate of 1 mL/min. The activity-containing fractions, which eluted with approximately 0.25 M (NH₄)₂SO₄, were pooled and dialyzed against 3 L of buffer for 8 h.

The concentrated enzyme solution was applied to a Superdex-200 HR10/30 column (1.0 × 30 cm; GE Healthcare) equilibrated with buffer containing 0.15 M NaCl, and the enzyme was eluted with the same buffer.

The enzyme solution was then applied to a RESOURCE Q column (0.5×5 cm) previously equilibrated with the same buffer and was eluted by FPLC with a linear gradient of 0 to 0.8 M NaCl in 35 mL of buffer at a flow rate of 1 mL/min. The activity-containing fractions, eluting with approximately 0.3 M NaCl, were collected.

The $(\text{NH}_4)_2\text{SO}_4$ concentration of the enzyme solution was adjusted to 1 M by the addition of solid $(\text{NH}_4)_2\text{SO}_4$, and the enzyme solution was applied to a HiTrap Butyl FF column (1.0×10 cm; GE Healthcare) previously equilibrated with the buffer containing 1 M $(\text{NH}_4)_2\text{SO}_4$. The enzyme was eluted by FPLC with a linear gradient of 1 to 0 M $(\text{NH}_4)_2\text{SO}_4$ in 230 mL of buffer at a flow rate of 2 mL/min. The activity-containing fractions, which eluted with approximately 0.3 M $(\text{NH}_4)_2\text{SO}_4$, were pooled, dialyzed against 3 L of the buffer for 8 h, and used as the purified enzyme.

Molecular weight determination.

SDS-PAGE was performed using a 4.5% acrylamide stacking gel and 12.5% acrylamide separation gel. For molecular weight determination of the enzyme subunit by SDS-PAGE, the following molecular weight standards were used: phosphorylase *b* ($M_r = 97,400$), bovine serum albumin (66,300), aldolase (42,400), carbonic anhydrase (30,000), trypsin inhibitor (20,100), and lysozyme (14,400).

The molecular weight of the native protein was determined by gel-permeation liquid chromatography on TSKgel Super-SW3000 (Tosoh, Japan) using glutamate dehydrogenase ($M_r = 290,000$), lactate dehydrogenase (142,000), enolase (67,000), myokinase (32,000), and cytochrome *c* (12,400) as molecular weight standards. The column was equilibrated and eluted with 100 mM potassium phosphate buffer (pH 6.7) containing 0.1 M Na_2SO_4 , and 0.05 % (w/v) NaN_3 at a flow rate of 0.35 mL/min.

The molecular weight of the recombinant enzyme was also estimated using a MALDI-TOF-MS (Voyager Biospectrometry, Applied Biosystems) in linear mode at 25-kV acceleration voltage, with sinapic acid as the matrix.

Amino acid sequence analysis.

The N-terminal and internal peptide sequences were determined as described previously (16) with a modification. After lysyl endopeptidase digestion of the enzyme, the peptides were separated by HPLC (L-2000 system; Hitachi-Hitec Corp., Tokyo, Japan) on a CAPCELL PAK C₁₈ MGIII column (4.6 × 250 mm; SHISEIDO, Tokyo, Japan). Edman degradation was performed at the Center for Instrumental Analysis at Hokkaido University.

Construction of the expression plasmid.

Total DNA was isolated from *Delftia* sp. HT23 using Isoplant II (Nippon Gene, Toyama, Japan). For the expression of *dthadh* in *E. coli*, a DNA fragment containing the open reading frame of *dthadh* was prepared by PCR, using *Delftia* sp. HT23 genomic DNA as a template and oligonucleotide sense (5'-ATGCGGATCCATGCAAGACACACTTCTGAC-3') and antisense primers (5'-ATATAAGCTTTCACCAGCCATGGAGCCGCT-3' (the underlined sequences are *Bam*HI and *Hind*III sites, respectively). The PCR mixture (50 µL) contained 10 pmol of each primer, 0.2 mM of each dNTP, and 1.25 U of PrimeSTAR HS DNA polymerase (Takara Bio, Ohtsu, Japan). The thermal cycler program was 98°C for 10 s, 55°C for 5 s, and 72°C for 1.5 min. The unique amplified band corresponding to about 1,100 bp was digested with *Bam*HI and *Hind*III, and then ligated into the *Bam*HI and *Hind*III sites of pQE30 (Qiagen, Hilden, Germany) to obtain pQE30dthadh. The expression vector was introduced into *E. coli* JM109 cells, and the nucleotide sequence of the insert

was confirmed.

Expression and purification of recombinant D-THA DH.

The transformed *E. coli* JM109 cells carrying pQE30dthadh were grown at 37°C in 50 mL of LB medium containing ampicillin. To induce gene expression, 0.1 mM IPTG was added to the culture medium when the absorbance at 600 nm reached 0.3. After cultivation for another 8 h at 37°C, cells were harvested by centrifugation. All purification procedures were carried out at 4°C in 10 mM Tris-HCl buffer (pH 8.0) containing 0.01 mM PLP, 0.1 mM MnCl₂, and 0.1 mM dithiothreitol. The cells (0.62 g wet weight) obtained from a 50-mL culture were disrupted with an ultrasonic oscillator. After centrifugation (8,000 × *g* for 15 min), the supernatant was applied to a HisTrap HP column (0.7 × 2.5 cm; GE Healthcare, UK) equilibrated with buffer supplemented with 20 mM imidazole. The enzyme was eluted by FPLC with a step-wise gradient of 20 to 500 mM imidazole. Fractions showing activity, which eluted with approximately 150 mM imidazole, were collected, dialyzed against the buffer, and used as the enzyme for characterization.

Site-directed mutagenesis.

A mutant enzyme, K43A, was prepared according to the protocol for a PrimeSTAR Mutagenesis Basal Kit (Takara Bio). The nucleotide substitutions were confirmed by DNA sequencing. The mutant enzyme was produced in *E. coli* JM109 cells and purified by the same procedure as that used for the wild-type recombinant enzyme.

EDTA-treated enzyme.

Recombinant D-THA DH was dialyzed against 10 mM Tris-HCl buffer (pH 8.0) containing 0.01 mM

PLP, 0.1 mM dithiothreitol, and 5 mM EDTA for 12 h at 4°C. To remove EDTA, the enzyme solution was further dialyzed against the same buffer without EDTA for 12 h at 4°C.

Nucleotide sequence accession number.

The nucleotide sequence of the *dthadh* gene of *Delftia* sp. HT23 has been deposited in the DDBJ/EMBL/GenBank database under accession number AB433986.

Bioinformatic analysis.

A homology search was performed with the FASTA program at DDBJ (<http://www.ddbj.nig.ac.jp/search/fasta-j.html>) (17). The amino acid sequence alignment was performed using ClustalW 1.83 and BOXSHADE 3.21 (18).

Results

Result of screening

D-THA-utilizing microorganisms were screened, and strains with high D-THA-degrading activity were isolated from soil. Four hundred thirty seven strains were isolated as D-THA-utilizing microorganisms, and 17 strains were selected by TLC as D-THA-degrading microorganisms. Most of these 17 strains showed 0.02 – 0.2 unit (mg protein)⁻¹ activity of D-THA DH, which produces oxalacetate from D-THA (8). Among these strains, strain HT23, which is rod-shaped, and gram-negative bacterium, was selected as the enzyme source, because HT23 showed the highest D-THA DH activity (around 0.2 unit (mg protein)⁻¹) among the strains tested.

The most variable region of the 16S rDNA sequence (460 bp) revealed 99.9% identity to *Delftia acidovorans* (19). These results indicate that strain HT23 belongs to the genus, *Delftia* sp. Thus, we designated this strain as *Delftia* sp. HT23.

Induction of the enzyme.

When *Delftia* sp. HT23 was cultured in the medium described above, the cell-free extract showed an enzyme activity of approximately 0.2 unit (mg protein)⁻¹. However, when DL-THA in the medium was replaced by an equal amount of D-serine, D-threonine, D-aspartate, or peptone, the cell-free extract showed no or only a trace [<0.02 unit (mg protein)⁻¹] of enzyme activity. These results suggest that the enzyme was induced by 3-hydroxyaspartate in the medium.

Purification and molecular weight determination of D-threo-3-hydroxyaspartate dehydratase.

With the purification procedures described above, the enzyme was purified approximately 116-fold to homogeneity, with approximately 1.4% recovery (Table I). The purified enzyme preparation gave a single band on SDS-PAGE (Fig. 1). Furthermore, by high-performance gel-permeation liquid chromatography, the enzyme gave a single symmetrical protein peak.

Using a calibrated TSKgel Super-SW3000 column, the relative molecular weight of the enzyme was estimated to be 36,000. By SDS-PAGE, the relative molecular weight of the subunit was estimated to be about 41,000, suggesting that the enzyme is a monomer.

Substrate specificity of the enzyme.

The enzyme showed broad specificity toward 3-hydroxyaspartate isomers. Table II shows the substrate specificity and kinetic parameters of D-THA DH. In addition, D- and L-serine reacted as

poor substrates for the enzyme. Normal hyperbolic kinetics were observed with all substrates tested.

The K_m , k_{cat} , and k_{cat}/K_m values, calculated from Lineweaver-Burk plots, are shown in Table II.

Alanine, serine, and aspartate racemase activities of the enzyme were investigated using a sensitive assay method. After a 12-h incubation of the enzyme with L-alanine, L-serine, or L-aspartate, no D-alanine, D-serine, or D-aspartate was observed by HPLC with fluorometric detection. Thus, any alanine, serine, or aspartate racemase activity of D-THA DH is below the detection limit of 5.0×10^{-2} pmol h⁻¹ (mg protein)⁻¹.

Effects of pH and temperature.

The optimal pH and temperature of the enzyme were 8.5 and 50°C, respectively.

Absorption spectrum.

The enzyme exhibited absorption maxima at 280 and 412 nm (Fig. 2). Solutions of the pure enzyme are distinctly yellow. These results suggest that the enzyme contains PLP as its prosthetic group.

Effects of chemicals.

Hydroxylamine or EDTA, each at a final concentration of 1 mM, was added to the standard reaction mixture, and enzyme activity was measured. The enzyme was strongly inhibited by hydroxylamine (91.2% inhibition), suggesting that PLP participates in the enzyme reaction, as in the L-threo-3-hydroxyaspartate dehydratase reaction (8–10). The enzyme was also modestly inhibited by EDTA (27% inhibition), suggesting that metal ions are involved in the enzyme reaction.

N-Terminal and internal amino acid sequence analysis.

Automated Edman degradation of the enzyme protein was performed with a pulsed liquid phase sequencer, giving an NH₂-terminal amino acid sequence of MQDTLLTLDTPAAVIDLDRMQXNIA, where X is an unidentified amino acid. The enzyme was digested with lysyl endopeptidase, and the peptides were separated by HPLC. One peptide was isolated, and the amino acid sequence was found to be RDRGTARQK. The two sequences were compared with sequences in a protein sequence database (nr-aa), using the sequence similarity search program FASTA (17). Both sequences were almost perfectly matched to the partial amino acid sequences of the putative alanine racemase of *Delftia acidovorans* SPH-1 (accession No. ABX33617), whose whole genome has been sequenced (<http://genome.jgi-psf.org/delac/delac.home.html>). These results strongly suggest that the enzyme purified from *Delftia* sp. HT23 is identical to or very closely related to the putative alanine racemase from *D. acidovorans* SPH-1.

Nucleotide sequence of the D-THA DH gene from Delftia sp. HT23.

Based on the sequence of the gene encoding *D. acidovorans* SPH-1 putative alanine racemase, a pair of specific PCR primers was synthesized. The PCR product was ligated into the vector pQE30, and the nucleotide sequence was determined.

The open reading frame was 1,143 bp long and encoded a protein of 380 amino acid residues, with a predicted molecular weight of 40,300. A FASTA search revealed that the deduced amino acid sequence was 99% identical with the putative alanine racemase from *D. acidovorans* SPH-1, described above. The amino acid sequence showed high identity with similar enzymes from various Gram-negative bacteria, including the putative alanine racemases from *Ralstonia solanacearum* (71%), *Burkholderia cenocepacia* (71%), and *Pseudomonas syringae* (68%), and the putative metal-activated pyridoxal enzyme from *Bordetella pertussis* (75%). The amino acid sequence of D-THA DH

had relatively low, but significant, similarity to two eukaryotic D-serine dehydratases, those from *Gallus gallus* (chicken, 36%) (20) and *S. cerevisiae* (26%) (21), as shown in the alignment in Fig. 4. The PROSITE database (22) (<http://au.expasy.org/prosite/>) could not predict the PLP-binding site of D-THA DH; however, the three enzymes share a common motif, R(P/A)HVKT, in their N-terminal regions. In D-serine dehydratase from *S. cerevisiae*, the lysine residue in this motif binds PLP (21). Thus, Lys43 of D-THA DH is probably a PLP-binding residue.

Characterization of the recombinant enzyme.

We purified recombinant His-tagged D-THA DH from *E. coli* cells and characterized its enzymatic properties. The molecular weight determined by SDS-PAGE analysis (41,000) and that determined by MALDI-TOF-MASS analysis (41,600) were in agreement with that calculated from the deduced amino acid sequence of the recombinant enzyme (40,900). The first 15 N-terminal amino acid residues sequenced in recombinant His-tagged enzyme perfectly matched the deduced amino acid sequence. Moreover, the purified recombinant enzyme showed high activity, with specific activity of around 20 units (mg protein)⁻¹ toward D-THA. From these results, we concluded that this protein is recombinant D-THA DH.

Effect of metal ions on the recombinant enzyme.

To investigate the role of metal ions in more detail, EDTA-treated recombinant enzyme was prepared as described in the Materials and methods. The EDTA-treated enzyme showed about 0.1 unit (mg protein)⁻¹ activity toward D-THA, which was about 0.5% activity of the non-EDTA-treated enzyme. However, when CoCl₂, MnCl₂, NiCl₂, ZnCl₂, CaCl₂, or FeCl₂ was added to the enzyme solution at various concentrations, the relative activity was restored or even activated as shown in Fig. 3. No

activity was detected when SnCl_2 or CuCl_2 was added to the enzyme solution, suggesting that Sn^{2+} and Cu^{2+} are inhibitors of this enzyme. These results indicate that divalent cations such as Co^{2+} , Mn^{2+} , Ni^{2+} , Ca^{2+} , Zn^{2+} , and Fe^{2+} are activators of this enzyme.

Identification of PLP-binding lysine residue.

To identify the PLP-interacting lysine residue of the enzyme, a mutant enzyme, K43A, was constructed and purified as described in the Materials and methods. The K43A mutant enzyme, in which Lys43 was replaced by alanine, showed no detectable activity (<0.01% of the activity toward D-THA) and lacked a large absorption maximum at 412 nm (Fig. 2). Thus, we concluded that Lys43 is most likely involved in both PLP binding and catalysis.

Discussion

In this report, an enzyme that acts on D-threo-3-hydroxyaspartate was isolated and purified to homogeneity for the first time. The microbial metabolism of 3-hydroxyaspartate was described by Kornberg and Morris in the 1960s (6). They reported two enzymes in *Micrococcus denitrificans* that act on 3-hydroxyaspartate: erythro-3-hydroxyaspartate aldolase (EC 4.1.3.14) (5, 6) and erythro-3-hydroxyaspartate dehydratase (EC 4.2.1.38) (7). However, these enzymes have not been purified to homogeneity or well characterized. More recently, the gene encoding 3-hydroxyaspartate aldolase was cloned and expressed in *E. coli* (23). In addition to these enzymes, L-THA DH, which acts only on L-THA, has been identified and characterized by our group (8–10), but enzymes acting on D-THA have not yet been reported. To our knowledge, the present study is the first to report an enzyme that catalyzes the deamination of D-THA. Therefore, we designated this enzyme D-threo-3-hydroxyaspartate dehydratase (D-THA DH), although it also has some activity toward L-EHA. Based

on this substrate specificity, the enzyme is clearly distinct from the *erythro*-3-hydroxyaspartate dehydratase (EC 4.2.1.38) reported by Gibbs and Morris (7), which acts only on L-EHA and not on D-THA.

D-THA DH had approximately 30% amino acid sequence similarity to two eukaryotic D-serine dehydratases, from chicken (20) and *S. cerevisiae* (21) (Fig. 4). Although D-THA DH acts on D-serine, the relative activity toward D-serine was only about 23% of the activity toward D-THA. Furthermore, the production of D-THA DH in *Delftia* sp. HT23 cells was not induced by D-serine, but was induced by D-THA in the medium. From these results, we concluded that this enzyme is a D-*threo*-3-hydroxyaspartate dehydratase, rather than a D-serine dehydratase.

The absorption spectrum of the purified enzyme revealed that D-THA DH contains PLP, as do other bacterial dehydratases, although the amino acid sequence of D-THA DH was not similar to that of L-THA DH, which catalyzes the same reaction of L-THA to oxaloacetate (9, 10). Despite catalyzing the dehydratase reaction of D-THA, D-THA DH does not belong to the family of serine/threonine dehydratases, which contains most bacterial dehydratases (24). Based on their folding patterns, the PLP-dependent enzymes are classified into five groups, designated fold types I to V (25). The L-THA DH enzymes of *Pseudomonas* sp. T62 and *S. cerevisiae* belong to the fold type II group (9, 10), whereas D-THA DH belongs to the fold type III group. Eukaryotic D-serine dehydratases, which are also fold-type III PLP-dependent enzymes, have been reported from chicken and from *S. cerevisiae* (20, 21). However, D-THA DH is the first example of a prokaryotic dehydratase belonging to the fold type III PLP-dependent enzyme family.

The high amino acid sequence identities between D-THA DH and several putative alanine racemases and putative metal-activated pyridoxal enzymes from various Gram-negative bacteria, including enzymes from *Bordetella pertussis* (75%), *Ralstonia solanacearum* (71%), *Burkholderia*

cenocepacia (71%), and *Pseudomonas syringae* (68%), suggest that these putative enzymes may have D-THA DH activity. D-THA DH is expected to be present in other organisms, especially the Gram-negative soil bacteria mentioned above.

Site-directed mutagenesis experiment revealed that lysine 43 is an important residue involved in PLP binding and catalysis (Fig. 2), however, the K43A mutant enzyme still has weak absorption peaks around 320 nm and 420 nm. The reason why K43A mutant enzyme has these absorption peaks remains unknown, however, non-covalently bound PLP may cause these absorption peaks as in the case of aspartate aminotransferase of *E. coli* (26). Another explanation is also possible; Lys residue other than 43 (for example, Lys46) may bind PLP weakly, and gave catalytically inactive enzyme-PLP complex with absorption around 320 nm, and 420 nm. But, further investigation is required to elucidate this phenomenon.

Like L-THA DH from *Pseudomonas* sp. T62, D-THA DH requires divalent cations for its activation (8, 9), but the activation pattern differed between the two enzymes. Both Co^{2+} and Zn^{2+} act as inhibitors of L-THA DH, whereas these divalent cations are activators of D-THA DH. The pattern of D-THA DH activation by divalent cations was similar to that of D-threonine aldolase from *Arthrobacter* sp. DK-38 (27, 28), which is also activated by Mn^{2+} , Co^{2+} , Ni^{2+} , Fe^{2+} , and Ca^{2+} . D-Threonine aldolase from *Arthrobacter* sp. DK-38 also belongs to the fold-type III PLP-dependent enzymes (28) and shares around 30% amino acid identity to D-THA DH (data not shown). D-Threonine aldolase from *Arthrobacter* sp. DK-38 can bind 1 mol of Mn^{2+} ion per mol of subunit (28), however, the amount of metal ion, which binds to D-THA DH, has not yet been determined.

Although the reason why activity of the EDTA-treated enzyme added with 100 μM MnCl_2 is higher than that of the non-EDTA-treated enzyme dialyzed against the same concentration of MnCl_2 is not clear (Fig. 3), the following explanation might be possible; the EDTA-treated enzyme is

somewhat more stable than the non-EDTA-treated enzyme, *i.e.*, the non-EDTA-treated enzyme (containing Mn^{2+}) might be partially inactivated during the dialysis, however, the EDTA-treated enzyme (containing no or very little amount of metal ions) may become more active than the non-EDTA-treated enzyme when Mn^{2+} is added just before activity measurement. Effect of the metal ions on stability of the enzyme, however, still needs to be elucidated.

Although the physiological function of this enzyme remains to be clarified, the dehydratase reaction catalyzed by D-THA DH may be one of the reactions enabling *Delftia* sp. HT23 to grow on medium containing D-THA as the sole carbon source. This hypothesis is also supported by the fact that D-THA DH is inducible in *Delftia* sp. HT23; however, the details have not been elucidated.

To analyze the detailed reaction mechanism and 3D structure of this enzyme, an efficient expression system is necessary. Unfortunately, the expression level of D-THA DH in *E. coli* was low. The specific activity of the cell-free extract of *E. coli* expressing the D-THA DH gene was about 0.2 unit (mg protein)⁻¹, which is almost equal to that of the cell-free extract of the original strain, *Delftia* sp. HT23. The reason for poor expression in *E. coli* is unknown; however, the high GC content of the gene (71.9%) is one possible explanation. Currently, we are trying to improve the expression level of D-THA DH in *E. coli*.

Funding

This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Sports, and Culture of Japan (KAKENHI, No. 19580074 to MW).

Conflict of interest

None declared.

Acknowledgments

The MALDI-TOF-MS analysis was carried out at the OPEN FACILITY, Hokkaido University Sousei Hall.

References

1. Balcar, V. J., Johnston, G. A., and Twitchin, B. (1977) Stereospecificity of the inhibition of L-glutamate and L-aspartate high affinity uptake in rat brain slices by *threo*-3-hydroxyaspartate. *J. Neurochem.* **28**, 1145-1146
2. Shigeri, Y., Shimamoto, K., Yasuda-Kamatani, Y., Seal, R. P., Yumoto, N., Nakajima, T., and Amara, S. G. (2001) Effects of *threo*- β -hydroxyaspartate derivatives on excitatory amino acid transporters (EAAT4 and EAAT5). *J. Neurochem.* **79**, 297-302
3. Kaneko, T. and Katsura, H. (1963) The synthesis of four optical isomers of β -hydroxyaspartic acid. *Bull. Chem. Soc. Jpn.* **36**, 899-903
4. Lebrun, B., Sakaitani, M., Shimamoto, K., Yasuda-Kamatani, Y., and Nakajima, T. (1997) New β -hydroxyaspartate derivatives are competitive blockers for the bovine glutamate/aspartate transporter. *J. Biol. Chem.* **272**, 20336-20339
5. Gibbs, R. G. and Morris, J. G. (1964) Assay and properties of β -hydroxyaspartate aldolase from *Micrococcus denitrificans*. *Biochim. Biophys. Acta* **85**, 501-503
6. Kornberg, H. L. and Morris, J. G. (1965) The utilization of glycollate by *Micrococcus denitrificans*: the β -hydroxyaspartate pathway. *Biochem. J.* **95**, 577-586
7. Gibbs, R. G. and Morris, J. G. (1965) Purification and properties of *erythro*- β -hydroxyaspartate dehydratase from *Micrococcus denitrificans*. *Biochem. J.* **97**, 547-554
8. Wada, M., Matsumoto, T., Nakamori, S., Sakamoto, M., Kataoka, M., Liu, J.Q., Itoh, N., Yamada, H., and Shimizu, S. (1999) Purification and characterization of a novel enzyme, L-*threo*-3-hydroxyaspartate dehydratase, from *Pseudomonas* sp. T62. *FEMS Microbiol. Lett.* **179**, 147-151

9. Murakami, T., Maeda, T., Yokota, A., and Wada, M. (2009) Gene cloning and expression of pyridoxal 5'-phosphate-dependent L-threo-3-hydroxyaspartate dehydratase from *Pseudomonas* sp. T62, and characterization of the recombinant enzyme. *J. Biochem.* **145**, 661-668
10. Wada, M., Nakamori, S., and Takagi, H. (2003) Serine racemase homologue of *Saccharomyces cerevisiae* has L-threo-3-hydroxyaspartate dehydratase activity. *FEMS Microbiol. Lett.* **225**, 189-193
11. Robinson, W. G. and Labow, R. (1971) D-Serine dehydratase (*Escherichia coli*). *Meth. Enzymol.* **17B**, 356-360.
12. Panizzutti, R., de Souza Leite, M., Pinheiro, C. M., and Meyer-Fernandes, J. R. (2006) The occurrence of free D-alanine and an alanine racemase activity in *Leishmania amazonensis*. *FEMS Microbiol. Lett.* **256**, 16-21
13. Uo, T., Yoshimura, T., Shimizu, S., and Esaki, N. (1998) Occurrence of pyridoxal 5'-phosphate-dependent serine racemase in silkworm, *Bombyx mori*. *Biochem. Biophys. Res. Commun.* **246**, 31-34
14. Shibata, K., Watanabe, T., Yoshikawa, H., Abe, K., Takahashi, S., Kera, Y., and Yamada, R. (2003) Purification and characterization of aspartate racemase from the bivalve mollusk *Scapharca broughtonii*. *Comp. Biochem. Physiol. Part B* **134**, 307-314
15. Bradford, M. M. (1976) A rapid and sensitive method for quantitation of microgram quantities of proteins utilizing the principle of protein-dye binding. *Anal. Biochem.* **72**, 248-254

16. Wada, M., Yoshizumi, A., Nakamori, S., and Shimizu, S. (1999) Purification and characterization of monovalent cation-activated levodione reductase from *Corynebacterium aquaticum* M-13. *Appl. Environ. Microbiol.* **65**, 4399-4403
17. Pearson, W. R., and Lipman, D. J. (1988) Improved tools for biological sequence comparison. *Proc. Natl. Acad. Sci. USA* **85**, 2444-2448
18. Thompson, J. D., Higgins, D. G., and Gibson, T. J. (1994) CLUSTAL W: Improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673-4680
19. Wen, A., Fegan, M., Hayward, C., Chakraborty, S., Sly, L.I. (1999) Phylogenetic relationships among members of the *Comamonadaceae*, and description of *Delftia acidovorans* (den Dooren de Jong 1926 and Tamaoka *et al.* 1987) gen. nov., comb. nov. *Int. J. Syst. Bacteriol.*, **49**, 567-576
20. Tanaka, H., Yamamoto, A., Ishida, T., and Horiike, K. (2008) D-Serine dehydratase from chicken kidney: a vertebral homologue of the cryptic enzyme from *Burkholderia cepacia*. *J. Biochem.* **143**, 49-57
21. Ito, T., Hemmi, H., Kataoka, K., Mukai, Y., and Yoshimura, T. (2008) A novel zinc-dependent D-serine dehydratase from *Saccharomyces cerevisiae*. *Biochem. J.* **409**, 399-406
22. Hulo, N., Bairoch, A., Bulliard, V., Cerutti, L., De Castro, E., Langendijk-Genevaux, P. S., Pagni, M., and Sigrist, C. J. A. (2006) The PROSITE database. *Nucleic Acids Res.* **34**, D227-D230

23. Liu, J. Q., Dairi, T., Itoh, N., Kataoka, M., and Shimizu, S. (2003) A novel enzyme, D-3-hydroxyaspartate aldolase from *Paracoccus denitrificans* IFO 13301: purification, characterization, and gene cloning. *Appl. Microbiol. Biotechnol.* **62**, 53-60
24. Eliot, A. C, Kirsch, J. F. (2004) Pyridoxal phosphate enzymes: mechanistic, structural, and evolutionary considerations. *Annu. Rev. Biochem.* **73**, 383-415
25. Schneider, G., Käck, H., and Lindqvist, Y. (2000) The manifold of vitamin B₆ dependent enzymes. *Structure* **8**, R1-6
26. Kuramitsu, S., Inoue, Y., Tanase, S., Morino, Y., and Kagamiyama, H. (1987) Substitution of an arginyl residue for the active site lysyl residue (Lys258) of aspartate aminotransferase. *Biochem. Biophys. Res. Commun.*, 146, 416-421
27. Kataoka, M., Ikemi, M., Morikawa, T., Miyoshi, T., Nishi, K., Wada, M., Yamada, H., and Shimizu, S. (1997) Isolation and characterization of D-threonine aldolase, a pyridoxal-5'-phosphate-dependent enzyme from *Arthrobacter* sp. DK-38. *Eur. J. Biochem.* **248**, 385-393
28. Liu, J.-Q., Dairi, T., Itoh, N., Kataoka, M., Shimizu, S., and Yamada, H. (1998) A novel metal-activated pyridoxal enzyme with a unique primary structure, low specificity D-threonine aldolase from *Arthrobacter* sp. strain DK-38. *J. Biol. Chem.* **273**, 16678-16685

Figure legends

Fig. 1 Molecular weight analysis of D-THA DH from *Delftia* sp. HT23.

SDS-PAGE analysis of D-THA DH. Lane (a), molecular weight markers; Lane (b), purified D-THA DH.

Fig. 2 Absorption spectrum of the recombinant enzyme.

Absorption spectra were obtained with a Beckman DU-800 spectrophotometer. The recombinant enzyme (0.2 mg/mL) was dissolved in 10 mM Tris-HCl buffer (pH 8.0) containing 0.01 mM PLP, 0.1 mM MnCl₂, and 0.1 mM dithiothreitol. Solid line, wild-type enzyme; dotted line, K43A mutant enzyme.

Fig. 3 Effect of metal ions on EDTA-treated D-THA DH. MnCl₂, CaCl₂, ZnCl₂, CoCl₂,

FeCl₂, and NiCl₂ were added to the reaction mixture at various concentrations, and incubated for 10 minutes. The activity of non-EDTA-treated enzyme, which dialyzed against 100 μM MnCl₂, was taken as 100%. Values are means of two independent experiments. Symbols: open circle, Mn²⁺; filled circle, Ca²⁺; open triangle, Zn²⁺; filled triangle, Co²⁺; open square, Fe²⁺; filled square, Ni²⁺.

Fig. 4 Multiple alignment of the amino acid sequences of D-THA DH from *Delftia* sp.

HT23 (D-THADH *Delftia*), D-serine dehydratase from chicken (DSD chicken), and D-

serine dehydratase from *S. cerevisiae* (DSD Yeast).

The alignment was generated with Clustal W 1.83 and BOXSHADE 3.21. The numbers on the left are the residue numbers for each sequence. White letters on a black background indicate identical residues, and white letters on a gray background indicate similar residues. The asterisk indicates the PLP-binding residue.

Table I. Purification of D-THA DH from *Delftia* sp. HT23.

Step	Total Protein (mg)	Total Activity (U)	Specific Activity (U/mg)	Purification (fold)	Yield (%)
Cell extract	1390	255	0.18	1.0	100
Ammonium Sulfate	346	129	0.37	2.0	50.5
Hi-Prep Q FF	73.5	109	1.5	8.0	42.6
Hi-Trap phenyl	7.92	58.0	7.3	39.8	22.8
Superdex-200	2.24	33.3	14.8	80.6	13.1
Resource Q	0.72	14.9	20.7	112.4	5.8
Hi-Trap butyl	0.17	3.6	21.3	115.8	1.4

Table II. **Kinetic parameters of D-THA DH.**

Substrate	k_{cat} (s^{-1})	K_{m} (mM)	$k_{\text{cat}}/K_{\text{m}}$
D-THA	10.93	0.42	25.96
L-THA	3.03	6.16	0.49
D-EHA	N.D.	-	-
L-EHA	8.68	0.16	54.25
D-Serine	0.89	0.15	5.91
L-Serine	0.18	38.70	0.0047

1 N.D., below the detection limit, *i.e.*, <0.01 $\mu\text{mol} / \text{min}/(\text{mg protein})$

↑ Fig. 1 Maeda et al.

↑ **Fig. 2 Maeda et al.**

D-THA DH Delftia	1	-----MQDTLLTLDTPAAVIDLDRMQRNIARMQQRMDAAG-----VRLRPHVKTSK*
DSD Chicken	1	-----MWLGALLDTPALPTIDRTTARRNABRMRECRALG-----VRLRPHVKTHK
DSD Yeast	1	MSDVLSQYKGCSSVRDLPTPNFVINEEKFDKNCCTMLNNVEKLSQECGVPFKFRAHVKTHK
D-THA DH Delftia	47	SVPVAAAQRAA-----GASGITVSTLKEAEQFFAAG-----TTDILYANSMAP-HR
DSD Chicken	49	TLEGGLLATGG-----TRRGIAVSTLAEARFFADGG-----FDDILLAYPVPT-AR
DSD Yeast	61	TAKGTLKQLGHGLPLAKRTTRAILVSTLKEAEELLNYQDRQCSYIDDITYSLPCCVPEF
D-THA DH Delftia	92	LPQALQLRRRGCDLKLIVDSVAAAQAIAAFGR-EQGEAFVWIEIDTDGHRSGVGADDTTP
DSD Chicken	94	LEECAGLAARRLDAFHVLLDRPEALASLRQRPL-CHGKRWLVLKLDGNGRAGVRPTDPA
DSD Yeast	121	TPLLSNLSRRRVNNEQVFVDNIEHLENLKNFGRPASGKKWSVFIKVDMGTKRAGLAFDSPE
D-THA DH Delftia	151	LLLAIGRTLHDG---GMRLGGVLTHAGSSYELDTPEALQALAERERAGCVQAAEALRAAG
DSD Chicken	153	ALELAQAIALNDAP-EEVTLVGVIYAHCGNTYGCSGADTIQAIARTTTNAVLSTFVAALRAAG
DSD Yeast	181	FLSLLKKLTSSEIKFVIEPYGFYAHAGHSYSSTINDTONLLMEEVKAVNSAAKVLCSVD
D-THA DH Delftia	208	LPCP---VVSVGSTP-----TALAASRLDGVTEVRAGVYVFFDLVMRNIGV
DSD Chicken	212	VPCP---QASIGSTPS-----CSHPPEMSQLTELHPGNYIFYDLQQTQLGS
DSD Yeast	241	PQFDPSKLTLSVGATPTNSNLKLDNKSTLVKFITTQLVSTLEIHCGRNYCMYDLQQVATGC
D-THA DH Delftia	251	CAAEDVALSVLATVIGHQADKGWAIVDAGWMMSRDRGTARQKQDFGYGQVCDLQGRVMP
DSD Chicken	256	CQPQDVATRVLTRVIGHYAHRGQLLVDCGWAALS LHAGAGQ---GPQGCAATDG--HP
DSD Yeast	301	VQDHELSGFVLGTVLSSYPSRGELLNTGVMCLTREASSIKGFGICADLEHVLKSESEFSR
D-THA DH Delftia	311	GEVLTGANQEHGILARADGAABADIATRFPLGTRLRLPNHACATGAQFPAYQALA-ADG
DSD Chicken	310	ELRLVGLTQEHGLLEHAGGQMDFG---RFPVGSVLALIPYHACATAAMHPVYVVE-ECK
DSD Yeast	361	ENYVARVSQEHGILRPIRNWNETTP---LKLGSKI AVLPPHACITMGQFPYYPVVNSEGI
D-THA DH Delftia	370	SVQTWERLHGW
DSD Chicken	366	VVALWHPVRGW
DSD Yeast	418	VNDVWLPFQKW

↑ Fig. 4 Maeda et al.