Title	Occurrence of all-cis-5,8,11,14,17,20,23-hexacosaeptaenoic acid (26:7n-3) in roughscale sole <i>Clidoderma asperrimum</i> flesh lipids
Author(s)	Fukuda, Yuuki; Ando, Yasuhiro
Citation	Fisheries Science, 77(5), 875-882 https://doi.org/10.1007/s12562-011-0387-y
Issue Date	2011-09
Doc URL	http://hdl.handle.net/2115/47167
Rights	© 2011 公益社団法人日本水産学会; © 2011 The Japanese Society of Fisheries Science; The final publication is available at www.springerlink.com
Type	article (author version)
File Information	Fukuda&Ando.pdf

[Instructions for use](#)

1 Occurrence of All-*cis*-5,8,11,14,17,20,23-Hexacosaeptaenoic Acid (26:7n-3) in the Roughscale
2 Sole *Clidoderma asperrimum* Flesh Lipids

7 Yuuki Fukuda and Yasuhiro Ando*

11 Division of Marine Life Sciences, Faculty of Fisheries Science, Hokkaido University,
12 Hakodate, Hokkaido 041-8611, Japan

18 *Correspondence should be addressed to:

19 Yasuhiro Ando

20 Division of Marine Life Sciences, Faculty of Fisheries Science, Hokkaido University,
21 3-1-1 Minato-cho, Hakodate, Hokkaido 041-8611, Japan

22 Phone: +81-138-40-8803

23 Fax: +81-138-40-8802

24 E-mail: ando@fish.hokudai.ac.jp

Abstract

Fatty acid analysis of roughscale sole *Clidoderma asperrimum* flesh lipids was carried out by gas chromatography. An unidentified peak appeared in the chromatogram in the elution region of $\geq C_{24}$ fatty acids. After enrichment by solvent partitioning, reversed-phase TLC, and argentation TLC, the peak component was subjected to structural analyses. The partially hydrogenated products after reaction with hydrazine hydrate gave seven isomers of *cis*-hexacosenoic acid (26:1). GC-MS analysis of their dimethyl disulfide (DMS) adducts identified the monounsaturates as 5-, 8-, 11-, 14-, 17-, 20-, and 23-26:1. The peak component was assigned to all-*cis*-5,8,11,14,17,20,23-hexacosaeptaenoic acid (26:7n-3). GC-MS analyses of the 4,4-dimethyloxazoline (DMOX) derivative and methyl ester confirmed this structure. This fatty acid is a rare, very long chain polyunsaturated fatty acid (VLCPUFA). The concentrations of the acid found in roughscale sole were 0.69 ± 0.34 % ($N=5$) of the total fatty acids in the flesh lipids. Roughscale sole appears to be characterized by the occurrence of 26:7n-3, which is lacking in popular sources of methylene-interrupted VLCPUFA, such as vertebrate retina, spermatozoa, and herring.

Keywords

Hexacosaeptaenoic acid, Very long chain polyunsaturated fatty acid, Fatty acid, Roughscale sole, *Clidoderma asperrimum*, GC, GC-MS

Introduction

Very long chain polyunsaturated fatty acids (VLCPUFA) occur in many species of animals, plants, and lower organisms [1–4]. Mammalian retinas [5,6], brains [7–9], testes [10], and spermatozoa [11–12] include methylene-interrupted VLCPUFA of the n-3 and n-6 series up to C34–C40. Aquatic organisms also contain VLCPUFA [13]. Baltic herring include those up to C28 [14–16], some of which are represented in the fatty acids of the predatory ringed seal [17]. Bottom-living flathead flounder is rich in 24:6n-3 [18,19], which presumably originated from their diet of brittle star [20–22]. Additionally, dinoflagellates and some species of microalgae include 28:8n-3 and 28:7n-6 [4], and even- and odd-numbered VLCPUFA up to C36 were observed in a species of dinoflagellate [23,24].

All-*cis*-5,8,11,14,17,20,23-hexacosaeptaenoic acid (26:7n-3) is also a VLCPUFA. However, this fatty acid is a very rare one among the VLCPUFA of n-3 series. To the best of our knowledge, 26:7n-3 has been found in only 6 species of microalgae (Cryptophyceae, Prymnesiophyceae and Dinophyceae) at less than 0.3% of the total fatty acid content [23–25]. This fatty acid also cannot be prepared from popular sources of n-3 and n-6 VLCPUFA standards such as retinas, sperm, and herring [1]. A detailed assignment of the structure has also not been reported.

In the present study, the fatty acids of roughscale sole *Clidoderma asperrimum*, i.e., ‘samegarei’ in Japanese, which is one of the edible fish in Japan, were investigated for VLCPUFA. The fatty acid analysis of the sole revealed the occurrence of 0.3%–1.3% of 26:7 in their flesh lipids. This paper reports the structural assignment of this fatty acid as 26:7n-3 along with their enrichment process and the fatty acid composition of the roughscale sole.

Materials and methods

Materials

Five individuals of roughscale sole caught in the Pacific waters off Hidaka, Hokkaido, Japan, and in the Sea of Okhotsk off Shari, Hokkaido, Japan, were purchased in stores in May 2009. Two of the Pacific samples were obtained as whole fish (females; body lengths, 39.8 and 41.2 cm; body weights, 2.4 and 2.8 kg) and another in frozen form without skin and viscera (body length, 34.8 cm). The two Okhotsk samples were obtained in frozen form without the head, skin, or viscera (lengths of trunk and tail, 24.0 and 25.8 cm). The flesh was removed, minced, and stored at -30°C before lipid extraction.

Fatty acid methyl esters

Total lipids were extracted from 150 g of the flesh by the method reported by Bligh and Dyer [26]. Fatty acid methyl esters were prepared from the lipids by transesterification with 7% BF_3 -methanol for 1 h at 100°C under a nitrogen atmosphere. Methyl esters were purified by thin-layer chromatography (TLC) on silica gel G plates (10×10 cm, 0.25 mm thickness; Analteck, Newark, USA) with hexane/diethyl ether (85:15, v/v) for development.

For quantity preparation, the total lipids were saponified by refluxing them with 1 M KOH in ethanol for 1 h; the unsaponifiable portion was extracted with diethyl ether. Following acidification of the aquatic phase using dilute HCl, the fatty acids were recovered by ether extraction. The fatty acids were converted to methyl esters by refluxing them with 7% BF_3 -methanol at 70°C for 14 min.

Fractionation of fatty acids

The fatty acid methyl esters were divided into two fractions obtained through a solvent partition method using a solvent system of 2,2,4-trimethylpentane and ethanol/water (1:1, v/v) containing 0.25 g/ml of silver nitrate [27,28].

The methyl esters were fractionated according to their partition number by reversed-phase

TLC (RP-TLC) on Partisil KC18F plates (20 × 20 cm, 0.2 mm thickness; Whatman, Maidstone, England) with acetonitrile/water (95:5, v/v) for double developments [29].

The methyl esters were fractionated according to the degree of unsaturation by argentation TLC (Ag-TLC) on 10% silver nitrate-impregnated layers of silica gel 60G (20 × 20 cm, 0.5 mm thickness; Merck, Darmstadt, Germany) with hexane/acetone (70:30, v/v) for double developments [29].

Derivatization for structural analysis

Partial hydrogenation of the polyunsaturated fatty acids was carried out using hydrazine hydrate [30,31]. A mixture of 5 mg of free fatty acids and 10% (v/v) hydrazine hydrate in methanol (5 ml) was stirred at 50°C for 7 h with aeration. The products, which were extracted with diethyl ether, were converted to methyl esters using 7% BF₃-methanol. The monounsaturated fatty acids were isolated from the products by Ag-TLC on 5% silver nitrate-impregnated silica gel 60G with hexane/acetone (95:5, v/v), and then fractionated according to the olefinic bond position by Ag-TLC on 15% silver nitrate-impregnated silica gel 60G with hexane/toluene (50:50, v/v) [32,33].

Dimethyl disulfide (DMDS) adducts of monounsaturated fatty acids were prepared following the procedure of Shibahara *et al.* [34,35]. The methyl esters were reacted with DMDS (1 ml) in the presence of catalytic I₂ (13 mg) for 1 h at 35°C before adding 30% aqueous NaHSO₃. The resulting adducts that were extracted by hexane/diethyl ether (50:50, v/v) were purified by TLC on a silica gel G plate with hexane/diethyl ether/acetic acid (80:20:1, v/v/v) for development.

4,4-Dimethyloxazoline (DMOX) derivatives of fatty acids were prepared by the procedure reported by Christie [36]. Acid chlorides formed from the free fatty acids by a reaction with oxalyl chloride were reacted with a 10 mg/ml solution of 2-amino-2-methyl-1-

propanol in dichloromethane (0.5 ml) for 1 h at room temperature. After the solvent was evaporated, trifluoroacetic anhydride was added to the residue and the mixture was left at 40°C for 1 h. The DMOX derivatives were purified by TLC on a silica gel G plate with hexane/diethyl ether/acetic acid (50:50:1, v/v/v) for development.

Instrumental analysis

The fatty acid methyl esters were analyzed by GC using a Shimadzu GC-18A gas chromatograph (Shimadzu, Kyoto, Japan) equipped with a flame ionization detector and a Restek FAMEWAX column (30 m × 0.32 mm i.d., 0.25 µm film thickness; Restek, Bellefonte, USA). The column temperature was programmed to either increase from 170 to 240°C at a rate of 4°C/min, or to remain isothermal at 240°C. The injector and detector temperatures were 240°C, and the carrier gas was helium (85 kPa). Peak area percentages were obtained using a Shimadzu C-R6A integrator.

The monounsaturated fatty acids isolated from the hydrazine hydrogenation products were analyzed using a Shimadzu GC-17A gas chromatograph equipped with a flame ionization detector and a SLB-IL100 column (60 m × 0.32 mm i.d., 0.26 µm film thickness; Supelco, Bellefonte, USA) [37]. The column temperature was isothermal at 200°C, the injector and detector temperatures were 240°C, and the carrier gas was helium (117.5 kPa). The peaks were monitored using a Shimadzu C-R3A integrator.

GC-MS analysis was carried out using an HP 6890 series gas chromatograph (Hewlett-Packard, Palo Alto, USA) linked to a JEOL JMS-700TZ mass spectrometer (JEOL, Tokyo, Japan). The latter was used in the electron impact mode at 70 eV with source temperatures of 240°C for methyl esters, 270°C for DMOX derivatives and 280°C for DMDS adducts. The GC was fitted with split/splitless injection. For the analyses of the methyl esters and DMOX derivatives, a DB-23 column (30 m × 0.25 mm i.d., 0.25 µm film thickness; Agilent, Santa

Clara, USA) was used. The column temperatures were programmed from 40°C to 230°C and from 40°C to 270°C at 20°C/min for the methyl esters and DMOX, respectively. For the analysis of the DMDS adducts, a Zebron ZB-1ms column (30 × 0.25 mm i.d., 0.25 µm film thickness; Phenomenex, Torrance, USA) was used. The column temperature was held at 40°C for 1 min then raised to 175°C at 40°C/min and then to either 265°C at 5°C/min or to 280°C at 20°C/min. Helium was used as the carrier gas.

Fourier transform infrared spectra were measured in CCl₄ solutions using a JASCO FT-IR 5300 spectrometer (JASCO, Tokyo, Japan).

Fatty acid composition determined by GC of methyl esters was represented by mean ± standard deviation of the five individuals of roughscale sole.

Results

GC and enrichment of 26:7

The GC analysis of the fatty acid methyl esters derived from roughscale sole flesh lipids showed two remarkable peaks, A and B, after elution of 24:1n-9 (Fig. 1). Their equivalent chain lengths (ECLs) on FAMEWAX at 240°C were 26.14 and 28.30, for peak A and B, respectively. The ECL of peak A was in fair agreement with that of 24:6n-3, which was previously found in flathead flounder [18,19] and brittle star [20,21]. The component of peak B was confirmed to be hexacosaeptaenoic acid (26:7) since the mass number of the molecular ion peak obtained under high resolution conditions agreed with that calculated for a 26:7 methyl ester; the *m/z* found was 396.30280 and the value calculated was 396.30283 for C₂₇H₄₀O₂.

For the structural analysis, the enrichment of 26:7 was conducted by the solvent partition procedure, RP-TLC, and 10% Ag-TLC in this order (Table 1). The concentration of 26:7, which started at 1.3% of the total fatty acids, was boosted to 10.4% by the solvent partition, 27.7% by RP-TLC, and finally 83.0% by Ag-TLC. A coexistent minor component of the final fraction was

24:6n-3 (17.0%).

Structure of 26:7

The infrared spectrum of the 26:7 concentrate showed absorptions at 1650 cm^{-1} and 710 cm^{-1} , but not that near 970 cm^{-1} which is characteristic of a *trans*-olefinic bond. Hydrazine hydrogenation yielded *n*-26:0 as a saturated fatty acid. These results indicate that the fatty acid is normal-chain 26:7 with all *cis*-geometry.

The monounsaturated fatty acids, which were produced by the partial hydrogenation of 26:7, separated into four fractions on the 15% Ag-TLC plate. The GC analyses of these fractions showed a total of seven peaks corresponding to 26:1 isomers with ECLs in the range of 26.41–27.23 on SLB-IL100 at 200°C. The mass spectra of the DMDS adducts of the 26:1 isomers gave a molecular ion at m/z 502 corresponding to the DMDS adduct of 26:1 methyl ester and a series of key fragment ions showing the olefinic bond position in 26:1 (Fig. 2). In Fig. 2a, the fragment ions at m/z 161 and 341 indicate cleavage between the methylthio-substituted carbons of C5 and C6. The fragment ion at m/z 129 was due to the loss of methanol (m/z 32) from the ion at m/z 161. A set of the fragment ions indicated the structure of 5-26:1. In the same manner, the other isomers were identified as 8-, 11-, 14-, 17-, 20- and 23-26:1, as shown in Figs. 2b-2g. Hydrazine reduces olefinic bonds without positional and geometrical isomerization of the remaining olefinic bonds [30]. The structure of 26:7 was assigned as all-*cis*-5,8,11,14,17,21,23-hexacosaeptaenoic acid (26:7n-3).

The mass spectrum of the DMOX derivative of 26:7 showed irregular intervals of m/z 12 between the maxima in the fragment ion peaks for each carbon atom as follows: C7 (m/z 180, intensity 19.0%)–C8 (m/z 192, 9.7%); C10 (m/z 220, 23.2%)–C11 (m/z 232, 15.0%); C13 (m/z 260, 16.5%)–C14 (m/z 272, 8.6%); C16 (m/z 300, 13.7%)–C17 (m/z 312, 7.3%); and C19 (m/z 340, 15.6%)–C20 (m/z 352, 9.8%) (Fig. 3a). These fragments indicate the occurrence of olefinic

bonds at the $\Delta 8$, 11, 14, 17, and 20 positions in 26:7 [29]. An olefinic bond at the $\Delta 5$ position was shown by the fragment ion due to cleavage at this position (m/z 152, 17.6%) accompanied by an intense odd-numbered peak at m/z 153 (32.3%) [38]. GC-MS analysis of the 26:7 methyl ester revealed fragment ions characteristic of n-3 and $\Delta 5$ series polyunsaturated fatty acids at m/z 108 (33.4%) and 180 (14.1%), respectively [29], confirming the structure of 26:7n-3 (Fig. 3b).

Fatty acids of the roughscale sole flesh lipids

The fatty acid composition of the roughscale sole flesh lipids is shown in Table 2. The major fatty acids (>5% of the total lipids) were 14:0, 16:0, 16:1n-7, 18:1n-9+18:1n-11, 20:1n-11+20:1n-13, and 20:5n-3. The proportions of these fatty acids were not very different from those previously reported for deep-sea flounders [39], in which the monounsaturated fatty acids were rich in the liver and flesh neutral lipids. The major highly unsaturated fatty acids of the roughscale sole were 20:5n-3, 22:5n-3, 22:6n-3, 24:6n-3, and 26:7n-3. The content of 26:7n-3 was $0.69 \pm 0.34\%$ of the total lipids, ranging from 0.33% to 1.26% among the five individuals. The lipid content of the flesh was $30.5 \pm 4.6\%$ on wet-weight base, much higher than the previous datum (5.6%) observed for smaller-sized roughscale sole (mean body length, 22.3 cm; mean body weight, 199.8 g) [39].

Discussion

In the present study, 26:7n-3 and 24:6n-3 were found in the roughscale sole flesh lipids as their VLCPUFA. Methylene-interrupted VLCPUFA of n-3 series are typical of vertebrate retinas (up to 36:6n-3) [5,6], spermatozoa (up to 34:6n-3 in chain length and 32:7n-3 in unsaturation) [11,12], and Baltic herring (generally up to 28:7n-3) [14–16]. These tissues or fish have been recognized as convenient sources of VLCPUFA standards for the n-3 and n-6 series [1]. One of

the shorter-chain VLCPUFA, 24:6n-3, was rich in flathead flounder [18,19] and brittle stars [20–22], and was also found in sea lilies [20], coelenterates [40,41], gorgonians [42], jellyfish [43], and gastropods [44]. Freshwater crustacea of the order Bathynellacea were reported to contain more than 50 VLCPUFA up to 40:8n-3 [45,46]. Marine dinoflagellates contain 28:8n-3 together with 28:7n-6 [23–25,47–53]. Fatty acids of the dinoflagellate *Amphidinium carterae* included even- and odd-numbered chain VLCPUFA up to 36:8n-3 [23,24].

As a tentatively identified component, 26:7n-3 were found in Dinophyceae (*Heterocapsa niei*, 0.2% of total fatty acids; and *Amphidinium carterae*, in trace amounts), Prymnesiophyceae (*Pavlova pinguis*, up to 0.3%), and Cryptophyceae (*Proteomonas sulcata* and *Phodomonas salina*, each in trace amounts) [25]. In the fatty acids of the dinoflagellate *Amphidinium carterae*, 26:7n-3 was found at a concentration of 3.2% in the heptaenoic + octaenoic acid concentrate [23], which corresponds to 0.064% of the total fatty acids. The same acid was also found at 1.40% in a concentrate of VLCPUFA with more than 3 olefinic bonds [24]. To the best of our knowledge, there has been no other report describing the occurrence of 26:7n-3 in nature. While the analogous 26:6n-3 was observed in vertebrate retina [5,6], murine testis [10], and ram spermatozoa [11], the fatty acids of these tissues were not reported to include 26:7n-3. Baltic herring [15], seal [17], and crustacea (Bathynellacea) [45,46] contain both analogous 26:6n-3 and homologous 28:7n-3, but not 26:7n-3.

Therefore, the roughscale sole flesh lipids are characterized by the occurrence of the rare VLCPUFA, 26:7n-3. This fatty acid seems to be formed from coexistent 24:6n-3 via two-carbon chain elongation followed by Δ^5 -desaturation. In GC analysis, unidentified minor peaks were observed between the peaks of 24:6n-3 and 26:7n-3 (Fig. 1). Roughscale sole preferentially feed on brittle star [54,55], which is usually rich in 24:6n-3 [20–22]. At this time, the occurrence of 26:7n-3 in brittle star is unknown. Although the concentrations of 26:7n-3 in the flesh lipids (0.3–1.3%) were not very high, there was no sample where 26:7n-3 was not found.

Acknowledgments

The authors wish to thank Ms. Seiko Oka, Instrumental Analysis Division, Equipment Management Center, Creative Research Institution, Hokkaido University, for the GC-MS analyses.

References

1. Řezanka T (1989) Very-long-chain fatty acids from the animal and plant kingdoms. *Prog Lipid Res* 28:147-187
2. Poulos A (1995) Very long chain fatty acids in higher animals-a review. *Lipids* 30:1-14
3. Řezanka T, Votruba J (2002) Chromatography of very long-chain fatty acids from animal and plant kingdoms. *Anal Chim Acta* 465:273-297
4. Řezanka T, Sigler K (2009) Odd-numbered very-long-chain fatty acids from the microbial, animal and plant kingdoms. *Prog Lipid Res* 48:206-238.
5. Aveldaño MI, Sprecher H (1987) Very long chain (C₂₄ to C₃₆) polyenoic fatty acids of the n-3 and n-6 series in dipolyunsaturated phosphatidylcholines from bovine retina. *J Biol Chem* 262:1180-1186
6. Aveldaño MI (1987) A novel group of very long chain polyenoic fatty acids in dipolyunsaturated phosphatidylcholines from vertebrate retina. *J Biol Chem* 262:1172-1179
7. Poulos A, Sharp P, Singh H, Johnson D, Fellenberg A, Pollard A (1986) Detection of a homologous series of C26-C38 polyenoic fatty acids in the brain of patients without peroxisomes (Zellweger's syndrome). *Biochem J* 235:607-610
8. Sharp P, Poulos A, Fellenberg A, Johnson D (1987) Structure and lipid distribution of polyenoic very-long-chain fatty acids in the brain of peroxisome-deficient patients

- (Zellweger syndrome). Biochem J 248:61-67
9. Poulos A, Sharp P, Johnson D, Easton C (1988) The occurrence of polyenoic very long chain fatty acids with greater than 32 carbon atoms in molecular species of phosphatidylcholine in normal and peroxisome-deficient (Zellweger's syndrome) brain. Biochem J 253:645-650
10. Furland NE, Maldonado EN, Aveldaño MI (2003) Very long chain PUFA in murine testicular triglycerides and cholesterol esters. Lipids 38:73-80
11. Poulos A, Sharp P, Johnson D, White I, Fellenberg A (1986) The occurrence of polyenoic fatty acids with greater than 22 carbon atoms in mammalian spermatozoa. Biochem J 240:891-895
12. Furland NE, Oresti GM, Antollini SS, Venturino A, Maldonado EN, Aneldaño MI (2007) Very long-chain polyunsaturated fatty acids are the major acyl groups of sphingomyelins and ceramides in the head of mammalian spermatozoa. J Biol Chem 282:18151-18161
13. Ackman RG (1989) Fatty acids. In: Ackman RG (ed) Marine biogenic lipids, fats, and oils, vol 1. CRC Press, Boca Raton, pp 103-137
14. Linko RR, Karinkanta H (1970) Fatty acids of long chain length in Baltic herring lipids. J Am Oil Chem Soc 47:42-46
15. Řezanka T (1990) Identification of very long polyenoic acids as picolinyl esters by Ag⁺ ion-exchange high-performance liquid chromatography, reversed-phase high-performance liquid chromatography and gas chromatography-mass spectrometry. J Chromatogr 513:344-348
16. Kallio H, Vauhkonen T, Linko RR (1991) Thin-layer silver ion chromatography and supercritical fluid chromatography of Baltic herring (*Clupea harengus membras*) triacylglycerols. J Agric Food Chem 39:1573-1577
17. Käkälä R, Ackman RG, Hyvärinen H (1995) Very long chain polyunsaturated fatty acids in

- the blubber of ringed seals (*Phoca hispida* sp.) from Lake Saimaa, Lake Ladoga, the Baltic Sea, and Spitsbergen. *Lipids* 30:725-731
18. Ota T, Chihara Y, Itabashi Y, Takagi T (1994) Occurrence of all-*cis*-6,9,12,15,18,21-tetracosahexaenoic acid in flatfish lipids. *Fish Sci* 60:171-175
19. Tomita Y, Ando Y (2009) Reinvestigation of positional distribution of tetracosahexaenoic acid in triacyl-*sn*-glycerols of flathead flounder flesh. *Fish Sci* 75:445-451
20. Takagi T, Kaneniwa M, Itabashi Y (1986) Fatty acids in Crinoidea and Ophiuroidea: occurrence of all-*cis*-6,9,12,15,18,21-tetracosahexaenoic acid. *Lipids* 21:430-433
21. Sato D, Ando Y, Tsujimoto R, Kawasaki K (2001) Identification of novel nonmethylene-interrupted fatty acids, 7*E*,13*E*-20:2, 7*E*,13*E*,17*Z*-20:3, 9*E*,15*E*,19*Z*-22:3, and 4*Z*,9*E*,15*E*,19*Z*-22:4, in Ophiuroidea (brittle star) lipids. *Lipids* 36:1371-1375
22. Mansour MP, Holdsworth DG, Forbes SE, Macleod CK, Volkman JK (2005) High contents of 24:6(n-3) and 20:1(n-13) fatty acids in the brittle star *Amphiura elandiformis* from Tasmanian coastal sediments. *Biochem Syst Ecol* 33:659-674
23. Řezanka T, Nedbalová L, Sigler K (2008) Identification of very-long-chain polyunsaturated fatty acids from *Amphidinium carterae* by atmospheric pressure chemical ionization liquid chromatography-mass spectroscopy. *Phytochemistry* 69:2391-2399
24. Řezanka T, Nedbalová L, Sigler K (2008) Odd-numbered very-long-chain polyunsaturated fatty acids from the dinoflagellate *Amphidinium carterae* identified by atmospheric pressure chemical ionization liquid chromatography-mass spectrometry. *Phytochemistry* 69:2849-2855
25. Mansour MP, Frampton DMF, Nichols PD, Volkman JK, Blackburn SI (2005) Lipid and fatty acid yield of nine stationary-phase microalgae: applications and unusual C₂₄-C₂₈ polyunsaturated fatty acids. *J Appl Phycol* 17:287-300
26. Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J*

Biochem Physiol 37:911-917

27. Christie WW (2003) Lipid analysis, 3rd edn. The Oily Press, Bridgwater
28. Peers KE, Coxon DT (1986) Simple enrichment procedure for the estimation of minor polyunsaturated fatty acids in food fats. J Food Technol 21:463-469
29. Christie WW, Han X (2010) Lipid Analysis, 4th edn. The Oily Press, Bridgwater
30. Ratnayake WMN, Grossert JS, Ackmen RG (1990) Studies on the mechanism of the hydrazine reduction reaction: applications to selected monoethylenic, diethylenic and triethylenic fatty acids of *cis* configurations. J Am Oil Chem Soc 67: 940-946
31. Ando Y, Ota T, Takagi T (1989) Japanese sardine oil as a source of 16:3(n-4) and 16:4(n-1) fatty acids. J Am Oil Chem Soc 66:1323-1325.
32. Gunstone FD, Ismail IA, Lie Ken Jie M (1967) Fatty acids, part 16. Thin layer and gas-liquid chromatographic properties of the *cis* and *trans* methyl octadecenoates and of some acetylenic esters. Chem Phys Lipids 1:376-385
33. Nikolova-Damyanova B (2003) Lipid analysis by silver ion chromatography. In: Adlof RO (ed) Advances in lipid methodology-five. The Oily Press, Bridgwater, pp 43-123
34. Shibahara A, Yamamoto K, Nakayama T, Kajimoto G (1985) Rapid determination of double bond positions in monounsaturated fatty acids by GC-MS and its application to fatty acid analysis. J Jpn Oil Chem Soc 34:618-625
35. Shibahara A, Yamamoto K, Kinoshita A, Anderson BL (2008) An improved method for preparing dimethyl disulfide adducts for GC/MS analysis. J Am Oil Chem Soc 85:93-94.
36. Christie WW (1998) Mass spectrometry of fatty acids with methylene-interrupted ene-yne systems. Chem Phys Lipids 94:35-41.
37. Ando Y, Sasaki T (2011) GC separation of *cis*-eicosenoic acid positional isomers on an ionic liquid SLB-IL100 stationary phase. J Am Oil Chem Soc 88, in press.
38. Spitzer V (1996) Structure analysis of fatty acids by gas chromatography-low resolution

electron impact mass spectrometry of their 4,4-dimethyloxazoline derivatives-a review.

Prog Lipid Res 35:387-408

39. Hayashi K, Yamada M (1975) The lipids of marine animals from various habitat depth-IV.

On the fatty acid composition of the neutral lipids in nine species of flatfishes. Bull Fac

Fish Hokkaido Univ 26:265-276

40. Vysotskii MV, Svetashev VI (1991) Identification, isolation and characterization of

tetracosapolyenoic acids in lipids of marine coelenterates. Biochim Biophys Acta

1083:161-165

41. Imbs AB, Yakovleva IM, Pham LQ (2010) Distribution of lipids and fatty acids in the

zooxanthellae and host of the soft coral *Sinularia* sp. Fish Sci 76:375-380

42. Carballeira NM, Sostre A, Rodríguez AD (1997) Phospholipid fatty acid composition of

goronians of the genus *Eunicea*: further identification of tetracosapolyenoic acids. Comp

Biochem Physiol 118B:257-260

43. Nichols PD, Danaher KT, Koslow JA (2003) Occurrence of high levels of

tetracosahexaenoic acid in the jellyfish *Aurelia* sp. Lipids 38:1207-1210

44. Go JV, Řezanka T, Srebnik M, Dembitsky VM (2002) Variability of fatty acid compositions

of marine and freshwater gastropods species from the littoral zone of the Red Sea,

Mediterranean Sea, and Sea of Galilee. Biochim Syst Ecol 30:819-835

45. Řezanka T, Dembitsky VM (1999) Very long chain polyunsaturated fatty acids in crustacea

of the order Bathynellacea. Biochim Syst Ecol 27:551-558

46. Řezanka T (2000) Analysis of very long chain polyunsaturated fatty acids using high-

performance liquid chromatography-atmospheric pressure chemical ionization mass

spectrometry. Biochim Syst Ecol 28:847-856

47. Mansour MP, Volkman JK, Holdsworth DG, Jackson AE, Blackburn SI (1999) Very long-

chain (C₂₈) highly unsaturated fatty acids in marine dinoflagellates. Phytochemistry

50:541-548

48. Mansour MP, Volkman JK, Jackson AE, Blackburn SI (1999) The fatty acid and sterol composition of five marine dinoflagellates. *J Phycol* 35:710-720
49. Van Pelt CK, Huang M-C, Tschanz CL, Brenna JT (1999) An octaene fatty acid, 4,7,10,13,16,19,22,25-octacosaoctaenoic acid (28:8n-3), found in marine oils. *J Lipid Res* 40:1501-1505
50. Mansour MP, Volkman JK, Blackburn SI (2003) The effect of growth phase on the lipid class, fatty acid and sterol composition in the marine dinoflagellate, *Gymnodinium* sp. in batch culture. *Phytochemistry* 63:145-153
51. Leblond JD, Evans TJ, Chapman PJ (2003) The biochemistry of dinoflagellate lipids, with particular reference to the fatty acid and sterol composition of a *Karenia brevis* bloom. *Phycologia* 42:324-331
52. Mansour MP (2005) Reversed-phase high-performance liquid chromatography purification of methyl esters of C16-C28 polyunsaturated fatty acids in microalgae, including octacosaoctaenoic acid [28:8(n-3)]. *J Chromatogr A* 1097:54-58
53. Mooney BD, Nichols PD, de Salas MF, Hallegraeff GM (2007) Lipid, fatty acid, and sterol composition of eight species of Kareniaceae (Dinophyta): chemotaxonomy and putative lipid phycotoxins. *J Phycol* 43:101-111.
54. Tokranov AM, Orlov AM (2003) On the distribution and biology of roughscale sole *Clidoderma asperrimum* (Temminck et Schlegel, 1846) in the Pacific waters off the northern Kuril Islands and southeastern Kamchatka. *Bull Sea Fish Inst* 159:67-80
55. Fujita T (1996) Bathymetric distribution of Ophiuroids (Echinodermata) off Sendai Bay, northern Japan, with notes on the diet of the roughscale sole *Clidoderma asperrimum* (Pisces, Pleuronectidae). *Mem Natn Sci Mus Tokyo* 29:209-222

Figure captions

Fig.1 Gas chromatogram of fatty acid methyl esters formed from roughscale sole flesh lipids (Restek FAMEWAX, 170 to 240°C at 4°C/min).

Fig. 2 Mass spectra of dimethyl disulfide (DMDS) adducts of 26:1 isomers formed by hydrazine hydrogenation of the roughscale sole 26:7 acid.

Fig. 3 Mass spectra of 4,4-dimethyloxazoline (DMOX) derivative (**a**) and methyl ester (**b**) of the roughscale sole 26:7 acid.

Fukuda & Ando
Figure 1

Fukuda & Ando
Figure 2

Table 1 Enrichment of 26:7 from the fatty acids of roughscale sole flesh lipids

Fatty acid	Intact ^a	Solvent partition	RP-TLC	Ag-TLC
Concentration (wt%)				
20:5n-3	7.7	48.4	0.4	ND
21:5n-3	0.2	1.1	0.5	ND
22:5n-3	0.7	2.0	6.8	ND
22:6n-3	2.0	15.7	ND	ND
24:6n-3	3.1	16.4	62.1	17.0
26:7n-3	1.3	10.4	27.7	83.0
Others	85.0	6.0	2.5	0.0

^a Fatty acid methyl esters prepared from a Pacific sample (41.2 cm, 2.8 kg).

Table 2 Fatty acid composition of the roughscale sole flesh lipids (wt%)

Fatty acid	Composition ^a			Fatty acid	Composition		
12:0	0.06	±	0.02	18:4n-3	0.37	±	0.06
14:0	5.97	±	0.64	18:4n-1	0.10	±	0.02
14:1n-5	0.28	±	0.03	20:0	0.14	±	0.03
iso-15:0	0.24	±	0.08	20:1n-11+n-13	6.50	±	1.38
anteiso-15:0	0.10	±	0.04	20:1n-9	2.63	±	0.34
15:0	0.31	±	0.01	20:1n-7	0.76	±	0.16
iso-16:0	0.12	±	0.03	20:2n-6	0.18	±	0.02
16:0	12.05	±	1.44	20:3n-6	0.02	±	0.02
16:1n-9	0.33	±	0.04	20:4n-6	0.83	±	0.13
16:1n-7	9.30	±	0.65	20:3n-3	0.10	±	0.02
16:1n-5	0.20	±	0.03	20:4n-3	0.23	±	0.03
iso-17:0	0.52	±	0.19	20:5n-3	8.31	±	0.46
anteiso-17:0	0.11	±	0.03	22:0	0.07	±	0.01
16:2n-4	0.15	±	0.04	22:1n-11+n-13	3.07	±	0.59
17:0	0.18	±	0.02	22:1n-9	0.93	±	0.14
16:3n-4	0.08	±	0.03	22:1n-7	0.22	±	0.05
17:1n-7	0.32	±	0.02	22:2n-6	0.05	±	0.03
16:4n-1	0.19	±	0.09	21:5n-3	0.24	±	0.05
18:0	2.12	±	0.13	22:5n-6	0.13	±	0.05
18:1n-13	0.69	±	0.15	22:5n-3	0.80	±	0.11
18:1n-9+n-11	23.54	±	1.48	22:6n-3	2.88	±	0.74
18:1n-7	4.34	±	0.25	24:1n-9	0.75	±	0.11
18:1n-5	0.60	±	0.07	24:6n-3	3.75	±	1.02
18:2n-6	0.33	±	0.05	26:7n-3	0.69	±	0.34
18:2n-4	0.13	±	0.04	Others	3.18	±	0.34
18:3n-6	0.02	±	0.02				
18:3n-3	0.15	±	0.03	Lipid content (%) ^b	30.5	±	4.5

^a Mean ± SD of the 5 individuals caught in the Pacific water and the Sea of Okhotsk around Hokkaido, Japan.

^b Determined by gravimetry (% on the wet-weight base).

サメガレイ *Clidoderma asperrimum* の筋肉脂質における全-*cis*-5,8,11,14,17,20,23-ヘキサ コサヘプタエン酸 (26:7n-3) の存在

福田悠紀・安藤靖浩（北大院水）

北海道産サメガレイの筋肉脂質の脂肪酸を GC で分析したところ、炭素数 24 以上の領域に未知のピークが出現した。この成分を濃縮後、数種の誘導体に変換して GC-MS に供した結果、同成分は全-*cis*-5,8,11,14,17,20,23-ヘキサコサヘプタエン酸 (26:7n-3) と同定された。総脂肪酸中の含有量は供試 5 個体の平均で $0.69 \pm 0.34\%$ であった。サメガレイは、n-3 系および n-6 系超長鎖ポリエン酸の給源として一般的な高等動物の網膜、精子、ニシン油などには見られない 26:7n-3 を含む点で特徴的である。