



Title	Herbivory in canopy gaps created by a typhoon varies by understory plant leaf phenology
Author(s)	Takafumi, Hino; Kawase, Satoru; Nakamura, Masahiro; Hiura, Tsutom
Citation	Ecological Entomology, 35(5), 576-585 https://doi.org/10.1111/j.1365-2311.2010.01216.x
Issue Date	2010-10
Doc URL	http://hdl.handle.net/2115/47185
Rights	The definitive version is available at onlinelibrary.wiley.com
Type	article (author version)
File Information	EE35-5_576-585.pdf



[Instructions for use](#)

**Herbivory in canopy gaps created by a typhoon varies by understory plant leaf
phenology**

Running title: Herbivory varies by plant leaf phenology

List of Authors: Hino Takafumi¹, Satoru Kawase¹, Masahiro Nakamura², Tsutomu Hiura¹.

Affiliation: 1 Tomakomai Research Station, Field Science Center for Northern

Biosphere, Hokkaido University, Tomakomai, Hokkaido, 053-0035, Japan.

2 Nakagawa Research Station, Field Science Center for Northern Biosphere, Hokkaido

University, Ootomai, Hokkaido, 098-2501, Japan.

Corresponding author: Hino Takafumi

Tomakomai Research Station, Field Science Center for Northern Biosphere,

Hokkaido University, Tomakomai 053-0035, Japan.

E-mail: takafumi.hi@gmail.com

Tel.: +81-144-332171

Fax: +81-144-332173

Abstract

1. Availabilities of light and soil nitrogen for understory plants vary by extent of canopy gap formation through typhoon disturbance. We predicted that variation in resource availability and herbivore abundance in canopy gaps would affect herbivory through variation in leaf traits among plant species. We studied six understory species that expand their leaves before or after canopy closure in deciduous forests. We measured the availabilities of light, soil nitrogen, soil water content, and herbivore abundance in 20 canopy gaps (28.3–607.6 m²) formed by a typhoon and in four undisturbed stands. We also measured leaf traits and herbivory on understory plants.

2. The availabilities of light and soil nitrogen increased with increasing gap size. However, soil water content did not. The abundance of herbivorous insects (such as Lepidoptera and Orthoptera) increased with increasing gap size.

3. Concentrations of condensed tannins, total phenolics, and nitrogen in leaves and the leaf mass per area increased in late leaf expansion species with increasing gap size, while none of the leaf traits varied by gap size in early leaf expansion species.

4. Herbivory increased on early leaf expansion species with increasing gap size, but decreased on late leaf expansion species. In these late leaf expansion species, total phenolics and C:N ratio had negative relationships with herbivory.

5. These results suggested that following typhoon disturbance, increased herbivory on early leaf expansion species can be explained by increased herbivore abundance, while decreased herbivory on late leaf expansion species can be explained by variation in leaf traits.

Key-words: deciduous forest, gap size, insect–plant interactions, resource availability, timing of leaf expansion

Introduction

Forest canopy gaps formed by natural disturbances produce variability in resource availabilities for the understory plant community. For example, light availability increases in canopy gaps (Carlton & Bazzaz, 1998; Denslow *et al.*, 1990; Oguchi *et al.*, 2006). This is also the case for soil nitrogen availability (Denslow *et al.*, 1998; Oguchi *et al.*, 2006; Prescott, 2002) because high soil water content and temperatures in canopy gaps accelerate decomposition of organic matter (Bormann *et al.*, 1974). Natural disturbances play major roles in biological interactions by increasing variability in resource availabilities (Connell, 1978; Hiura, 1995; Menge & Sutherland, 1987; Pickett & White, 1985; Sousa, 1984). Variations in resource availability following natural disturbance alter insect–plant interactions in understory vegetation because herbivory in canopy gaps may increase (Chacon & Armesto, 2006; Dudt & Shure, 1994; Hunter & Forkner, 1999), decrease (Schowalter & Ganio, 1999; Shure & Wilson, 1993), and/or may not vary (Shure & Wilson, 1993). Responses of herbivory to canopy gaps remain largely undetermined.

Leaf traits, such as concentrations of secondary metabolites, toughness, and leaf nitrogen content, alter feeding behaviors of herbivorous insects (e.g., Bryant *et al.*, 1987; Coley, 1983; Coley & Barone, 1996; Feeny, 1970) and may be influenced by

variation in resource availability (Bryant *et al.*, 1983, 1987; Close *et al.*, 2003; Close & McArthur, 2002; Nakamura *et al.*, 2008; Yamasaki & Kikuzawa, 2003). Secondary metabolites (e.g., condensed tannins and total phenolics) in plants often act as herbivore deterrents (Bryant *et al.*, 1987; Coley & Barone, 1996; Feeny, 1970), and their concentrations in plant tissues increase when light availability is elevated (Bryant *et al.*, 1983, 1987; Close *et al.*, 2003; Close & McArthur, 2002). This increase in secondary metabolite concentrations is explained by two non-mutually exclusive hypotheses. First, the carbon-nutrient balance (CNB) hypothesis explains increasing secondary metabolite concentrations as a function of the balance between carbon and nitrogen (C:N ratio) within a plant (Bryant *et al.*, 1983, 1987), although the generality of this hypothesis is debated (Hamilton *et al.*, 2001; Koricheva, 2002; Nakamura *et al.*, 2008). Second, taking no account of the C:N ratio, the photoinhibition hypothesis explains increasing secondary metabolite concentrations in terms of photodamage risk reduction (Close *et al.*, 2003; Close & McArthur, 2002). Secondary metabolites often function as chemical deterrents of grazers, while the leaf mass per area (LMA) is closely correlated with toughness and physical defense against herbivores (Choong, 1996; Coley, 1983). In contrast, elevated leaf nitrogen content is an attractant for herbivores (Coley, 1983). All of these leaf traits increase under high light and/or high soil nitrogen availability (Mattson, 1980; Nakamura *et al.*, 2008; Nordin *et al.*, 1998; Oguchi *et al.*, 2006;

Yamasaki & Kikuzawa, 2003). Thus, canopy gaps may affect herbivory by causing variation in leaf traits.

Physiological responses (e.g., photosynthetic adaptation, nitrogen-use efficiency) to seasonal resource variation vary among understory plant species with differing leaf expansion phenologies (Kudo *et al.*, 2008; Rothstein & Zak, 2001a,b; Skillman *et al.*, 1996). In deciduous forests, understory plants have two distinct strategies of leaf expansion timing for avoiding or tolerating shade from canopy trees in summer (Neufeld & Young, 2003; Uemura, 1994). Early leaf expansion species open their leaves before canopy trees, and thus avoid shading early in the season. This phenology allows expanded leaves to capture a large proportion of current-year photosynthate before canopy closure (Ida & Kudo, 2008). In contrast, late leaf expansion species have shade-tolerating strategies because the leaves open after canopy closure (Neufeld & Young, 2003). If leaf trait responses to canopy gaps differ between early and late leaf expansion species, responses of herbivory to canopy gaps likely depend on leaf phenology.

Herbivorous insects on understory plants are more abundant in canopy gaps than under closed canopy (Basset *et al.*, 2001; Jokimaki *et al.*, 1998; Louda & Rodman, 1996; Richards & Coley, 2007; Sipura & Tahvanainen, 2000). Elevated light availability and temperature in the gaps promote faster insect development and immigration of

grazers from forest shade (Cappuccino & Root, 1992; Courtney & Courtney, 1982; Sipura & Tahvanainen, 2000; Weiss et al., 1988). Increased herbivore abundance in canopy gaps probably increases feeding pressure on understory plants. Thus, both leaf traits and herbivore abundances should be considered as important determinants of herbivory in canopy gaps (Louda & Rodman, 1996).

In this study, we focused on the ways in which canopy gaps formed by a typhoon affect herbivory on understory plants through disturbance-induced variations in leaf traits and herbivore abundances. We studied leaf traits and herbivory in six perennial understory species (differing in leaf expansion timing) among various canopy gap sizes (28.3–607.6 m²) created by a typhoon in a deciduous forest. In particular, we addressed the following questions: Do leaf trait responses (C:N ratio, LMA and concentrations of condensed tannins, total phenolics and nitrogen) to canopy gaps differ between early and late leaf expansion species? Do herbivore abundances respond to canopy gaps? Does herbivory in canopy gaps differ between early and late leaf expansion species?

Methods

STUDY SITE

This study was conducted in Tomakomai Experimental Forest (TOEF), Hokkaido University (42°41' N, 141°36' E; area: 2,715 ha), Hokkaido, Japan. A large part of TOEF is located on flat land, with a gradient of <math><5^\circ</math>. Mean monthly temperatures range from -3.2 to 19.1°C, and annual precipitation is 1450 mm. Snow cover reaches a depth of 50 cm from December to March. Minimum of 350 vascular plant species have been recorded in TOEF (Kudo & Yoshimi, 1916). The dominant canopy trees in natural stands are *Quercus crispula* (Fagaceae), *Acer mono* (Aceraceae), *Sorbus alnifolia* (Rosaceae), and *Tilia japonica* (Tiliaceae) (Hiura, 2001). The forest formed on 2-m-deep porous volcanic ash, which accumulated after the eruptions of Mt. Tarumae between 1669 and 1739 (Shibata et al., 1998).

Two typhoons in 1954 and 2004 were major natural disturbances from 1904 to 2004 in TOEF (Hiura et al., 2004; Takafumi & Hiura, 2009). The typhoon in 2004 disturbed about 32% of the forest area in TOEF (Hiura et al., 2004). To determine the effects of canopy gaps created by the typhoon on resource availabilities for understory plants, herbivore abundances, leaf traits, and herbivory on understory plants, we set up

20 study plots within varying gap sizes (28.3–607.6 m²), and four undisturbed study plots under full canopy (total: 24 plots) in July 2005. The study plots had not been disturbed by harvesting over the previous 50 years (Takafumi & Hiura, 2009). We measured the lengths and widths of gaps with a steel tape and calculated areas of gaps assuming they were elliptical.

RESOURCE AVAILABILITY

To evaluate the availability of light in canopy gaps, we captured a hemispherical photographic image to measure canopy openness above each study plot. The hemispherical photographs were taken with a digital camera (Coolpix 990; Nikon, Tokyo, Japan) equipped with a fish-eye lens (FC-E8; Nikon). The top element of the lens was positioned 0.5 m above the ground. The hemispherical images were analyzed with CanopOn2 (free software programmed by A. Takenaka available at <http://www.takenaka-akio.cool.ne.jp/etc/canopon2/>) software based on a standard overcast sky condition.

To evaluate the availabilities of soil nitrogen in canopy gaps, N mineralization (NH₄⁺) and nitrification (NO₃⁻) in intact soil cores were determined in each study plot following the methods of Oguchi *et al.* (2006). Sampling points in each study plot under

a canopy gap were set at the center of the gap and at four midpoints between the center and the gap edges, avoiding pits and mounds (total: five sampling points per plot), when measurements were started in July 2005. In undisturbed study plots (no canopy gaps), sampling points were set at the center and four corners of a 5 × 5-m quadrat located at the center of each plot. Three cores (5.0 cm diameter × 5.0 cm long) were extracted from each sampling point (total: 15 cores per plot). One core from each sampling point was used for the analysis of initial soil nitrogen availability. The other two cores were reinserted into their holes in the plot and incubated in the field using the resin technique (Distefano & Gholz, 1986). The resin (active anion exchange resin, Amberlite IRA-400 and active cation exchange resin, Amberlite IR-120B; Rohm and Haas, Philadelphia, PA, USA) wrapped in fine mesh nylon was connected to the base of each soil core sample in polyvinyl chloride tubes. The resin packs collected ions leaching through soil core samples. At the end of measurement periods (September 2005, September 2006), cores were collected. Afterward, a solution of NO_3^- and NH_4^+ was extracted and analyzed with an autoanalyzer (AACS-4; BL-TEC Inc., Osaka, Japan). The availabilities of NO_3^- and NH_4^+ were calculated from the differences between start and endpoints of the measurement periods.

To evaluate soil water content (percentage by volume) in canopy gaps, we used time domain reflectometry (Trime-FM2/P2; IMKO, Ettlingen, Germany) (Topp & Davis,

1985; Topp et al., 1980) in August and November 2005, and in August 2006. We measured the soil water contents after 5 -6 days after a rainfall event. A pair of 5-cm-long reflectometry rods was installed in the soil near each resin bag.

UNDERSTORY PLANT SPECIES WITH DIFFERENT TIMINGS OF LEAF EXPANSION

To determine the effects of canopy gaps on understory plant species with different timings of leaf expansion, we observed four dominant herb species (*Trillium tschonoskii* Maxim., *Smilacina japonica* A. Gray, *Polygonatum odoratum* var. *maximowiczii* Koidz, *Desmodium podocarpum* DC.) and two canopy tree species (*Acer mono* Maxim., *Quercus crispula* Blume) in the understory. In late April, *T. tschonoskii* and *S. japonica* expand their leaves as soon as snow melts and before canopy closure. In early May, the other herb species, *P. odoratum* var. *maximowiczii* and *D. podocarpum*, expand leaves immediately following the start of canopy tree leaf expansion. In late May, seedlings (<130 cm tall) of two canopy tree species (*A. mono*, *Q. crispula*) expand leaves following canopy closure.

LEAF TRAITS OF UNDERSTORY PLANTS

To evaluate the leaf traits of understory plants in canopy gaps, we randomly sampled one leaf having the least herbivory (<10% leaf loss) from each of six individuals of each plant species in early August 2006. When less than six individuals of a plant species occurred in a plot, we randomly sampled one leaf from each of the individuals present (>2 individuals in all cases). Afterward, we punched five leaf disks (6 mm in diameter) out of each leaf, and the sampled disks were oven-dried for 72 h at 40°C. The LMA was calculated for each leaf. Using these punched disks, we measured concentrations of carbon and nitrogen using a CN analyzer (NC-900; Sumitomo, Osaka, Japan), and the C:N ratios were calculated. Condensed tannins and total phenolics in leaves were measured following Julkunen-Tiitto (1985). Since herbivory was evaluated by assessing the percentage of leaf area consumed, we also calculated condensed tannins, total phenolics, and nitrogen on unit-area bases.

HERBIVORE ABUNDANCE

To evaluate herbivore abundance in canopy gaps, herbivorous chewing insects were collected by hand from understory plants over 30-min periods in single 5 × 5-m quadrats in each plot in late July 2006. We sorted insects into orders (Lepidoptera,

Orthoptera, Coleoptera) and measured the dry mass of each order in each plot after oven-drying for 48 h at 60°C.

HERBIVORY ON UNDERSTORY PLANTS

To evaluate herbivory in canopy gaps, 100 leaves were randomly selected from each plant species in each plot. We assessed percentages of consumed leaf areas by chewing herbivore insects in late July 2006. We transformed these percentages into seven classes of herbivory as follows: 1: 0%, 2: 1–5%, 3: 6–10%, 4: 11–25%, 5: 26–50%, 6: 51–75%, 7: 76–100%. When we had less than 100 leaves of a plant species in a plot, we assessed herbivory on all leaves present (>30 leaves in all cases). Leaves grazed by mammals and snails were omitted because we hardly observed leaves grazed by these herbivores.

DATA ANALYSIS

The availabilities of light and soil nitrogen, soil water content, herbivore abundance, and leaf traits of understory plants were analyzed using generalized linear models with gap size as the explanatory variable. We analyzed herbivory by gap size using ordinal logistic models (= proportional odds models) (McCullagh, 1980) as each class of

consumed leaf area was an ordinal category. The relationships between herbivory and leaf traits were also analyzed with ordinal logistic models. We selected best-fit models from linear and log-linear models using Akaike's information criterion (AIC). We used the likelihood ratio test to determine whether the data supported selected models over a null model. All statistical analyses were conducted with R Version 2.4.1. software (R development Core Team, 2006).

Results

RESOURCE AVAILABILITY

Canopy openness increased significantly from 7.8% to 38.3% with increasing gap size (gap size: 0–607.6 m²) (Table 1). Nitrate in soil between July 2005 and October 2006 increased significantly with increasing gap size. Only NH₄⁺ mineralized in soil between July and October in 2005 (i.e., not at other times) increased significantly with increasing gap size. However, soil water content did not vary by gap size during the measurement period ($P > 0.05$).

LEAF TRAITS OF UNDERSTORY PLANTS

The responses of leaf traits of understory plants to canopy gaps differed between early and late leaf expansion species. Most leaf traits varied in late leaf expansion species by gap size, but no significant variation in leaf traits of early leaf expansion species was observed as gap size increased (Table 2). In all late leaf expansion species, condensed tannin and total phenolic concentrations in leaves, and LMA increased with increasing gap size. With increasing gap size, the C:N ratio increased in *D. podocarpum*, did not vary in *A. mono*, and decreased in *Q. crispula*. Leaf nitrogen content did not vary in *D. podocarpum*, but increased in *A. mono* and *Q. crispula* with increasing gap size.

HERBIVORY ON UNDERSTORY PLANTS

Herbivory responses to canopy gaps differed between early and late leaf expansion species. Herbivory increased on early leaf expansion species (*T. tschonoskii* and *S. japonica*) with increasing gap size, but decreased on late leaf expansion species (*D. podocarpum* and *Q. crispula*) (Table 3). In addition, the total dry mass of herbivorous insects increased with increasing gap size (Fig. 1), particularly in the orders Lepidoptera and Orthoptera.

RELATIONSHIP BETWEEN HERBIVORY AND LEAF TRAITS

Among leaf traits of late leaf expansion species that varied by canopy gap size, the C:N ratio in *D. podocarpum* and total phenolics in *Q. crispula* had negative relationships with herbivory (Table 4).

Discussion

This study clearly showed that with increasing gap size, concomitant increases occurred in the availabilities of plant resources (light and soil nitrogen) and herbivore abundance. Furthermore, most leaf traits varied by gap size in late leaf expansion species but not in early leaf expansion species. Herbivory increased on early leaf expansion species with increasing gap size but decreased in late leaf expansion species.

CHANGE IN RESOURCE AVAILABILITY IN CANOPY GAPS

The availability of soil nitrogen increases in canopy gaps (Denslow *et al.*, 1998; Oguchi *et al.*, 2006; Prescott, 2002) through accelerated decomposition of organic matter

resulting from elevated soil water contents and temperatures (Bormann *et al.*, 1974). We found that availabilities of NO_3^- and NH_4^+ increased with increasing gap size, but soil water content did not, probably because this forest is well drained (being formed on 2-m-deep porous volcanic ash with a very shallow soil layer) (Shibata *et al.*, 1998). In general, the soil temperature increases in canopy gaps (Gray *et al.*, 2002; Naidu & DeLucia, 1997, 1998; Phillips & Shure, 1990; Ritter, 2005; Wright *et al.*, 1998). For example, Phillips & Shure (1990) reported that soil temperature increased with increasing gap size (160–20000 m^2) in a southern Appalachian deciduous forest (USA). Although our study did not measure soil temperature, the high soil nitrogen availability might have been caused by increased soil temperatures in canopy gaps.

DIFFERENT LEAF TRAIT RESPONSES AMONG PLANTS DIFFERING IN TIMING OF LEAF EXPANSION

Understory plants with different leaf expansion phenologies have different physiological responses to changing light availability (Kudo *et al.*, 2008; Rothstein & Zak, 2001a; Skillman *et al.*, 1996). We found that concentrations of total phenolics, condensed tannins, and nitrogen in leaves, and the LMA increased in late leaf expansion species with increasing gap size, but none of these traits varied by gap size in early leaf

expansion species. Late leaf expansion species usually grow under limited light conditions after canopy closure, but canopy gaps provided illumination for these species, which may have led to variation in the leaf traits. Two non-mutually exclusive hypotheses propose different mechanisms by which higher light availability may lead to increased secondary metabolite concentrations. First, the CNB hypothesis explains that the balance of carbon and nitrogen within a plant (C:N ratio) regulates the allocation of resources to primary and secondary metabolites (Bryant *et al.*, 1983, 1987). Under high light conditions, excess C is used preferentially to synthesize C-based secondary metabolites (Bryant *et al.*, 1983; Coley *et al.*, 1985). We found that in *D. podocarpum*, both the C:N ratio and secondary metabolites (condensed tannins and total phenolics) increased with increasing gap size, but N content did not; thus responses of this species alone support the CNB hypothesis. The photoinhibition hypothesis explains that plants are induced to produce secondary metabolites (regardless of C:N ratio) as a way of reducing the risk of photodamage by oxidants to photosynthesis (Close *et al.*, 2003; Close & McArthur, 2002). Plants usually increase carotenoids under high light conditions to protect the photosystems (Demmig-Adams *et al.*, 1996; Havaux, 1998).

Previous studies have shown that total phenolics and condensed tannins also had similar function (Hagerman *et al.*, 1998; Kuhlmann & Muller, 2009; Mole *et al.*, 1988; Treutter, 2006). We found that secondary metabolite (total phenolics and condensed tannins)

concentrations increased in all late leaf expansion species with increasing gap size. Thus, the photoinhibition hypothesis may agree with all our results for later leaf expansion species. Similar to secondary metabolite concentrations, the LMA and leaf nitrogen increased in late leaf expansion species with increasing gap size. Studies have reported that higher light availability increases the LMA and leaf nitrogen under canopy gaps due to an elevation of photosynthetic production (Oguchi *et al.*, 2006; Poorter *et al.*, 2009). We found that soil nitrogen increased with increasing gap size, which may have led to increases in leaf nitrogen in late leaf expansion species.

In contrast to late leaf expansion species, no leaf trait variation with increasing gap size was observed in early leaf expansion species. Since these species produce the largest proportion of current-year photosynthate before canopy closure (Ida & Kudo, 2008), higher light availability in canopy gaps would not affect concentrations of secondary metabolites and nitrogen in leaves, or the LMA.

Alternatively, the first four species to expand their leaves are herbs, while the two late leaf expansion species are canopy tree species. Canopy gaps are essential for growth and recruitment of most of canopy trees species (Denslow, 1987; Pickett & White, 1985). Thus, not only different leaf expansion phenologies but also different life forms may explain the different responses of leaf traits to canopy gaps.

INCREASED HERBIVORE ABUNDANCE IN CANOPY GAPS

Herbivorous insects on understory plants are more abundant in canopy gaps than under closed canopy because the high light availability and warmer condition lead to faster insect development and immigration (Basset *et al.*, 2001; Jokimaki *et al.*, 1998; Louda & Rodman, 1996; Richards & Coley, 2007; Sipura & Tahvanainen, 2000). We found that the abundance of herbivorous insects, such as Lepidoptera and Orthoptera, increased with increasing gap size. This is probably because lepidopteran larvae prefer the edges of canopy gaps (Cappuccino & Martin, 1997; Cappuccino & Root, 1992; Courtney & Courtney, 1982), and they migrate from the canopy to the understory as canopy leaf quality degrades (leaf toughness and tannin increased, while the nitrogen and water contents decreased) in late season (Murakami & Wada, 1997). These lepidopteran larvae feed canopy trees and understory plants (Murakami & Wada, 1997), thus, most of these Lepidoptera larvae may be generalists. Furthermore, the majority of orthopterans are open landscape species, and their abundances are also elevated in canopy gaps compared to closed canopy (Bouget & Duelli, 2004; Richards & Coley, 2007; Shure & Phillips, 1991).

DIFFERING HERBIVORY RESPONSES BETWEEN PLANTS WITH DIFFERENT

TIMINGS OF LEAF EXPANSION

Herbivory rates may be determined by a combination of variation in leaf traits and herbivore abundances (Louda & Rodman, 1996). We showed that herbivory increased on early leaf expansion species (*T. tschonoskii* and *S. japonica*) with increasing gap size, but decreased on late leaf expansion species (*D. podocarpum* and *Q. crispula*). None of the leaf traits varied in early leaf expansion species in canopy gaps, although herbivore abundance increased. Thus, increased herbivory on early leaf expansion species in canopy gaps may be explained by increased herbivore abundance. In contrast, most leaf traits in late leaf expansion species varied by gap size. Among these traits, the C:N ratio in *D. podocarpum* and total phenolics in *Q. crispula* had negative relationships with herbivory. However, measured secondary metabolite (condensed tannins and total phenolics) concentrations in *D. podocarpum* were not negatively related to herbivory. Thus, other (as yet unmeasured) secondary metabolites may be responsible for decreased herbivory in canopy gaps. Decreased herbivory on late leaf expansion species can be explained by increased secondary metabolite concentrations in canopy gaps. These results suggest that important determinants of herbivory in canopy gaps differed between early and late leaf expansion species.

In conclusion, we clearly showed that following typhoon disturbance, increased

herbivory on early leaf expansion species can be explained by increased herbivore abundance; decreased herbivory on late leaf expansion species in canopy gaps can be explained by variation in leaf traits. Several studies have reported that understory plants with different timings of leaf expansion have different physiological characteristics (Kudo *et al.*, 2008; Rothstein & Zak, 2001a,b; Skillman *et al.*, 1996). Our findings emphasize that initial responses of herbivory to canopy gaps formed by the typhoon also varied between plants with different timings of leaf expansion. To consider both leaf traits and herbivore abundance as important determinant of herbivory, we can understand how herbivory responds to canopy gaps.

Acknowledgements

We thank staff members and graduate students at Tomakomai Research Station, Hokkaido University, for their support during the study. We also thank Drs. R. Oguchi, H. Shibata, M. Ozawa, K. Fukuzawa, and Y. Nakagawa for soil analysis. This work was partly supported by grants from the Ministry of Education, Culture, Sports, Science and Technology, Japan (no. 15208014) and from the Ministry of Environment to T. Hiura.

References

- Basset, Y., Charles, E., Hammond, D.S., & Brown, V.K. (2001) Short-term effects of canopy openness on insect herbivores in a rain forest in Guyana. *Journal of Applied Ecology*, 38, 1045-1058.
- Bormann, F.H., Likens, G.E., Siccama, T.G., Pierce, R.S., & Eaton, J.S. (1974) The export of nutrients and recovery of stable conditions following deforestation at Hubbard Brook. *Ecological Monographs*, 44, 255-277.
- Bouget, C. & Duelli, P. (2004) The effects of windthrow on forest insect communities: a literature review. *Biological Conservation*, 118, 281-299.
- Bryant, J.P., Chapin, F.S., & Klein, D.R. (1983) Carbon nutrient balance of boreal plants in relation to vertebrate herbivory. *Oikos*, 40, 357-368.
- Bryant, J.P., Clausen, T.P., Reichardt, P.B., McCarthy, M.C., & Werner, R.A. (1987) Effect of nitrogen fertilization upon the secondary chemistry and nutritional value of quaking aspen (*Populus tremuloides* Michx.) leaves for the large aspen tortrix (*Choristoneura conflictana* (Walker)). *Oecologia*, 73, 513-517.
- Cappuccino, N. & Martin, M.A. (1997) The birch tube-maker *Acrobasis betulella* in a fragmented habitat: The importance of patch isolation and edges. *Oecologia*, 110, 69-76.
- Cappuccino, N. & Root, R.B. (1992) The significance of host patch edges to the

- colonization and development of *Corythucha marmorata* (Hemiptera, Tingidae).
Ecological Entomology, 17, 109-113.
- Carlton, G.C. & Bazzaz, F.A. (1998) Resource congruence and forest regeneration following an experimental hurricane blowdown. *Ecology*, 79, 1305-1319.
- Chacon, P. & Armesto, J.J. (2006) Do carbon-based defences reduce foliar damage? Habitat-related effects on tree seedling performance in a temperate rainforest of Chiloe Island, Chile. *Oecologia*, 146, 555-565.
- Choong, M.F. (1996) What makes a leaf tough and how this affects the pattern of *Castanopsis fissa* leaf consumption by caterpillars. *Functional Ecology*, 10, 668-674.
- Close, D., McArthur, C., Paterson, S., Fitzgerald, H., Walsh, A., & Kincade, T. (2003) Photoinhibition: A link between effects of the environment on eucalypt leaf chemistry and herbivory. *Ecology*, 84, 2952-2966.
- Close, D.C. & McArthur, C. (2002) Rethinking the role of many plant phenolics - protection from photodamage not herbivores? *Oikos*, 99, 166-172.
- Coley, P.D. (1983) Herbivory and defensive characteristics of tree species in a lowland tropical forest. *Ecological Monographs*, 53, 209-233.
- Coley, P.D. & Barone, J.A. (1996) Herbivory and plant defenses in tropical forests. *Annual Review of Ecology and Systematics*, 27, 305-335.
- Coley, P.D., Bryant, J.P., & Chapin, F.S. (1985) Resource availability and plant

- antiherbivore defense. *Science*, 230, 895-899.
- Connell, J.H. (1978) Diversity in tropical rain forests and coral reefs. *Science*, 199, 1302-1310.
- Courtney, S.P. & Courtney, S. (1982) The 'edge-effect' in butterfly oviposition: causality in *Anthocharis cardamines* and related species. *Ecological Entomology*, 7, 131-137.
- Demmig-Adams, B., Gilmore, A.M., & Adams, W.W. (1996) Carotenoids .3. In vivo functions of carotenoids in higher plants. *The Federation of American Societies for Experimental Biology Journal*, 10, 403-412.
- Denslow, J.S. (1987) Tropical rain-forest gaps and tree species-diversity. *Annual Review of Ecology and Systematics*, 18, 431-451.
- Denslow, J.S., Ellison, A.M., & Sanford, R.E. (1998) Treefall gap size effects on above- and below-ground processes in a tropical wet forest. *Journal of Ecology*, 86, 597-609.
- Denslow, J.S., Schultz, J.C., Vitousek, P.M., & Strain, B.R. (1990) Growth-responses of tropical shrubs to treefall gap environments. *Ecology*, 71, 165-179.
- Distefano, J.F. & Gholz, H.L. (1986) A proposed use of ion exchange resins to measure nitrogen mineralization and nitrification in intact soil cores. *Communications in Soil Science and Plant Analysis*, 17, 989-998.
- Dudt, J.F. & Shure, D.J. (1994) The influence of light and nutrients on foliar phenolics

and insect herbivory. *Ecology*, 75, 86-98.

Feeny, P. (1970) Seasonal changes in oak leaf tannin and nutrients as a cause of spring feeding by winter moth caterpillars. *Ecology*, 51, 565-581.

Gray, A.N., Spies, T.A., & Easter, M.J. (2002) Microclimatic and soil moisture responses to gap formation in coastal Douglas-fir forests. *Canadian Journal of Forest Research*, 32, 332-343.

Hagerman, A.E., Riedl, K.M., Jones, G.A., Sovik, K.N., Ritchard, N.T., Hartzfeld, P.W., & Riechel, T.L. (1998) High molecular weight plant polyphenolics (tannins) as biological antioxidants. *Journal of Agricultural and Food Chemistry*, 46, 1887-1892.

Hamilton, J.G., Zangerl, A.R., DeLucia, E.H., & Berenbaum, M.R. (2001) The carbon-nutrient balance hypothesis: its rise and fall. *Ecology Letters*, 4, 86-95.

Havaux, M. (1998) Carotenoids as membrane stabilizers in chloroplasts. *Trends in Plant Science*, 3, 147-151.

Hiura, T. (1995) Gap formation and species-diversity in Japanese beech forests - a test of the intermediate disturbance hypothesis on a geographic scale. *Oecologia*, 104, 265-271.

Hiura, T. (2001) Stochasticity of species assemblage of canopy trees and understory plants in a temperate secondary forest created by major disturbances. *Ecological Research*, 16, 887-893.

- Hiura, T., Okuda, A., Okuyama, S., & Takafumi, H. (2004) Report on the damage by the Typhoon 18 of 2004 and future problem in the Tomakomai Experimental Forest (in Japanese). *Technical Report for Boreal Forest Conservation*, 23, 24-26.
- Hunter, M.D. & Forkner, R.E. (1999) Hurricane damage influences foliar polyphenolics and subsequent herbivory on surviving trees. *Ecology*, 80, 2676-2682.
- Ida, T.Y. & Kudo, G. (2008) Timing of canopy closure influences carbon translocation and seed production of an understorey herb, *Trillium apetalon* (Trilliaceae). *Annals of Botany*, 101, 435-446.
- Jokimaki, J., Huhta, E., Itamies, J., & Rahko, P. (1998) Distribution of arthropods in relation to forest patch size, edge, and stand characteristics. *Canadian Journal of Forest Research*, 28, 1068-1072.
- Julkunen-Tiitto, R. (1985) Phenolic constituents in the leaves of northern willows: methods for the analysis of certain phenolics. *Journal of Agricultural and Food Chemistry*, 33, 213-217.
- Koricheva, J. (2002) The carbon-nutrient balance hypothesis is dead; long live the carbon-nutrient balance hypothesis? *Oikos*, 98, 537-539.
- Kudo, G., Ida, T.Y., & Tani, T. (2008) Linkages between phenology, pollination, photosynthesis, and reproduction in deciduous forest understory plants. *Ecology*, 89, 321-331.

- Kudo, Y. & Yoshimi, T. (1916) Flora of Tomakomai Experimental Forest, Hokkaido University. *Research Bulletin of Tohoku Emperor University Forests*, 3, 1-62.
- Kuhlmann, F. & Muller, C. (2009) Development-dependent effects of UV radiation exposure on broccoli plants and interactions with herbivorous insects. *Environmental and Experimental Botany*, 66, 61-68.
- Louda, S.M. & Rodman, J.E. (1996) Insect herbivory as a major factor in the shade distribution of a native crucifer (*Cardamine cordifolia* A. Gray, bittercress). *Journal of Ecology*, 84, 229-237.
- Mattson, W.J. (1980) Herbivory in relation to plant nitrogen content. *Annual Review of Ecology and Systematics*, 11, 119-161.
- McCullagh, P. (1980) Regression models for ordinal data. *Journal of the Royal Statistical Society Series B-Methodological*, 42, 109-142.
- Menge, B.A. & Sutherland, J.P. (1987) Community regulation - variation in disturbance, competition, and predation in relation to environmental-stress and recruitment. *American Naturalist*, 130, 730-757.
- Mole, S., Ross, J.A.M., & Waterman, P.G. (1988) Light-Induced Variation in Phenolic Levels in Foliage of Rain-Forest Plants .1. Chemical-Changes. *Journal of Chemical Ecology*, 14, 1-21.
- Murakami, M. & Wada, N. (1997) Difference in leaf quality between canopy trees and

- seedlings affects migration and survival of spring-feeding moth larvae. *Canadian Journal of Forest Research*, 27, 1351-1356.
- Naidu, S.L. & DeLucia, E.H. (1997) Acclimation of shade-developed leaves on saplings exposed to late-season canopy gaps. *Tree Physiology*, 17, 367-376.
- Naidu, S.L. & DeLucia, E.H. (1998) Physiological and morphological acclimation of shade-grown tree seedlings to late-season canopy gap formation. *Plant Ecology*, 138, 27-40.
- Nakamura, M., Hina, T., Nabeshima, E., & Hiura, T. (2008) Do spatial variation in leaf traits and herbivory within a canopy respond to selective cutting and fertilization? *Canadian Journal of Forest Research*, 38, 1603-1610
- Neufeld, H.S. & Young, R.D. (2003) Ecophysiology of the herbaceous layer in temperate deciduous forests. In *The Herbaceous Layer in Forests of Eastern North America*. (eds F.S. Gilliam & M.R. Roberts), pp. 38-90. Oxford University Press, New York.
- Nordin, A., Nasholm, T., & Ericson, L. (1998) Effects of simulated N deposition on understorey vegetation of a boreal coniferous forest. *Functional Ecology*, 12, 691-699.
- Oguchi, R., Hikosaka, K., Hiura, T., & Hirose, T. (2006) Leaf anatomy and light acclimation in woody seedlings after gap formation in a cool-temperate deciduous

- forest. *Oecologia*, 149, 571-582.
- Phillips, D.L. & Shure, D.J. (1990) Patch-size effects on early succession in southern Appalachian forests. *Ecology*, 71, 204-212.
- Pickett, S.T.A. & White, P.S. (1985) The ecology of natural disturbance and patch dynamics. Academic press, New York.
- Poorter, H., Niinemets, U., Poorter, L., Wright, I.J., & Villar, R. (2009) Causes and consequences of variation in leaf mass per area (LMA): a meta-analysis. *New Phytologist*, 182, 565-588.
- Prescott, C.E. (2002) The influence of the forest canopy on nutrient cycling. *Tree Physiology*, 22, 1193-1200.
- R development Core Team (2006) R: a language and environment for statistical computing. R foundation for statistical computing, Vienna, Austria. ISBN 3-900051-07-0. <http://www.R-project.org>.
- Richards, L.A. & Coley, P.D. (2007) Seasonal and habitat differences affect the impact of food and predation on herbivores: a comparison between gaps and understory of a tropical forest. *Oikos*, 116, 31-40.
- Ritter, E. (2005) Litter decomposition and nitrogen mineralization in newly formed gaps in a Danish beech (*Fagus sylvatica*) forest. *Soil Biology & Biochemistry*, 37, 1237-1247.

- Rothstein, D.E. & Zak, D.R. (2001a) Photosynthetic adaptation and acclimation to exploit seasonal periods of direct irradiance in three temperate, deciduous-forest herbs. *Functional Ecology*, 15, 722-731.
- Rothstein, D.E. & Zak, D.R. (2001b) Relationships between plant nitrogen economy and life history in three deciduous-forest herbs. *Journal of Ecology*, 89, 385-394.
- Schowalter, T.D. & Ganio, L.M. (1999) Invertebrate communities in a tropical rain forest canopy in Puerto Rico following Hurricane Hugo. *Ecological Entomology*, 24, 191-201.
- Shibata, H., Kirikae, M., Tanaka, Y., Sakuma, T., & Hatano, R. (1998) Proton budgets of forest ecosystems on volcanogenous regosols in Hokkaido, northern Japan. *Water Air and Soil Pollution*, 105, 63-72.
- Shure, D.J. & Phillips, D.L. (1991) Patch size of forest openings and arthropod populations. *Oecologia*, 86, 325-334.
- Shure, D.J. & Wilson, L.A. (1993) Patch-size effects on plant phenolics in successional openings of the southern Appalachians. *Ecology*, 74, 55-67.
- Sipura, M. & Tahvanainen, J. (2000) Shading enhances the quality of willow leaves to leaf beetles - but does it matter? *Oikos*, 91, 550-558.
- Skillman, J.B., Strain, B.R., & Osmond, C.B. (1996) Contrasting patterns of photosynthetic acclimation and photoinhibition in two evergreen herbs from a winter

deciduous forest. *Oecologia*, 107, 446-455.

Sousa, W.P. (1984) The role of disturbance in natural communities. *Annual Review of Ecology and Systematics*, 15, 353-391.

Takafumi, H. & Hiura, T. (2009) Effects of disturbance history and environmental factors on the diversity and productivity of understory vegetation in a cool-temperate forest in Japan. *Forest Ecology and Management*, 257, 843-57.

Topp, G.C. & Davis, J.L. (1985) Measurement of soil-water content using time-domain reflectometry (TDR) - a field-evaluation. *Soil Science Society of America Journal*, 49, 19-24.

Topp, G.C., Davis, J.L., & Annan, A.P. (1980) Electromagnetic determination of soil water content: measurements in coaxial transmission lines. *Water Resources Research*, 16, 574-582.

Treutter, D. (2006) Significance of flavonoids in plant resistance: a review. *Environmental Chemistry Letters*, 4, 147-157.

Uemura, S. (1994) Patterns of leaf phenology in forest understory. *Canadian Journal of Botany*, 72, 409-414.

Weiss, S.B., Murphy, D.D., & White, R.R. (1988) Sun, slope, and butterflies: topographic determinants of habitat quality for *Euphydryas editha*. *Ecology*, 69, 1486-1496.

Wright, E.F., Coates, K.D., & Bartemucci, P. (1998) Regeneration from seed of six tree species in the interior cedar-hemlock forests of British Columbia as affected by substrate and canopy gap position. *Canadian Journal of Forest Research*, 28, 1352-1364.

Yamasaki, M. & Kikuzawa, K. (2003) Temporal and spatial variations in leaf herbivory within a canopy of *Fagus crenata*. *Oecologia*, 137, 226-232.

Figure legend

Fig. 1. Relationships between dry mass (g) of herbivorous insects and gap size (m²).

Symbols represent collected data, and lines indicate the best-fit and significant models ($P < 0.05$). The best-fit models were selected from among linear and log-linear models using the AIC. All selected models were linear and significant. Significance of differences was determined by the likelihood ratio test ($P < 0.05$). Proportions of total herbivorous insect dry mass were Lepidoptera (81.8%), Orthoptera (15.0%), and Coleoptera (3.2%).




Table 1. Relationships between gap size (m²) and canopy openness, soil nitrogen, and soil water content. Values are means \pm SE ($n = 24$) of plots. Coefficients, P -values for best-fit significant models are shown. The best-fit models were selected among linear and log-linear models using the AIC.

Environment factors	Measurement	Average	Coefficients	P	Selected model
Canopy openness (%)	Aug. 2005	19.71 \pm 1.51	2.97	****	Log-linear
NO ₃ ⁻ -N (mg/cm ³)	Jul. 2005 (content of initial)	0.09 \pm 0.01	0.01	***	Log-linear
	Jul. 2005-Oct. 2005	83.22 \pm 7.15	9.60	****	Log-linear
	Oct. 2005-Oct. 2006	165.37 \pm 19.39	22.56	***	Log-linear
NH ₄ ⁺ -N (mg/cm ³)	Jul. 2005 (content of initial)	0.41 \pm 0.02	-	-	Null
	Jul. 2005-Oct. 2005	1.53 \pm 0.23	0.32	****	Log-linear
	Oct. 2005-Oct. 2006	4.86 \pm 0.65	-	-	Null
Soil water content (%)	Aug. 2005	19.68 \pm 0.89	-	-	Null
	Nov. 2005	26.78 \pm 0.51	-	-	Null
	Aug. 2006	25.88 \pm 0.52	-	-	Null

P : The likelihood ratio test was used to test the difference in deviance between the selected model and the null model.

: $p < 0.01$, *: $p < 0.001$.

Table2. Relationships between gap size (m²) and concentrations of condensed tannins and total phenolics, LMA, C:N ratio, and nitrogen content. Values are means ± SE (n = 24) of plots. Coefficients, *P*-values and best fit significant models are shown. The best fit models were selected from linear and log-linear models using the AIC.

Species name	Leaf expansion	Leaf traits	Average	Coefficients	<i>P</i>	Selected model
<i>Trillium tschonoskii</i>	1st	Condensed tannins (mg/mm ²)	0.022 ± 0.001	0.001	n.s.	Log-linear
		Total phenolics (mg/mm ²)	6.560 ± 0.327	0.003	n.s.	Linear
		LMA (mg/mm ²)	221.033 ± 4.497	3.687	n.s.	Log-linear
		CN ratio	13.476 ± 0.176	0.180	n.s.	Log-linear
		N (mg/mm ²)	7.000 ± 0.134	-	-	Null
<i>Smilacina japonica</i>	2nd	Condensed tannins (mg/mm ²)	0.103 ± 0.007	-	-	Null
		Total phenolics (mg/mm ²)	5.950 ± 0.310	-	-	Null
		LMA (mg/mm ²)	218.366 ± 7.447	6.192	n.s.	Log-linear
		CN ratio	15.796 ± 0.415	-	-	Null
		N (mg/mm ²)	5.800 ± 0.237	0.197	n.s.	Log-linear
<i>Polygonatum odoratum</i> var. <i>maximowiczii</i>	3rd	Condensed tannins (mg/mm ²)	0.903 ± 0.088	-	-	Null
		Total phenolics (mg/mm ²)	13.350 ± 1.504	-	-	Null
		LMA (mg/mm ²)	160.032 ± 7.388	3.525	n.s.	Log-linear
		CN ratio	12.384 ± 0.381	-	-	Null
		N (mg/mm ²)	5.800 ± 0.347	0.002	n.s.	Linear
<i>Desmodium podocarpum</i>	4th	Condensed tannins (mg/mm ²)	0.078 ± 0.008	0.002	****	Linear
		Total phenolics (mg/mm ²)	3.760 ± 0.180	0.039	****	Linear
		LMA (mg/mm ²)	209.746 ± 4.804	0.141	***	Linear
		CN ratio	12.384 ± 0.381	0.414	*	Log-linear
		N (mg/mm ²)	7.100 ± 0.191	0.003	n.s.	Linear
<i>Acer mono</i>	5th	Condensed tannins (mg/mm ²)	0.905 ± 0.115	0.094	*	Log-linear
		Total phenolics (mg/mm ²)	22.130 ± 1.077	1.402	***	Log-linear
		LMA (mg/mm ²)	301.445 ± 9.978	15.787	****	Log-linear
		CN ratio	12.412 ± 0.135	-	-	Null
		N (mg/mm ²)	6.900 ± 0.237	0.345	****	Log-linear
<i>Quercus crispula</i>	6th	Condensed tannins (mg/mm ²)	0.517 ± 0.076	0.002	***	Linear
		Total phenolics (mg/mm ²)	33.270 ± 1.721	1.882	**	Log-linear
		LMA (mg/mm ²)	338.024 ± 10.854	0.208	****	Linear
		CN ratio	19.709 ± 0.242	-0.003	*	Log-linear
		N (mg/mm ²)	8.500 ± 0.340	0.007	****	Linear

P: The likelihood ratio test was used to test the difference in deviance between the selected model and the null model.

*: *P* < 0.1, **: *P* < 0.05, ***: *P* < 0.01, ****: *P* < 0.001.

P-values were corrected for multiple pairwise testing with the sequential Bonferroni technique.

Intercepts and coefficients of tannins and total phenolics content, LMA, and nitrogen content are 10⁴ times.

Table 3. Relationships between herbivory and gap size (m²).
 Values shown are coefficients and *P*-values for best-fit and significant models.
 The best-fit models were selected among linear model and log-linear models using the AIC.

Species name	Leaf expansion	Coefficients	<i>P</i>	Selected model
<i>Trillium tschonoskii</i>	1st	0.072	***	Log-linear
<i>Smilacina japonica</i>	2nd	0.137	****	Log-linear
<i>Polygonatum odoratum</i> var. <i>maximowiczii</i>	3rd	-	-	Null
<i>Desmodium podocarpum</i>	4th	-0.001	***	Linear
<i>Acer mono</i>	5th	-	-	Null
<i>Quercus crispula</i>	6th	-0.056	***	Linear

P: The likelihood ratio test was used to test the difference in deviance between the selected model and the null model.

: $p < 0.01$, *: $p < 0.001$.

Table 4. Relationships between herbivory and concentrations of condensed tannins and total phenolics, LMA, C:N ratio, and nitrogen content. Values shown are coefficients and *P*-values for best-fit and significant models. The best-fit models were selected from among linear model and log-linear models using the AIC.

Species name	Leaf expansion	Leaf traits	Coefficients	<i>P</i>	Selected model
<i>Trillium tschonoskii</i>	1st	Condensed tannins (mg/mm ²)	-	-	Null
		Total phenolics (mg/mm ²)	927.7	n.s.	Log-linear
		LMA (mg/mm ²)	-	-	Null
		CN ratio	0.1	n.s.	Linear
		N (mg/mm ²)	-	-	Null
<i>Smilacina japonica</i>	2nd	Condensed tannins (mg/mm ²)	-	-	Null
		Total phenolics (mg/mm ²)	-	-	Null
		LMA (mg/mm ²)	-	-	Null
		CN ratio	-	-	Null
		N (mg/mm ²)	-	-	Null
<i>Polygonatum odoratum</i> var. <i>maximowiczii</i>	3rd	Condensed tannins (mg/mm ²)	-	-	Null
		Total phenolics (mg/mm ²)	-	-	Null
		LMA (mg/mm ²)	-165.9	****	Linear
		CN ratio	-0.3	*	Linear
		N (mg/mm ²)	-	-	Null
<i>Desmodium podocarpum</i>	4th	Condensed tannins (mg/mm ²)	-2884.1	n.s.	Log-linear
		Total phenolics (mg/mm ²)	-181.6	n.s.	Log-linear
		LMA (mg/mm ²)	-	-	Null
		CN ratio	-0.3	****	Linear
		N (mg/mm ²)	585.3	*	Log-linear
<i>Acer mono</i>	5th	Condensed tannins (mg/mm ²)	-	-	Null
		Total phenolics (mg/mm ²)	-	-	Null
		LMA (mg/mm ²)	-	-	Null
		CN ratio	-1.5	***	Log-linear
		N (mg/mm ²)	-	-	Null
<i>Quercus crispula</i>	6th	Condensed tannins (mg/mm ²)	-	-	Null
		Total phenolics (mg/mm ²)	-159.0	**	Log-linear
		LMA (mg/mm ²)	-	-	Null
		CN ratio	-3.7	****	Log-linear
		N (mg/mm ²)	1055.8	n.s.	Log-linear

P: The likelihood ratio test was used to test the difference in deviance between the selected model and the null model.

*: *p*<0.1, **: *p*< 0.05, ***: *p*<0.01, ****: *p*< 0.001.

P-values were corrected for multiple pairwise testing with the sequential Bonferroni technique.