



Title	Diazepam-Induced Increases of Synaptic Efficacy in the Hippocampal-Medial Prefrontal Cortex Pathway Are Associated With Its Anxiolytic-like Effect in Rats
Author(s)	Shikanai, Hiroki; Izumi, Takeshi; Matsumoto, Machiko; Togashi, Hiroko; Yamaguchi, Taku; Yoshida, Takayuki; Yoshioka, Mitsuhiro
Citation	Journal of Pharmacological Sciences, 114(3), 341-346 <a href="https://doi.org/10.1254/jphs.10182FP">https://doi.org/10.1254/jphs.10182FP</a>
Issue Date	2010
Doc URL	<a href="http://hdl.handle.net/2115/47360">http://hdl.handle.net/2115/47360</a>
Type	article (author version)
File Information	JPS114-3_341-346.pdf



[Instructions for use](#)

**Diazepam-induced increases of synaptic efficacy in the hippocampal-mPFC pathway are associated with its anxiolytic effect.**

Hiroki Shikanai<sup>1)</sup>, Takeshi Izumi<sup>1)</sup>, Machiko Matsumoto<sup>2)</sup>, Hiroko Togashi<sup>2)</sup>, Taku Yamaguchi<sup>1)</sup>, Takayuki Yoshida<sup>1)</sup>, Mitsuhiro Yoshioka<sup>1)</sup>

1) Department of Neuropharmacology, Hokkaido University Graduate School of Medicine, Kita-15, Nishi-7, Kita-ku, Sapporo 060-8638, Japan

2) Department of Pharmacology, Faculty of Pharmaceutical Sciences, Health Sciences University of Hokkaido, Ishikari-Tobetsu 061-0293, Japan

**Abstract.**

The medial prefrontal cortex (mPFC) has recently been shown to be an important brain region for emotional function as well as cognitive ability. In previous experiments, we studied the population spike amplitude (PSA) in the mPFC induced by stimulation of the CA1/subicular region as an index of synaptic efficacy in the hippocampal–mPFC pathway. In the present study, we investigated the relationship between the anxiolytic effect of diazepam and the changes of synaptic efficacy in this pathway. In contextual fear conditioning tests, diazepam (0.1 mg/kg) was not effective for fear-related freezing behavior. At a dose of 0.5 mg/kg, diazepam decreased freezing behavior 20 min after administration, with no discernible effect 30 min after administration. In electrophysiological experiments, 0.1 mg/kg diazepam had no effect on the PSA in the mPFC. In contrast, 0.5 mg/kg diazepam increased the PSA in the mPFC within 30 min of administration; however, this PSA increase was attenuated over the 30-min period. Based on these results, we propose that the diazepam-induced PSA increase in the mPFC is associated with its anxiolytic-like effect.

**Keywords:** electrophysiology, hippocampal – medial prefrontal cortex (mPFC) pathway, behavior, anxiety, prefrontal cortex

## **Introduction**

Functional brain imaging techniques, such as positron emission tomography and functional magnetic resonance imaging, are used to investigate the pathologies of psychiatric diseases. Investigations of brain activity in patients with psychiatric disorders (e.g., schizophrenia, depression, and anxiety disorders) have revealed a dysfunction in the frontal cerebral cortex, which has been termed hypofrontality. However, this phenomenon alone cannot explain the etiology of these psychiatric disorders.

When we carry out a pharmacological investigation in rats that focuses on the prefrontal cortex, it is important to consider whether or not the rodent prefrontal cortex is comparable to that in primates. In the past, prefrontal cortex functionality was thought to be unique to primates. However, recent tracer and lesion studies have revealed that rats have a functional prefrontal cortex that includes not only features of the medial and orbital areas that are observed in primates but also includes several features of the primate dorsolateral prefrontal cortex (1).

Previous studies have indicated that the medial prefrontal cortex (mPFC) shares mutual projections with the amygdala complex (2). In addition, previous studies have indicated that the mPFC receives input from the hippocampus in rats (3). Both of these neural circuits are important for emotional regulation. Previous studies have also suggested that the mPFC has inhibitory control over the amygdala during fear-related

behavior. For example, an mPFC lesion (4) in conjunction with inhibition of protein synthesis in the mPFC (5) blocked the extinction of conditioned fear. Furthermore, electrical stimulation of the mPFC inhibited the expression of conditioned fear (6, 7). Previous studies have also reported that the extracellular concentration of dopamine and serotonin (5-hydroxytryptamine: 5-HT) in the mPFC increased during freezing behavior as the expression of conditioned fear (8, 9) and that a selective serotonin reuptake inhibitor (SSRI), which increased 5-HT release in the mPFC, reduced conditioned fear-induced freezing behavior (10). These behavioral and neurochemical studies suggest that the mPFC plays an important role in the inhibitory regulation of fear-related emotions.

In contrast to the amygdala, the mPFC and several hippocampal structures are necessary for the execution of memory-related behaviors. The CA1/subicular region of the ventral hippocampus plays an important role in the acquisition of spatial memory (11). Furthermore, this region is connected to the mPFC (3), and this connection could contribute to the encoding of spatial information (12). Taking these results together, we hypothesized that the connections in the hippocampal–mPFC pathway may be necessary for memory-dependent emotional regulation.

We have studied the population spike amplitude (PSA) in the mPFC that is induced by stimulation of the hippocampus because it is a measure of synaptic efficacy in this pathway (Fig. 1). Electrophysiological experiments in rats showed that stimulation of the CA1/subicular region produces negative-going field potentials in the mPFC. Because the excitatory response of the mPFC was blocked by a selective alpha-amino-3-hydroxyl-5-methylisoxazole-4-propionate (AMPA) receptor antagonist, this negative-going peak reflects the excitatory input from the hippocampus to the

mPFC via glutamatergic neurons (3). In the previous experiment, we studied the effect of an SSRI on synaptic efficacy in the mPFC. A single administration of the SSRI fluvoxamine induced an increase of the PSA in the mPFC (13).

Based on these studies, we hypothesized that the synaptic efficacy in the hippocampal–mPFC pathway may be involved in emotional regulation. To test this hypothesis, we investigated the effect of diazepam, an anxiolytic-like agent that is widely used in clinical practice and laboratory research (14), on synaptic efficacy in the hippocampal–mPFC pathway.

## **Materials and Methods**

### *Animals*

Male Wistar rats (280 – 410 g and 10 – 14-week-old) were used for these experiments (Slc:Wistar/ST; Shizuoka Laboratory Animal Center, Hamamatsu). The animals were housed in a 12-h light-dark cycle (lights on from 19:00 – 7:00) at  $21 \pm 2^\circ\text{C}$  and were given free access to food and water. All experiments in the present study were conducted in accordance with the standards established by the Guide for the Care and Use of Laboratory Animals of Hokkaido University.

### *Drugs*

Diazepam (0.1, 0.5, or 1.0 mg/kg; Wako Pure Chemical Industries, Ltd., Osaka) or flumazenil (10 or 30 mg/kg; Tocris, Hung Road, Bristol, UK) suspended in 1% arabic gum (Nacalai Tesque, Kyoto) were intraperitoneally (i.p.) injected. Urethane (20%; Sigma, St. Louis, MO, USA) was dissolved in 0.9% saline and injected i.p. Lidocaine was purchased from Astra Zeneca (Osaka), and the formaldehyde neutral buffer solution

was purchased from Sigma Aldrich (Tokyo).

#### *Contextual conditioned fear test*

A chamber (30 × 30 × 27 cm) constructed from aluminum (side walls) and Plexiglass (rear wall, ceiling, and front door) was used to evaluate fear-related behavior. The floor of the chamber consisted of 27 stainless steel rods wired to a shock generator (Medical Agent, Kyoto) that delivered foot shocks. On day 1, the rats were exposed to the chamber for 5 min and were given five foot shocks (2 s, 0.5 mA, at 30-s intervals). Thirty seconds after the final shock, freezing behavior was counted at 5-s intervals for 5 min. Freezing was defined as the absence of all movement, with the exception of respiration and whisker movements. After an observation of freezing, the rats were returned to their home cages. On day two, vehicle (arabic gum, 2.5 ml/kg, i.p.) or diazepam (0.1 or 0.5 mg/kg, i.p.) was injected. At 20 or 30 min after the injection, the rats were re-exposed to the chamber, and freezing behavior was counted at 5-s intervals for 10 min. The freezing time was calculated as a percentage of the total time.

#### *Measurement of the PSA*

Each rat was anesthetized with urethane (1.0 g/kg, i.p.) prior to insertion of a tracheal cannula. Next, each rat was fixed in a stereotaxic frame (Narishige Scientific Instrument Lab., Tokyo). The stereotaxic coordinates using bregma and lambda in the same horizontal plane) were derived from the Paxinos and Watson's brain atlas. In addition to urethane, the local anesthetic lidocaine was used during the scalp dissection. For each rats, the body temperature was maintained at 37°C using a thermoregulated heating pad (TMO35-1; Takasago Ltd., Kanagawa). Bipolar stimulating electrodes

(diameter: 100  $\mu\text{m}$ , resistance: 20  $\text{k}\Omega$ , material: stainless steel; Eiko Science, Tokyo) with a tip separation of 500  $\mu\text{m}$  were placed in the CA1/subicular region of the ventral hippocampus (Fig. 2: 6.0-mm posterior to the bregma, 5.6-mm lateral, and 4.0 – 7.0-mm ventral from the cortical surface) according to Paxinos and Watson's brain atlas. A recording electrode (diameter: 100  $\mu\text{m}$ , resistance: 20  $\text{k}\Omega$ , material: stainless steel; Eiko Science, Tokyo) was lowered into the mPFC (Fig. 2: 3.3-mm anterior to the bregma, 0.8 mm lateral, and 3.3-mm ventral from the cortical surface). Electrode positions were optimized to record the maximal field response, which was evoked at a frequency of 0.1 Hz (250- $\mu\text{s}$  duration) with the first negative peak latency of  $20.05 \pm 0.5$  ms (Fig. 1). Field responses were amplified and monitored with an oscilloscope (VC-10; Nihon Kohden, Tokyo). The evoked field potentials were converted from analog to digital data and were averaged with a data analyzing system (Power Lab System; ADI Instruments Pty., Ltd., Castle Hill, Australia). Then, the PSA of the first negative peak was measured (Fig. 1). After electrode insertion, the input/output characterization of each rat was assessed. The test stimuli were delivered every 30 s at an intensity that produced a 60% maximal response that was induced by the maximal stimulation intensity (100 – 640  $\mu\text{A}$ ). The vehicle or diazepam was injected after sampling the data for 30 min, and changes in the PSA were observed for 60 min. Seven successive responses were averaged and collected every 5 min throughout the experiment. After the experiments, the brains were removed and fixed in a formaldehyde neutral buffer solution, followed by confirmation of a proper electrode insertion location using a microscope (Fig. 2). The evoked potentials were expressed as a percentage of the baseline level, which was determined immediately prior to drug administration.

### *Data analyses*

All of the results are given as the mean  $\pm$  S.E.M. To compare the two groups, Student's t-test was used. For multiple group comparisons, a one-way ANOVA (followed by a Tukey post-hoc test) was used. The statistical significance was set at  $P < 0.05$ .

## **Results**

### *Dose- and time-dependent effects of diazepam on fear related behavior*

Fear-conditioning test was performed to confirm the optimal dose for the anxiolytic-like effect of diazepam. Twenty minutes after administration, 0.1 mg/kg diazepam had no effect on the freezing behavior, whereas 0.5 mg/kg diazepam significantly reduced the freezing behavior compared to that of the control group [one-way ANOVA,  $F(3,19) = 4.00$ ,  $P < 0.05$ ; post-hoc test, control vs. diazepam (0.5 mg/kg),  $P = 0.021$ ] (Fig. 3A1)]. However, 1.0 mg/kg of diazepam did not reduce the freezing behavior. This effect is likely due to the sedative effects of diazepam. In contrast to the 20-min time point, 30 min after administration, 0.5 mg/kg diazepam did not reduce freezing behavior [Student's t-test,  $F(1,9) = 0.78$ ,  $P > 0.05$ ] (Fig. 3B)]. In addition, concurrent administration of 0.5 mg/kg diazepam and 30 mg/kg flumazenil, which is a benzodiazepine-receptor antagonist, blocked the effect of 0.5 mg/kg of diazepam on freezing [one-way ANOVA,  $F(2,11) = 3.61$ ,  $P < 0.05$ ; post-hoc test, diazepam (0.5 mg/kg) vs. diazepam (0.5 mg/kg) + flumazenil (30 mg/kg),  $P = 0.021$ ] (Fig. 3A2)], whereas 10 and 30 mg/kg flumazenil alone had no effect on freezing behavior (data not shown).

### *Effect of diazepam on the PSA in the hippocampal–mPFC pathway*

We examined the effects of diazepam on the synaptic efficacy of the hippocampal–mPFC



pathway. Systemic administration of 0.1 mg/kg diazepam had no effect on the PSA in this pathway, whereas 0.5 mg/kg diazepam significantly increased the PSA [maximum responses:  $167.9 \pm 29.9\%$ ; one-way ANOVA,  $F(2,12) = 4.40$ ,  $P < 0.05$ ; post-hoc test, control vs. diazepam (0.5 mg/kg),  $P = 0.010$ ]. The PSA value returned to the basal level after 20 min, and the increase of the PSA was significantly inhibited by the concurrent administration of 0.5 mg/kg diazepam and 10 mg/kg flumazenil (Fig. 4). We did not perform the electrophysiological experiment at 1.0 mg/kg diazepam because sedative effects of diazepam occurred in the conditioned fear tests at this dose.

## Discussion

In the present study, we investigated the effect of diazepam, which is widely used clinically as an anxiolytic drug, on conditioned fear, an animal model of anxiety. At a dose of 0.1 mg/kg, diazepam had no effect on conditioned fear-induced freezing behavior. At a dose of 0.5 mg/kg, diazepam significantly reduced freezing 20 min after administration; however, 30 min after administration, 0.5 mg/kg diazepam failed to reduce freezing. In addition, 1.0 mg/kg of diazepam did not reduce freezing. This effect was likely due to the sedative effect at this dose (Fig. 3A1).

In electrophysiological experiments, 0.1 mg/kg diazepam had no effect on the PSA in the hippocampal–mPFC pathway. However, 0.5 mg/kg diazepam increased the PSA at 20 min after administration, and the PSA returned these results, it is conceivable that the effect of diazepam on the synaptic efficacy in the hippocampal–mPFC pathway is similar to its anxiolytic-like effect in a time and dose-dependent manner. In previous pharmacokinetic studies in rats, after a single intraperitoneal injection, diazepam was rapidly absorbed, and its blood concentration

reached the maximum in 10 min (15). In addition, the diazepam was rapidly cleared from the plasma and brain, with an overall half-life of 0.88 and 0.89 h, respectively (16). Specific blood concentrations may be needed for the anxiolytic and electrophysiological effects of diazepam.

A wide consensus exists that the amygdala mediates fear learning (17). In addition, the hippocampus has bidirectional connections with the amygdala. The majority of the projections from the hippocampus to the amygdala originate at the ventral hippocampus (18). Furthermore, several studies have indicated that the hippocampus is involved in the regulation of anxiety and fear. The expression of contextual conditioned fear has been shown to be abolished by a hippocampal-lesion after conditioning (19). The lesions of the ventral hippocampal efferents to the amygdala have been shown to prevent contextual fear conditioning (20), suggesting that ventral hippocampus–amygdala interactions mediate fear learning. It has been reported that the hippocampus CA1 field unidirectionally projected to the mPFC (3, 21), and this pathway has been speculated to be relevant to spatial short-term memory and working memory (12). Additionally, the mPFC has been implicated in fear memory extinction. For example, electric lesions or chemical inactivation of the mPFC have been shown to impair the extinction process (4, 22), and fear extinction-induced c-Fos expression in the mPFC has been shown to be blocked by an infusion of a protein synthesis inhibitor into the mPFC (5).

Recent studies have also reported the inhibitory control of the mPFC on the amygdala in fear extinction. Tracer studies have revealed the distribution of the mPFC projections to the amygdala (2, 23). Given that most of the mPFC projections to the amygdala are excitatory (24, 25), it has been proposed that the mPFC inhibits the

amygdala by activation of inhibitory interneurons within the amygdala (26, 27). It was also reported that mPFC stimulation can suppress the firing of pyramidal cells in the amygdala, which respond to a fear-conditioned stimulus (28, 29).

Synaptic plasticity in the hippocampal–mPFC pathway has been assessed by the formation of long-term potentiation (LTP), which depends on the activity of the specific afferent. Acute stress has been shown to inhibit LTP formation in the hippocampal–mPFC pathway (30). Furthermore, our previous study indicated that the PSA in this pathway was reduced during exposure to the fearconditioned stimulus and that this reduction was attenuated in parallel with the extinction of conditioned fear (31). Thus, we inferred that the hippocampal–mPFC pathway was important for the regulation of emotion, especially during fear extinction.

The mechanism of the anxiolytic-like action of diazepam commonly involves the enhancement of the inhibition of neurons via gamma aminobutyric acid A (GABAA) receptors (32). The GABAA receptor has a benzodiazepine binding site, and upon benzodiazepine binding, the frequency of chloride channel opening increases, resulting in the inactivation of neurons. Because of this mechanism, it is plausible that the PSA decreases as a consequence of diazepam administration. In fact, it has been reported that a local injection of the GABAA agonist muscimol causes a PSA decrease in the hippocampal– mPFC pathway (33).

Unexpectedly, the PSA actually increased in the present experiment. We hypothesized that diazepam did not directly act on the mPFC, but, instead, acted on other brain regions that modulated the hippocampal–mPFC pathway. For example, we previously reported that LTP formation in this pathway was augmented by a 5-HTergic lesion of the whole brain using a 5-HT neurotoxin, 5,7-DHT (34). From these results, it is

impossible to determine if diazepam acted on the region that suppressed the mPFC, such as the 5-HTnergic neurons of the raphe, resulting in the increase of the PSA in the hippocampal–mPFC pathway. Therefore, diazepam indirectly activated mPFC neurons; as a result of this effect, neurons in the amygdala were inhibited, and a reduction in freezing behavior was observed.

We believe that the effect of diazepam in the present study was an anxiolytic-like effect, not an amnesic effect, although it has been reported that diazepam could lead to anterograde amnesia. However, by observing animal behavior, we could not infer whether the effect of diazepam on conditioned freezing was derived from its anxiolytic-like effect or from its amnesic effect. We previously reported that another type of anxiolytic agent, fluvoxamine, also increased the PSA in the hippocampal–mPFC pathway. In addition, a clinical study indicated that the anxiolytic effect of fluvoxamine was without amnesia (35). From these studies, there is support for the idea that diazepam decreased freezing through its anxiolytic-like effect, which is the same effect from an SSRI

In conclusion, the anxiolytic agent diazepam enhances synaptic efficacy in the hippocampal–mPFC pathway *in vivo*. Based on the results from previous studies, we speculate that the anxiolytic-like effect of diazepam is derived from the enhancement of the suppressive activity of the mPFC on the amygdala.

## References

1. Uylings HB, Groenewegen HJ, Kolb B. Do rats have a prefrontal cortex? *Behav Brain Res.* 2003;146:3-17

2. McDonald AJ, Mascagni F, Guo L. Projection of the medial and lateral prefrontal cortices in the amygdala: a Phaseolus vulgaris leucoagglutinin study in the rat. *Neuroscience*. 1996;71:55-75
3. Jay TM, Glowinski J, Thierry AM. Selectivity of the hippocampal projection to the prelimbic area of the prefrontal cortex in the rat. *Brain Res*. 1989;505:337-340
4. Morgan MA, Ledoux JE. Differential contribution of dorsal and ventral medial prefrontal cortex to the acquisition and extinction of conditioned fear in rats. *Behav Neurosci*. 1995;109:681-688
5. Santini E, Ge H, Ren K, Peña de Ortiz S, Quirk GJ. Consolidation of fear extinction requires protein synthesis in the medial prefrontal cortex. *J Neurosci*. 2004;24:5704-5710
6. Milad MR, Vidal-Gonzalez I, Quirk GJ. Electrical stimulation of medial prefrontal cortex reduces conditioned fear in a temporally specific manner. *Behav Neurosci*. 2004;118:389-394
7. Quirk GJ, Lithtik E, Pelletier JG, Pare D. Stimulation of medial prefrontal cortex decreases the responsiveness of central amygdala output neurons. *J Neurosci*. 2003;23:8800-8807
8. Yoshioka M, Matsumoto M, Togashi H, Saito H. Effects of Conditioned Fear

Stress on 5-HT Release in the Rat Prefrontal Cortex. *Pharmacol Biochem Behav.* 1995;51:515-519

9. Yoshioka M, Matsumoto M, Togashi H, Saito H. Effect of conditioned fear stress on dopamine release in the rat prefrontal cortex. *Neurosci Lett.* 1996;209:201-203

10. Hashimoto S, Inoue T, Koyama T. Serotonin reuptake inhibitor reduces conditioned fear stress-induced freezing behavior in rats. *Psychopharmacology (Berl).* 1996;123:182-186

11. Floresco SB, Seamans JK, Phillips AG. Differential effects of lidocaine infusions into the ventral CA1/subiculum or the nucleus accumbens on the acquisition and retention of spatial information. *Behav Brain Res.* 1996;81:163-171

12. Floresco SB, Seamans JK, Phillips AG. Selective roles for hippocampal, prefrontal cortical, and ventral striatal circuits in radial-arm maze tasks with or without a delay. *J Neurosci.* 1997;17:1880-1890

13. Ohashi S, Matsumoto M, Otani H, Mori K, Togashi H, Ueno K, Kaku A, Yoshioka M. Changes in synaptic plasticity in the rat hippocampo-medial prefrontal cortex pathway induced by repeated treatments with fluvoxamine. *Brain Res.* 2002;949:131-138

14. Nishino T, Takeuchi T, Takechi K, Kamei C. Evaluation of anxiolytic-like

effects of some short-acting benzodiazepine hypnotics in mice. *J Pharmacol Sci.* 2008;107:349-354

15. Walker MC, Tong X, Brown S, Shorvon SD, Patsalos PN. Comparison of single- and repeated- dose pharmacokinetics of diazepam. *Epilepsia.* 1998;39:283-289

16. Friedman H, Abernethy DR, Greenblatt DJ, Shader RI. The pharmacokinetics of diazepam and demethyldiazepam in rat brain and plasma. *Psychopharmacology (Berl).* 1986;88:267-270

17. LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci.* 2000;23:155-184

18. Pikkarainen M, Rönkkö S, Savander V, Insausti R, Pitkänen A. Projections from the lateral, basal, and accessory basal nuclei of the amygdala to the hippocampal formation in rat. *J Comp Neurol.* 1999;403:229-260

19. Kim JJ, Fanselow MS. Modality-specific retrograde amnesia of fear. *Science.* 1992;256:675-677

20. Maren S, Fanselow MS. Synaptic plasticity in the basolateral amygdala induced by hippocampal formation stimulation in vivo. *J Neurosci.* 1995;15:7548-7564

21. Jay TM, Thierry AM, Wiklund L, Glowinski J. Excitatory Amino Acid Pathway from the Hippocampus to the Prefrontal Cortex. Contribution of AMPA Receptors in

Hippocampo-prefrontal Cortex Transmission. *Eur J Neurosci.* 1992;4:1285-1295

22. Morrow BA, Elsworth JD, Rasmusson AM, Roth RH. The role of mesoprefrontal dopamine neurons in the acquisition and expression of conditioned fear in the rat. *Neuroscience.* 1999;92:553-564

23. Vertes RP. Differential projections of the infralimbic and prelimbic cortex in the rat. *Synapse,* 2004;51:32-58

24. Smith Y, Paré JF, Paré D. Differential innervation of parvalbumin-immunoreactive interneurons of the basolateral amygdaloid complex by cortical and intrinsic inputs. *J Comp Neurol.* 2000;416:496-508

25. Ohashi S, Togashi H, Matsumoto M, Mori K, Ueno K, Yoshioka M. Changes in Synaptic Properties in Cortical-Limbic Communications Induced by Repeated Treatments With Fluvoxamine in Rats. *J Pharmacol Sci.* 2003;92:100-107

26. Rosenkranz JA, Grace AA. Modulation of basolateral amygdala neuronal firing and afferent drive by dopamine receptor activation in vivo. *J Neurosci.* 1999;19:11027-11039

27. Rosenkranz JA, Grace AA. Cellular mechanisms of infralimbic and prelimbic prefrontal cortical inhibition and dopaminergic modulation of basolateral amygdala neurons in vivo. *J Neurosci.* 2002 1;22:324-337



28. Quirk GJ, Gehlert DR. Inhibition of the amygdala: key to pathological states? *Ann N Y Acad Sci.* 2003;985:263-272
29. Rosenkranz JA, Moore H, Grace AA. The prefrontal cortex regulates lateral amygdala neuronal plasticity and responses to previously conditioned stimuli. *J Neurosci.* 2003;23:11054-11064
30. Rocher C, Spedding M, Munoz C, Jay TM. Acute stress-induced changes in hippocampal/prefrontal circuits in rats: effects of antidepressants. *Cereb Cortex.* 2004;14:224-229
31. Koseki H, Matsumoto M, Togashi H, Miura Y, Fukushima K, Yoshioka M. Alteration of synaptic transmission in the hippocampal-mPFC pathway during extinction trials of context-dependent fear memory in juvenile rat stress models. *Synapse.* 2009;63:805-813
32. Nakamichi N, Takarada T, Yoneda Y. Neurogenesis mediated by gamma-aminobutylic acid and glutamate signaling. *J Pharmacol Sci.* 2009;110:133-149
33. Takita M, Kuramochi M, Izaki Y, Ohtomi M. In vivo temporal property of GABAergic neural transmission in collateral feed-forward inhibition system of hippocampal-prefrontal pathway. *Brain Res.* 2007;1150:69-73

34. Ohashi S, Matsumoto M, Togashi H, Ueno K, Yoshioka M. The serotonergic modulation of synaptic plasticity in the rat hippocampo-medial prefrontal cortex pathway. *Neurosci Lett.* 2003;342:179-182

35. Hindmarch I. Cognitive toxicity of pharmacotherapeutic agents used in social anxiety disorder. *Int J Clin Pract.* 2009;63:1085-1094

### **Figure legends**

Figure 1. Typical wave-form of an evoked potential in the mPFC after stimulation of the CA1/subicular region. We assessed the depth of the first negative peak in this wave form as the synaptic efficacy in the mPFC and evaluated this as the population spike amplitude (PSA).

Figure 2. Schematic representation of the locations of the tips of the recording electrodes in the mPFC and the stimulating electrodes in the CA1/subicular region in the ventral hippocampus (open square: control group; solid square: drug administration group).

Figure. 3. Effects of diazepam on the experiment evaluating conditioned fear-induced freezing behavior. A1) At 20 min after administration, diazepam (0.5 mg/kg) reduced the freezing behavior of rats in fear conditioning [one-way ANOVA,  $F(3,19) = 4.00$ ,  $P < 0.05$ ; posthoc test, control vs. DZP (0.5 mg/kg),  $P = 0.021$ ]. A2) Moreover, concurrent administration of diazepam (0.5 mg/kg) and flumazenil (30 mg/kg) did not reduce the

freezing behavior [one-way ANOVA,  $F(2,11) = 3.61$ ,  $P < 0.05$ ; post-hoc test, DZP (0.5 mg/kg) vs. DZP (0.5 mg/kg) + flumazenil (30 mg/kg),  $P = 0.021$ ]. B) At 30 min after administration, diazepam (0.5 mg/kg) did not reduce the freezing behavior of rats in fear conditioning [Student's t-test,  $F(1,9) = 0.78$ ,  $P > 0.05$ ]. The parenthetic digits are the numbers of rats in these experiments. \* $P < 0.05$ : control group vs. DZP (0.5 mg/kg) group, # $P < 0.05$ : DZP (0.5 mg/kg) group vs. DZP (0.5 mg/kg) + flumazenil (30 mg/kg) group. DZP, diazepam.

Figure. 4. Effects of diazepam on the time course of the population spike amplitude (PSA) changes (A) and the AUC from 0 to 60 min in the mPFC (B). Diazepam (0.5 mg/kg) increased the PSA in the mPFC. Inset of panel A indicates a representative wave-form of the DZP (0.5 mg/kg) group. The parenthetic digits are sample numbers in these experiments. \* $P < 0.05$ : control group vs. DZP (0.5 mg/kg) group, one-way ANOVA,  $F(2,12) = 4.40$ ,  $P < 0.05$ ; post-hoc test, control vs. DZP (0.5 mg/kg),  $P = 0.010$ . # $P < 0.05$ : DZP (0.5 mg/kg) group vs. DZP (0.5 mg/kg) + flumazenil (10 mg/kg) group, Student's t-test  $F(1,7) = 4.58$ ,  $P < 0.05$ . DZP, diazepam.

Figures  
Figure 1.




Figure 2.




Figure 3A.




Figure 3B.




Figure 4.

