




Title	International Collaborative Study on Ecology and Evolution in Plant-Insect Interactions
Author(s)	Utsumi, Shunsuke
Citation	北方圏の環境研究に関するシンポジウム. 2011年10月31日（月）. 北海道大学学術交流会館 小講堂. Northern Environmental Research Symposium (Hokkaido-Finland Days: A Bridge for Northern Cooperation). Monday, 31 October, 2011. Hokkaido University Conference Hall.
Issue Date	2011-10-31
Doc URL	http://hdl.handle.net/2115/47652
Type	conference presentation
Note	Session 2.1: Biodiversity and Environmental Protection in the North
File Information	2-1-5_utsumi.pdf



[Instructions for use](#)



Northern Environmental Research Symposium
Sustainability Week 2011 @ Sapporo, Oct 31

International Collaborative Study on Ecology and Evolution in Plant-Insect Interactions

Shunsuke Utsumi
University of Tokyo

Research Fellow of Japan Society for the Promotion of Science


International collaboration in ecological and evolutionary research

Ecosystems have

- ◆ Complexity
- ◆ Variety
- ◆ Generality

Across boarder problems

- ◆ Biodiversity loss
- ◆ Species invasion
- ◆ Climate change



International collaboration offers many opportunities and advantages



- 

Collaboration between Japan and Finland to explore general patterns



Prof Heikki Roininen
(University of Eastern Finland)

Funded by Finnish Government Scholarship, Grant-in-Aid for Scientific Research for JSPS fellowship

Research Question

How multiple species influences evolutionary trajectories of a community member?

[Background]

Organisms are embedded in complex interactions in multiple species community

However, researchers have overwhelmingly focused on coevolution between 2 species (e.g., arms race: plant's defense – insect's counterdefense)

Study organism

Willow leaf beetle

Plagiodera versicolora (Chrysomelidae)

*Distribution: Europe, America, Asia

*Specialist on willow species




Larva



Adult




Regrowth response in Willows



[Regrowth Response to Herbivory]

Willows produce new lateral shoots in response to attack by some herbivore species. As a result, a period of new leaf production is drastically extended.

Evolution of Preference between leaf-age types in Japan




The leaf beetle populations evolutionarily develop diverse preference according to local regrowth intensity of willows


Utsumi *et al.* (2009) *Ecol Lett* **12**: 920-929

Hypothesis

1. Herbivore community structure affects plant regrowth response
2. The leaf beetle populations develop diverse preference according to local regrowth intensity.
3. There is a similar pattern between Japan and Finland even when host plant species are different.



Host plant: *Salix eriocarpa*




Host plant: *Salix caprea*

Why Finland?

- ◆ Willows are very common throughout Finland.
- ◆ Willows are widely distributed in the north hemisphere.
- ◆ The north limit of *Plagiodera versicolora* distribution.
- ◆ Long history of ecological research for insect-willow interactions.


Herbivore species diversity governs the expression of willows' regrowth response



Significant difference
in species composition among sites
[Permutation MANOVA]
JPN: $P < 0.001$, FIN: $P < 0.001$

Higher species diversity
→ Greater plant regrowth intensity



Leaf beetle populations develop diverse preference according to local regrowth intensity



Greater plant regrowth intensity
→ Stronger preference for new leaves

Significant difference
in preference among populations
[ANOVA]
JPN: $P < 0.01$, FIN: $P < 0.05$

Population Genetics in Japan and Finland




Genetic differentiation was detected among populations within a country (mtDNA COI)


[Exact test]: JPN: $P < 0.0001$, FIN: $P < 0.0001$

Experimentally estimated fitness landscape of the leaf beetle

Without Aphids




With Aphids



1. Herbivore species diversity modifies evolutionary trajectory of a community member
2. There is a similar pattern between Japan and Finland even when host plant species are different.

International Collaborative Study



Novelty and Generality

(→...toward sharing concerns for biodiversity loss)

Collaboration between Japan and US to address biological invasion



Prof Tim Craig
(University of Minnesota, Duluth)

Funded by JSPS Core-to-Core program, Grant-in-Aid for Scientific Research for JSPS fellowship

Study organisms

Solidago altissima (Asteraceae)

Uroleucon nigrotuberculatum
(Hemiptera: Aphidae)



Invasive species from US to Japan

Motivation

- ◆ Studying exotic species in both their introduced and native is important.
- ◆ Biological and ecological knowledge of exotic species are often accumulated in the native range institute.


Research Question

- ◆ How exotic plant genotypic diversity contributes to population size of exotic herbivores?
- ◆ How exotic plant genotypic diversity contributes to spatial dynamics of exotic herbivores?


First we examined these questions in a **native habitat** because the US collaborator maintains a variety of *Solidago* genotypes in a common garden)

Genetic variation and spatial spreading dynamics of exotic species are important issues in biological invasion...

Genotypic variation in plant resistance to the aphid



Plant genotypic diversity increased aphid population size



**Greater than a sum of single genotype effects (expected)
= non-additive (synergistic) effect of diverse genotypes**

Mechanism: Source-sink dynamics

Local density-dependent processes determine aphid abundance under isolated condition (monoculture)

(Diverse genotype plot)


A surplus over carrying capacity on a susceptible genotype (source)



moving to a bad quality genotype (sink) where reproductive success is low (walking by wingless morphs)




Synergistic increase in population size



In Japan?

- ◆ Genotypic diversity and composition might be different between native and introduced ranges - **undergoing**
- ◆ Insect movement manner might be different between a native and introduced range due to biotic and abiotic factors - **undergoing**
- ◆ Genetics and spatial spreading dynamics in Japan differ from US?

International Collaborative Study in Invasion Biology



Accumulation of knowledge about differences in genetics, population dynamics, and species interactions between a native and introduced habitat

→ **Condition dependence** in evolutionary and ecologically dynamics (invasive species often show rapid adaptation in an introduced habitat)

→ Insights for adaptive management



**Thanks for your attention
and amazing memories in Finland**