

## HOKKAIDO UNIVERSITY

Title	Dermatan sulfate epimerase 2 is the predominant isozyme in the formation of the chondroitin sulfate/dermatan sulfate hybrid structure in postnatal developing mouse brain			
Author(s)	Akatsu, Chizuru; Mizumoto, Shuji; Kaneiwa, Tomoyuki; Maccarana, Marco; Malmström, Anders; Yamada, Shuhei; Sugahara, Kazuyuki			
Citation	Glycobiology, 21(5), 565-574 https://doi.org/10.1093/glycob/cwq208			
Issue Date	2011-05			
Doc URL	http://hdl.handle.net/2115/47770			
Rights	This is a pre-copy-editing, author-produced PDF of an article accepted for publication in Glycobiology following peer review. The definitive publisher-authenticated version Glycobiology (2011) 21 (5): 565-574 is available online at: http://glycob.oxfordjournals.org/content/21/5/565			
Туре	article (author version)			
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.			
File Information	Supplementary_Data.pdf (Supplementary Data)			



## Supplementary Table I. Disaccharide composition of the CS/DS chains from thalamus and cerebral cortex.

The CS/DS chains in the GAG-peptide preparations from thalamus and cerebral cortex of five mice (7W) were treated with a mixture of CSases ABC and AC-I, or a mixture of CSases AC-I and AC-II. The content of each disaccharide unit was determined by the methods described in the legend to **Fig. 6**.

CS/DS disaccharide unit	Thalamus		Cerebral cortex	
	pmol/mg of acetone powder	(mol%)	pmol/mg of acetone powder	(mol%)
0	18	3.6	38	9.8
С	19	3.8	10	2.6
А	443	89	330	85
iA	ND	_	ND	—
D	1.5	0.30	0.60	0.15
iD	5.5	1.1	1.3	0.34
В	0.77	0.16	0.64	0.17
iB	2.0	0.40	0.63	0.16
Е	6.4	1.3	6.3	1.6
iE	ND	_	ND	
Total CS/DS disaccharide	496	100	387	100

ND, not detected (<0.1 pmol/mg).

-, not occurring.