Title	Mutagenicity of secondary oxidation products of 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate (8-hydroxy-2'-deoxyguanosine 5'-triphosphate)
Author(s)	Hori, Mika; Suzuki, Tetsuya; Minakawa, Noriaki; Matsuda, Akira; Harashima, Hideyoshi; Kamiya, Hiroyuki
Citation	Mutation Research/Fundamental and Molecular Mechanisms of Mutagenesis, 714(1-2), 11-16 https://doi.org/10.1016/j.mrfmmm.2011.05.015
Issue Date	2011-09-01
Doc URL	http://hdl.handle.net/2115/48349
Type	article (author version)
File Information	MR714-1-2_11-16.pdf

[Instructions for use](#)

**Mutagenicity of secondary oxidation products of
8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate (8-hydroxy-2'-
deoxyguanosine 5'-triphosphate)**

**Mika Hori^{a,1}, Tetsuya Suzuki^{a,2}, Noriaki Minakawa^{a,b}, Akira Matsuda^a,
Hideyoshi Harashima^a, and Hiroyuki Kamiya^{a,c,*}**

^aFaculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6,
Kita-ku, Sapporo 060-0812, Japan

^bGraduate School of Pharmaceutical Sciences, The University of Tokushima,
1-78-1 Sho-machi, Tokushima 770-8505, Japan

^cGraduate School of Science and Engineering, Ehime University, 2-5 Bunkyo-cho,
Matsuyama 790-8577, Japan

¹Present address: Division of Cancer Development System, National Cancer Center
Research Institute, 5-1-1 Tsukiji, Chuo-ku, Tokyo 104-0045, Japan

²Present address: Division of Genetics and Mutagenesis, National Institute of
Health Sciences, 1-18-1 Kamiyoga, Setagaya-ku, Tokyo 158-8501, Japan

*Corresponding author: Tel: +81-89-927-9609, Fax: +81-89-927-9590, E-mail:
hirokam@ehime-u.ac.jp (Hiroyuki Kamiya)

ABSTRACT

8-Oxo-7,8-dihydroguanine (8-hydroxyguanine) is oxidized more easily than normal nucleobases, which can produce spiroiminodihydantoin (Sp) and guanidinohydantoin (Gh). These secondary oxidation products of 8-oxo-7,8-dihydroguanine are highly mutagenic when formed within DNA. To evaluate the mutagenicity of the corresponding oxidation products of 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate (8-hydroxy-2'-deoxyguanosine 5'-triphosphate) in the nucleotide pool, *Escherichia coli* cells deficient in the *mutT* gene were treated with H₂O₂, and the induced mutations were analyzed. Moreover, the 2'-deoxyriboside 5'-triphosphate derivatives of Sp and Gh were also introduced into competent *E. coli* cells. The H₂O₂ treatment of *mutT E. coli* cells resulted in increases of G:C → T:A and A:T → T:A mutations. However, the incorporation of exogenous Sp and Gh 2'-deoxyribonucleotides did not significantly increase the mutation frequency. These results suggested that the oxidation product(s) of 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate induces G:C → T:A and A:T → T:A mutations, and that the 2'-deoxyriboside 5'-triphosphate derivatives of Sp and Gh exhibit quite weak mutagenicity, in contrast to the bases in DNA.

Key words: Oxidized nucleotide, 8-Oxo-7,8-dihydroguanine, 8-Hydroxyguanine, Secondary oxidation product, Spiroiminodihydantoin, Guanidinohydantoin

Abbreviations: G^O, 8-oxo-7,8-dihydroguanine; Sp, spiroiminodihydantoin; Gh, guanidinohydantoin; dG^OTP, 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate; wt, wild-type; dSpTP, spiroiminodihydantoin-2'-deoxyriboside-5'-triphosphate; dGhTP, guanidinohydantoin-2'-deoxyriboside-5'-triphosphate; pol, polymerase

1. Introduction

Reactive oxygen species, such as superoxide, hydrogen peroxide, hydroxyl radical, and singlet oxygen, are produced through normal cellular metabolism, and by various environmental mutagens and dietary factors [1-4]. Oxidatively damaged nucleobases in DNA are the focus of considerable interest, due to their implications in mutagenesis, carcinogenesis, aging, and neurodegeneration [5-7]. One of the major oxidized nucleobases is 8-oxo-7,8-dihydroguanine (G^O , also known as 8-hydroxyguanine), and 100–500 G^O residues are generated per cell per day [8-11]. The G^O base is susceptible to further oxidation, due to its lower redox potential (0.5 V lower than G) [12]. Spiroiminodihydantoin (Sp) and guanidinohydantoin (Gh) are two major secondary oxidation products of G^O [13-17]. Sp and Gh in single-stranded DNA are both >98% mutagenic and cause $G \rightarrow C$ and $G \rightarrow T$ transversions, in contrast to the ~3% mutagenicity of G^O under the same conditions [18].

Nucleobase oxidation occurs in the cellular nucleotide pool as well as in DNA, and the oxidized 2'-deoxyribonucleoside 5'-triphosphates induce mutagenic events [19]. 8-Oxo-7,8-dihydro-2'-deoxyguanosine 5'-triphosphate (dG^O TP, 8-hydroxy-2'-deoxyguanosine 5'-triphosphate) induces $A:T \rightarrow C:G$ transversion mutations in living cells [20,21]. The facts that *Escherichia coli* MutT (NudA) catalyzes the hydrolysis of dG^O TP *in vitro* [22] and that the mutation frequency in *mutT*-deficient strains is 10- to 130-fold higher than that in wild-type (wt) strains

[23-25] indicate the importance of endogenous dG^oTP as a mutagen in cells. This oxidized form of dGTP may be further oxidized to spiroiminodihydantoin-2'-deoxyriboside 5'-triphosphate (dSpTP) and guanidinohydantoin-2'-deoxyriboside 5'-triphosphate (dGhTP) under physiological conditions (Fig. 1). Thus, it is important to examine the mutagenicity of the secondary oxidation products of dG^oTP, particularly dSpTP and dGhTP, *in vivo*.

In this study, *mutT E. coli* cells, in which endogenous dG^oTP is expected to spontaneously accumulate, were treated with H₂O₂, and the induced mutations were analyzed. In addition, dSpTP and dGhTP (Fig. 1) were introduced into competent *E. coli* cells. Although the H₂O₂ treatment of *mutT E. coli* cells caused an increase in G:C → T:A and A:T → T:A transversions, the mutation frequency was not increased when dSpTP and dGhTP were incorporated into *E. coli* cells. These results suggested that the oxidation product(s) of dG^oTP induces G:C → T:A and A:T → T:A mutations. However, dSpTP and dGhTP do not seem to contribute to these induced mutations.

2. Materials and Methods

2.1. Bacterial strains

The AB1157 *E. coli* strain (*F*: *thr*-1, *leuB*6, *thi*-1, *lacY*1, *galK*2, *ara*-14, *xyl*-5, *mtl*-1, *proA*2, *his*-4, *argE*3, *rpsL*31, *tsx*-33, *supE*44, *flaND*) was obtained

from the National Institute of Genetics, Stocks Research Center (Mishima, Japan). The MK601 (AB1157 but *leu*⁺) and MK602 (AB1157 but *leu*⁺ *mutT*) strains were kindly provided by Prof. Yusaku Nakabeppu of Kyushu University [24].

2.2. Synthesis of oxidized 2'-deoxyribonucleoside 5'-triphosphates

dSpTP (the two isomers, dSpTP1 and dSpTP2) and dGhTP were synthesized by the oxidation of dGTP with Rose Bengal plus light at pH 8.0 and pH 4.4, respectively, according to the literature [26]. dG^oTP and 2-hydroxy-2'-deoxyadenosine 5'-triphosphate were synthesized as described previously [27,28].

2.3. MutT, Orf135, and Orf17 proteins

The MutT gene fragment was amplified by KOD DNA polymerase (pol) (Toyobo, Osaka, Japan) from a lysate of AB1157 *E. coli* cells, with the primers 5'-dTAGGTTTTGGATCCATGAAAAAGCTGC and 5'-dCGCCTTAGTTCGACCTACAGA (Life Technologies Japan, Tokyo, Japan), in which the underlined sequences correspond to BamH I and Sal I sites, respectively. The fragment was inserted into the pGEX-6P-3 plasmid (GE Healthcare Bio-Sciences, Piscataway, New Jersey, USA), which had been digested with the two restriction enzymes, as described previously [29]. This manipulation generated the gene encoding the GST-MutT fusion protein.

The MutT, Orf135, and Orf17 proteins without the GST-tag were purified after PreScission protease treatment, as described previously [29,30].

2.4. Mutant frequencies upon H₂O₂ treatment

A single colony (MK601 or MK602) taken from an LB agar plate was inoculated into 7 mL of LB medium. The *E. coli* culture was incubated at 37°C until the turbidity at 570 nm reached 0.2, and then 1 mL of the culture was transferred to a microtube. After H₂O₂ or H₂O (10 µL) was added, the culture was incubated at 37°C for 30 min. The culture was then centrifuged at 13,000 X *g* for 2 min at room temperature. The pellet thus obtained was resuspended in 1 mL of prewarmed LB medium, cultured at 37°C for 2 h, and then placed on ice. A portion of the suspension was diluted with ice-cold LB medium and transferred onto an LB agar plate without rifampicin (the titer plate), which was incubated at 37°C for 12 h. Another portion of the suspension was transferred onto an LB agar plate containing rifampicin (100 µg/mL) (the selection plate), which was incubated at 37°C for 20 h. The mutant frequency was calculated according to the numbers of colonies on the titer and selection plates.

2.5. Introduction of oxidized 2'-deoxyribonucleotides

A single colony (AB1157) taken from an LB agar plate was inoculated into 7 mL of LB medium. The *E. coli* culture was incubated at 37°C until the

turbidity at 570 nm reached 0.6. The culture was placed on ice for 10 min, and was then divided into 500- μ L aliquots in microtubes, which were centrifuged at 13,000 $\times g$ for 2 min at 4°C. Competent cells were prepared by a treatment with 0.1 M calcium chloride. To 50 μ L of the *E. coli* suspension, a 2'-deoxyribonucleotide solution or H₂O (2.5 μ L) was added, and then the mixture was placed on ice for 2 h. Prewarmed SOC medium (450 μ L) was then added, and cells were incubated at 37°C for 1 h. The *rpoB* mutant frequency was calculated as described above.

2.6. *MutT/Orf135/Orf17* assay

The MutT activities were assayed in a reaction mixture (50 μ L) containing 20 mM Tris-HCl, pH 7.5, 8 mM MgCl₂, 5 mM dithiothreitol, 4% glycerol, and the nucleotide substrates (20 μ M). Following a preincubation at 30°C for 2 min, the mixtures were incubated at 30°C for 10 min with the MutT protein (20 nM). The Orf135 activities were assayed in a reaction mixture (50 μ L) containing 50 mM Tris-HCl, pH 9.0, 5 mM MgCl₂, 1 mM dithiothreitol, and the nucleotide substrates (20 μ M). Following a preincubation at 37°C for 5 min, the mixtures were incubated at 37°C for 10 min with the Orf135 protein (15 nM). The Orf17 activities were assayed in a reaction mixture (50 μ L) containing 50 mM Tris-HCl, pH 9.0, 10 mM MgCl₂, 1 mM dithiothreitol, 1% glycerol, and the nucleotide substrates (20 μ M). Following a preincubation at 37°C for 5 min, the mixtures were incubated at 37°C for 15 min with the Orf17 protein (400 nM). dG^oTP, 5-methyl-dCTP, and dATP

were used as “standard” substrates for the MutT, Orf135, and Orf17 proteins, respectively. Reactions were terminated by the addition of EDTA. All samples were injected into a TSK-gel DEAE-2SW column (ϕ 4.6 mm X 250 mm, Tosoh, Tokyo, Japan), with isocratic elution by 20% CH₃CN, 75 mM sodium phosphate buffer, pH 7.0, at a flow rate of 1 ml/min. The amounts of the 2'-deoxyribonucleoside triphosphates and their hydrolyzed products were quantitated by measuring the area of UV absorbance (230 nm).

2.7. Statistical analysis

The statistical significance of the values was examined by the Student's *t*-test (mutant frequency and survival ratio) or the χ^2 test for independence (mutation spectrum). Levels of $P < 0.05$ were considered to be significant.

3. Results

3.1. Hydrogen peroxide-induced mutations in *mutT* cells

The MutT protein seems to play a crucial role in the removal of dG^oTP from the nucleotide pool, and this oxidized form of dGTP is expected to accumulate in *mutT E. coli* cells [22,24]. Indeed, the A:T→C:G mutation, the typical mutation induced by dG^oTP, is frequently observed in *mutT* cells [23-25]. Thus, the H₂O₂ treatment of *mutT* cells would produce secondary oxidation

product(s) of dG^oTP in the nucleotide pool. First, we analyzed the mutations induced by H₂O₂ in *mutT* cells, to examine whether the secondary oxidation product(s) are mutagenic. Wt (MK601) and *mutT* (MK602) cells were treated with H₂O₂, and the survival ratios and the mutant frequencies for the chromosomal *rpoB* gene were calculated (Table 1 and Fig. 2).

The survival ratios of both types of cells upon H₂O₂ treatment were similar (Table 1). The survival ratios of the wt cells were 8.8% and 1.9%, and those of the *mutT* cells were 11.9% and 4.3%, upon treatments with 1.2 and 3.5 mM H₂O₂, respectively. Interestingly, the *mutT* cells seemed to be more resistant to H₂O₂ (3 and 3.5 mM).

The treatment with 1.2 mM H₂O₂ increased the *rpoB* mutant frequency from 1.7×10^{-8} to 7.4×10^{-7} in the wt strain (Fig. 2). The mutant frequency decreased to 3.1×10^{-7} with the 3.5 mM H₂O₂ treatment. Thus, the mutants corresponding to the differences in the frequencies, 7.2×10^{-7} and 2.9×10^{-7} , were induced by 1.2 and 3.5 mM H₂O₂, respectively. Meanwhile, the background (0 mM) mutant frequency was higher in the *mutT* cells (1.3×10^{-6}), and the H₂O₂ treatment further promoted the mutant frequency to $4\text{--}6 \times 10^{-6}$. The treatment enhanced the mutation formation in the *mutT* cells relative to the wt cells, since a greater number of mutants, corresponding to the difference in the mutant frequencies ($\sim 4 \times 10^{-6}$), was induced by H₂O₂ in the *mutT* cells. These results suggested that the greater efficiency of mutation induction by H₂O₂ in the *mutT*

cells is due to increased cellular accumulation of the 2'-deoxyribonucleotide MutT substrates, dG^oTP and its diphosphate derivative [22,31], and their further oxidation products.

We then analyzed the mutation spectra in the *rpoB* gene in each strain (Tables 2 and 3). A:T → C:G transversions (85%) were observed in the *mutT* cells without the H₂O₂ treatment, as expected. The ratio of the A:T → C:G transversions decreased with the H₂O₂ treatment (statistically significant, $P < 0.01$ for 0 and 1.2 mM, $P < 0.005$ for 0 and 3.5 mM) and they constituted only 32% of the total mutations upon the treatment with 3.5 mM H₂O₂. We multiplied the *rpoB* mutant frequencies by the ratio of the A:T → C:G transversion mutations to the total colonies analyzed, and expressed them as the mutation frequencies. The *rpoB* mutant frequencies were 1.3×10^{-6} , 6.0×10^{-6} , and 5.8×10^{-6} for the 0, 1.2, and 3.5 mM H₂O₂ treatments, respectively. Thus, the A:T → C:G transversion occurred with efficiencies of 1.2×10^{-6} , 2.7×10^{-6} , and 2.1×10^{-6} for the 0, 1.2, and 3.5 mM H₂O₂ treatments, respectively. These results indicated that the H₂O₂ treatment increases the A:T → C:G transversion mutation, and that other types of mutations are also more efficiently induced.

G:C → T:A and A:T → T:A transversions represented ~40% of the total mutations in the *mutT* cells treated with 3.5 mM H₂O₂, although these mutations were undetected in the control (0 mM) experiment. The increase in G:C → T:A and A:T → T:A mutations suggested that the oxidation product(s) of dG^oTP

produced by the H_2O_2 treatment induce these types of mutations. $\text{G:C} \rightarrow \text{A:T}$ transition was also detected in the *mutT* cells treated with 1.2 mM H_2O_2 , although this mutation seemed to be decreased by 3.5 mM H_2O_2 treatment. This result suggested that secondary oxidation product(s) responsible for the transition mutation are further oxidized/degraded.

A mutational hotspot exists at position 1538 in the *mutT* cells, and the $\text{A:T} \rightarrow \text{C:G}$ transversion mutation at this position was detected for 17 of the total of 19 single-base substitutions (0 mM, Table 3). The $\text{A:T} \rightarrow \text{C:G}$ mutation at position 1538 was decreased, depending on the H_2O_2 concentration, and only three cases among the total of 18 single-base substitutions were detected for the 3.5 mM H_2O_2 treatment. These results also supported the interpretation that the intracellular $\text{dG}^\text{O}\text{TP}$ was further oxidized by the H_2O_2 treatment.

3.2. Mutagenicity of *dSpTP* and *dGhTP*

Various damaged bases have been identified as secondary oxidation products of G^O [32]. Among them, the Sp and Gh bases in DNA are highly mutagenic and induce transversion mutations in *E. coli* [18]. *dSpTP* and *dGhTP* are the oxidation products of $\text{dG}^\text{O}\text{TP}$ (Fig. 1). In contrast to $\text{dG}^\text{O}\text{TP}$, these base structure are quite different from that of G and they might be poor substrates for DNA pols. However, the Sp and Gh bases in DNA are not absolutely blocking lesions [18], suggesting the possibility of incorporation of *dSpTP* and *dGhTP* by DNA pols.

Thus, we examined their mutagenicity as the candidates that might be responsible for the mutations observed for the H₂O₂-treated *mutT* cells.

It was previously demonstrated that the direct incorporation of dG^oTP, added to a CaCl₂-treated *E. coli* suspension, induced chromosomal gene mutations [20,33]. Therefore, we examined the mutagenicity of dSpTP1, dSpTP2, and dGhTP in *E. coli* cells by the direct incorporation method. These 2'-deoxyribonucleotides were synthesized by the oxidation of dGTP, according to the published method [26]. Note that the two stereo-isomers of dSpTP, dSpTP1 and dSpTP2, were separately purified, as described above [26], although their configurations have not been determined. dGhTP was purified as the mixture of two isomers in equilibrium. dSpTP and dGhTP were introduced into wt *E. coli* cells, and the mutant frequencies for the chromosomal *rpoB* gene were calculated. The numbers of colonies on the titer plates were very similar (data not shown), indicating the minimal, if any, cytotoxicity of dSpTP and dGhTP.

Unexpectedly, the mutagenicities of dSpTP1 and dGhTP were low, in contrast to the high mutagenic potentials of Sp and Gh in DNA (Fig. 3) [18]. No mutagenicity by dSpTP2 was detected (data not shown). In contrast, the *rpoB* mutant frequencies upon dG^oTP and 2-hydroxy-2'-deoxyadenosine 5'-triphosphate treatments (12.5 nmol) were 2.4×10^{-7} and 3.1×10^{-7} , respectively. Thus, the mutagenicities of dSpTP and dGhTP were at least one order of magnitude lower

than that of dG^oTP in *E. coli*. Moreover, the possibility that they contributed to the H₂O₂-induced mutations observed in *mutT* cells is excluded.

3.3. *dSpTP and dGhTP are not hydrolyzed by bacterial MutT-type enzymes*

The low mutagenic potentials of dSpTP and dGhTP might be due to their efficient degradation by *E. coli* MutT-type protein(s). We finally examined the possibility that dSpTP and dGhTP are good substrates for the *E. coli* MutT-type proteins, MutT, Orf135 (NudG), and Orf17 (NudB, NtpA). dSpTP1, dSpTP2, and dGhTP were incubated with recombinant MutT, and the reaction mixtures were analyzed by anion-exchange HPLC [34]. The MutT protein produced a significant amount of the monophosphate derivative of dG^oTP, a known substrate of MutT (Fig. 4A). In contrast, little, if any, degradation of dSpTP1, dSpTP2, and dGhTP was observed (Fig. 4C and E, and data not shown). Thus, they are unlikely to be degraded with high efficiencies by the MutT protein in *E. coli* cells. Meanwhile, the 2'-deoxyribonucleotides of Sp and Gh might be degraded by the other MutT-type nucleotide pool sanitization enzymes, Orf135 and Orf17, in *E. coli* cells. To examine this possibility, dSpTP1, dSpTP2, and dGhTP were incubated with the Orf135 and Orf17 proteins *in vitro*. However, their hydrolysis was not observed under our experimental conditions (data not shown). Therefore, the weak mutagenicity of dSpTP1, dSpTP2, and dGhTP could not be explained by their efficient hydrolysis by the three MutT-type proteins.

4. Discussion

The objective of this study was to examine the mutagenicity of secondary oxidation products of dG^oTP. We treated *mutT E. coli* cells with H₂O₂ to oxidize intracellular dG^oTP, and examined the mutation spectrum. As shown in Table 2, the treatment caused an increase in G:C → T:A and A:T → T:A transversions. These results suggested that the oxidation product(s) of dG^oTP cause G:C → T:A and A:T → T:A mutations, since the degree of the increase in mutant frequency was much higher in the *mutT* strain than the wt strain (Fig. 2). G:C → T:A mutations could be generated by the incorporation of damaged dGTP opposite C in the template DNA, and dATP insertion opposite the damaged G base in DNA during the next round of replication. In addition, the incorporation of damaged dGTP opposite G in the template DNA, and dTTP insertion opposite the damaged G base in DNA, could induce the same type of mutation. A:T → T:A mutations could be generated by the incorporation of damaged dGTP opposite T in the template DNA and dATP insertion opposite the damaged G base in DNA, and by the incorporation of damaged dGTP opposite A in the template DNA, followed by dTTP insertion opposite the damaged G base in DNA.

Alternatively, dG^oTP could be incorporated opposite C and A, and further oxidation of G^o in DNA might induce the mutations we observed. If this were the

case, then one would expect $G^O:C$ to be converted to $X:C$ (X represents oxidized G^O), and subsequent dATP incorporation opposite X would result in $G:C \rightarrow T:A$ mutations. Likewise, if $G^O:A$ were converted to $X:A$, then subsequent dTTP incorporation opposite X would induce $A:T \rightarrow T:A$ mutations. Studies using plasmid DNA modified with G^O at a predetermined site might be required to examine these possibilities.

Previously, we examined the effects of the UvrA- and UvrB-deficiencies on the mutations induced in *mutT* cells treated with H_2O_2 [35]. The deficiency in the *uvrA* or *uvrB* gene resulted in fewer mutations caused by H_2O_2 -treatment of the *mutT* cells. We assumed that the formation of d G^O TP by the oxidation of dGTP would increase under the oxidative stress produced by the exposure to H_2O_2 in *mutT* cells. However, based on the results shown in Tables 2 and 3, the increased amount of d G^O TP does not explain the entire increase in the mutant frequency induced by the H_2O_2 treatment. The reduced mutant frequencies in the *uvrA/mutT* and *uvrB/mutT* cells, at least in part, could be due to the involvement of UvrA and UvrB in the metabolism of the putative secondary oxidation product(s) of d G^O TP.

The secondary oxidation products, Sp and Gh, are highly mutagenic when generated in DNA [18]. Conversely, no mutagenicities of dSpTP and dGhTP were observed (Fig. 3). Based on the assumption that they are eliminated from the nucleotide pool by MutT-type enzymes, we incubated these damaged 2'-deoxyribonucleotides with MutT, Orf135, and Orf17 *in vitro*. However, no

hydrolysis products were detected under our experimental conditions (Fig. 4 and data not shown). Thus, the absence of mutagenicity by dSpTP and dGhTP might be attributable to their inability to be incorporated by the *E. coli* DNA pol(s). Alternatively, they might be incorporated in a highly “error-free” manner by DNA pol(s). For example, no mutation is induced if dGhTP is incorporated opposite G (the first round of replication) and dGTP is inserted opposite Gh (the next round of replication). Experiments using various bacterial DNA pols, including the specialized DNA pols (DNA pols II, IV, and V), would be required to understand the discrepancy.

As shown in Table 1, the survival ratios of the H₂O₂ treated wt and *mutT* cells were similar. However, the cell death ratios of the *mutT* cells were less than those of the wt cells, in the cases of the 3 and 3.5 mM H₂O₂ treatments. One possible explanation is that the intracellular H₂O₂ concentration was lower in the *mutT* cells than in the wt cells, for an unknown reason(s) triggered by the MutT-deficiency. In contrast, the resistance to H₂O₂ is difficult to explain by the amount of cytotoxic, oxidized 2'-deoxyribonucleotide(s) and/or DNA lesion(s) in the *mutT* cells. However, it might be due to a putative stress-induced response that confers resistance to reactive oxygen species.

In this study, H₂O₂-induced mutations in *mutT* cells were analyzed, and the lack of mutagenicity by dSpTP and dGhTP was demonstrated. Our results suggested that the secondary oxidation product(s) of dG^oTP, except for dSpTP and

dGhTP, are mutagenic. The other secondary oxidation products should be examined to reveal the mechanism(s) of the promoted mutations observed in the H₂O₂-treated *mutT* cells.

Conflict of interest statement

The authors declare that there are no conflicts of interest.

Acknowledgements

We thank Profs. Naoki Kamo and Seiji Miyauchi for their assistance in the synthesis of dSpTP and dGhTP. We thank Prof. Yusaku Nakabeppu for providing *E. coli* strains. This work was supported in part by Grants-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology of Japan, the Takeda Science Foundation, and by fellowships and grants from Research Fellowships of the Japan Society for the Promotion of Science to M. H.

References

- [1] B.N. Ames, Dietary carcinogens and anticarcinogens, *Science* 221 (1983) 1256-1264.
- [2] B.N. Ames, L.S. Gold. Endogenous mutagens and the causes of aging and cancer, *Mutat. Res.* 250 (1991) 3-16.
- [3] E.S. Henle, S. Linn, Formation, prevention, and repair of DNA damage by iron/hydrogen peroxide, *J. Biol. Chem.* 272 (1997) 19095-19098.
- [4] Y. Hiraku, M. Murata, S. Kawanishi, Role of oxidative DNA damage in dietary carcinogenesis, *Genes Environ.* 28 (2006) 127-140.
- [5] H. Kamiya, Mutagenic potentials of damaged nucleic acids produced by reactive oxygen/nitrogen species: Approaches using synthetic oligonucleotides and nucleotides, *Nucleic Acids Res.* 31 (2003) 517-531.
- [6] K.B. Beckman, B.N. Ames, Endogenous oxidative damage of mtDNA, *Mutat. Res.* 424 (1999) 51-58.
- [7] D. Wang, D.A. Kreutzer, J.M. Essigmann, Mutagenicity and repair of oxidative DNA damage: insights from studies using defined lesions, *Mutat. Res.* 400 (1998) 99-115.
- [8] H. Kasai, S. Nishimura, Hydroxylation of deoxyguanosine at the C-8 position by ascorbic acid and other reducing agents, *Nucleic Acids Res.* 12 (1984) 2137-2145.

- [9] T. Lindahl, Instability and decay of the primary structure of DNA, *Nature* 362 (1993) 709-715.
- [10] H. Kasai, Analysis of a form of oxidative DNA damage, 8-hydroxy-2'-deoxyguanosine, as a marker of cellular oxidative stress during carcinogenesis, *Mutat. Res.* 387 (1997) 147-163.
- [11] H. Kamiya, Mutagenicities of 8-hydroxyguanine and 2-hydroxyadenine produced by reactive oxygen species, *Biol. Pharm. Bull.* 27 (2004) 475-479.
- [12] S. Steenken, S.V. Jovanovic, M. Bietti, K. Bernhard, The trap depth (in DNA) of 8-oxo-7,8-dihydro-2'-deoxyguanosine as derived from electron-transfer equilibria in aqueous solution, *J. Am. Chem. Soc.* 122 (2000) 2373-2374.
- [13] W. Luo, J.G. Muller, E.M. Rachlin, C.J. Burrows, Characterization of spiroiminodihydantoin as a product of one-electron oxidation of 8-Oxo-7,8-dihydroguanosine, *Org. Lett.* 2 (2000) 613-616.
- [14] W. Luo, J.G. Muller, E.M. Rachlin, C.J. Burrows, Characterization of hydantoin products from one-electron oxidation of 8-oxo-7,8-dihydroguanosine in a nucleoside model, *Chem. Res. Toxicol.* 14 (2001) 927-938.
- [15] T. Suzuki, T. Inoue, M. Inukai, Formation of spiroiminodihydantoin nucleosides from 8-oxo-7,8-dihydro-2'-deoxyguanosine by ultraviolet light, *Genes Environ.* 32 (2010) 31-36.

- [16] T. Suzuki, Formation of spiroiminodihydantoin nucleoside from 8-oxo-7,8-dihydro-2'-deoxyguanosine by nitric oxide under aerobic conditions, *Bioorg. Med. Chem. Lett.* 19 (2009) 4944-4947.
- [17] K. Kino, M. Morikawa, T. Kobayashi, T. Kobayashi, R. Komori, Y. Sei, H. Miyazawa, The oxidation of 8-oxo-7,8-dihydroguanine by iodine, *Bioorg. Med. Chem. Lett.* 20 (2010) 3818-3820.
- [18] P.T. Henderson, J.C. Delaney, J.G. Muller, W.L. Neeley, S.R. Tannenbaum, C.J. Burrows, J.M. Essigmann, The hydantoin lesions formed from oxidation of 7,8-dihydro-8-oxoguanine are potent sources of replication errors in vivo, *Biochemistry* 42 (2003) 9257-9262.
- [19] H. Kamiya, Mutations induced by oxidized DNA precursors and their prevention by nucleotide pool sanitization enzymes, *Genes Environ.* 29 (2007) 133-140.
- [20] M. Inoue, H. Kamiya, K. Fujikawa, Y. Ootsuyama, N. Murata-Kamiya, T. Osaki, K. Yasumoto, H. Kasai, Induction of chromosomal gene mutations in *Escherichia coli* by direct incorporation of oxidatively damaged nucleotides, *J. Biol. Chem.* 273 (1998) 11069-11074.
- [21] K. Satou, K. Kawai, H. Kasai, H. Harashima, H. Kamiya, Mutagenic effects of 8-hydroxy-dGTP in live mammalian cells, *Free Radic. Biol. Med.* 42 (2007) 1552-1560.

- [22] H. Maki, M. Sekiguchi, MutT protein specifically hydrolyses a potent mutagenic substrate for DNA synthesis, *Nature* 355 (1992) 273-275.
- [23] C. Yanofsky, E.C. Cox, V. Horn, The unusual mutagenic specificity of an *E. coli* mutator gene, *Proc. Natl. Acad. Sci. USA.* 55 (1966) 274–281.
- [24] T. Tajiri, H. Maki, M. Sekiguchi, Functional cooperation of MutT, MutM and MutY proteins in preventing mutations caused by spontaneous oxidation of guanine nucleotide in *Escherichia coli*, *Mutat. Res.* 336 (1995) 257-267.
- [25] H. Kamiya, C. Ishiguro, H. Harashima, Increased A:T→C:G mutations in the *mutT* strain upon 8-hydroxy-dGTP treatment: Direct evidence for MutT in the prevention of mutations by oxidized dGTP, *J. Biochem.* 136 (2004) 1359-1362.
- [26] Y. Ye, J.G. Muller, C.J. Burrows, Synthesis and characterization of the oxidized dGTP lesions spiroiminodihydantoin-2'-deoxynucleoside-5'-triphosphate and guanidinohydantoin-2'-deoxynucleoside-5'-triphosphate, *J. Org. Chem.* 71 (2006) 2181-2184.
- [27] K.C. Cheng, D.S. Cahill, H. Kasai, S. Nishimura, L.A. Loeb, 8-Hydroxyguanine, an abundant form of oxidative DNA damage, causes G→T and A→C substitutions, *J. Biol. Chem.* 267 (1992) 166-172.
- [28] H. Kamiya, H. Kasai, Formation of 2-Hydroxydeoxyadenosine triphosphate, an oxidatively damaged nucleotide, and its incorporation by DNA polymerases, *J. Biol. Chem.* 270 (1995) 19446-19450.

- [29] H. Kamiya, N. Murata-Kamiya, E. Iida, H. Harashima, Hydrolysis of oxidized Nucleotides by the *Escherichia coli* Orf135 protein, *Biochem. Biophys. Res. Commun.* 288 (2001) 499-502.
- [30] M. Hori, K. Fujikawa, H. Kasai, H. Harashima, H. Kamiya, Dual hydrolysis of diphosphate and triphosphate derivatives of oxidized deoxyadenosine by Orf17 (NtpA), a MutT-type enzyme, *DNA Repair (Amst.)* 4 (2005) 33-39.
- [31] R. Ito, H. Hayakawa, M. Sekiguchi, T. Ishibashi, Multiple enzyme activities of *Escherichia coli* MutT protein for sanitization of DNA and RNA precursor pools, *Biochemistry* 44 (2005) 6670-6674.
- [32] W.L. Neeley, J.M. Essigmann, Mechanisms of formation, genotoxicity, and mutation of guanine oxidation products, *Chem. Res. Toxicol.* 19 (2006) 491-505.
- [33] H. Kamiya, E. Iida, N. Murata-Kamiya, Y. Yamamoto, T. Miki, H. Harashima, Suppression of spontaneous and hydrogen peroxide-induced mutations by a MutT-type nucleotide pool sanitization enzyme, the *Escherichia coli* Orf135 protein, *Genes Cells* 8 (2003) 941-950.
- [34] K. Fujikawa, H. Kamiya, H. Yakushiji, Y. Fujii, Y. Nakabeppu, H. Kasai, The oxidized forms of dATP are substrates for the human MutT homologue, the hMTH1 protein, *J. Biol. Chem.* 274 (1999) 18201-18205.
- [35] M. Hori, C. Ishiguro, T. Suzuki, N. Nakagawa, T. Nunoshiba, S. Kuramitsu, K. Yamamoto, H. Kasai, H. Harashima, H. Kamiya, UvrA and UvrB enhance

mutations induced by oxidized deoxyribonucleotides, DNA Repair (Amst.) 6
(2007) 1786-1793.

Figure Legends

Fig. 1. Structures of dSpTP and dGhTP. P represents a phosphate group.

Fig. 2. Frequency of *rpoB* mutants upon H₂O₂ treatment. A single colony taken from an agar plate was inoculated into LB medium and incubated at 37 °C. The *E. coli* culture was then treated with water or H₂O₂, and was incubated at 37 °C for 30 min. After centrifugation, the pellet was resuspended in LB medium and was incubated at 37 °C for 2 hr. The *rpoB* mutant frequency was calculated according to the numbers of colonies on the titer and selection plates, as described in the Materials and Methods section. Open diamonds, the wt strain; closed squares, the *mutT* strain. Experiments were repeated at least three times, except for wt, 1.6 mM (n=1), and data are expressed as means $\pm$ standard deviation. Asterisks (**) indicate a significant difference versus the wt strain, with $P < 0.01$.

Fig. 3. Mutant frequency in wt *E. coli* cells treated with damaged 2'-deoxyribonucleotides. After adding the 2'-deoxyribonucleotides to competent *E. coli* suspensions, the *rpoB* mutant frequency was calculated according to the numbers of colonies on the titer and selection plates, as described in the Materials and Methods section. The 2'-deoxyribonucleotides that were added are shown below each bar. Experiments were repeated at least six times, and the data are

expressed as means $\pm$ standard deviation. The *rpoB* mutant frequencies were 2.4×10^{-7} and 3.1×10^{-7} when *E. coli* cells were treated with dG^oTP and 2-hydroxy-2'-deoxyadenosine 5'-triphosphate treatments (12.5 nmol), respectively.

Fig. 4 HPLC analysis of reaction mixtures of oxidized 2'-deoxyribonucleotides and MutT. Each nucleotide (20 μ M) was incubated with 20 nM of the MutT protein at 30°C for 10 min. (A) dG^oTP, postincubation, (B) dSpTP1, preincubation, (C) dSpTP1, postincubation, (D) dGhTP, preincubation, and (E) dGhTP, postincubation. The reaction mixtures were analyzed by anion-exchange HPLC under the conditions described in section 2.6. dG^oMP in panel A represents 8-oxo-7,8-dihydro-2'-deoxyguanosine 5'-monophosphate.

dSpTP

dGhTP

Fig. 1

Fig. 2

Fig. 3

A

Fig. 4

B

Fig. 4

C

Fig. 4

D

Fig. 4

E

Fig. 4

Table 1. H₂O₂-induced death in wt and *mutT* cells

[H₂O₂] (mM)	wt	<i>mutT</i>
0.0	100.0	100.0
0.4	73.9 (27.0)	74.5 (8.8)
0.8	16.4 (14.7)	24.9 (13.7)
1.2	8.8 (4.5)	11.9 (6.5)
1.6	4.4	6.5 (3.1)
2.0	5.8 (3.3)	6.3 (3.9)
3.0	2.8 (2.1)	6.9 (5.2)*
3.5	1.9 (0.7)	4.3 (2.2)**

Relative *E. coli* colonies on the titer plates to that of 0.0 mM are shown as percentage. Experiments were repeated at least three times except for wt, 1.6 mM, and the data are expressed as the mean. Numbers in parentheses represent standard deviations. Asterisks indicate a significant difference versus the wt strain, with *P*<0.05 (*) or *P*<0.01 (**).

Table 2. Mutations induced by H₂O₂ in wt and *mutT* strains

	wt			<i>mutT</i>		
	0 mM	1.2 mM	3.5 mM	0 mM	1.2 mM	3.5 mM
Single base substitution						
Transition						
A:T to G:C	7 (32)	2 (10)	0 (0)	1 (5)	0 (0)	0 (0)
G:C to A:T	9 (41)	9 (43)	11 (48)	0 (0)	5 (25)	2 (9)
Transversion						
A:T to T:A	0 (0)	0 (0)	2 (9)	0 (0)	0 (0)	3 (14)
A:T to C:G	2 (9)	0 (0)	1 (4)	17 (85)	9 (45)	7 (32)
G:C to T:A	1 (5)	6 (29)	4 (17)	0 (0)	4 (20)	5 (23)
G:C to C:G	0 (0)	4 (19)	3 (13)	1 (5)	0 (0)	1 (5)
Insertion/deletion	0 (0)	0 (0)	0 (0)	0 (0)	1 (5)	2 (9)
Others	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
No mutation	3 (14)	0 (0)	2 (9)	1 (5)	1 (5)	2 (9)
Total mutation	22 (100)	21 (100)	23 (100)	20 (100)	20 (100)	22 (100)
Total colonies analyzed	22	21	23	19	20	19

A portion of the *rpoB* gene (nucleotides 1519-1725) was analyzed. All data are represented as cases found (%).

Table 3. The distribution of single-base substitutions induced by H₂O₂ in wt and mutT strains.

wt			mutT		
0 mM	1.2 mM	3.5 mM	0 mM	1.2 mM	3.5 mM
1534 T to C (1)	1535 C to A (1)	1535 C to T (1)	1536 T to C (1)	1537 C to A (2)	1535 C to A (1)
1535 C to T (1)	1546 G to A (2)	1538 A to C (1)	1538 A to C (17) ^a	1538 A to C (6)	1537 C to A (2)
1546 G to A (1)	1546 G to T (1)	1546 G to A (2)	1559 C to G (1) ^a	1546 G to A (1)	1538 A to T (1)
1546 G to T (1)	1547 A to G (2)	1546 G to T (2)		1546 G to T (1)	1538 A to C (3)
1547 A to G (5)	1565 C to T (1)	1547 A to T (1)		1576 C to T (3)	1545 G to A (1)
1552 A to G (1)	1576 C to T (4)	1576 C to T (5)		1576 C to A (1)	1571 T to A (2) ^b
1576 C to T (5)	1576 C to A (4)	1576 C to A (2)		1691 C to T (1)	1576 C to A (1) ^b
1592 C to T (2)	1576 C to G (3)	1576 C to G (3)		1714 A to C (1)	1576 C to G (1)
1715 T to G (2)	1592 C to T (1)	1691 C to T (2)		1715 T to G (2)	1592 C to T (1)
	1691 C to T (1)	1714 A to T (1)			1592 C to A (1)
	1716 C to G (1)	1721 C to T (1)			1714 A to C (3)
					1715 T to G (1)

A portion of the *rpoB* gene (nucleotides 1519-1725) was analyzed. The sequence of the sense strand is shown. The numbers of colonies are shown on the right side in parentheses.

^aOne colony contained two base substitutions.

^bOne colony contained two base substitutions.