



Title	Mechanism for propofol inhibition of Na <sup>+</sup> , K <sup>+</sup> -ATPase activity in rat brain
Author(s)	Hase, Yuri; Deyama, Yoshiaki; Yoshimura, Yoshitaka; Suzuki, Kuniaki; Fukushima, Kazuaki
Citation	北海道歯学雑誌, 32(2), 147-155
Issue Date	2012-03
Doc URL	<a href="http://hdl.handle.net/2115/48709">http://hdl.handle.net/2115/48709</a>
Type	article
File Information	05-hase_gencho.pdf



[Instructions for use](#)

## ORIGINAL

Mechanism for propofol inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in rat brain

Yuri Hase<sup>1</sup>, Yoshiaki Deyama<sup>2</sup>, Yoshitaka Yoshimura<sup>2</sup>,  
Kuniaki Suzuki<sup>2</sup> and Kazuaki Fukushima<sup>1</sup>

**ABSTRACT** : Propofol is one of the most widely used intravenous anesthetics, however the mechanism of the anesthetic effect is not fully understood.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase is an enzyme present in all animal cell membranes and plays essential roles for the maintenance of neuronal excitability. There is a report of propofol inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity, but the mechanism is not clearly established. To study the mechanism for propofol inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase purified from whole brains of rats, the effects of propofol on  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity,  $\text{Na}^+$ -ATPase, and  $\text{K}^+$ - $\beta$ NPPase activities, which are partial reactions of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase were examined.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase and  $\text{Na}^+$ -ATPase activities decreased depending on the concentration of propofol, and were completely inhibited at 1.03 mM. Propofol decreased the maximum activity of  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ , and ATP-dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity depending on its concentration, and changed the half maximal concentration for  $\text{Na}^+$ ,  $\text{K}^+$ , and ATP, but not for  $\text{Mg}^{2+}$ . Propofol also decreased the maximum activities of  $\text{Na}^+$ -ATPase and  $\text{K}^+$ - $\beta$ NPPase, suggesting that propofol inhibits  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity by affecting the whole reaction process of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase. The inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity by propofol was reversible by dilution of its concentration. These results suggest that propofol reversibly inhibits  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in a mixed-type inhibition pattern.

**Key Words** : Intravenous anesthetic, propofol,  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase

## Introduction

Propofol is a short-acting intravenous anesthetic with mainly a sedative effect, widely used for the induction and maintenance of general anesthesia, as well as for sedation of respiratory management in intensive care units and for intravenous sedation in dental therapy. The exact mechanism of general anesthetics including propofol has not been revealed yet, though various studies have been reported [1-7]. Propofol has been reported to have many pharmacological effects [8] : 1) It reduces cerebral blood flow, cerebral metabolic rate, and intracranial pressure. 2) It acts as an antioxidant. 3) It reduces ischemic neuronal injury in animal models of transient global or focal cerebral ischemia. 4) It activates  $\gamma$ -aminobutyric acid ( $\text{GABA}_A$ ) receptors directly [8-12].

$\text{Na}^+$ ,  $\text{K}^+$ -ATPase is an enzyme present in all animal

cell membranes, which translocates sodium and potassium ions across the cell membrane, utilizing the chemical energy of hydrolysis of ATP.  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase plays important physiological roles [13], and the activity of this enzyme is very sensitive to the influence of various bioregulators, such as cardiac steroids, transition and heavy metals, as well as metal complexes [14, 15]. We hypothesize that the inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity by propofol may be related to the state of anesthesia or side effects caused by propofol.

Kutchai *et al.* [16] reported that 73-800  $\mu\text{M}$  propofol inhibited  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in canine renal medulla, and its  $\text{IC}_{50}$  was  $127 \pm 13 \mu\text{M}$ , however, details still remain to be elucidated. The aim of this study was to investigate the inhibition mechanism of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity by propofol. We studied the propofol inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity and the effects of propofol on

<sup>1</sup>Department of Dental Anesthesiology, Hokkaido University Graduate School of Dental Medicine, Sapporo, Hokkaido 060-8586, Japan

<sup>2</sup>Department of Molecular Cell Pharmacology, Hokkaido University Graduate School of Dental Medicine, Sapporo, Hokkaido 060-8586, Japan

$\text{Na}^+$ -,  $\text{K}^+$ -,  $\text{Mg}^{2+}$ -, and ATP- dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity. We also studied the effects of propofol on partial reactions of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase, and the reversibility of the inhibition by dilution of propofol concentration. We show that propofol reversibly inhibits  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity in a mixed-type inhibition pattern.

## Materials and Methods

### 1. Enzyme preparation

The animals and tissue specimens were treated in accordance with the Guidelines of the Experimental Animal Committee, Hokkaido University Graduate School of Dental Medicine. First, microsomes were prepared from the whole brains of rats and the purification of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase from microsome was accomplished according to Jorgensen's method [17] with some modifications [18]. The microsome was treated with 0.55 mg of sodium dodecyl sulfate (SDS) per milligram of microsomal protein, and centrifuged on a glycerol density gradient. After centrifugation, we recovered two layers (white cloud and medium) and pellet (button). The protein concentration was estimated by using the Bio-Rad Protein Assay (Bio-Rad Laboratories, CA) according to the manufacture's protocol with bovine serum albumin as a standard. The specific activities of white cloud, medium, and button were 3.11, 4.63, and 0.8  $\mu\text{mol}/\text{min}/\text{mg}$  protein.

### 2. $\text{Na}^+$ , $\text{K}^+$ -ATPase assay

$\text{Na}^+$ ,  $\text{K}^+$ -ATPase and  $\text{Na}^+$ -ATPase activities were determined by the measurement of inorganic phosphate production according to Chifflet's method [19].  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity was assayed in a total volume 300  $\mu\text{l}$  of reaction mixture containing the enzyme (1  $\mu\text{g}$  white cloud), 25 mM sucrose, 0.1 mM ethylenediaminetetraacetic acid (EDTA), 50 mM tris-HCl at pH 7.41, 160 mM NaCl, 16 mM KCl and 5 mM  $\text{MgCl}_2$ . After pre-incubation, the reaction was started by the addition of 50  $\mu\text{l}$  of 30 mM ATP, allowed to proceed for 30 minutes at 37°C, and then the reaction was stopped by the addition of 12% SDS. In brief, 0.6 ml of the solution containing 3% ascorbic acid, 0.5 N HCl and 0.5% ammonium molybdate was added to the 0.6 ml reaction mixture with SDS, which was left for 3-10 minutes at room temperature. Then, 0.9 ml of a solution containing 2% sodium citrate, 2% sodium metaarsenite and 2% acetic acid was added to the mixture, which was then incubated for 10 minutes at 37°C. The developed color was read at 850 nm spectrophotometrically

with a Hitachi U-2000 spectrophotometer. The results are expressed as the mean percentage of enzyme activity relative to the corresponding control value, the data derived from at least 3 experiments, each experiment using 3 samples of propofol.

#### 1) Concentration-dependent inhibition of $\text{Na}^+$ , $\text{K}^+$ -ATPase activity by propofol

Various concentrations of propofol were added to the above reaction mixture and the effects were observed by assaying  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

#### 2) The effect of propofol on the affinities of $\text{Na}^+$ , $\text{K}^+$ -ATPase for $\text{Na}^+$ , $\text{K}^+$ , $\text{Mg}^{2+}$ , or ATP.

$\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity was measured in the reaction mixture containing the enzyme with different concentrations of  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$ , or ATP. The effects of 0.14, 0.26, or 0.35 mM propofol and dimethyl sulfoxide (DMSO) used as a solvent were observed on the  $\text{Na}^+$ -,  $\text{K}^+$ -,  $\text{Mg}^{2+}$ - or ATP-concentration dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

#### 3. The effect of propofol on the affinity of $\text{Na}^+$ -ATPase activity for $\text{Na}^+$ .

To reveal at which stage of the reaction mechanism propofol had an effect on  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase,  $\text{Na}^+$ -ATPase activity, the anterior half partial reaction of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase, was measured.  $\text{Na}^+$ -ATPase activity was measured in the reaction mixture containing the enzyme (5.25  $\mu\text{g}$  medium), 25 mM sucrose, 0.1 mM EDTA, 50 mM tris-HCl at pH 7.41 and 5 mM  $\text{MgCl}_2$ , changing the concentration of  $\text{Na}^+$  from 0-20 mM with and without propofol and DMSO. The reaction time was decided to be 1 hour because of weak ATP hydrolysis of  $\text{Na}^+$ -ATPase.  $\text{Na}^+$ -ATPase activity was also determined by measurement of inorganic phosphate.

#### 4. The effect of propofol on the affinity of $\text{K}^+$ - $\rho\text{NPPase}$ activity for $\text{K}^+$ .

$\text{K}^+$ - $\rho\text{NPPase}$  activity, posterior half partial reaction of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase, was measured in the reaction mixture containing the enzyme (15  $\mu\text{g}$  medium), 25 mM sucrose, 0.1 mM EDTA, 50 mM tris-HCl at pH 7.41 and 5 mM  $\text{MgCl}_2$ , changing the concentration of  $\text{K}^+$  from 0-20 mM with and without propofol and DMSO. The reaction was started by the addition of 200  $\mu\text{l}$  of 16 mM  $\rho\text{NPP}$ , allowed to proceed for 30 minutes at 37°C, and then it was stopped by adding 2 ml of 2% SDS and 1.25%  $\text{Na}_2\text{CO}_3$ . The

developed color was read at 420 nm spectrophotometrically.

### 5. Reversibility of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity inhibited by propofol.

Whether the inhibition by propofol was reversible or not was examined as below. At first, the reaction mixture containing the enzyme (1 μg white cloud), 25 mM sucrose, 0.1 mM EDTA, 50 mM tris-HCl at pH 7.41, 160 mM NaCl, 16 mM KCl and 5 mM MgCl<sub>2</sub> was pre-incubated with 1.03 mM propofol and 10% DMSO at room temperature or on ice for 30 minutes, and then the propofol and DMSO concentration was reduced by dilution to the concentrations described in Fig. 9. Then Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was measured as described above.

### 6. Drugs and chemicals

All the drugs and chemicals used in this study were obtained from Wako Pure (Osaka, Japan). 2, 6-Diisopropylphenol, which is the active ingredient of propofol, has high lipid solubility and little water solubility. Therefore propofol is clinically used as a propofol injection with additives like soybean oil and triglyceride to make an emulsion. However, it is impossible to use an emulsion in this study as turbidity disturbs the measurement of absorbance. We dissolved 2, 6-diisopropylphenol into DMSO as a 0.2% solution and then this solution was diluted with water.

### 7. Statistics

The data are expressed using the means of at least 3 independent experiments. Statistical assessment of the data was examined by a Student's *t*-test. Differences were considered to be statistically significant when  $P < 0.05$  and  $P < 0.01$ .

## Results

### 1. Concentration-dependent inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity by propofol

Specific activity of Na<sup>+</sup>, K<sup>+</sup>-ATPase used for ATPase assay was 3.11–4.63 μmol/mg protein/min. Fig. 1 shows the inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity in the presence of various concentrations of propofol. The activity decreased depending on the concentration of propofol, and was inhibited completely at 1.03 mM. The concentration that caused half-maximal inhibition (IC<sub>50</sub>) of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was 0.26 mM and the Hill coefficient *n* was 1.11, analyzed by the Hill equation.




Fig. 1 Concentration-dependent inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity by propofol.

Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was measured in the presence of various concentrations of propofol. The activity decreased depending on the concentration of propofol, and was inhibited completely at 1.03 mM. The concentration that caused half-maximal inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was 0.26 mM. Data have a mean ± SEM ( $n \geq 3$ ).

### 2. The effect of propofol on the ATP concentration-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

Fig. 2a shows the effect of propofol on the ATP concentration dependency of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity. In the absence of DMSO and propofol (◇), the activity increased, depending on the ATP concentration. A double-reciprocal plot of ATP concentration versus ATPase activity resulted in two straight lines (Fig. 2b). By extrapolation of each line to x and y axes, high (0.12 mM) and low (0.48 mM) Km values and corresponding maximum activities (Vmax) were calculated (Fig. 2b and Table 1). Similar experiments were done in the presence of 0.14 (△), 0.26 (○) and 0.35 mM (×) propofol or DMSO (□) as controls for propofol solvent (Fig. 2a). The results were analyzed as shown in Fig. 2b, and Km and Vmax values for high and low affinity sites were summarized in Table 1. Propofol decreased Vmax values for both sites depending on its concentration. However, it increased affinities for ATP for both high and low ATP affinity sites (Table 1).

### 3. The effect of propofol on the Na<sup>+</sup> concentration-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

Fig. 3 shows the effect of propofol on the Na<sup>+</sup> concentration dependency of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity. In the absence of DMSO and propofol (◇), the activity increased, depending on the Na<sup>+</sup> concentration. The results were analyzed by the Hill equation, and the Na<sup>+</sup> concentration that caused half-maximal activation ([S]<sub>0.5</sub>), maximum activity (Vmax) of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity,




Fig. 2a The effect of propofol on the ATP concentration-dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

ATP concentration dependency of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity was measured in the absence of DMSO and propofol ( $\diamond$ ), or in the presence of 2.5% DMSO without propofol ( $\square$ ); 1.3% DMSO and 0.14 mM propofol ( $\triangle$ ); 2.5% DMSO and 0.26 mM propofol ( $\circ$ ); 3.3% DMSO and 0.35 mM propofol ( $\times$ ). The results were analyzed as shown in Fig. 2b, and  $K_m$  and  $V_{max}$  values for high and low affinity sites are summarized in Table 1. Data have a mean  $\pm$  SEM ( $n \geq 3$ ).

\* $P < 0.05$  and \*\* $P < 0.01$  compared with two groups of each by a Student's  $t$ -test. ( $\diamond$ ) vs ( $\square$ );  $P < 0.01$ , ( $\triangle$ ) vs ( $\circ$ );  $P < 0.05$ , ( $\circ$ ) vs ( $\times$ );  $P < 0.01$ .




Fig. 2b Double-reciprocal plot of ATP concentration versus ATPase activity.

The results obtained in the experiments of Fig. 2 a were analyzed by a double-reciprocal plot. The double-reciprocal plot of ATP concentration versus  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity resulted in two straight lines. By extrapolation of each line to  $x$  and  $y$  axes, high and low  $K_m$  values and corresponding maximum activities ( $V_{max}$ ) were calculated. The concentration of ATP was shown as molar concentration multiplied by  $10^4$  M. Reaction velocity was expressed as  $\mu\text{mol Pi released/mg protein per min}$ . The  $K_m$  and  $V_{max}$  values for high and low affinity sites calculated from each experiment are summarized in Table 1.

Table 1 Effects of propofol on ATP-dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

DMSO (%)	Propofol (mM)	High affinity site		Low affinity site	
		$K_m$ (mM)	$V_{max}$ ( $\mu\text{mol/mg/min}$ )	$K_m$ (mM)	$V_{max}$ ( $\mu\text{mol/mg/min}$ )
0	0	0.12	1.82	0.48	4.45
2.5	0	0.35	2.6	0.67	3.82
1.3	0.14	0.12	1.63	0.51	3.99
2.5	0.26	0.06	0.9	0.27	2.22
3.3	0.35	0.02	0.8	0.24	1.76

$K_m$  and  $V_{max}$  values were calculated as shown in Fig. 2b using double-reciprocal plot.




Fig. 3 The effect of propofol on the  $\text{Na}^+$  concentration-dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

$\text{Na}^+$  concentration dependency of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity was measured in the absence of DMSO and propofol ( $\diamond$ ), or in the presence of 2.5% DMSO without propofol ( $\square$ ); 1.3% DMSO and 0.14 mM propofol ( $\triangle$ ); 2.5% DMSO and 0.26 mM propofol ( $\circ$ ); 3.3% DMSO and 0.35 mM Propofol ( $\times$ ). The results were analyzed by the Hill equation, and the  $\text{Na}^+$  concentration that caused half-maximal activation ( $[\text{S}]_{0.5}$ ), maximum activity ( $V_{max}$ ) of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity, and the Hill coefficient  $n$  are summarized in Table 2. Data have a mean  $\pm$  SEM ( $n \geq 3$ ).

\* $P < 0.05$  and \*\* $P < 0.01$  compared with two groups of each. ( $\triangle$ ) vs ( $\circ$ );  $P < 0.01$ , ( $\circ$ ) vs ( $\times$ );  $P < 0.05$ .

and the Hill coefficient  $n$  were calculated as 12.57 mM, 2.26  $\mu\text{mol/mg protein/min}$  and 1.71, respectively (Table 2). Similar experiments were done in the presence of 0.14 ( $\triangle$ ), 0.26 ( $\circ$ ) and 0.35 mM ( $\times$ ) propofol or DMSO ( $\square$ ) (Fig. 3). The results were analyzed by the Hill equation, and  $[\text{S}]_{0.5}$ ,  $V_{max}$  and the Hill coefficient  $n$  are summarized in Table 2. DMSO did not change the  $[\text{S}]_{0.5}$  and  $V_{max}$  significantly, but propofol decreased both  $V_{max}$  and affinities for  $\text{Na}^+$ .

**Table 2** Effects of propofol on Na<sup>+</sup>-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

DMSO (%)	Propofol (mM)	Hill coefficient	[S] <sub>0.5</sub> (mM)	Vmax (μmol/mg/min)
0	0	1.71	12.57	2.26
2.5	0	1.91	11.91	2.37
1.3	0.14	1.86	21.16	1.87
2.5	0.26	2.04	21.65	1.03
3.3	0.35	2.14	19.67	0.69

Concentration-response data of Fig. 3 were fitted by nonlinear regression to the Hill equation.


[S]<sub>0.5</sub> = the concentration with half maximal activity. Vmax = the maximum activity.

#### 4. The effect of propofol on the K<sup>+</sup> concentration-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

Fig. 4 shows the effect of propofol on the K<sup>+</sup> concentration dependency of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity. In the absence of DMSO and propofol (◇), the activity increased, depending on the K<sup>+</sup> concentration. The results were analyzed by the Hill equation, and the [S]<sub>0.5</sub> for K<sup>+</sup>, Vmax and the Hill coefficient n were calculated as 1.5 mM, 2.19 μmol/mg protein/min and 1.53, respectively (Table 3). Similar experiments were done in the presence of 0.14 (△), 0.26 (○) and 0.35 mM (×) propofol or DMSO (□) (Fig. 4). The results were analyzed by the Hill equation, and [S]<sub>0.5</sub>, Vmax and the Hill coefficient n are summarized in Table 3. DMSO did not change the [S]<sub>0.5</sub> and slightly increased Vmax, but propofol decreased the Vmax and increased the affinity for K<sup>+</sup>.

#### 5. The effect of propofol on the Mg<sup>2+</sup> concentration-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

Fig. 5 shows the effect of propofol on the Mg<sup>2+</sup> concentration dependency of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity. In the absence of DMSO and propofol (◇), the activity increased, depending on the Mg<sup>2+</sup> concentration. The results were analyzed by the Hill equation, and the [S]<sub>0.5</sub> for Mg<sup>2+</sup>, Vmax and the Hill coefficient n were calculated as 0.43 mM, 2.23 μmol/mg protein/min and 1.15, respectively (Table 4). Similar experiments were done in the presence of 0.14 (△), 0.26 (○) and 0.35 mM (×) propofol or DMSO (□) (Fig. 5). The results were analyzed by the Hill equation, and [S]<sub>0.5</sub>, Vmax and the Hill coefficient n are summarized in Table 4. DMSO did not change the [S]<sub>0.5</sub> significantly but increased Vmax. Propofol also did not change the [S]<sub>0.5</sub> significantly but decreased the Vmax, depending on its concentration.

**Fig. 4** The effect of propofol on the K<sup>+</sup> concentration-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

K<sup>+</sup> concentration dependency of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was measured in the absence of DMSO and propofol (◇), or in the presence of 2.5% DMSO without propofol (□); 1.3% DMSO and 0.14 mM propofol (△); 2.5% DMSO and 0.26 mM propofol (○); 3.3% DMSO and 0.35 mM propofol (×). The results were analyzed by the Hill equation, and the [S]<sub>0.5</sub> for K<sup>+</sup>, Vmax, and the Hill coefficient n are summarized in Table 3. Data have a mean ± SEM (n ≥ 3).

\*\*P < 0.01 compared with two groups of each. (◇) vs (△); P < 0.01, (△) vs (○); P < 0.01, (○) vs (×); P < 0.01.

**Table 3** Effects of propofol on K<sup>+</sup>-activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

DMSO (%)	Propofol (mM)	Hill coefficient	[S] <sub>0.5</sub> (mM)	Vmax (μmol/mg/min)
0	0	1.53	1.5	2.19
2.5	0	1.54	1.5	2.53
1.3	0.14	1.5	1.09	1.78
2.5	0.26	1.67	0.99	1
3.3	0.35	1.75	0.96	0.21

Concentration-response data of Fig. 4 were fitted by nonlinear regression to the Hill equation like Table 2.

**Table 4** Effects of propofol on Mg<sup>2+</sup>-dependent activation of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

DMSO (%)	Propofol (mM)	Hill coefficient	[S] <sub>0.5</sub> (mM)	Vmax (μmol/mg/min)
0	0	1.15	0.43	2.23
2.5	0	1.37	0.47	2.7
1.3	0.14	1.13	0.44	1.59
2.5	0.26	1.51	0.5	1.25
3.3	0.35	1.61	0.49	1.01

Concentration-response data of Fig. 5 were fitted by nonlinear regression to the Hill equation like Table 2, and 3.




Fig. 5 The effect of propofol on the  $Mg^{2+}$  concentration-dependent activation of  $Na^+$ ,  $K^+$ -ATPase activity.

$Mg^{2+}$  concentration dependency of  $Na^+$ ,  $K^+$ -ATPase activity was measured in the absence of DMSO and propofol ( $\diamond$ ), or in the presence of 2.5% DMSO without propofol ( $\square$ ); 1.3% DMSO and 0.14 mM propofol ( $\triangle$ ); 2.5% DMSO and 0.26 mM propofol ( $\circ$ ); 3.3% DMSO and 0.35 mM propofol ( $\times$ ). The results were analyzed by the Hill equation, and the  $[S]_{0.5}$  for  $Mg^{2+}$ ,  $V_{max}$ , and the Hill coefficient  $n$  are summarized in Table 4. Data have a mean  $\pm$ SEM ( $n \geq 3$ ).

\* $P < 0.05$  compared with two groups of each. ( $\square$ ) vs ( $\diamond$ );  $P < 0.05$ , ( $\diamond$ ) vs ( $\triangle$ );  $P < 0.05$ , ( $\triangle$ ) vs ( $\circ$ );  $P < 0.05$ , ( $\circ$ ) vs ( $\times$ );  $P < 0.05$ .

## 6. Concentration-dependent inhibition of $Na^+$ -ATPase activity by propofol

Fig. 6 shows the inhibition of  $Na^+$ -ATPase activity in the presence of various concentrations of propofol. The data obtained by  $Na^+$ -ATPase activity tended to scatter compared with  $Na^+$ ,  $K^+$ -ATPase, because of the untableteness of  $Na^+$ -ATPase. The activity decreased, depending on the concentration of propofol, and was inhibited completely at 1.03 mM.  $IC_{50}$  of  $Na^+$ -ATPase inhibition was about 0.26 mM (Fig.6).

## 7. The effect of propofol on the $Na^+$ concentration-dependent activation of $Na^+$ -ATPase activity.

Fig. 7 shows the effect of propofol on the  $Na^+$  concentration dependency of  $Na^+$ -ATPase activity. In the absence of DMSO and propofol ( $\diamond$ ), the activity increased, depending on the  $Na^+$  concentration ( $[S]_{0.5} \approx 2$  mM), and  $V_{max}$  was attained at 5 mM. Similar experiments were done in the presence of 0.14 mM propofol ( $\square$ ) or DMSO ( $\triangle$ ) (Fig. 7).  $Na^+$ -ATPase activity also increased, depending on the concentration of  $Na^+$ , and  $V_{max}$  was attained at 5 mM. Both DMSO and propofol decreased the  $V_{max}$ , but DMSO decreased  $[S]_{0.5}$




Fig. 6 Concentration-dependent inhibition of  $Na^+$ -ATPase activity by propofol.

$Na^+$ -ATPase activity was measured in the presence of various concentrations of propofol. The activity decreased depending on the concentration of propofol, and was inhibited completely at 1.03 mM. The concentration that caused half-maximal inhibition of  $Na^+$ -ATPase activity was 0.26 mM. Data have a mean  $\pm$ SEM ( $n \geq 3$ ).





Fig. 7 The effect of propofol on the  $Na^+$  concentration-dependent activation of  $Na^+$ -ATPase activity.

$Na^+$  concentration dependency of  $Na^+$ -ATPase activity was measured in the absence of DMSO and propofol ( $\diamond$ ), or in the presence of 2.5% DMSO without propofol ( $\square$ ); 1.3% DMSO and 0.14mM propofol ( $\triangle$ ). Data have a mean  $\pm$ SEM ( $n \geq 3$ ).

for  $Na^+$  ( $[S]_{0.5} \approx 1$  mM), and propofol increased it ( $[S]_{0.5} \approx 3$  mM).

## 8. The effect of propofol on the $K^+$ concentration-dependent activation of $K^+$ -pNPPase activity.

Fig. 8 shows the effect of propofol on the  $K^+$  concentration dependency of  $K^+$ -pNPPase activity. In the absence of DMSO and propofol ( $\diamond$ ), the activity increased, depending on the  $K^+$  concentration. The results were analyzed by the Hill equation, and the  $[S]_{0.5}$  for  $K^+$ ,  $V_{max}$  and the Hill coefficient  $n$  were calculated as 2.43 mM, 908 nmol/mg protein/min, and 1.86 respectively (Table 5). Similar experiments were done in the presence of 0.14 ( $\triangle$ ), 0.26 ( $\circ$ ) and 0.35 mM ( $\times$ )



**Fig. 8** The effect of propofol on the K<sup>+</sup> concentration-dependent activation of K<sup>+</sup>-pNPPase activity.

K<sup>+</sup> concentration dependency of K<sup>+</sup>-pNPPase activity was measured in the absence of DMSO and propofol (◇), or in the presence of 2.5% DMSO without propofol (□); 1.3% DMSO and 0.14mM propofol (△); 2.5% DMSO and 0.26 mM propofol (○); 3.3% DMSO and 0.35 mM propofol (×). The results were analyzed by the Hill equation, and the [S]<sub>0.5</sub> for K<sup>+</sup>, V<sub>max</sub>, and the Hill coefficient n are summarized in Table 5. Data have a mean ± SEM (n≥3).

\*P<0.05 and \*\*P<0.01 compared with two groups of each. (□) vs (◇); P<0.05, (◇) vs (△); P<0.01, (○) vs (×); P<0.01.

**Table 5** Effects of propofol on K<sup>+</sup>-dependent activation of K<sup>+</sup>-pNPPase activity.


DMSO (%)	Propofol (mM)	Hill coefficient	[S] <sub>0.5</sub> (mM)	V <sub>max</sub> (nmol/mg/min)
0	0	1.86	2.43	908
2.5	0	1.85	2.1	1041
1.3	0.14	1.97	1.62	415
2.5	0.26	2.49	2.04	372
3.3	0.35	1.9	2.61	163

Concentration-response data of Fig. 8 were fitted by nonlinear regression to the Hill equation like Table 2-4.

propofol or DMSO (□) (Fig. 8). The results were analyzed by the Hill equation, and [S]<sub>0.5</sub>, V<sub>max</sub>, and the Hill coefficient n were summarized in Table 5. DMSO slightly decreased the [S]<sub>0.5</sub> and increased V<sub>max</sub>. Propofol decreased the V<sub>max</sub>, depending on its concentration but increased the affinity for K<sup>+</sup> at 0.14 (△) and 0.26 (○) mM, and then decreased it at 0.35 mM (×).

### 9. Reversibility of propofol inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity by decreasing its concentration.

As described in Materials and Methods, Na<sup>+</sup>, K<sup>+</sup>-ATPase was exposed to 1.03 mM propofol at which concentration the activity was completely inhibited, and



**Fig. 9** Reversibility of propofol inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity by decreasing its concentration.

At first, Na<sup>+</sup>, K<sup>+</sup>-ATPase was exposed to 1.03 mM propofol and 10% DMSO (open bar) or 10% DMSO without propofol as a control for solvent (closed bar). In the presence of 1.03 mM propofol and 10% DMSO (open bar), Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was inhibited completely (See also Fig. 1 and text). In the presence of 10% DMSO (closed bar), about 35% of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was inhibited. Then propofol and/or DMSO concentration was reduced to the final concentrations shown on the graph by dilution and then Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was measured. The activity was recovered depending on the degree of dilution and was about 60% at 0.05 mM propofol. Data have a mean ± SEM (n≥3).

\*\*P<0.01 compared with two groups.

then after dilution of propofol Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was measured (Fig. 9). Na<sup>+</sup>, K<sup>+</sup>-ATPase activity was recovered to about 60% at 0.05 mM and 40% at 0.17 mM propofol after dilution. The extent of activity recovery was dependent on the propofol concentration after dilution. A difference in the temperature, being on ice or at room temperature during preincubation did not affect the results.

## Discussion

### 1. Mechanism for propofol inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity

There is a report about the effect of propofol on Na<sup>+</sup>, K<sup>+</sup>-ATPase activity by Kutchai *et al.* [16] which showed propofol inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity depending on its concentration. However, they did not study the inhibition mechanism. As Na<sup>+</sup>, K<sup>+</sup>-ATPase plays essential roles for the maintenance of neuronal excitability, its inhibition may be related to a change in brain function. For this reason, we studied the inhibition mechanism for propofol inhibition of Na<sup>+</sup>, K<sup>+</sup>-ATPase activity.

ATP hydrolysis by Na<sup>+</sup>, K<sup>+</sup>-ATPase requires the presence of Na<sup>+</sup>, K<sup>+</sup> and Mg<sup>2+</sup> [13, 20]. In the Post-Albers reaction sequence, the enzyme binds ATP in the presence of Na<sup>+</sup> and Mg<sup>2+</sup>, hydrolyze ATP and forms a

phosphorylated enzyme (EP) with  $\gamma$ -phosphate of ATP as a reaction intermediate. Then  $\text{Na}^+$ -bound EP (called E1P) releases  $\text{Na}^+$  to the outside of the cell and changes the conformation to E2P, which is sensitive to  $\text{K}^+$ . In the absence of  $\text{K}^+$ , E2P is dephosphorylated spontaneously and releases phosphate. This reaction is called  $\text{Na}^+$ -ATPase activity, as a partial reaction of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase. In the presence of  $\text{K}^+$ ,  $\text{K}^+$  binds to E2P and accelerates the dephosphorylation. After the release of phosphate  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase forms a  $\text{K}^+$ -bound enzyme (KE2). KE2 hydrolyzes *p*-nitrophenyl phosphate (*p*NPP) and this activity is called  $\text{K}^+$ -*p*NPPase as a partial reaction of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity [21]. To release  $\text{K}^+$  from KE2 and start a new cycle, relatively high concentrations of ATP is necessary (low affinity site), however low concentrations of ATP is enough to form EP (high affinity site) [22].  $\text{Mg}^{2+}$  is essential for an ATPase reaction. As expected from these reaction sequences,  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and ATP modulates the  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

In order to study at which step propofol inhibits  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity, we examined the effect of propofol on  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and ATP dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity. Propofol decreased all  $V_{\text{max}}$  values for  $\text{Na}^+$  (Fig 3 and Table 2),  $\text{K}^+$  (Fig 4 and Table 3),  $\text{Mg}^{2+}$  (Fig 5 and Table 4) and ATP (Fig 2 and Table 1) dependent activation of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity. These results suggest that propofol affects the whole reaction sequence of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase, not a specific step. Propofol increased the affinity of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase for both high and low ATP and  $\text{K}^+$ , decreased it for  $\text{Na}^+$  and did not change it for  $\text{Mg}^{2+}$  significantly. In brief, propofol decreased the  $V_{\text{max}}$  and also changed the affinities of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase for  $\text{Na}^+$ ,  $\text{K}^+$ ,  $\text{Mg}^{2+}$  and ATP. These results suggest that the inhibition pattern of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase by propofol is a mixed-type, but not competitive or non-competitive.

We studied the effects of propofol on  $\text{Na}^+$ -ATPase (Fig 7) and  $\text{K}^+$ -*p*NPPase (Fig 8 and Table 5) activities, which are partial reactions of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity. Propofol decreased the  $V_{\text{max}}$  and the affinity for  $\text{Na}^+$  of  $\text{Na}^+$ -ATPase activity. Propofol also decreased the  $V_{\text{max}}$  and changed the affinity for  $\text{K}^+$  of  $\text{K}^+$ -*p*NPPase activity. These results support the above suggestions obtained from the effect of propofol on  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity.

As far as we know, this is the first report about the inhibition mechanism for  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase by propofol.

## 2. Reversibility of propofol inhibition of $\text{Na}^+$ , $\text{K}^+$ -ATPase activity

By diluting propofol concentration, the  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity which was inhibited nearly completely by 1.03 mM propofol, was recovered depending on the degree of dilution (Fig. 9). This result suggests that the effect of propofol is reversible *in vitro*, and we may suppose that the inhibition of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity by propofol is also reversible *in vivo*, which is necessary to wake a human from anesthesia.

## 3. Propofol concentration for the inhibition of $\text{Na}^+$ , $\text{K}^+$ -ATPase activity and anesthesia.

Propofol inhibited  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase and  $\text{Na}^+$ -ATPase activities depending on its concentration (Figs. 1 and 6). For complete inhibition, 1.03 mM propofol was necessary, however approximately 29% of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase or 24% of  $\text{Na}^+$ -ATPase activity was inhibited in the presence of 65  $\mu\text{M}$  propofol.

Though there are some discrepancies about propofol concentrations necessary for anesthesia, Kazama *et al.* [23] measured the propofol concentrations of plasma obtained from the patients during the time that the patient was under anesthesia using high-performance liquid chromatography. They reported that the propofol concentrations at which 50% of the patients did not respond to skin incisions, peritoneum incisions or abdominal wall retractions, were 72, 96 or 109  $\mu\text{M}$ , respectively. These results suggest that almost 30% of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase activity may be inhibited at the range of plasma concentrations found by Kazama *et al.* during anesthesia and that this inhibition may be related to the anesthesia. Higuchi *et al.* [24] reported that propofol suppressed a hyperpolarization-activated inward current by approximately 10–20% in similar concentrations, supporting our results. The role of  $\text{Na}^+$ ,  $\text{K}^+$ -ATPase inhibition for the state of anesthesia remains to be studied.

## References

- 1) Reves JG, Glass PSA, Lubarsky DA, McEvoy MD, Martinez-Ruiz R : Intravenous anesthetics. edited by Miller RD, Miller's anesthesia seventh edition, 719–728, Churchill Livingstone Elsevier, Philadelphia, 2010.
- 2) Bovill JG : Mechanism of intravenous anesthesia. edited by White PF, Textbook of intravenous anesthesia, 27–46, Williams and Wilkins, Philadelphia,

- 1997.
- 3) Trapani G, Altomare C, Sanna E, Biggio G, Liso G : Propofol in anesthesia. Mechanism of action, structure-activity relationships, and drug delivery. *Curr Med Chem* 7 : 249-271, 2000.
  - 4) Krasowski MD, Nishikawa K, Nikolaeva N, Lin A, Harrison NL : Methionine 286 in transmembrane domain 3 of the GABA<sub>A</sub> receptor  $\beta$  subunit controls a binding cavity for propofol and other alkylphenol general anesthetics. *Neuropharmacology* 41 : 952-964, 2001.
  - 5) Lingamaneni R, Birch ML, Hemmings HC : Widespread inhibition of sodium channel-dependent glutamate release from isolated nerve terminals by isoflurane and propofol. *Anesthesiology* 95 : 1460-1466, 2001.
  - 6) Pain L, Jeltsch H, Lehmann O, Lazarus C, Laalou FZ, Cassel JC : Central cholinergic depletion induced by 192 IgG-Saporin alleviates the sedative effects of propofol in rats. *Br J Anaesth* 85 : 869-873, 2000.
  - 7) Antognini JF, Wang XW, Piercy M, Carstens E : Propofol directly depresses lumbar dorsal horn neuronal responses to noxious stimulation in goats. *Can J Anesth* 47 : 273-279, 2000.
  - 8) Kotani Y, Shimazawa M, Yoshimura S, Iwama T, Hara H : Review The experimental and clinical pharmacology of propofol, an anesthetic agent with neuroprotective properties. *CNS Neurosci & Ther* 14 : 95-106, 2008.
  - 9) Hara M, Kai Y, Ikemoto Y : Enhancement by propofol of the  $\gamma$ -aminobutyric acid<sub>A</sub> response in dissociated hippocampal pyramidal neurons of the rat. *Anesthesiology* 81 : 988-994, 1994.
  - 10) Fagerlund MJ, Sjodin J, Krupp J, Dabrowski MA : Reduced effect of propofol at human  $\alpha 1 \beta 2$  (N289M)  $\gamma 2$  and  $\alpha 2 \beta 3$  (N290M)  $\gamma 2$  mutant GABA<sub>A</sub> receptors. *Br J Anaesth* 104 : 472-481, 2010.
  - 11) Eghbali M, Gage PW, Birnir B : Effects of propofol on GABA<sub>A</sub> channel conductance in rat-cultured hippocampal neurons. *Eur J Pharmacol* 468 : 75-82, 2003.
  - 12) Buggy DJ, Nicol B, Rowbotham DJ, Lambert DG : Effects of intravenous anesthetic agents on glutamate release. A role for GABA<sub>A</sub> receptor-mediated inhibition. *Anesthesiology* 92 : 1067-1073, 2000.
  - 13) Kaplan JH : Biochemistry of the Na, K-ATPase. *Annu Rev Biochem* 71 : 511-535, 2002.
  - 14) Vasic V, Krinulovic K, Krstic D, Momic T, Horvat A : ATPases as multi-response sensing system for various organic and inorganic analytes. *Monatshefte fur Chemie* 135 : 605-614, 2004.
  - 15) Krstic D, Krinulovic K, Spasojevic-Tisma V, Joksic G, Momic T, Vasic V : Effects of digoxin and ditoxin on the enzymatic activity and kinetic parameters of Na<sup>+</sup> / K<sup>+</sup>-ATPase. *J Enz Inhibit Med Chem* 19 : 409-415, 2004.
  - 16) Kutchai H, Geddis LM, Farley RA : Effects of general anaesthetics on the activity of the Na, K-ATPase of canine renal medulla. *Pharmacol Res* 40 : 469-473, 1999.
  - 17) Jorgensen PL : Purification of Na<sup>+</sup>, K<sup>+</sup>-ATPase. Enzyme sources, preparative problems, and preparation from mammalian kidney. *Methods Enzymol* 156 : 29-42, 1988.
  - 18) Vasallo PM, Post RL : Calcium ion as a probe of the monovalent cation center of sodium, potassium ATPase. *J Biol Chem* 261 : 16957-16962, 1986.
  - 19) Chifflet S, Torriglia A, Chiesa R, Tolosa S : A method for the determination of inorganic phosphate in the presence of labile organic phosphate and high concentrations of protein : application to lens ATPases. *Anal Biochem* 168 : 1-4, 1988.
  - 20) Horisberger JD : Recent insights into the structure and mechanism of the sodium pump. *Physiology* 19 : 377-387, 2004.
  - 21) Taniguchi K, Suzuki K, Iida S : Conformational change accompanying transition of ADP-sensitive phosphoenzyme to potassium-sensitive phosphoenzyme of (Na<sup>+</sup>, K<sup>+</sup>) -ATPase modified with *N*-[*p*-(2-Benzimidazolyl) phenyl]maleimide. *J Biol Chem* 257 : 10659-10667, 1982.
  - 22) Neufeld AH, Levy HM : A second ouabain-sensitive sodium-dependent adenosine triphosphatase in brain microsomes. *J Biol Chem* 244 : 6493-6497, 1969.
  - 23) Kazama T, Ikeda K, Morita K : The pharmacodynamic interaction between propofol and fentanyl with respect to the suppression of somatic or hemodynamic responses to skin incision, peritoneum incision, and abdominal wall retraction. *Anesthesiology* 89 : 894-906, 1998.
  - 24) Higuchi H, Funahashi M, Miyawaki T, Mitoh Y, Kohjitani A, Shimada M, Matuo R : Suppression of the hyperpolarization-activated inward current contributes to the inhibitory actions of propofol on rat CA1 and CA3 pyramidal neurons. *Neurosci Res* 45 : 459-472, 2003.