Title	Successful Kidney Transplantation Ameliorates Arterial Stiffness in End-Stage Renal Disease Patients
Author(s)	Hotta, Kiyohiko; Harada, Hiroshi; Sasaki, Hajime; Iwami, Daiki; Fukuzawa, Nobuyuki; Morita, Ken; Seki, Toshimori; Togashi, Masaki; Nonomura, Katsuya
Citation	Transplantation Proceedings, 44(3), 684-686 https://doi.org/10.1016/j.transproceed.2011.11.016
Issue Date	2012-04
Doc URL	http://hdl.handle.net/2115/49270
Type	article (author version)
File Information	TP44-3_684-686.pdf

[Instructions for use](#)

Title page

Successful kidney transplantation ameliorates arterial stiffness in end-stage renal disease patients

Kiyohiko Hotta^{1,2}, hotta1125@mac.com

Hiroshi Harada², hirohara80@mac.com

Hajime Sasaki^{1,2}, hajimes1207@hotmail.com

Daiki Iwami¹, iwamidaiki@ybb.ne.jp

Nobuyuki Fukuzawa¹, fukuzawa1972@mac.com

Ken Morita¹, kenordic@mac.com

Toshimori Seki³, uroseki@jcom.home.ne.jp

Masaki Togashi³, masaki.togashi@doc.city.sapporo.jp

Katsuya Nonomura¹, k-nonno@med.hokudai.ac.jp

¹ Department of Renal and Genitourinary Surgery, Hokkaido University Graduate School of Medicine, Sapporo, Japan.

² Department of Kidney Transplant Surgery, Sapporo City General Hospital, Sapporo, Japan.

³ Department of Urology, Sapporo City General Hospital, Sapporo, Japan.

Correspondence: *Kiyohiko Hotta, Department of Kidney Transplant Surgery, Sapporo City General Hospital, kita-11 Nishi-13 Chuoku, Sapporo, Hokkaido, 060-8604, Japan. Phone: 81-11-726-2211; Fax: 81-11-726-7912; E-mail: hotta1125@mac.com*

Two tables and one figure are included.

Abstract

PURPOSE: Successful kidney transplantation (KTx) can ameliorate bodily damage caused by end-stage renal disease (ESRD). Arterial stiffness (AS) is one of the critical factors, that shorten the survival of patients due to cardiovascular events. KTx may reduce AS as well; however, this has not been investigated well. We therefore conducted a retrospective study using non-invasive pulse wave velocity (PWV), which is a useful index of aortic damage.

PATIENTS and METHODS: Fifty-eight consecutive kidney recipients (34 men, 24 women) were enrolled in this study. Mean age at transplantation was 40.5 ± 12.3 years and the dialysis period was 73.1 ± 95.8 months. The brachial-ankle PWV was measured preoperatively and six months postoperatively. First, we investigated the relationship between the PWV and the other parameters related to AS. Second, we studied the pre- to post-transplant change in PWV to evaluate the amelioration of AS after successful kidney transplantation.

RESULTS: PWV showed significant positive correlations with age, systolic blood pressure (BP), diastolic BP and abdominal aortic calcification index. After successful KTx, PWV significantly decreased. ($P < 0.01$) In addition, systolic and diastolic BP significantly decreased. ($P < 0.01$ and $P < 0.05$, respectively) **CONCLUSION:** Successful KTx ameliorates AS in ESRD patients. This might explain the improved cardiovascular prognosis of ESRD patients who undergo kidney transplantation.

Introduction

Chronic kidney disease (CKD) is one of the main risk factor for cardiovascular events. Arterial stiffness (AS) is accelerated in patients with end stage renal disease (ESRD) and leads to cardiovascular events.¹ Patients on dialysis suffer higher rates of cardiovascular morbidity and mortality than the general population. Successful kidney transplantation (KTx) can reduce the cardiovascular risk and improve cardiovascular survival for dialysis patients.^{2,3} Artery damage is one of the major factors determining the high prevalence of cardiovascular disease (CVD) in patients with ESRD.⁴ KTx may ameliorate arterial stiffness as well; however, this observation is not completely confirmed.

In clinical studies, pulse wave velocity (PWV) is a highly reproducible index of arterial rigidity. Repeatability studies have made it quiet feasible to investigate arterial stiffness in cardiovascular epidemiological studies.⁵⁻⁷ PWV is an independent predictive parameter for overall and cardiovascular mortality and a strong predictor of prognosis in patients with ESRD.⁸ In this study, we investigated the effect of kidney transplantation on arterial stiffness using noninvasive PWV, which is a useful index of aortic damage.

Methods

Patients

Fifty-eight consecutive kidney recipients (34 men, 24 women) were enrolled in this study. Mean

age at transplantation was 40.5 ± 12.3 years and the dialysis period was 73.1 ± 95.8 months. They were maintained with a quadruple immunosuppression comprised of tacrolimus, mycophenolate mofetil, methylprednisolone, and basiliximab. The inclusion requirement was stable renal function for six months after kidney transplantation.

Vascular parameters

First, patients rested in a supine position for five minutes. Oscillometric measurement of the pulse volume was recorded with a volume-plethysmographic apparatus (Colin Co. Ltd., Komaki, Japan), that simultaneously recorded brachial-ankle PWV (baPWV) and blood pressure (BP). The baPWV and BP were measured preoperatively and six months postoperatively. We also measured the abdominal aortic calcification index (ACI) calculated by abdominal CT scan preoperatively. We divided the CT scan images of the aorta in all slices from the diaphragm to the aortic bifurcation into 12 deltaic sectors and counted the number of calcified sectors. Then we calculated the ACI using the formula: $ACI (\%) = \text{sum total of calcified sectors} / \text{sum total of all sectors}$.

Biochemical parameters

Data for total cholesterol (TC), triglyceride (TG), hemoglobin A1c (HbA1c), serum calcium (Ca) and serum phosphorus (P) were obtained at the time of the preoperative baPWV examination.

Statistical Analysis

Statistical analysis was performed with Prism (version 5). Continuous data were summarized as mean $\pm$ standard deviation. All tests were two-sided and $P < 0.05$ was considered statistically significant.

Results

The clinical characteristics and biochemical parameters of the subjects are shown in Table 1. The mean baPWV was 15.9 ± 4.5 m/s in all patients. First, we investigated the relationship between the preoperative baPWV and the parameters related to arterial stiffness (AS), such as age, body mass index (BMI), BP, duration of pre-transplant dialysis, lipid profile (TC, TG), HbA1c, Ca, P, Ca x P and ACI. PWV showed significant positive correlations with age ($P < 0.001$), dialysis period ($P < 0.05$), systolic BP ($P < 0.001$), diastolic BP ($P < 0.001$) and ACI ($P < 0.001$). The other parameters did not correlate with PWV. (Fig. 1)

Second, we studied the difference between pre- and post-transplant vascular parameters, including baPWV, BP and ACI, to evaluate the amelioration of AS after successful kidney transplantation. After successful kidney transplantation, baPWV significantly decreased in all patients ($P < 0.01$). In addition, systolic and diastolic BP significantly decreased ($P < 0.01$ and $P < 0.05$, respectively). However, ACI did not change after kidney transplantation. (Table. 2)

Discussion

Successful kidney transplantation ameliorates arterial stiffness and reduces cardiovascular risk.

However, there is little information on the amelioration of AS after transplantation. In this study, using noninvasive baPWV, we demonstrated the amelioration of arterial stiffness six months after successful KTx.

It has previously been shown that aortic PWV, measured between the carotid and femoral arteries, is an independent predictive parameter for overall and cardiovascular mortality in patients with ESRD.⁸ Several studies examined the change of aortic PWV after KTx. Covic et al. found significant improvement of aortic PWV after transplantation.⁹ In contrast, another studies reported that KTx did not decrease aortic PWV.^{10,11} Recently, brachial-ankle PWV, which can be measured easily and noninvasively, was described. baPWV is also a useful parameter to estimate aortic damage, like aortic PWV.^{12,13} Therefore, we used noninvasive baPWV in this study as a parameter of arterial stiffness and demonstrated improvement after KTx. Similarly, in a cohort of 36 transplant recipients, Zoungas et al. investigated the reduction of peripheral PWV one year after KTx. They described that the difference of improvement between aortic and peripheral PWV may reflect the propensity of distal muscular arteries to be more responsive to changes in volume state, sympathetic innervation and neurohumoral-mediated vasomotor tone.¹⁴

There are several reports to evaluating the amelioration of arterial stiffness after KTx using other parameters. Hornum et al reported that the aortic augmentation index (AIX) and nitroglycerin-induced vasodilation were improved one year after KTx.¹⁵ Similarly, Zoungas et al.

also suggested that KTx significantly reduced the AIX.¹⁴ These data suggested that successful KTx ameliorated arterial stiffness in ESRD patients.

In contrast, several studies reported no reduction of arterial stiffness after KTx. Zoungas et al. examined the change of carotid artery intima-media thickness (IMT). However, there was no significant change in it at one year after KTx.¹⁴ In this study, we examined the aortic calcification in 29 KTx recipients. There was no significant change in the ACI six months after KTx. Sharon et al. also reported that there was no amelioration of coronary artery calcification (CAC) after KTx. However, they found that KTx appeared to slow down or arrest CAC, whereas CAC progressed in hemodialysis patients.¹⁶ Larger numbers of patients and a longer period of observation will be needed to confirm the improvements of these variables; however, it is at least clear that KTx does not cause severe arterial damage to progress.

In conclusion, using noninvasive baPWV, we demonstrated that successful KTx ameliorates arterial stiffness. This might explain the improved cardiovascular prognosis of ESRD patients who undergo kidney transplantation.

References

1. Go AS, Chertow GM, Fan D, *et al.* Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *N Engl J Med* 351: 1296 2004
2. Wolfe RA, Ashby VB, Milford EL, *et al.* Comparison of mortality in all patients on

- dialysis, patients on dialysis awaiting transplantation, and recipients of a first cadaveric transplant. *N Engl J Med* 341: 1725 1999
3. Meier-Kriesche HU, Schold JD, Srinivas TR, *et al.* Kidney transplantation halts cardiovascular disease progression in patients with end-stage renal disease. *Am J Transplant* 4: 1662 2004
 4. Covic A, Gusbeth-Tatomir P, Goldsmith DJ. Arterial stiffness in renal patients: an update. *Am J Kidney Dis* 45: 965 2005
 5. Blacher J, Asmar R, Djane S, *et al.* Aortic pulse wave velocity as a marker of cardiovascular risk in hypertensive patients. *Hypertension* 33: 1111 1999
 6. Willum-Hansen T, Staessen JA, Torp-Pedersen C, *et al.* Prognostic value of aortic pulse wave velocity as index of arterial stiffness in the general population. *Circulation* 113: 664 2006
 7. Laurent S, Boutouyrie P, Asmar R, *et al.* Aortic stiffness is an independent predictor of all-cause and cardiovascular mortality in hypertensive patients. *Hypertension* 37: 1236 2001
 8. Blacher J, Guerin AP, Pannier B, *et al.* Impact of aortic stiffness on survival in end-stage renal disease. *Circulation* 99: 2434 1999
 9. Covic A, Goldsmith DJ, Gusbeth-Tatomir P, *et al.* Successful renal transplantation

decreases aortic stiffness and increases vascular reactivity in dialysis patients.

Transplantation 76: 1573 2003

10. Bachelet-Rousseau C, Kearney-Schwartz A, Frimat L, *et al.* Evolution of arterial stiffness after kidney transplantation. *Nephrol Dial Transplant*: 2011
11. Pan CR, Schmaderer C, Roos M, *et al.* Comparing aortic stiffness in kidney transplant recipients, hemodialysis patients, and patients with chronic renal failure. *Clin Transplant* 25: E463 2011
12. Yamashina A, Tomiyama H, Takeda K, *et al.* Validity, reproducibility, and clinical significance of noninvasive brachial-ankle pulse wave velocity measurement. *Hypertens Res* 25: 359 2002
13. Nakamura U, Iwase M, Nohara S, *et al.* Usefulness of brachial-ankle pulse wave velocity measurement: correlation with abdominal aortic calcification. *Hypertens Res* 26: 163 2003
14. Zoungas S, Kerr PG, Chadban S, *et al.* Arterial function after successful renal transplantation. *Kidney Int* 65: 1882 2004
15. Hornum M, Clausen P, Idorn T, *et al.* Kidney transplantation improves arterial function measured by pulse wave analysis and endothelium-independent dilatation in uraemic patients despite deterioration of glucose metabolism. *Nephrol Dial Transplant* 26: 2370

2011

16. Moe SM, O'Neill KD, Reslerova M, *et al.* Natural history of vascular calcification in dialysis and transplant patients. *Nephrol Dial Transplant* 19: 2387 2004

Figure legend

Fig 1. Correlation between brachial-ankle pulse wave velocity (PWV) and the parameters related to arterial stiffness. baPWV show significant positive correlations with age ($P<0.001$), dialysis period ($P<0.05$), systolic BP ($P<0.001$), diastolic BP ($P<0.001$) and abdominal aortic calcification index (ACI) ($P<0.001$).

Fig 1

Table 1. The clinical characteristics and biochemical parameters in 58 transplant recipients

	n = 58
Age	40.5 $\pm$ 12.3
Male gender	34 (59%)
Dialysis duration (months)	73.1 $\pm$ 95.8
Body mass index (Kg/m ²)	21.0 $\pm$ 3.3
Systolic BP (mmHg)	134.4 $\pm$ 19.4
Diastolic BP (mmHg)	83.4 $\pm$ 11.5
Hemoglobin A1c (%)	4.9 $\pm$ 0.5
Total cholesterol (mg/dl)	162.2 $\pm$ 37.7
Triglyceride (mg/dl)	118.6 $\pm$ 81.6
Ca (mg/dl)	9.3 $\pm$ 1.0
P (mg/dl)	5.4 $\pm$ 1.7
Ca $\times$ P	50.6 $\pm$ 17.5
Aortic calcification index (%)	10.9 $\pm$ 16.3
Pulse wave velocity (m/sec)	15.9 $\pm$ 4.5

BP: blood pressure, Ca: serum calcium
P: serum phosphorus

Table 2. Comparison of vascular parameters pre- and post transplantation

	n	Pre KTx	Post KTx	P value
PWV (m/sec)	58	15.9 $\pm$ 4.5	14.3 $\pm$ 2.6	P<0.01
SBP (mmHg)	58	137.4 $\pm$ 19.4	127.6 $\pm$ 26.9	P<0.01
DBP (mmHg)	58	83.4 $\pm$ 11.5	79.5 $\pm$ 12.4	P<0.05
ACI (%)	29	10.1 $\pm$ 13.8	10.37 $\pm$ 14.9	NS

KTx: kidney transplantation, PWV: pulse wave velocity,
 SBP: systolic blood pressure, DBP: diastolic blood pressure
 ACI: aortic calcification index