Title	Experimental studies of surface reactions among OH radicals that yield H ₂ O and CO ₂ at 40-60 K
Author(s)	Oba, Yasuhiro; Watanabe, Naoki; Kouchi, Akira; Hama, Tetsuya; Pirronello, Valerio
Citation	Physical Chemistry Chemical Physics, 13(35), 15792-15797 https://doi.org/10.1039/c1cp20596j
Issue Date	2011-09-21
Doc URL	http://hdl.handle.net/2115/49372
Rights	Phys. Chem. Chem. Phys., 2011, 13(35), 15792-15797 - Reproduced by permission of the PCCP Owner Societies
Type	article (author version)
File Information	PCCP13-35_15792-15797.pdf

[Instructions for use](#)

Experimental studies of surface reactions among OH radicals that yield H₂O and CO₂ at 40–60 K

Yasuhiro Oba,^{*a} Naoki Watanabe,^a Akira Kouchi,^a Tetsuya Hama,^a and Valerio Pirronello^b

Received Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

First published on the web Xth XXXXXXXXXX 200X

DOI: 10.1039/b000000x

We investigated the OH-related formation routes of two astrophysically important molecules, H₂O and CO₂, under relatively warm astrophysical conditions. OH radicals, together with other neutral species such as H, O, H₂, and O₂, were produced in H₂O microwave-discharge plasma and cooled to 100 K before being deposited on an Al substrate at 40–60 K. H₂O formed at 40 and 50 K, but not at 60 K. Taking the experimental conditions into account, a possible route of H₂O formation is via reactions involving OH + OH, which yield H₂O₂ as the main reaction product. The present study is the first to show experimentally that surface reactions of two OH radicals can yield H₂O at low temperatures. The products' branching ratio was 0.2 and 0.8 for H₂O and H₂O₂, respectively. When CO was co-deposited with neutral species that formed in the H₂O plasma, CO₂ was formed at 40–60 K. H₂CO₃ formed at 40 and 50 K. The present results may suggest that chemical reactions related to OH radicals are effective at yielding various molecules in relatively warm astrophysical environments, such as protostars.

1 Introduction

It has been demonstrated theoretically that hydroxyl (OH) radicals are related to various chemical reactions that occur on solid surfaces in astrophysical environments.^{1,2} It is also well known that OH radicals play a significant role in photon-induced chemical reactions involving H₂O ices, because they are easily produced by the photolysis of solid H₂O at various temperatures.^{3–5} Previous experiments demonstrated experimentally that chemical reactions induced by UV photons in astrophysically relevant ices can yield various complex species.^{6,7} In those experiments, various reactions may occur due to energy input from photons, even if they are highly endothermic and have large activation barriers. In contrast, nonenergetic surface reactions are simple and suitable for understanding the elementary process of radical–molecule reactions. In our previous studies, nonenergetic OH radicals were obtained by dissociating H₂O in microwave-induced plasma, followed by cooling to 100 K before the reaction.^{8,9} Therefore, we are able to investigate the OH-related formation pathways of various molecules under astrophysically relevant conditions.

Solid H₂O is ubiquitous in icy grain mantles in dense molecular clouds.¹⁰ Previous studies have demonstrated experimentally that solid H₂O can be formed at ~10 K by the sequential addition of H to O,^{11,12} O₂,^{13,14} or O₃.^{15,16} Cup-

pen et al.¹⁷ recently proposed that hydrogen addition to HO₂ radicals can also yield H₂O on a low-temperature surface. Although H₂O formation routes involving atomic H have been extensively investigated experimentally, few studies have focused on OH-related formation routes of H₂O under astrophysically relevant conditions. In fact, OH radicals should form as an intermediate species during the sequential addition of H to O, O₂, or O₃.¹⁸ However, OH-related routes cannot be distinguished from other H₂O formation routes that have been shown to occur concurrently under experimental conditions.^{11–16} For example, Dulieu et al.¹² investigated reactions of atomic D (instead of H) with atomic O. In their experiment, atomic O was obtained by dissociating molecular O₂ with a dissociation efficiency of ~40%, which means that ~60% of molecular O₂ remained nondissociated and, therefore, could react with D. Although they estimated the efficiency of the O + D (OD + D) reaction from the temperature-programmed desorption spectra of the reaction products, the latter were not monitored in situ; no direct evidence was presented for the efficiency of the reaction. Therefore, if the OH-related formation routes of H₂O are understood in terms of their efficiency, the chemical-reaction network under astrophysically relevant conditions can be better understood.

Carbon dioxide (CO₂) is one of the most abundant solid components in icy grain mantles within molecular clouds. Recent astronomical observations by the Spitzer Space Telescope have revealed that the column density of solid CO₂ is high and almost constant relative to H₂O.^{19,20} It has been demonstrated theoretically that gas-phase reactions are unsuitable for the formation of the abundant CO₂ typically observed² and

^aInstitute of Low Temperature Science, Hokkaido University, N19W8, Kita-ku, Sapporo, Hokkaido 060-0819, JAPAN. Fax: +81 11 706 7142; Tel: +81 11 706 5477; E-mail: oba@lowtem.hokudai.ac.jp

^bDMFCI, Università di Catania, 95125 Catania, Sicily, ITALY

experimentally that abundant CO₂ can be formed by energetic irradiation, such as by UV or ions, of pure solid CO, H₂O/CO binary ices, and other astrophysically relevant ices containing CO.^{21–24} We recently reported that solid CO₂ can be formed by nonenergetic (*i.e.*, without external energetic sources such as photons and ions) surface reactions between CO and OH radicals at 10 and 20 K.⁸ More recently, CO₂ formation through the same reaction (CO + OH) was reported by other groups.^{25,26} These findings indicate that CO₂ formation can occur inside dense molecular clouds, where the UV field is relatively weak, which is in agreement with astronomical observations.^{20,27,28} However, it remains unknown whether nonenergetic reactions yield CO₂ at higher temperatures.

In the present paper, we discuss the formation routes of H₂O and other molecules (*e.g.*, O₃ and H₂O₂) that form on an aluminum (Al) substrate when neutral species in H₂O discharge plasma (H, O, OH, H₂, and O₂) are deposited at 40–60 K. At these temperatures, H₂O formation via H-related reactions (*e.g.*, hydrogenation to O and O₂) is expected to occur less frequently because of very short residence time of H atom at the surface, which means that OH-related routes of H₂O formation are better studied in the present experiment. This is the first report to investigate H₂O formation pathway through surface reactions of two OH radicals (OH + OH) at low temperatures. We also performed experiments on the co-deposition of plasma-derived neutral species with CO at the same temperatures, to show the efficiency of CO₂ formation at up to 60 K.

2 Experimental settings

Experiments were performed using the Apparatus for Surface Reaction in Astrophysics (ASURA) system. Details of this apparatus and experimental conditions have been described previously.^{8,9,29} Briefly, OH radicals, together with H atoms, were produced in an H₂O microwave-discharge plasma³⁰ in a Pyrex tube. H₂O dissociation may also lead to the direct formation of H₂ and O. These species are denoted as H₂O fragments in this paper. H₂O fragments were transferred via a series of poly(tetrafluoroethylene) and Al tubes, cooled to 100 K in the Al tube which was connected to another He refrigerator, and impinged on the substrate. The OH radicals are thought to be excited when formed in plasma, but after many collisions with the cold aluminum pipe, they are thermalized to 100 K and should already be vibrationally and electronically in the ground state, as expected in the gas-phase collision,³¹ before reacting on the cold substrate (40–60 K). The deposition rate of OH radicals could not be measured directly, but can be roughly estimated from the rates of H₂O₂ formation and H₂O deposition when the microwave source is turned off,^{8,9} yielding a value of approximately 1.2×10^{13} radicals cm⁻² s⁻¹. CO molecules were co-deposited with the H₂O

fragments onto the Al substrate for 120 min. The deposition rate of CO molecules was 1.4×10^{13} molecules cm⁻² s⁻¹. The reaction products were monitored in situ by infrared (IR) reflection-absorption spectroscopy with a resolution of 4 cm⁻¹ in the spectral range between 4000 and 700 cm⁻¹. The desorbed species from the substrate were monitored using a quadrupole mass spectrometer (QMS). A temperature for each experiment was chosen between 40 and 60 K, and the temperature of the Al substrate was maintained at that temperature throughout the experiment.

3 Results and Discussion

3.1 Deposition of H₂O fragments

When the H₂O fragments were solely deposited onto the Al substrate at 40 K, we observed the formation of H₂O₂ (~3300, 2849, and 1410 cm⁻¹), H₂O (~3300 and 1645 cm⁻¹), and O₃ (1040 cm⁻¹) in the IR spectrum of the reaction product (see Fig. 1; hereafter, this experiment is denoted as the blank experiment). Detailed assignments of these molecules have been reported elsewhere.^{32–34} The formation of these molecules was also observed at 50 K, but not at 60 K (data not shown). Because their desorption temperatures are higher than 60 K, we conclude that H₂O₂, H₂O, and O₃ were not formed in the gas phase but on the Al substrate. Trace amounts of CO₂ were observed in the spectrum, but CO₂ was not a reaction product, as explained previously.^{8,9}

QMS analyses revealed that molecular oxygen (O₂) was formed when the microwave source was turned on. However, we cannot deduce whether O₂ was formed on the Al surface or in the radical source. Brunet et al.³⁵ proposed that O₂ is formed on the inner wall of the radical source when H₂O is dissociated in microwave-induced plasma. Nevertheless, we believe that O₂ was partly formed by the following surface reactions:

Given that atomic O is present on the Al surface, these two reactions are expected to occur. The presence of atomic O is supported by the formation of O₃, as explained below.

For H₂O₂ formation, we considered the occurrence of the following reactions under the present experimental conditions:

Reaction (3) may be the predominant route of H₂O₂ formation, because it has no activation barrier.³⁶ It has been demonstrated experimentally that reactions (4) and (5) have essentially no barriers¹³ and are effective at yielding H₂O₂, even at

40 K;³⁷ hydrogenation of O₂ is likely to contribute to H₂O₂ formation. However, when O₂ was co-deposited with H₂O fragments onto the Al substrate at 10 K, the column density of the subsequently formed H₂O and H₂O₂ was almost the same as that in the blank experiment at the same temperature. This finding indicates that reactions (4) and (5) are very minor at 10 K and should also be at 40 K. Therefore, we assume that very little atomic H is contained in the H₂O fragments, a large fraction of atomic H is released into the gas phase prior to any reaction, and/or most of the atomic H reacted with itself to form H₂ on the surface.

Cuppen et al.¹⁷ proposed another H₂O₂ formation pathway from HO₂:

They also proposed that HO₂ radicals can dissociate into H + O₂ in low-temperature ices without energy inputs:

However, these reactions are not expected to occur under the present experimental conditions because reactions (6) and (7) are highly endothermic, with large barriers of 13100 and 24000 K, respectively.³⁶

The following are possible routes of H₂O formation:^{18,38}

Reactions (8)–(10) have no barriers, while reactions (11) and (12) have barriers of ~2000 K³⁹ and ~2100 K,³⁸ respectively. At lower temperatures (~10 K), barrierless reactions related to atomic H—reactions (8) and (9)—are effective because of the sufficiently long residence time of atomic H.^{12,17} Reaction (11) can also occur because of a tunneling effect.¹³ Reaction (12) has yet to be investigated experimentally but is expected to occur at temperatures as low as 10 K through quantum tunneling.⁴⁰ Under the present experimental conditions, however, reactions related to atomic H—reactions (8), (9), and (11)—are not expected to occur as mentioned earlier. Moreover, the residence time of H₂ on the Al substrate at 40 K would be too short for reaction (12) to occur through quantum tunneling.⁴¹ Therefore, the predominant route leading to H₂O formation is most likely reaction (10) in this experiment.

In addition to H₂O formation pathways, one may assume that the observed H₂O includes nondissociated H₂O from the radical source. However, little H₂O was observed in the blank experiment at 60 K (data not shown). We assume that dissociation fraction of H₂O is near unity and/or most H₂O from the

source was adsorbed on the inner wall of the aluminum pipe, which was cooled to 100 K.

We proposed above that reactions of two OH radicals (OH + OH) are solely responsible for the formation of both H₂O via reaction (10) and H₂O₂ via reaction (3). In addition, the following reactions have been studied in the gas phase:⁴²

However, reactions (13) and (14) are not expected to occur under the present experimental conditions, because these routes have large barriers of more than 25000 K.⁴²

Next, we calculated the column densities of H₂O and H₂O₂ that formed in the blank experiment at 40 K, using the band strengths for the peaks at 1645 and 1410 cm⁻¹, respectively.¹³ The column density of H₂O₂ was four times higher than that of H₂O, indicating that the branching ratios for the formation of H₂O and H₂O₂ by OH + OH were 0.2 and 0.8 at 40 K, respectively. At 50 K, on the other hand, we could not calculate these column densities because of the low signal-to-noise ratio of the IR spectrum.

The formation pathway of O₃ is much simpler than that of other molecules, and is limited to the following reaction:⁴

3.2 Codeposition of H₂O fragments with CO

When H₂O fragments were codeposited with CO at 40 K, we observed an increase in the CO₂ peak intensity at 2353 cm⁻¹ compared with the blank experiment (Fig. 1). It has been proposed that CO₂ forms by the following reaction routes under the present experimental conditions:⁸

Neither trans- nor cis-HOCO radicals were identified in the reaction products at 40–60 K, probably because HOCO radicals are highly reactive at these temperatures.^{8,43} Other routes to the formation of CO₂ such as CO + O or HOCO + H are unlikely to occur under the present experimental conditions as already mentioned in our previous study.⁸

In addition to CO₂, new peaks appeared at around 1750, 1500, and 1300 cm⁻¹, which were derived from carbonic acid (H₂CO₃) formed via the following reactions:⁹

Further results and discussion of the formation of H₂CO₃ have been published elsewhere.⁹ The formation of HCO, H₂CO,

CH₃OH, and HCOOH was not positively identified in the IR spectra of the reaction products, indicating that the hydrogenation of CO and HOCO did not occur under the present experimental conditions.

Note that with increasing substrate temperature, the formation of H₂O₂, H₂O, and O₃ becomes less effective, while CO₂ formed even at 60 K (Fig. 2). H₂CO₃ formed at 50 K, but negligible amounts were produced at 60 K (Fig. 2a). These results imply that once HOCO radicals are formed, they can reside on the surface, even at 50 K, and further yield CO₂ + H or react with additional OH radicals to yield H₂CO₃. At 60 K, the HOCO radicals may rapidly yield CO₂ through reactions (16)–(18). Fig. 2a shows that little or no H₂O₂ formation occurred at 60 K, although CO₂ was formed. This result indicates that the actual deposition rate of OH radicals could be lower than the estimated value (1.2×10^{13} radicals cm⁻² s⁻¹) and hence the actual one is much lower than that of CO (1.4×10^{13} molecules cm⁻² s⁻¹); OH has little chance to react with another OH before desorbing from the Al substrate. Nevertheless, our results clearly indicate that formation of CO₂ quickly occurred before CO and OH desorb from the Al substrate in spite of their very short residence time on the substrate at 60 K.

The peak position and shape of the CO₂ asymmetric stretch varied with substrate temperature (Fig. 2b). At 40 K, a strong peak appeared at 2353 cm⁻¹, while at 50 K, the peak appeared at 2363 cm⁻¹, concurrent with the appearance of a small shoulder at 2377 cm⁻¹. At 60 K, a strong peak appeared at 2380 cm⁻¹, with two small shoulders at 2345 and 2362 cm⁻¹. It has been reported that the asymmetric-stretch mode of pure, solid CO₂ falls at 2344 cm⁻¹ at 10 K, and its position changes little during warming up to 80 K.⁴⁴ These findings are clearly inconsistent with our experimental results. Therefore, a different explanation is required for the temperature dependence of the peak positions and shapes in Fig. 2b. Previous studies have revealed that the peak positions for the asymmetric mode of CO₂ crystals vary with size and conformation.^{45–48} For CO₂ crystals, their so-called transverse optical and longitudinal optical modes fall at 2344 and 2381 cm⁻¹, respectively,⁴⁷ which is consistent with the peak positions of CO₂ formed at 60 K (Fig. 2b). We assume that the band-shifting and splitting feature seen at higher temperatures is related to the degree of crystallinity and conformation of CO₂ formed at each temperatures. At 50 and 60 K, the CO₂ peak shape and position could also be perturbed by co-existing H₂O₂, O₃, or H₂CO₃ on the Al substrate. In view of molecular-vibrational spectroscopy, it may be of value to investigate the properties of the crystalline CO₂ formed in the present study; however, this is beyond the scope of our study.

The column density of CO₂ increased linearly with experimental time at each temperature (Fig. 3). The band strength for CO₂ (7.6×10^{-17} cm molecules⁻¹) was derived from Ger-

akines et al.³² The variation in CO₂ column density with time is clearly different from that in the same experiment at lower temperatures (10 and 20 K), where the column density reaches saturation with time.⁸ We reported previously that as the reaction proceeds, there is an increasing number of surface sites where CO and OH can be adsorbed, resulting in a decrease in the frequency of these species encountering each other on the surface at 10 and 20 K.⁸ In addition, the mobility of CO and OH show an increase with increasing temperature. In this case, the surface morphology constrains the reaction frequency to a much smaller extent, although the residence time on the surface should decrease for both species. Consequently, the CO₂ column density showed a linear increase with time (Fig. 3).

CO₂ formation becomes less effective with increasing temperature (Fig. 3). The conversion factor of CO into CO₂ was calculated to be 1.4%, 0.8%, and 0.3% at 40, 50, and 60 K, respectively. This temperature dependence is explained by the shorter residence times of CO and OH on the surface at higher temperatures. Although the conversion factor is not large compared with that for the same reaction at 10 K (which yielded values up to 10 %),⁸ the present results are of use because, to the best of our knowledge, this is the first experimental confirmation of the formation of solid CO₂ at temperatures as high as 60 K.

4 Summary

We investigated the reactions of H₂O fragments (H, O, OH, H₂, and O₂) on an Al substrate at 40–60 K, revealing that the formation of H₂O₂, H₂O, and O₃ occurred on the substrate and not in the gas phase. We propose that the predominant formation route of H₂O₂ and H₂O was a reaction between two OH radicals, where the branching ratios of the products were 0.8 for H₂O₂ and 0.2 for H₂O. The present study is the first to show experimentally that reactions of two OH radicals can yield H₂O under astrophysically relevant conditions.

When CO and H₂O fragments were co-deposited onto the Al substrate, CO₂ was formed by surface reactions of CO with nonenergetic OH radicals that formed in H₂O discharge plasma, even at 60 K. The conversion factor of CO to CO₂ was up to 1.4% under the present experimental conditions. H₂CO₃ formation was observed at 40 and 50 K, but was negligible at 60 K. HOCO radicals were not identified in IR spectra of the reaction products, unlike for the same experiment at 10 K.⁸ However, the formation of CO₂ and H₂CO₃ indicates that HOCO radicals were actually formed at 40–60 K, although they would have been rapidly consumed by further reactions to yield these molecules. These results may suggest that CO₂ and H₂CO₃ form via nonenergetic processes in relatively warm environments such as protostars.

Acknowledgement

We thank Dr. H. Hidaka for fruitful discussions. This work was partly supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (JSPS) and by a research fellowship from JSPS for Young Scientists (Y.O.).

References

- 1 L. B. d'Hendecourt, L. J. Allamandola and J. M. Greenberg, *Astron. Astrophys.*, 1985, **152**, 130.
- 2 T. I. Hasegawa, E. Herbst and C. M. Leung, *Astrophys. J. Suppl. Ser.*, 1992, **82**, 167.
- 3 T. Hama, A. Yabushita, M. Yokoyama, M. Kawasaki and N. Watanabe, *J. Chem. Phys.*, 2009, **131**, 114511.
- 4 P. D. Cooper, M. H. Moore and R. L. Hudson, *Icarus*, 2008, **194**, 379.
- 5 P. A. Gerakines, W. A. Schutte and P. Ehrenfreund, *Astron. Astrophys.*, 1996, **312**, 289.
- 6 L. J. Allamandola, S. A. Sandford and G. J. Valero, *Icarus*, 1988, **76**, 225.
- 7 K. I. Oberg, R. T. Garrod, E. F. van Dishoeck and H. Linnartz, *Astron. Astrophys.*, 2009, **504**, 891.
- 8 Y. Oba, N. Watanabe, A. Kouchi, T. Hama and V. Pirronello, *Astrophys. J.*, 2010, **712**, L174.
- 9 Y. Oba, N. Watanabe, A. Kouchi, T. Hama and V. Pirronello, *Astrophys. J.*, 2010, **722**, 1598.
- 10 E. L. Gibb, D. C. B. Whittet, A. C. A. Boogert and A. G. G. M. Tielens, *Astrophys. J.*, 2004, **151**, 35.
- 11 K. Hiraoka, T. Miyagoshi, T. Takayama, K. Yamamoto and Y. Kihara, *Astrophys. J.*, 1998, **498**, 710.
- 12 F. Dulieu, L. Amiaud, E. Congiu, J. H. Fillion, E. Matar, A. Momeni, V. Pirronello and J. L. Lemaire, *Astron. Astrophys.*, 2010, **512**, A30.
- 13 N. Miyauchi, H. Hidaka, T. Chigai, A. Nagaoka, N. Watanabe and A. Kouchi, *Chem. Phys. Lett.*, 2008, **456**, 27.
- 14 S. Ioppolo, H. M. Cuppen, C. Romanzin, E. F. van Dishoeck and H. Linnartz, *Astrophys. J.*, 2008, **686**, 1474.
- 15 H. Mokrane, H. Chaabouni, M. Accolla, E. Congiu, F. Dulieu, M. Chehrouri and J. L. Lemaire, *Astrophys. J.*, 2009, **705**, L195.
- 16 C. Romanzin, S. Ioppolo, H. M. Cuppen, E. F. van Dishoeck and H. Linnartz, *Phys. Chem. Chem. Phys.*, 2011, **134**, 084504.
- 17 H. M. Cuppen, S. Ioppolo, C. Romanzin and H. Linnartz, *Phys. Chem. Chem. Phys.*, 2010, **12**, 12077.
- 18 A. G. G. M. Tielens and W. Hagen, *Astron. Astrophys.*, 1982, **114**, 245.
- 19 K. M. Pontoppidan, A. C. A. Boogert, H. J. Fraser, E. F. van Dishoeck, G. A. Blake, F. Lahuis, K. I. Oberg, N. J. Evans and C. Salyk, *Astrophys. J.*, 2008, **678**, 1005.
- 20 D. C. B. Whittet, S. S. Shenoy, E. A. Bergin, J. E. Chiar, P. A. Gerakines, E. L. Gibb, G. J. Melnick, and D. A. Neufeld, *Astrophys. J.*, 2007, **655**, 332.
- 21 M. E. Palumbo, G. A. Baratta, J. R. Brucato, A. C. Castorina, M. A. Satorre and G. Strazzulla, *Astron. Astrophys.*, 1998, **334**, 247.
- 22 N. Watanabe and A. Kouchi, *Astrophys. J.*, 2002, **567**, 651.
- 23 M. J. Loeffler, G. A. Baratta, M. E. Palumbo, G. Strazzulla, and R. A. Baragiola, *Astron. Astrophys.*, 2005, **435**, 587.
- 24 S. Ioppolo, M. E. Palumbo, G. A. Baratta, and V. Menella, *Astron. Astrophys.*, 2009, **493**, 1017.
- 25 S. Ioppolo, Y. van Boheemen, H. M. Cuppen, E. F. van Dishoeck, and H. Linnartz, *Mon. Not. R. Astron. Soc.*, 2011, **413**, 2281.
- 26 J. A. Noble, F. Dulieu, E. Congiu, and H. J. Fraser, arXiv:1104.0031v1, 2011.
- 27 E. A. Bergin, G. J. Melnick, P. A. Gerakines, D. A. Neufeld and D. C. B. Whittet, *Astrophys. J.*, 2005, **627**, L33.
- 28 C. Knez, A. C. A. Boogert, K. M. Pontoppidan, J. Kessler-Silacci, E. F. van Dishoeck, N. J. Evans II, J.-C. Aigereau, G. A. Blake and F. Lahuis, *Astrophys. J.*, 2005, **635**, L145.
- 29 A. Nagaoka, N. Watanabe and A. Kouchi, *J. Phys. Chem. A*, 2007, **111**, 3016.
- 30 E. A. H. Timmermans, J. Jonkers, I. A. J. Thomas, A. Rodero, M. C. Quintero, A. Sola, A. Gamero and J. A. M. van der Mullen, *Spectroc. Acta Pt. B-Atom. Spectr.*, 1998, **53**, 1553.
- 31 N. Kohno, M. Izumi, H. Kohguchi and K. Yamasaki, *J. Phys. Chem. A*, 2011, **115**, 4867.
- 32 P. A. Gerakines, W. A. Schutte, J. M. Greenberg and E. F. van Dishoeck, *Astron. Astrophys.*, 1995, **296**, 810.
- 33 W. Zheng, D. Jewitt and R. I. Kaiser, *Phys. Chem. Chem. Phys.*, 2007, **9**, 2556.
- 34 C. J. Bennett and R. I. Kaiser, *Astrophys. J.*, 2005, **635**, 1362.
- 35 P. E. Brunet, X. Deglise, and P. A. Giguere, *Can. J. Chem.*, 1970, **48**, 2042.
- 36 W. Tsang and R. F. Hampson, *J. Phys. Chem. Ref. Data*, 1986, **15**, 1087.
- 37 Y. Oba, N. Miyauchi, H. Hidaka, T. Chigai, N. Watanabe and A. Kouchi, *Astrophys. J.*, 2009, **701**, 464.
- 38 R. Atkinson, D. L. Baulch, R. A. Cox, J. N. Crowley, R. F. Hampson, R. G. Hynes, M. E. Jenkin, M. J. Rossi and J. Troe, *Atmos. Chem. Phys.*, 2004, **4**, 1461.
- 39 H. Koussa, M. Bahri, N. Jaidane and Z. Ben Lakhdar, *J. Mol. Struct.: THEOCHEM*, 2006, **770**, 149.

- 40 H. M. Cuppen and E. Herbst, *Astrophys. J.*, 2007, **668**, 294.
- 41 N. Watanabe and A. Kouchi, *Prog. Surf. Sci.*, 2008, **83**, 439.
- 42 S. P. Karkach and V. I. Osherov, *J. Chem. Phys.*, 1999, **110**, 11918.
- 43 D. E. Milligan and M. E. Jacox, *J. Chem. Phys.*, 1971, **54**, 927.
- 44 S. A. Sandford and L. J. Allamandola, *Astrophys. J.*, 1990, **355**, 357.
- 45 G. E. Ewing and D. T. Sheng, *J. Phys. Chem.*, 1988, **92**, 4063.
- 46 F. Fleyfel and J. P. Devlin, *J. Phys. Chem.*, 1989, **93**, 7292.
- 47 M. A. Ovchinnikov and C. A. Wight, *J. Chem. Phys.*, 1993, **99**, 3374.
- 48 M. K. Kunzmann, R. Signorell, M. Taraschewski and S. Bauerecker, *Phys. Chem. Chem. Phys.*, 2001, **3**, 3742.

Fig. 1 FTIR spectra obtained after (a) deposition of the H₂O fragments and (b) codeposition of the H₂O fragments with CO at 40 K. Vertical lines are provided for guidance.

Fig. 3 Variations in CO₂ column density with experimental time at 40, 50, and 60 K.

Fig. 2 FTIR spectra obtained after codeposition of the H_2O fragments with CO: (a) 4000–700 cm^{-1} and (b) 2400–2300 cm^{-1} . From top to the bottom: 40, 50, and 60 K. For peak assignments, refer to Fig. 1.