Title	Highly efficient enzymatic synthesis of 3'-deoxyapionucleic acid (apioNA) having the four natural nucleobases
Author(s)	Kataoka, Mayumi; Kouda, Yasuo; Sato, Kousuke; Minakawa, Noriaki; Matsuda, Akira
Citation	Chemical Communications, 47(30), 8700-8702 https://doi.org/10.1039/c1cc12980e
Issue Date	2011-08-14
Doc URL	http://hdl.handle.net/2115/49422
Rights	Chem. Commun., 2011, 47, 8700-8702 - Reproduced by permission of The Royal Society of Chemistry (RSC)
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	CC47-30_8700-8702.pdf

[Instructions for use](#)

Cite this: DOI: 10.1039/c0xx00000x

www.rsc.org/xxxxxx

ARTICLE TYPE

Highly efficient enzymatic synthesis of 3'-deoxyapionucleic acid (apioNA) having the four natural nucleobases

Mayumi Kataoka, Yasuo Kouda, Kousuke Sato, Noriaki Minakawa and Akira Matsuda*

Received (in XXX, XXX) Xth XXXXXXXXXX 20XX, Accepted Xth XXXXXXXXXX 20XX

DOI: 10.1039/b000000x

The synthesis of the 3'-deoxyapionucleoside 3''-triphosphates (apioNTPs) having the four natural nucleobases and their enzymatic incorporation into a DNA-DNA primer-template have been tried. Terminator DNA polymerase was shown to incorporate these apioNTPs effectively giving 43mer DNA-apioNA chimera.

Since nucleic acid aptamers are relatively easily isolated from iterative rounds of selection for binding to various targets with high affinities and selectivities, they are expected to be not only useful therapeutic agents but also biological tools.¹ Therapeutic aptamers can act as nucleic acid antibodies against protein targets;² they can also transport oligonucleotide therapeutic agents into target cells through receptor-mediated endocytosis.³ However, because the unmodified oligonucleotides selected are spontaneously degraded before reaching the target, time-consuming post-selection modifications for stabilizing them against nucleases abundant in biological fluids are usually required.⁴ Although isolation of fully-modified aptamers during the selection process is desired, it is nevertheless difficult to achieve because of the high substrate specificity of DNA/RNA polymerases required to incorporate modified nucleoside triphosphates, which are necessary to impart nuclease-resistance after incorporation into the oligonucleotides.⁵ Therefore, the development of a system that incorporates nuclease-resistant chemically modified nucleoside triphosphates having four natural nucleobases still remains a challenge.

Recently, α -L-threofuranosyl nucleoside 3'-triphosphates (tNTPs) have been incorporated into a DNA-DNA primer-template by DNA polymerases, such as Terminator DNA polymerase, to afford DNA-TNA [α -L-threose-(3'→2') nucleic acid] chimeras up to a 75mer (25mer DNA + 50mer TNA) with >20% yield within 24 h, although the chemical structures of tNTPs are rather different from those of the natural 2'-deoxyribonucleoside 5'-triphosphates (dNTPs).⁶ Other sugar-modified nucleoside triphosphates such as GNA⁷, FNA⁸, LNA/BN⁹ and HNA¹⁰, were synthesized and their

Figure 1. Structures of DNA, TNA, INA, apioNA (A) and apioNTP (B).

incorporation into DNA by DNA polymerases was also investigated. These results have inspired us to further investigate other sugar-modified nucleoside triphosphates, which could be incorporated into the desired length of oligonucleotides, since oligonucleotides containing these sugar-modified nucleosides would be expected to be nuclease-resistant. Although TNAs are linked together through their 3'- and 2'-hydroxyl groups by a phosphodiester bond and their repeating unit consists of a backbone (5-bonds) one atom shorter than the natural DNA phosphodiester backbone (6-bonds) (Figure 1), tNTPs acted as substrates of DNA polymerases, but not well enough to be able to isolate aptamers. Therefore, we designed two new candidates that could be inserted and elongated much more efficiently by DNA polymerases. These are 2'-deoxy-2'-isorebionucleoside 5'-triphosphates (iNTPs) and 3'-deoxyapionucleoside 3''-triphosphates (apioNTPs), both of which have a 6-bond phosphodiester backbone similar to DNA although they are regioisomers. We and others have examined the incorporation of iNTPs into DNA by DNA polymerases.¹¹ However, the shape of the iNTPs and local conformational changes of the duplex containing 2'-deoxy-2'-isorebionucleosides disrupted the elongation reaction to give the full length DNA-INA product. Therefore, in this paper, we examined the enzymatic insertion and elongation reactions of apioNTPs, which are one-carbon longer homologues of the tNTPs at the 3' position.

The synthesis of the 3'-deoxyapionucleosides having the four natural nucleobases was carried out using the previous methods¹² with a slight modification, and their 3''-triphosphates (apioNTPs) were prepared (see supporting

Faculty of Pharmaceutical Sciences, Hokkaido University, Sapporo, Hokkaido 060-0812, Japan. E-mail: matuda@pharm.hokudai.ac.jp; Fax: (+81) 11-706-4980; Tel: (+81) 11-706-3228

Figure 2. A) Sequence of the primer-template complex and the apioNA elongation product. Elongated apioNA is shown in bold letters. B) Primer extension experiments: The primer was 5'-FAM labeled and annealed with the DNA template. Polymerization reactions were performed with 0.8 μ M primer-template and 200 μ M all four apioNTPs by incubating for 1 h with 1.0 μ L of DNA polymerase [Klenow (exo-) 0.9 U/ μ L, exTaq 0.5 U/ μ L, Tfl 0.5 U/ μ L, Pfu 0.25 U/ μ L, KOD Dash 0.25 U/ μ L, Vent (exo-) 0.2 U/ μ L, Deep Vent (exo-) 0.2 U/ μ L, Therminator 0.2 U/ μ L, and Therminator II 0.2 U/ μ L] under conditions optimal for each enzyme.

information). In the presence of all four apioNTPs, the primer extension reaction of a 5'-FAM labeled DNA primer (20mer) and a DNA template (27mer) was examined using 9 different DNA polymerases (Figure 2). As shown in Figure 2B, most of the DNA polymerases were unable to elongate the primer with the apioNTPs, but Therminator and Therminator II DNA polymerases incorporated three or four apioNTPs (lanes 10 and 11). Mn^{2+} is known to relax the substrate specificity of many DNA polymerases, possibly by allowing it to enhance its binding to the β - and γ -phosphates of the dNTPs.¹³ Addition of Mn^{2+} (1.25 mM) to the enzyme reaction mixture was found to show a dramatic effect in the elongation reaction by the polymerase (Figure 2C). Klenow (exo-), Pfu, KOD Dash, Vent (exo-), Deep Vent (exo-) and Therminator (lanes 3, 6-10) gave longer elongation product than Mn (-) reactions (Figure 2B), and especially Therminator revealed full length – 1 product as a major product. Therefore, reaction conditions of Therminator were optimized relative to the reaction temperatures and the enzyme concentrations (Figures 3 and S1). The reaction temperature sometimes influences the primer extension reactions for modified triphosphates.^{14,6d} Therefore, we investigated the reaction under lower temperatures than those optimized in the presence of Mn^{2+} . Interestingly full length bands were detected when the reactions were performed at 64 and 54 $^{\circ}$ C (lanes 4 and 5, respectively), but not at 74 $^{\circ}$ C. At even lower temperatures, e.g., 44 and 34 $^{\circ}$ C, the elongation reaction was accelerated to

afford the full length product as essentially one band (lanes 6 and 7). The enzyme concentration was also optimized in Figure S1, the higher concentration of Therminator DNA polymerase (0.2 U/ μ L; 1.7 μ M) was needed for efficient elongation of apioNA.

Figure 3. Primer extension reactions with various reaction temperatures. The reactions were performed with same reaction conditions in general method (see supporting information) except for reaction temperature; 74, 64, 54, 44, and 34 $^{\circ}$ C (lanes 3-7). Lane 1; primer, Lane 2; control.

Since the mobility of the full length bands was slightly different from that of its DNA counterpart on their denaturing polyacrylamide gel when the reactions were performed using dNTPs or apioNTPs (Figure 3), the full length band (27mer) from the apioNTPs was confirmed by MALDI-TOF mass spectroscopy at m/z 8401.8 (Figure S2). Consequently, we found the optimized primer extension reaction conditions, which were 0.8 μ M primer-template duplex and 1.7 μ M (0.2 U/ μ L) Therminator DNA polymerase in the ThermoPol buffer containing 1.25 mM $MnCl_2$ at 44 $^{\circ}$ C for 1 h.

In order to evaluate the substrate ability of dNTP and apioNTP quantitatively, we determined the kinetic parameters (K_m = the Michaelis constant, V_{max} = the maximum rate of the enzyme reaction and V_{max}/K_m = the insertion efficiency) of every triphosphate at various concentrations in the presence or absence of Mn^{2+} , and the data are shown in Table 1. In the absence of Mn^{2+} , the relative efficiency of incorporation of dTTP into the primer was about twice that of the apioTTP (100 vs 46). The difference is presumably a reflection of their K_m values (0.022 vs 0.058 μ M), but not that of their V_{max} values (7.4 vs 9.4 % $\cdot$ min⁻¹). Other triphosphates showed a similar effect with thymine nucleotides, namely relatively larger differences in the K_m values (1.5-3.5 times) and smaller differences in the V_{max} values (1.2-1.3 times) between dNTPs and apioNTPs, respectively. These quantitative analyses revealed that Therminator incorporated the apioNTPs into the DNA/DNA primer/template duplex with 2-5 times lower relative efficiency than the dNTPs. The effect of the addition of Mn^{2+} was next investigated by using single nucleotide insertion reactions. As can be seen from Table 1, the polymerase incorporated apioTTP with almost the same efficiency in the presence or absence of Mn^{2+} (relative efficiency; 48 vs 46), and also with similar K_m (0.086 vs 0.058 μ M) and V_{max} (14.0 vs 9.4 % $\cdot$ min⁻¹) values. Other nucleotides showed similar results with little difference in K_m (0.4-1.6 times) and V_{max} (0.4-1.5 times) values. Therefore, the addition of Mn^{2+} to the enzyme reaction mixture did not show any dramatic effects on the single nucleotide insertion reaction of the apioNTP.

It is obvious that the elongation reactions of the apioNTPs

Table 1. The kinetic parameters of dNTPs and apioNTPs in the absence

NTP (MnCl ₂)	K _m (μM)	V _{max} (%·min ⁻¹)	V _{max} /K _m (%·min ⁻¹ ·M ⁻¹)	relative efficiency
dTTP (–)	0.022 ± 0.0075	7.4 ± 1.7	3.5 × 10 ⁸	100
apioTTP (–)	0.058 ± 0.025	9.4 ± 1.0	1.6 × 10 ⁸	46
(+)	0.086 ± 0.020	14.0 ± 1.9	1.7 × 10 ⁸	48
dCTP (–)	0.040 ± 0.012	5.9 ± 0.74	1.5 × 10 ⁸	100
apioCTP (–)	0.090 ± 0.020	4.8 ± 1.4	5.4 × 10 ⁷	35
(+)	0.063 ± 0.0033	4.5 ± 1.4	7.1 × 10 ⁷	46
dATP (–)	0.022 ± 0.0078	7.1 ± 0.82	3.4 × 10 ⁸	100
apioATP (–)	0.078 ± 0.0015	5.5 ± 0.54	7.1 × 10 ⁷	21
(+)	0.031 ± 0.007	2.4 ± 0.14	8.0 × 10 ⁷	24
dGTP (–)	0.036 ± 0.012	8.3 ± 1.6	2.4 × 10 ⁸	100
apioGTP (–)	0.047 ± 0.022	6.1 ± 0.92	1.6 × 10 ⁸	69
(+)	0.075 ± 0.046	5.8 ± 0.71	1.3 × 10 ⁸	56

or presence of Mn²⁺ with Terminator DNA polymerase^[a]

[a] Assay conditions: Reactions were initiated by adding 1 [L of a dNTP or apioNTP solution to 9 [L of the reaction mixture containing a 0.8 [M 5'-FAM labeled primer-template complex, 42.5 nM (0.005 U/[L) Terminator DNA polymerase in the ThermoPol buffer (see supporting information) in the absence or presence of 1.25 mM MnCl₂ incubating for 3 min at 74 °C, and quenching with the addition of 10 [L loading buffer.

by Terminator were evidently accelerated in the presence of Mn²⁺ as shown in Figure 2. Therefore, we hypothesized that Mn²⁺ influenced the elongation reaction after incorporation of several apionucleotides into the DNA primer. To examine this hypothesis, a chimera oligonucleotide primer (5'-FAM 20mer DNA + 4mer apioNA) was synthesized by the usual phosphoramidite method except for the coupling time of the apioNA (see supporting information). Single nucleotide insertion reactions using apioATP were then performed using a steady-state method in the presence or absence of Mn²⁺. Although V_{max} shows almost the same values (3.3 vs 3.0 %·min⁻¹), the K_m values were different by one order of magnitude with or without addition of Mn²⁺ (0.044 vs 0.403 μM) as shown in Table 2. Consequently, the relative efficiency increased over 10 fold when Mn²⁺ was added to the reaction mixture.

Table 2. The kinetic parameters of apioATP in the absence or presence of Mn²⁺ with Terminator DNA polymerase^[a]

NTP (MnCl ₂)	K _m (μM)	V _{max} (%·min ⁻¹)	V _{max} /K _m (%·min ⁻¹ ·M ⁻¹)	relative efficiency
apioATP (–)	0.403 ± 0.014	3.0 ± 0.11	7.4 × 10 ⁶	100
(+)	0.044 ± 0.007	3.3 ± 0.16	7.5 × 10 ⁷	1014

[a] Assay conditions: The reactions were performed with the same reaction conditions described in Table 1 except for 1.7 [M (0.2 U/[L) Terminator DNA polymerase.

Finally, a longer elongation polymerase reaction by Terminator was carried out using a 21mer DNA primer and a 43mer DNA template. In this case, the reactions under lower temperature are also effective for longer elongation in the presence of Mn²⁺ (Figure S3). Consequently, a longer elongation polymerase reaction (22mer apioNA elongation) at

lower temperatures, such as 44 and 34 °C, would appear to be more efficient for full length product formation by Terminator DNA polymerase containing 1.25 mM MnCl₂ within 1 h.

In summary, we have synthesized apioNTPs, the one-carbon longer homologue of the α-L-threofuranosyl nucleoside 3'-triphosphates (tNTPs) at the 3'-position having the four natural bases (T, C, A and G) and also identified Terminator DNA polymerase as a polymerase that is capable of highly efficient enzymatic apioNA synthesis. Moreover we found that Mn²⁺ and lower temperatures are important for longer elongation of apioNA but are not required for a single nucleotide insertion reaction into a DNA-DNA duplex (Figure 3 and Table 1). A longer elongation polymerase reaction (22mer apioNA elongation) has also been achieved at 34 or 44 °C within 1 h. Therefore, the apioNTPs are some of the most effective sugar-modified triphosphates for enzymatic polymerization by DNA polymerase. Studies of apioNA templated apioNA synthesis by Terminator DNA polymerase are now under investigation.

Notes and references

† Electronic Supplementary Information (ESI) available: Experimental details of synthesis, ¹H and ³¹P NMR of apioNTPs, ODN synthesis, elongation reactions, single nucleotide insertion reactions, and MALDI-TOF mass spectrum of full length 27mer elongated product. See DOI: 10.1039/b000000x/

- D. H. J. Bunka, P. G. Stockley, *Nat. Rev. Microbiol.* 2006, **4**, 588.
- (a) A. D. Keefe, S. Pai, A. Ellington, *Nat. Rev. Drug Discovery* 2010, **9**, 537; (b) P. R. Bouchard, R. M. Hutabarat, K. M. Thompson, *Annu. Rev. Pharmacol. Toxicol.* 2010, **50**, 237.
- (a) N. S. Que-Gewirth, B. A. Sullenger, *Gene Ther.* 2007, **14**, 283; (b) A. C. Yan, M. Levy, *RNA Biol.* 2009, **6**, 316.
- A. D. Keefe, S. T. Cload, *Curr. Opin. Chem. Biol.* 2008, **12**, 448.
- (a) P. E. Burmeister, S. D. Lewis, R. F. Silva, J. R. Preiss, L. R. Horwitz, P. S. Pendergrast, T. G. McCauley, J. C. Kurz, D. M. Epstein, C. Wilson, A. D. Keef, *Chem. Biol.* 2005, **12**, 25; (b) N. Minakawa, M. Sanji, Y. Kato, A. Matsuda, *Bioorg. Med. Chem.* 2008, **16**, 9450.
- (a) J. C. Chaput, J. W. Szostak, *J. Am. Chem. Soc.* 2003, **125**, 9274; (b) A. Horhota, K. Zou, J. K. Ichida, B. Yu, L. W. McLaughlin, J. W. Szostak, J. C. Chaput, *J. Am. Chem. Soc.* 2005, **127**, 7427; (c) V. Kempeneers, K. Vastmans, J. Rozenski, P. Herdewijn, *Nucleic Acids Res.* 2003, **31**, 6221; (d) J. K. Ichida, K. Zou, A. Horhota, B. Yu, L. W. McLaughlin, J. W. Szostak, *J. Am. Chem. Soc.* 2005, **127**, 2802.
- (a) A. T. Horhota, J. W. Szostak, L. W. McLaughlin, *Org. Lett.* 2006, **8**, 5345; (b) J. J. Chen, C.-H. Tsai, X. Cai, A. T. Horhota, L. W. McLaughlin, J. W. Szostak, *PLoS ONE* 2009, **4**, 1.
- B. D. Heuberger, C. Switzer, *J. Am. Chem. Soc.* 2008, **130**, 412.
- (a) R. N. Veedu, B. Vester, J. Wengel, *ChemBioChem* 2007, **8**, 490; (b) R. N. Veedu, B. Vester, J. Wengel, *J. Am. Chem. Soc.* 2008, **130**, 8124; (c) R. N. Veedu, B. Vester, J. Wengel, *Nucleosides Nucleotides Nucleic Acids* 2007, **26**, 1207.
- (a) K. Vastmans, S. Pochet, A. Peys, L. Kerremans, A. van Aerschot, C. Hendrix, P. Marlière, P. Herdewijn, *Biochemistry* 2000, **39**, 12757; (b) M. Renders, M. Abramov, M. Froeyen, P. Herdewijn, *Chem. Eur. J.* 2009, **15**, 5463.
- (a) C. Jiang, B. Li, Z. Guan, Z. Yang, L. Zhang, L. Zhang, *Bioorg. Med. Chem.* 2007, **15**, 3019; (b) T. Ogino, K. Sato, A. Matsuda, *ChemBioChem* 2010, **11**, 2597.
- D. Z. Jin, S. H. Kwon, H. R. Moon, P. Gunaga, H. O. Kim, D.-K. Kim, M. W. Chun, L. S. Jeong, *Bioorg. Med. Chem.* 2004, **12**, 1101.
- (a) R. C. Cadwell, G. F. Joyce, *PCR Methods Appl.* 1992, **2**, 28; (b) S. Tabor, C. C. Richardson, *Proc. Natl. Acad. Sci. USA* 1989, **86**, 4076.

-
- 14 J. K. Ichida, A. Horhota, K. Zou, L. W. MaLaughlin, J. W. Szostak, *Nucleic Acids Res.* 2005, **33**, 5219.