Title	Effect of liquid whey feeding on fecal microbiota of mature and growing pigs
Author(s)	Kobayashi, Yasuo; Itoh, Aya; Miyawaki, Kanae; Koike, Satoshi; Iwabuchi, Osamu; Iimura, Yuji; Kobashi, Yuri; Kawashima, Tomoyuki; Wakamatsu, Junichi; Hattori, Akihito; Murakami, Hiroshi; Morimatsu, Fumiki; Nakaebisu, Takashi; Hishinuma, Takeshi
Citation	Animal Science Journal, 82(4), 607-615 https://doi.org/10.1111/j.1740-0929.2011.00876.x
Issue Date	2011-08
Doc URL	http://hdl.handle.net/2115/49681
Rights	The definitive version is available at wileyonlinelibrary.com
Type	article (author version)
File Information	ASJ82-4_607-615.pdf

[Instructions for use](#)

ORIGINAL ARTICLE

Title: Effect of liquid whey feeding on fecal microbiota of mature and growing pigs

Authors: Yasuo KOBAYASHI¹, Aya ITOH¹, Kanae MIYAWAKI¹, Satoshi KOIKE¹, Osamu IWABUCHI², Yuji IIMURA², Yuri KOBASHI³, Tomoyuki KAWASHIMA³, Junichi WAKAMATSU¹, Akihito HATTORI¹, Hiroshi MURAKAMI⁴, Fumiki MORIMATSU⁴, Takashi NAKAEBISU⁵ and Takeshi HISHINUMA⁵

Institute, address, country:

¹Research Faculty of Agriculture, Hokkaido University, Kita, Sapporo 060-8589, Japan

²Interfarm Co., Ltd., Abashiri, 099-3008, Japan

³National Institute of Livestock and Grassland Science, Tsukuba 305-0901, Japan

⁴Nippon Meat Packers, Inc., R&D Center, Tsukuba 300-2648, Japan

⁵Japan Livestock Industry Association, Minato, Tokyo 105-0001, Japan

Running title: FECAL MICROBIOTA OF WHEY-FED PIGS

Correspondence: Yasuo Kobayashi, Research Faculty of Agriculture, Hokkaido University, Kita, Sapporo 060-8589, Japan.

Phone: +81-11-706-2476; Facsimile: +81-11-706-2476; Email: kyas@anim.agr.hokudai.ac.jp

ABSTRACT

The effect of liquid whey feeding on fecal bacteria and their metabolites was assessed in five pregnant sows and 66 growing pigs. Sows were fed a control diet for 4 weeks (control period) followed by the same diet but with whey feeding (5 L/day/pig) for 4 weeks (whey period). One group of growing pigs was given 267 L of whey per pig (whey group), while the other group was not (control group). In both cases, liquid whey was given separately from control diet. Sows in the whey period had feces showing lower pH, lower ammonia concentration, and larger population sizes of total bacteria, lactobacilli, and bifidobacteria. The bacterial gene library analysis indicated that *Mitsuokella* and *Megasphaera* were more frequently detected, while *Clostridium disporicum* were detected less frequently in the whey period. Feces from whey-fed growing pigs showed lower pH than that from control pigs in the early stage of growing. Also, larger populations of total bacteria, lactobacilli, and bifidobacteria were recorded in the whey group. From the analysis of bacterial gene library, the detection frequency of *Lactobacillus reuteri* tended to be higher in the whey group. These results indicate that whey feeding influences the hindgut microbiota of pigs, possibly leading to a fermentation shift that is favorable for animal health.

Key words: fermentation, hindgut bacteria, liquid whey, pig, 16S rRNA gene.

INTRODUCTION

Liquid whey is an attractive feed resource for domestic animals, especially pigs, because liquid feeding is being introduced worldwide with the advantage that feed ingredients are given directly without additional costly processes such as drying. Liquid whey could be considered as a prebiotic feed as well as a protein source because it contains 5% lactose, which is utilized by beneficial intestinal bacteria such as lactobacilli and bifidobacteria (Coppa et al. 2006). Growth promotion of these lactate-producing bacteria in the pig intestine could improve intestinal health through lactate production and the resultant reduction of intestinal pH, which indirectly prevents the growth of detrimental bacteria including pathogenic *Salmonella* and *Escherichia coli* (Wells et al. 2005). Also, the intestinal immune system could be promoted by selective proliferation of beneficial bacteria (Mitsuoka 2002). Therefore, evaluation of the potency of liquid whey as a prebiotic feed from the viewpoint of intestinal microbiology would be important. In addition, because liquid whey contains lactic acid bacteria that are blended as cheese starters by manufacturers, these bacteria also need to be assessed as possible probiotics for pigs. Furthermore, liquid whey that is rich in proteins, vitamins, and minerals is considered to be a nutritionally favorable material as a feed ingredient (Leibbrandt & Benevenga 1991).

Although liquid whey feeding is popular on swine farms and contributes to dairy byproduct consumption all over the world, its merits have not been fully scientifically evaluated. The usefulness of liquid whey feeding in terms of the growth promotion of pigs was initially indicated by Maswaure and Mandisodza (1995). We have also reported the improved growth performance of pigs as the result of liquid whey feeding

on a commercial farm and suggested the importance of clarifying some factors involved in this improvement (Kobashi et al. 2009). Intestinal microbiota is a primary factor in modulating pig nutrition and health. Therefore, the aim of the present study was to determine how liquid whey feeding influences the intestinal microbiota of pigs.

MATERIALS AND METHODS

Animals, feeding, and sampling

All the experimental pigs were kept in a commercial farm (Inter Farm co. Ltd, Abashiri, Japan) according to the farm's manual for animal management. Samplings were carried out, following Act on Welfare and Management of Animals (2005) and Guidelines for Animal Experiment, Hokkaido University (2007).

Five sows (LWD) in 7th wk of pregnancy were individually housed in stalls and fed an antibiotic-free commercial formula feed with free access to water. The feed (Pigfighter 73; Kumiai Feed Manufacturing, Ohta, Japan) consisted of 62% cereals, 17% oil meals, 14% bran, and 7% other ingredients and contained 15% crude protein (CP) and 73% total digestible nutrients (TDN). Sows were assigned to the formula feed for 4 weeks (control period) followed by the same diet but with additional feeding of liquid whey at the level of 5 L /day/pig for 4 weeks (whey period). Liquid whey was fed separately from the formula feed, using a wet feeder. Fresh feces samples were taken from the rectum by grab sampling at the end of the control period and at the 14th and 28th days of the whey period. Feces samples were immediately frozen at -30°C, shipped to the laboratory, and stored at -80°C until analysis.

Sixty-six piglets (LWD) from nine litters were used. The animals, their feeding, and

their management were the same as described by Kobashi et al. (2009). In brief, the animals were weaned at 28 days of age (7.8 ± 1.3 kg in body weight), divided into two groups to erase hereditary and sex interference, and given a commercial formula feed (*see below*) with and without additional supply of liquid whey during the period from 29 to 173 days of age (whey group vs. control group). Whey was given separately from the formula feed as described for sows, but its supply was gradually increased as pigs grew, *i.e.* 0.79, 1.5 and 2.0 L/day/pig were fed during the periods of 29-65, 66-104 and 105-173 days of age, respectively. The total supply of liquid whey in the whey group was 267 L/pig. The formula feed for pigs was the starter A (Uruoi Pro; 22.5% CP and 89% TDN), followed by the starter B (Manpuku IF; 18.5% CP and 81.0% TDN), the grower diet (Grower Phase; 16% CP and 79.5% TDN) and then the finisher diet (Finisher Wonder Rich 78IF; 12.5% CP and 78.0% TDN). All the formula feeds were purchased from Nippon Formula Feed, Tomakomai, Japan. Morantel citrate, colistin sulphate, and avilamycin were blended into the starter feed that was given to all pigs until 65 days of age, while no such antibiotics were offered thereafter. At 43, 65, 104, and 139 days of age, fresh feces samples from six pigs randomly selected from each group were taken and stored as described above for sows. This was due to difficulty in continuous sampling from specific individuals over the experimental period.

The experimental whey just before given to the animals was sampled at the time for fecal sampling (twice in whey period for sows and four times for growing pigs) and subjected to microbial analysis.

Microbial analysis

DNA extraction

Bacterial DNA was extracted from feces and whey samples by the repeated bead beating plus column (RBB+C) method (Yu & Morrison 2004). In brief, thawed feces was weighed and placed in a sterilized 2-mL screw-capped tube containing 0.4 g of glass beads (diameter: 100 to 500 μ m; Sigma Chemicals, St. Louis, MO, USA). The feces (0.25 g) was mixed with 1.0 mL of lysis buffer (500 mmol/L NaCl, 50 mmol/L Tris-HCl [pH 8.0], 50 mmol/L EDTA, and 4% sodium dodecyl sulfate). DNA was extracted by shaking the tube horizontally. The extracted DNA was purified by QIAamp columns (from the QIAamp DNA Stool Mini Kit; Qiagen, Hilden, Germany). The DNA was spectrophotometrically quantified by a plate reader (ARVO MX / Light; PerkinElmer, Waltham, MA, USA).

Real-time polymerase chain reaction

For polymerase chain reaction (PCR) quantification of lactobacilli, the genus-specific primers Lac1f (5'-AGC AGT AGG GAA TCT TCC A-3') (Walter et al. 2001) and Lab677r (5'-CAC CGC TAC ACA TGG AG-3') (Heilig et al. 2002) were used. For bifidobacteria, the genus-specific primers Bif164f (5'-GGG TGG TAA TGC CGG ATG-3') and Bif662r (5'-CCA CCG TTA CAC CGG GAA-3') (Kok et al. 1996) were employed. Standards for real-time PCR were constructed by amplifying the targeted regions of 16S rDNA of *Lactobacillus acidophilus* CH2 and *Bifidobacterium breve* ATCC15700, followed by cloning with *E. coli*. Both assays were validated for accuracy and reproducibility as described in Koike et al. (2007). Total bacteria were quantified according to the method of Koike et al. (2007).

Denaturing gradient gel electrophoresis

DNA (500 ng) isolated from feces of sows was used as a template to amplify the V3 region of bacterial 16S rDNA by PCR with primers 2 and 3 as described by Muyzer et

al. (1993). In brief, touchdown PCR was performed using the rTaq DNA polymerase system (TOYOBO, Osaka, Japan) and the PCR conditions described by Muyzer et al. (1993). The integrity of the PCR products was visually checked by electrophoresis on a 2% agarose gel. The PCR products and the denaturing gradient gel electrophoresis (DGGE) marker (NipponGene, Tokyo, Japan) were separated by the Decode Universal Mutation Detection System (Bio-Rad Laboratories, Hercules, CA, USA) using a 6% (v/v) polyacrylamide DGGE gel with a 30% to 70% gradient of denaturant. Electrophoresis was performed at 90 V for 16 h in TAE buffer (pH 7.4) at a controlled temperature of 60°C. The gel was stained with SYBR green or silver nitrate and the DGGE gel image was scanned using a LumiVision PRO 400EX image analyzer (Aisin Seiki Co., Ltd., Kariya, Japan). The DGGE band profiles obtained were analyzed by clustering via the unweighted pair group method with mathematical averages (UPGMA) using BioNumerics software (Applied Maths, Inc., Austin, TX, USA).

Bacterial 16S rDNA library

Primers 27F (5'-AGA GTT TGA TCM TGG CTC AG-3') and 1525R (5'-AAG GAG GTG WTC CAR CC -3') were used for amplifying bacterial 16S rDNA to construct its library. Templates used for library construction were the mixture of DNA from whey samples, feces of 5 sows at the end of control and whey periods, and also feces of 6 growing pigs from control and whey groups at 43 days of age. Polymerase chain reaction conditions were as follows: initial denaturation at 94°C for 5 min, then 20 amplification cycles of denaturation at 94°C for 0.5 min, annealing at 58°C for 0.5 min, and extension at 72°C for 1.5 min. A final extension was performed at 72°C for 7 min. The PCR product was purified by agarose electrophoresis and isolated using a QIAEX II Gel Extract Kit (Qiagen, Hilden, Germany). The purified product was then ligated

with pGEM T Easy Vector (Promega, San Louis Obispo, CA, USA) and introduced into *E. coli* JM109. White colonies developed on LB plates with ampicillin, isopropyl beta-D-1-thiogalactopyranoside and X-gal were employed for plasmid isolation using a QIAprep Spin Miniprep Kit (Qiagen) for the following sequencing analysis (Dragon Genomics Center, Takara Bio Inc., Yokkaichi, Japan).

The sequences were compared to those available in the GenBank database using the DDBJ BLAST program (<http://www.ddbj.nig.ac.jp/Welcome-e.html>). The 16S rDNA clone libraries from the control and whey periods (groups) were compared using the LIBSHUFF computer program (Singleton et al. 2001). The DNADIST program of PHYLIP using the Juke-Cantor model was used to generate the distance matrix analyzed by LIBSHUFF. Sequence identity of 97% was used as a criterion to define whether each sequence belonged to a certain species. The sequences (ca. 1500 bp) obtained were deposited in the DDBJ nucleotide sequence database under the accession numbers AB506124 through AB506454.

Chemical analysis

Each thawed feces sample (0.1 g) was dispersed into sterilized saline (0.5 mL). After measuring the pH value with an electrode (Horiba, Kyoto, Japan), this solution was centrifuged at $1,000\times g$ for 5 min. The supernatant was used for analyzing short chain fatty acids (SCFA) by gas-liquid chromatography (Suto 1970), lactate (L- and D-lactate) with a lactic acid assay kit (Megazyme, Wicklow, Ireland), and ammonia nitrogen by a phenol-hypochlorite method (Weatherburn et al. 1967).

Statistical analysis

The data on bacterial abundance and chemical parameters of feces from sows were subjected to one way variance analysis of SPSS software (Version 16.0 J, Tokyo), where period was a fixed effect. When significance was detected, multiple comparisons were made by Bonferroni's method. The data from growing pigs at each time point were compared between control and whey groups by Student's t-test. Statistical significance was defined at $P \leq 0.05$.

RESULTS AND DISCUSSION

Liquid whey as a tested feed

The experimental whey just before given to pigs showed variations in pH (3.7-4.9) and bacterial abundance (lactobacilli, $\times 10^{7-8}$ 16S rDNA copy/mL; bifidobacteria, $\times 10^{5-6}$ 16S rDNA copy /mL) due to season and storage duration. **Table 1** shows the bacteria detected in the 16S rDNA mini-library. They belonged to Lactobacillales (50%), Proteobacteria (36%), and Rhodospirillales (14%). Nineteen sequences showing more than 97% identity were identified as those of 8 known species. Of these known bacteria, some species of lactate-producing bacteria such as *Lactobacillus helveticus*, *Streptococcus thermophilus*, and *Lactococcus lactis* were considered as starter cultures blended by a cheese manufacturer. Other bacteria were considered as later contaminants during transportation and/or storage. Because all of these species were not detected from the feces of either mature or growing pigs in the present study (*see below*), they may be regarded as transient microorganisms in the gastro-intestinal tract of the experimental pigs. However, there might be a possibility that these organisms act as probiotics to stimulate indigenous bacteria without notable proliferation of themselves

[Table 1]

(Ohashi et al. 2007). The possibility is to be further described below.

[Figure 1]

Effects on mature pigs

Floral changes in sows' feces revealed by PCR/DGGE analysis are shown in **Figure 1**. Clustering of banding patterns divided fecal community structures into three groups: those that were from sows in the control period and those from the second and fourth weeks of the whey period. In particular, samples clustered together from the fourth week of the whey period were located apart from the other samples, suggesting that liquid whey feeding gradually alters fecal microbiota and significant alteration occurs within 4 weeks of whey feeding.

This floral alteration was again evidenced by the analysis of bacterial 16S rDNA libraries as shown in **Table 2**. The bacterial community retrieved by the library analysis was significantly different between the control and whey periods, shown by the LIBSHUFF test ($P < 0.002$). Although Firmicutes occupied both libraries, their constituents were different. While the most dominant constituent in the control period was clostridial cluster IV (26.4%), it was replaced by cluster IX (29.5%) in the whey period.

[Table 2]

[Table 3]

[Table 4]

Although the numbers of known bacterial species (sharing more than 97% 16S rDNA sequence identity) were 24 in the control period and 21 in the whey period, the detection frequency of individual species was different between the control and whey periods (**Table 3**). For example, no sequence belonging to *Mitsuokella* or *Megasphaera* of cluster IX was detected in the control period, but *Mitsuokella* was detected at 7.5% frequency and *Megasphaera* was detected at 3.0% frequency in the whey period. *Mitsuokella jalaludinii* has been considered as a useful bacterium for producing lactate

and degrading phytate in the porcine intestine (Lan et al. 2002). *Megasphaera elsedenii*, a butyrate producer, is also useful for promoting intestinal health (Hashizume et al. 2003), because butyrate stimulates epithelial proliferation in pigs (Cummings et al. 2004). Meanwhile, *Clostridium disporicum* belonging to cluster I was decreased from 1.2% to nil with liquid whey feeding, suggesting the usefulness of liquid whey to expel this opportunistic bacterium (Woo et al. 2005) from intestinal tracts. However, these bacteria need to be examined in more quantitative analyses.

Quantitative PCR revealed that liquid whey feeding significantly increased ($P<0.05$) the numbers of total bacteria, lactobacilli, and bifidobacteria in 2 to 4 weeks compared with the control period, even though the increase in the percentages of lactobacilli and bifidobacteria (relative abundance of each group in 16SrDNA copy number) was not significant (**Table 4**). These results indicate that liquid whey could stimulate the growth of intestinal bacteria and that the stimulation is not absolutely selective for lactic acid bacteria. In fact, the above library analysis does not support the selective growth of lactobacilli and bifidobacteria by liquid whey feeding, because no bifidobacteria were detected and no lactobacilli showed an increase in detection frequency with whey feeding (**Table 3**). Wells et al. (2005) also pointed out that growth stimulation by lactose-rich skim milk is not highly selective for hindgut bacteria in pigs.

Chemical parameters, shown in **Table 4**, did not reflect the above-mentioned alteration of microbiota. While fecal pH was significantly reduced with 2 weeks of whey feeding, the total SCFA level was unchanged and the lactate level decreased. The SCFA proportion did not significantly differ among the samples. Ammonia tended to be consistently but not significantly lower in the whey period. Overall, the variations in chemical parameters did not correlate with those of microbial parameters, which made

speculation difficult. However, one possible explanation for reduced pH without increase of SCFA and lactate may be the accumulation of succinate. Although succinate is rarely detected in the hindgut of pigs under normal condition, it would be accumulated if the number of succinate-utilizing bacteria is limited and succinate-producing bacteria are abundantly present. Bacteria increased with whey feeding in the present study may be dominated by active producers of succinate including lactobacilli, some of which are considered as causative agents of succinate accumulation in pigs (Tsukahara & Ushida 2002). However, the possibility remains to be evidenced, because succinate production and the related bacteria were not assessed in the present study. The tendency for ammonia level to be lowered with whey feeding may imply the increased cell proliferation of hindgut bacteria which is partly supported by the higher number of bifidobacteria, lactobacilli and total bacteria in feces.

Effects on growing pigs

Bacterial phyla and groups within a phylum detected in the 16S rDNA library for growing pigs are indicated in **Table 5**. All sequences fell into Firmicutes and about half belonged to Lactobacillales (43.7% in the control group and 50.6% in the whey group). Two libraries from the control and whey groups were not significantly different by the LIBSHUFF test ($P=0.747$ and 0.806).

Table 6 shows the number of known bacterial species identified in the bacterial 16S rDNA library. Thirty-eight sequences were identified as known species in the control group and 49 sequences were identified in the whey group. Of these, *Lactobacillus reuteri* showed a higher detection frequency in whey-fed pigs (6.5% in the control group vs. 16.7% in the whey group). This bacterium is known to improve

[Table 5]

[Table 6]

[Table 7]

feed conversion by expelling the parasitic *Cryptosporidium parvum* from pig intestine (Casas & Dobrogosz, 1997). Other beneficial actions of *L. reuteri*, including the production of reuterin, a bacteriocin, and activation of the immune system (Morita & Masaoka, 2005), could promote host animal health. Because improvements in feed conversion and survivability of whey-fed growing pigs in the present study were previously reported (Kobashi et al. 2009), the higher abundance of *L. reuteri* in whey-fed pigs in the early stage of growing might have contributed to these improvements.

Table 7 shows the changes in bacterial quantity and chemical parameters in the feces of growing pigs taken at four different time points. Quantitative PCR revealed that whey-fed pigs had higher ($P<0.05$) numbers of total bacteria, lactobacilli, and bifidobacteria at 43 days of age in comparison with control pigs, even though the relative percentage of each bacterial group was not significantly different between control and whey-fed pigs. Any other changes in the bacterial parameters at other time points were not significant. Fecal pH showed a lower value ($P<0.05$) at 43 days in whey-fed pigs without alterations in total SCFA and lactate concentration. The results are not straightforward to be discussed, because fecal concentrations of organic acid do not necessarily reflect bacterial numbers and activities. The inconsistent association of pH with acid concentration may be again excused by the possibility of succinate accumulation as done for the data of matured pigs, which remains to be proven. No particular variations were observed for SCFA molar proportion and ammonia in relation to liquid whey feeding throughout the experimental period.

The fact that fecal bacteria responded to liquid whey only in the early stage of growth may suggest that the efficacy of liquid whey feeding depends on the age of pigs,

their original microbiota, and the amount of whey fed. The relative abundance of lactobacilli was much higher at 43 days of age (71.5%) than at 104 days (43.3%) and 139 days (12.7%), implying that the original microbiota at 43 days of age potentiate the effect of liquid whey feeding. Similarly, it is suggested that prebiotics may have a greater impact on fecal bacterial populations in younger infants than in older infants (Nakamura et al. 2009). The amount of whey fed to growing pigs in the present study varied from 0.79 to 2.0 L/days/pig in accordance with animal growth. This was approximately equivalent to 8% of body weight at 43 days and 3% of body weight at 139 days of age. Therefore, the difference in actual whey intake would have differentiated the efficacy of liquid whey.

The flora of growing pigs was apparently different from that of sows, which was supported by the LIBSHUFF test ($P < 0.05$). In fact, except for *Lactobacillus amylovorus*, no species of bacteria was shared between mature sows and growing pigs in the present study. Therefore, it is assumed that liquid whey could promote the growth of specific bacteria that varied between growing pigs (e.g., *L. reuteri*) and sows (e.g., *Mitsuokella* and *Megasphaera*). None of the bacteria detected from the experimental pigs originated from the experimental whey. This is again indicative of a possibility that bacteria included in the whey do not multiply in the intestinal tract of pigs but may exert their actions by interacting with indigenous beneficial bacteria as suggested by Ohashi et al. (2007). They demonstrated an increase of indigenous lactobacilli in the porcine cecum with feeding of a probiotic yoghurt strain of *Lactobacillus* without detecting the strain itself. Therefore, it is not definitive at present whether liquid whey can be a probiotic, or a prebiotic as a lactose carrier, or just a material for delivering other milk-originating nutrients such as proteins, minerals, and vitamins (Leibbrandt & Benevenga 1991;

Maswaure & Mandisodza 1995). Although whey can be all these beneficial materials for pigs, it should be clarified how the benefits are associated with each other. Then, analytical approach is to be argued, *e.g.* unsuitableness of fecal concentration of SCFA as a predictor of production rate has been mentioned (Sakata et al. 2003).

CONCLUSION

Liquid whey feeding altered the fecal microbiota of pigs, particularly mature pigs. Lactobacilli, bifidobacteria, and other beneficial bacterial species showed higher abundances in whey-fed pigs, while one opportunistic species disappeared. Therefore, whey is considered to be useful for promoting animal health and nutrition. Because all speculations made in the present paper are based on the analysis of a limited number of animals and gene sequences, those are to be evidenced by more extensive dataset.

ACKNOWLEDGEMENTS

The authors appreciate the financial support to the present study from the Northern Advancement Center for Science and Technology (the basic research development project 2007-2) and from the Japan Livestock Industry Association (the use development of cheese whey, 2007)

REFERENCES

- Casas IA, Dobrogosz WJ. 1997 *Lactobacillus reuteri* : Overview of a new probiotic for human and animals. *Microecology and Therapy* **26**, 221-231.
- Coppa GV, Zampini L, Galeazzi T, Gabrielli O. 2006. Prebiotics in human milk: a

- review. *Digestive and Liver Disease* **38** (Suppl 2), S291-S294.
- Cummings JH, Rombeau JL, Sakata T. 2004. Physiological and clinical aspects of short-chain fatty acids. p.289. Cambridge University Press, Cambridge, UK.
- Hashizume K, Tsukahara T, Yamada K, Koyama H, Ushida K. 2003. *Megasphaera elsdenii* JCM1772T normalizes hyperlactate production in the large intestine of fructooligosaccharide-fed rats by stimulating butyrate production. *Journal of Nutrition* **133**, 3187-90.
- Heilig HG, Zoetendal EG, Vaughan EE, Marteau P, Akkermans AD, de Vos WM. 2002. Molecular diversity of *Lactobacillus* spp. and other lactic acid bacteria in the human intestine as determined by specific amplification of 16S ribosomal DNA. *Applied and Environmental Microbiology* **68**, 114-123.
- Kobashi Y, Ishiguro T, Wakamatsu J, Okumura T, Takahagi Y, Iwabuchi O, Iinuma Y, Kawashima T, Kobayashi Y, Hattori A, Murakami H, Morimatsu F. 2009. Effects of liquid whey supplement on the productivity of pigs in a commercial pig farm. *Nihon Chikusan Gakkaiho* **80**, 443-450.
- Koike S, Yabuki H, Kobayashi Y. 2007. Validation and application of real-time polymerase chain reaction assays for representative rumen bacteria. *Animal Science Journal* **78**, 135-141.
- Kok RG, de Waal A, Schut F, Welling GW, Weenk G, Hellingwerf KJ. 1996. Specific detection and analysis of a probiotic Bifidobacterium strain in infant feces. *Applied and Environmental Microbiology* **62**, 3668-3672.
- Lan GQ, Abdullah N, Jalaludin S, Ho YW. 2002. Optimization of carbon and nitrogen sources for phytase production by *Mitsuokella jalaludinii*, a new rumen bacterial species. *Letters in Applied Microbiology* **35**, 157-161.

- Leibbrandt VD, Benevenga NJ. 1991. Utilization of liquid whey in feeding swine. In: Miller ER, Ullrey DE (eds), *Swine Nutrition*, pp.559-571. Butterworth-Heinemann. Boston, USA.
- Mansware SM, Mandisodza KT. 1995. An evaluation of the performance of weaner pigs fed diets incorporating fresh sweet liquid whey. *Animal Feed Science and Technology* **54**, 193-201.
- Mitsuoka T. 2002. Prebiotics and intestinal flora. *Bioscience and Microflora* **21**, 3-12.
- Morita H, Masaoka T. 2005. Lactic acid bacteria. In: Itoh K (ed), *Probiotics and Biogenics*, pp.215-226. NTS Co. Ltd., Tokyo..
- Muyzer G, de Waal EC, Uitterlinden AG. 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology* **59**, 695–700.
- Nakamura N, Gaskins HR, Collier CT, Nava GM, Rai D, Petschow B, Russell WM, Harris Mackie RI, Wampler JL, Walker DC. 2009. Molecular ecological analysis of fecal bacterial populations from term infants fed formula supplemented with selected blends of prebiotics. *Applied and Environmental Microbiology* **75**, 1121-1128.
- Ohashi Y, Tokunaga M, Taketomo N, Ushida K. 2007. Stimulation of indigenous lactobacilli by fermented milk prepared with probiotic bacterium, *Lactobacillus delbrueckii* subsp. *bulgaricus* strain 2038, in the pigs. *Journal of Nutritional Science and Vitaminology* **53**, 82-86.
- Sakata T, Kokima T, Fujieda M, Takahashi M, Michiba T. 2003. Influences of probiotic bacteria on organic acid production by pig caecal bacteria *in vitro*. *Proceedings of*

Nutrition Society **62**, 73-80.

- Singleton DR, Furlong MA, Rathbun SL, Whitman WB. 2001. Quantitative comparisons of 16S rDNA sequence libraries from environmental samples. *Applied and Environmental Microbiology* **67**, 4373–4376.
- Suto T. 1973. Examination of the rumen. In: Nakamura R, Yonemura T, Suto T (eds), *Methods in the Clinical Examination of the Bovine*, pp.39-42. Nosan Gyoson Bunka Kyokai, Tokyo.
- Tsukahara T, Ushida K. 2002. Succinate accumulation in pig large intestine during antibiotic-associated diarrhea and the constitution of succinate-producing flora. *Journal of General and Applied Microbiology* **48**, 143-154.
- Walter J, Hertel C, Tannock GW, Lis CM, Munro K, Hammes WP. 2001. Detection of *Lactobacillus*, *Pediococcus*, *Leuconostoc*, and *Weissella* species in human feces by using group-specific PCR primers and denaturing gradient gel electrophoresis. *Applied and Environmental Microbiology* **67**, 2578-2585.
- Weatherburn MW. 1967. Phenol-hypochlorite reaction for determination of ammonia. *Analytical Chemistry* **39**, 971–974.
- Wells JE, Yen JT, Miller DN. 2005. Impact of dried skim milk in production diets on *Lactobacillus* and pathogenic bacterial shedding in growing-finishing swine. *Journal of Applied Microbiology* **99**, 400–407.
- Woo PCY, Lau SKP, Chan KM, Fung AMY, Tang BSF, Yuen KY. 2005. *Clostridium bacteriaemia* characterized by 16S ribosomal RNA gene sequencing. *Journal of Clinical Pathology* **58**, 301-307.
- Yu Z, Morrison M. 2004. Improved extraction of PCR-quality community DNA from digesta and fecal samples. *Biotechniques* **36**, 808-812.

Figure 1 Fecal bacterial community determined by denatured gradient gel electrophoresis and its similarity between sows with and without liquid whey feeding.

¹ Similarity (%) = $2NAB/(NA+NB)$

NA and NB are the number of band detected in sample A and B, respectively, while NAB is the number of band detected both in sample A and B .

² 0wk (Control), 2wk and 4wk are samples taken prior to, 2weeks and 4weeks after whey feeding, respectively.

Table 1 Bacteria identified as known species in 16S rDNA mini-library prepared from whey fed to experimental pigs ¹

Species	No. of identified	(%)	Accession No.
<i>Streptococcus thermophilus</i>	1	(4.8) ²	AB506443
<i>Lactobacillus helveticus</i>	4	(19.0)	AB506440, AB506444, AB506448, AB506452
<i>Lactococcus piscium</i>	3	(14.3)	AB506435, AB506438, AB506451
<i>Lactococcus lactis</i>	2	(9.5)	AB506446, AB506447
<i>Rahnella aquatilis</i>	5	(23.8)	AB506439, AB506442, AB506445, AB506449, AB506454
<i>Pantoea agglomerans</i>	1	(4.8)	AB506441
<i>Acetobacter fabarum</i>	2	(9.5)	AB506434, AB506436
<i>Acetobacter orientalis</i>	1	(4.8)	AB506450
Total identified	19	(90.5)	
Total unidentified	2	(9.5)	AB506437, AB506453

¹ Share sequence identity at 97% or higher level with known bacteria

² Numbers of parenthesis are % of total detected.

Table 2 Bacteria detected in 16S rDNA library prepared from feces of sows with and without liquid whey feeding

Phyla/Groups	Control ¹		Whey ¹	
Firmicutes				
Clostridiales cluster I	13	(14.3) ²	7	(9.0)
III	1	(1.1)	0	(0)
IV	24	(26.4)	16	(20.5)
IX	9	(9.9)	23	(29.5)
XI	8	(8.8)	7	(9.0)
XIVa	17	(18.7)	13	(16.7)
Lactobacillales	14	(15.4)	5	(6.4)
Proteobacteria	4	(4.4)	0	(0)
Others	1	(1.1)	7	(9.0)
Total	91	(100)	78	(100)
<i>P</i> value ³	0.001		0.002	

¹ Prepared from feces taken prior to and 4wk after whey feeding

² Numbers in parenthesis are % of total identified.

³ Represents statistical significance in difference between two libraries.

Table 3 Bacteria identified as known species in 16S rDNA library prepared from feces of sows with and without liquid whey feeding ¹

Species	Control		Whey		Accession No.
<i>Clostridium sardiniense</i>	3	(3.7) ²	1	(1.5)	AB506285, AB506293, AB506322, AB506425
<i>Clostridium butyricum</i>	2	(2.4)	2	(3.0)	AB506292, AB506318, AB506380, AB506433
<i>Clostridium beijerinckii</i>	1	(1.2)	0	(0)	AB506291
<i>Clostridium puniceum</i>	0	(0)	1	(1.5)	AB506414
<i>Ruminococcus flavefaciens</i>	1	(1.2)	0	(0)	AB506290
<i>Mitsuokella jalaludinii</i>	0	(0)	5	(7.5)	AB506375, AB506376, AB506403, AB506405, AB506409,
<i>Megasphaera elsdenii</i>	0	(0)	1	(1.5)	AB506398
<i>Megasphaera hominis</i>	0	(0)	1	(1.5)	AB506427
<i>Clostridium glycolicum</i>	4	(4.9)	3	(4.5)	AB506297, AB506307, AB506321, AB506338, AB506377, AB506379, AB506390
<i>Clostridium disporicum</i>	1	(1.2)	0	(0)	AB506357
<i>Clostridium baratii</i>	0	(0)	1	(1.5)	AB506374
<i>Coprococcus eutactus</i>	0	(0)	1	(1.5)	AB506372
<i>Roseburia faecalis</i>	0	(0)	1	(1.5)	AB506423
<i>Kurthia gibsonii</i>	1	(1.2)	1	(1.5)	AB506298, AB506406
<i>Lactobacillus ruminis</i>	2	(2.4)	0	(0)	AB506300, AB506301
<i>Lactobacillus amylovorus</i>	4	(4.9)	3	(4.5)	AB506320, AB506325, AB506331, AB506343, AB506389, AB506395, AB506401
<i>Bacillus bhargavae</i>	1	(1.2)	0	(0)	AB506347
<i>Acinetobacter seohaensis</i>	2	(2.4)	0	(0)	AB506287, AB506310
<i>Acinetobacter calcoaceticus</i>	1	(1.2)	0	(0)	AB506359
<i>Eshcherichia coli</i>	1	(1.2)	0	(0)	AB506289
Total identified	24	(29.3)	21	(31.3)	
					AB506286, AB506288, AB506294, AB506295, AB506296, AB506299, AB506302 -AB506306, AB506308, AB506309, AB506311-AB506317, AB506319, AB506323, AB506324, AB506326 - AB506330, AB506332 - AB506337, AB506339, AB506340, AB506341, AB506342, AB506344-AB506346, AB506348-AB506356, AB506358, AB506360-AB506371, AB506373, AB506378, AB506381-AB506388, AB506391 -AB506394, AB506396, AB506399, AB506400, AB506402, AB506404, AB506407 -AB506413, AB506415 - AB506422, AB506424, AB506426, AB506428 - AB506432
Total unidentified	58	(70.7)	46	(68.7)	

¹ Share sequence identity at 97% or higher level with known bacteria.

² Numbers in parenthesis are % of total detected.

Table 4 Fecal characteristics of sows with and without liquid whey feeding ¹

Microbial and chemical parameters quantified	Whey feeding for		
	0wk (Control)	2wk	4wk
Total bacteria (log 16S rDNA copy/g feces)	12.07 ± 0.07 ^a	12.32 ± 0.16 ^{ab}	12.29 ± 0.17 ^b
Total lactobacilli (log 16S rDNA copy/g feces)	9.22 ± 0.63 ^a	9.47 ± 0.23 ^{ab}	9.77 ± 0.36 ^b
Total bifidobacteria (log 16S rDNA copy/g feces)	7.95 ± 0.48 ^a	8.47 ± 0.27 ^b	8.60 ± 0.66 ^{ab}
Lactobacilli (relative % of total bacteria)	0.14 ± 0.40	0.14 ± 0.13	0.30 ± 0.15
Bifidobacteria (relative % of total bacteria)	0.01 ± 0.01	0.01 ± 0.01	0.02 ± 0.06
pH	7.49 ± 0.36 ^a	6.92 ± 0.29 ^b	7.37 ± 0.20 ^{ab}
Total SCFA (mmol/g feces)	0.73 ± 0.21	0.69 ± 0.23	0.62 ± 0.18
Acetate (%)	63.2 ± 7.4	58.9 ± 2.9	58.7 ± 2.7
Propionate (%)	21.8 ± 3.7	24.7 ± 1.1	23.8 ± 2.1
n-Butyrate (%)	8.2 ± 3.1	9.6 ± 1.0	9.6 ± 1.7
Total lactate	7.45 ± 1.95 ^a	5.01 ± 0.75 ^b	4.48 ± 0.38 ^b
L-lactate (mmol/g feces)	4.21 ± 1.56	3.26 ± 0.64	2.96 ± 0.69
D-Lactate (mmol/g feces)	3.24 ± 0.76 ^a	1.75 ± 0.70 ^b	1.53 ± 0.60 ^b
Ammonia (mgN/g feces)	11.6 ± 5.0	6.9 ± 0.8	8.4 ± 1.9

¹ Feces from 5 sows at 0 (Control), 2 and 4 wk after starting liquid whey feeding was employed for analysis.

^{a, b}: Values with different superscripts differ significantly at *P* <0.05.

Table 5 Bacteria detected in 16S rDNA library prepared from feces of growing pigs with and without liquid whey feeding

Phyla / Groups	Control		Whey	
Firmicutes				
Clostridiales clustε I	3	(3.4) ¹	2	(2.3)
III	0	(0)	1	(1.1)
IV	14	(16.1)	6	(6.9)
IX	1	(1.1)	2	(2.3)
XI	3	(3.4)	3	(3.4)
XIVa	28	(32.2)	28	(32.2)
Lactobacillales	38	(43.7)	44	(50.6)
Others	0	(0)	1	(1.1)
Total	87	(100)	87	(100)
P value ²	0.747		0.806	

¹ Numbers in parenthesis are % of total identified.

² Represents statistical significance of difference between two libraries.

Table 6 Bacteria identified as known species in 16S rDNA library prepared from feces of growing pigs with and without liquid whey feeding ¹

Species	Control	Whey	Accession No.
<i>Clostridium bartlettii</i>	1 (1.3) ²	2 (2.4)	AB506166, AB506207, AB506252
<i>Coprococcus catus</i>	2 (2.6)	1 (1.2)	AB506150, AB506190, AB506281
<i>Ruminococcus obeum</i>	1 (1.3)	4 (4.8)	AB506135, AB506208, AB506219, AB506224, AB506258
<i>Roseburia inulinivorans</i>	1 (1.3)	0 (0)	AB506159
<i>Eubacterium rectale</i>	1 (1.3)	0 (0)	AB506124
<i>Eubacterium halii</i>	0 (0)	1 (1.2)	AB506265
<i>Lactobacillus amylovorus</i>	12 (15.6)	12 (14.3)	AB506133, AB506134, AB506141, AB506152, AB506157, AB506170, AB506173, AB506175, AB506179, AB506180, AB506187, AB506189, AB506203, AB506204, AB506210, AB506212, AB506220, AB506239, AB506240, AB506243, AB506247, AB506254, AB506280, AB506284
<i>Lactobacillus reuteri</i>	5 (6.5)	14 (16.7)	AB506155, AB506158, AB506160, AB506194, AB506196, AB506214, AB506216, AB506218, AB506228, AB506230, AB506231, AB506233, AB506235, AB506259, AB506263, AB506264, AB506268, AB506273,
<i>Lactobacillus gasseri</i>	1 (1.3)	0 (0)	AB506144
<i>Lactobacillus acidophilus johnsonii</i>	3 (3.9)	3 (3.6)	AB506143, AB506156, AB506176, AB506221, AB506251, AB506283
<i>Streptococcus alactolyticus</i>	11 (14.3)	12 (14.3)	AB506125, AB506137, AB506142, AB506145, AB506148, AB506149, AB506163, AB506164, AB506168, AB506183, AB506195, AB506213, AB506223, AB506225, AB506226, AB506234, AB506236, AB506238, AB506241, AB506245, AB506249, AB506257, AB506282,
Total identified	38 (49.4)	49 (58.3)	AB506126 - AB506132, AB506136, AB506138-AB506140, AB506146, AB506147, AB506151, AB506153, AB506154, AB506161, AB506162, AB506165 - AB506167, AB506169, AB506172, AB506174, AB506177, AB506178, AB506181, AB506182, AB506184 - AB506186, AB506188, AB506191 - AB506193, AB506197 - AB506202, AB506205, AB506206, AB506209, AB506211, AB506215, AB506217, AB506222, AB506227, AB506229, AB506232, AB506237, AB506242, AB506244, AB506246, AB506248, AB506250, AB506253, AB506255, AB506256, AB506260,-AB506262, AB506266, AB506267, AB506269 - AB506276, AB506278, AB506279
Total unidentified	39 (50.6)	35 (41.7)	

¹ Share sequence identity at 97% or higher level with known bacteria.

² Numbers in parenthesis are % of total detected.

Table 7 Fecal characteristics of growing pigs with and without liquid whey feeding ¹

Microbial and chemical parameters quantified	Days of age							
	43		65		104		139	
	Control	Whey	Control	Whey	Control	Whey	Control	Whey
Total bacteria (log 16S rDNA copy/g feces)	10.83 ± 0.34	11.74 ± 0.25 *	nd	nd	11.22 ± #####	11.40 ± 0.28	11.51 ± 0.22	11.36 ± 0.18
Total lactobacilli (log 16S rDNA copy/g feces)	10.59 ± 0.26	11.52 ± 0.25 *	nd	nd	10.76 ± #####	11.05 ± 0.36	10.47 ± 0.30	10.49 ± 0.30
Total bifidobacteria (log 16S rDNA copy/g feces)	7.96 ± 0.36	8.41 ± 0.16 *	nd	nd	9.01 ± #####	8.99 ± 0.48	8.66 ± 0.21	9.28 ± 0.42
Lactobacilli (relative % of total bacteria)	65.4 ± 35.1	77.3 ± 59.2	nd	nd	38.1 ± #####	48.5 ± 19.6	10.7 ± 5.8	14.6 ± 5.6
Bifidobacteria (relative % of total bacteria)	0.2 ± 0.2	0.1 ± 0.1	nd	nd	1.4 ± 2.2	0.6 ± 0.6	0.1 ± 0.1	1.6 ± 2.5
pH	6.41 ± 0.23	6.21 ± 0.31 *	6.64 ± 0.21	6.72 ± 0.15	6.75 ± 0.36	6.67 ± 0.33	6.72 ± 0.31	6.92 ± 0.35
Total SCFA (mmol/g feces)	0.18 ± 0.05	0.16 ± 0.04	0.16 ± 0.03	0.15 ± 0.04	0.16 ± 0.05	0.17 ± 0.03	0.17 ± 0.04	0.18 ± 0.05
Acetate (%)	56.7 ± 5.2	56.0 ± 3.6	52.6 ± 5.5	53.6 ± 3.5	51.0 ± 6.9	50.6 ± 5.7	53.1 ± 3.8	55.2 ± 3.8
Propionate (%)	25.2 ± 2.6	25.9 ± 2.1	22.7 ± 1.4	23.2 ± 1.2	24.0 ± 3.3	25.5 ± 1.7	24.2 ± 2.3	22.7 ± 2.2
n-Butyrate (%)	10.9 ± 2.3	11.8 ± 2.5	15.9 ± 3.1	14.5 ± 2.2	15.2 ± 3.9	14.8 ± 4.2	12.8 ± 1.8	12.3 ± 2.2
Total lactate (mmol/g feces)	10.47 ± 3.69	8.99 ± 2.17	5.66 ± 3.99	5.39 ± 2.11	10.82 ± #####	8.00 ± 3.76	8.52 ± 2.44	8.10 ± 2.16
L-lactate (mmol/g feces)	3.83 ± 1.95	2.97 ± 1.31	1.90 ± 1.33	2.19 ± 0.92	4.80 ± #####	3.08 ± 1.67 *	3.08 ± 1.03	2.99 ± 1.33 *
D-Lactate (mmol/g feces)	6.64 ± 1.90	6.02 ± 1.10	3.76 ± 3.00	3.20 ± 1.72	6.02 ± #####	4.92 ± 2.25	5.44 ± 1.44	5.11 ± 1.03
Ammonia (mgN/g feces)	0.77 ± 0.31	0.79 ± 0.29	0.94 ± 0.27	0.77 ± 0.34	0.63 ± #####	0.61 ± 0.19	0.65 ± 0.22	0.58 ± 0.23

¹ Feces from 6 pigs of each group was employed for analysis. * Significantly different from control at $P < 0.05$. nd, not determined; SCFA, short-chain fatty acids