Title	Prediction of maturational competence of feline oocytes using supravital staining of cumulus cells by propidium iodide
Author(s)	Uchikura, Kenzo; Nagano, Masashi; Hishinuma, Mitsugu
Citation	Zygote, 20(4), 333-337 https://doi.org/10.1017/S096719941100027X
Issue Date	2011-05
Doc URL	http://hdl.handle.net/2115/49780
Rights	Cambridge University Press
Type	article
File Information	Zygote2011.pdf

[Instructions for use](#)

Prediction of maturational competence of feline oocytes using supravital staining of cumulus cells by propidium iodide

Kenzo Uchikura^{2,3}, Masashi Nagano¹ and Mitsugu Hishinuma²

Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University; School of Veterinary Medicine, Faculty of Agriculture, Tottori University; The United Graduate School of Veterinary Science, Yamaguchi University Japan

Date submitted: 03.12.10. Date accepted: 02.03.11

Summary

We examined the relationship between integrity of cumulus cells and nuclear maturation rate after *in vitro* culture to determine a non-invasive prediction of the maturational competence of feline oocytes. Feline cumulus–oocyte complexes (COCs) were collected from either small (400–800 µm) or large (≥800 µm) follicles. Immediately after collection, cumulus cells were evaluated morphologically (thickness of cumulus cell layers) and stained with propidium iodide (PI), which penetrates only non-viable cells. Cumulus cells without PI staining were judged as having good membrane integrity. After evaluation, COCs were cultured for 30 h and their nuclear maturation rate was determined. The nuclear maturation rate of oocytes derived from large follicles (89.8%) was higher ($p < 0.05$) than that from small follicles (60.8%). There was no difference in the maturation rate of oocytes from follicles with the same size regardless of cumulus morphology. In contrast, oocytes that had cumulus cells with good membrane integrity showed a higher maturation rate (93.8%) than oocytes with poor cumulus integrity (76.9%) in large follicles ($p < 0.05$). We conclude that evaluation of membrane integrity of cumulus cells by propidium iodide staining can be used to predict the maturational competence of oocytes.

Keywords: Cat, Follicle, IVM, Membrane integrity, Propidium iodide

Introduction

The domestic cat is an important experimental model for advancing assisted reproductive technologies that may be useful for conservation of non-domestic felidae. Development of reliable methods for *in vitro* maturation (IVM), *in vitro* fertilization (IVF) and *in vitro* culture (IVC) of domestic cat oocytes is a prerequisite for their subsequent application for preservation of

endangered species. Research on *in vitro* production (IVP) of domestic cat embryos has increased rapidly (Pope, 2000; Pope *et al.*, 2006, 2009; Spinaci *et al.*, 2007; Nagano *et al.*, 2008; Naoi *et al.*, 2008; Uchikura *et al.*, 2010). Ideally, for efficient IVP of feline embryos, oocytes with high maturational competence should be selected non-invasively before maturational culture.

It is well established that oocytes surrounded by several layers of cumulus cells and with evenly granulated dark ooplasm have higher *in vitro* developmental competence than oocytes with irregularly granulated ooplasm and fewer cumulus cell layers (Pope *et al.*, 1997; Wood & Wildt, 1997; Katska-Ksiazkiewicz *et al.*, 2003; Naoi *et al.*, 2008). Also, *in vitro* maturation and developmental competence of oocytes are influenced by season (Spindler & Wildt, 1999; Freistedt *et al.*, 2001) and ovarian cyclicity status (Johnston *et al.*, 1989; Freistedt *et al.*, 2001; Karja *et al.*, 2002; Naoi *et al.*, 2008), even if oocytes are selected using similar criteria for oocyte and cumulus cell morphologies.

Oocytes and cumulus cells grow and develop in a highly coordinated and mutually dependent manner.

¹All correspondence to: Masashi Nagano. Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060–0818, Japan. Tel./Fax: +81 11 706 5232. e-mail: mnaga@vetmed.hokudai.ac.jp

²Department of Theriogenology, School of Veterinary Medicine, Faculty of Agriculture, Tottori University, 4–101 Koyama-Minami, Tottori 680–8553, Japan.

³The United Graduate School of Veterinary Science, Yamaguchi University, 1677–1 Yoshida, Yamaguchi 753–8515, Japan; 1 Laboratory of Theriogenology, Department of Veterinary Clinical Sciences, Graduate School of Veterinary Medicine, Hokkaido University, Sapporo 060–0818, Japan.

Cumulus cells support oocyte growth and maturation by supplying nutrients and growth factors (Tanghe *et al.*, 2002). Recently, researchers attempted to use the expression of cumulus genes as a marker for non-invasive assessment of oocyte quality in cattle (Assidi *et al.*, 2008; Bettgowda *et al.*, 2008) and humans (Assou *et al.*, 2008, 2010; Anderson *et al.*, 2009). However, this method requires a biopsy of the cumulus cells, and it is laborious, complicated and expensive. There is another method for easily assessing the quality of cumulus cells using propidium iodide (PI) staining. PI does not penetrate intact cell membrane, but can penetrate only cells with non-intact membranes that have lost their integrity. In a study that evaluated the membrane integrity of cumulus cells of cryopreserved murine oocytes at the germinal vesicle stage by PI staining, differences in developmental potential appeared to depend on the assumption that the proportion of oocytes with good cumulus integrity (without PI staining) was similar to the proportion of embryos developing to the blastocyst stage after IVF/IVC (Ruppert-Lingham *et al.*, 2006).

Therefore, in the present study, for non-invasive prediction of the maturational competence of feline oocytes, we examined the morphology of cumulus investments and stained cumulus cells immediately after collection using PI, and subsequently investigated nuclear maturation rate after IVM culture. In domestic cats, it was reported that follicles $\geq 800 \mu\text{m}$ in diameter expressed the luteinizing hormone (LH) receptor in granulosa cells (Saint-Dizier *et al.*, 2007). Therefore, we collected oocytes from small (400–800 μm) and large follicles ($\geq 800 \mu\text{m}$), and the diameter of oocytes was also examined.

Materials and methods

Animals for ovarian collection

Ovaries of eight cats were obtained following ovario-hysterectomy at the veterinary teaching hospital of Tottori University and a local veterinary clinic in Tottori city and kept in normal saline at room temperature before delivery to the laboratory within 1 h after collection.

Antral follicle isolation and oocyte recovery

Upon arrival in the laboratory, ovaries were transferred to isolation medium: TCM-199 (Invitrogen) containing 0.1% polyvinyl alcohol (Sigma-Aldrich), 25 mM HEPES (Sigma-Aldrich), 0.85 mg/ml NaHCO_3 (Kanto Chemical Co., Inc.) and 50 $\mu\text{g}/\text{ml}$ gentamicin sulfate (Sigma-Aldrich) (pH 7.4) (Harada *et al.*, 1997). Ovaries were cut with scissors along their long axis, the

ovarian medulla was removed and the cortex was cut into small pieces. It was difficult to identify the antrum in follicles that were $<400 \mu\text{m}$ in diameter. The antral follicles that were $\geq 400 \mu\text{m}$ in diameter were isolated using an 18-gauge needle under a stereomicroscope and transferred to fresh isolation medium. Using an ocular micrometer, follicular diameter was measured as the mean length of two perpendicular axes. The follicles were divided into two groups according to their diameter: 400–800 (small) and $\geq 800 \mu\text{m}$ (large). The follicles were then dissected with a 25-gauge needle under a stereomicroscope. Cumulus–oocyte complexes (COCs) were collected and their morphological characteristics were determined under a stereomicroscope ($\times 15$ –50). Oocytes surrounded by cumulus cells and having evenly granulated black ooplasm were defined as morphologically normal and used for the experiment.

Evaluation of membrane integrity of cumulus cells by propidium iodide staining and morphology

The method of evaluation of membrane integrity of cumulus cells was in accordance with that of a previous report (Ruppert-Lingham *et al.*, 2006) with slight modification. Briefly, COCs were incubated for 10 min in the dark at 39°C in 3 ml of isolation medium supplemented with 0.1 mg/ml PI (Sigma-Aldrich). The COCs were then washed twice in isolation medium and transferred into a 10- μl droplet of isolation medium covered with paraffin oil. Stained COCs were observed under an inverted microscope equipped with a fluorescence system (TS100-F, Nikon) using an appropriate filter (G-2A, Nikon). Cumulus cells with a damaged cell membrane fluoresced red. Each COC was scored for membrane integrity of the cumulus cells using the following scoring system (Ruppert-Lingham *et al.*, 2006) with slight modification. The COCs were defined as either grade 1: less than one-quarter of cumulus cells were stained by PI; grade 2: one-quarter to one-half of cumulus cells were stained; or grade 3: more than one-half of cumulus cells were stained. These oocytes were also divided into three categories by the morphology of their surrounding cumulus investments: categories A: at least three layers; B: less than three layers; or C, only corona radiata. Images of COCs were taken using a digital camera (DP-11, Olympus) attached to an inverted microscope and the diameter of ooplasm was measured as the mean length of two perpendicular axes.

In vitro maturational culture and assessment of nuclear maturation

After evaluation of membrane integrity and morphology, COCs were washed once in maturation medium, which was composed of 25 mM HEPES-

buffered TCM 199 (cat. no. 12340, Earle's salt, Invitrogen) supplemented with 3 mg/ml fatty-acid-free BSA (Sigma-Aldrich), 0.02 units/ml follicle stimulating hormone (F2293, from porcine pituitary, Sigma-Aldrich), 1 µg/ml estradiol-17β (Sigma-Aldrich), 0.2 mM sodium pyruvate (Sigma-Aldrich) and 50 µg/ml gentamicin sulfate (Nagano *et al.*, 2008). They were then cultured individually in 10-µl droplets of maturation medium covered with paraffin oil at 39°C in a humidified atmosphere of 5% CO₂ in air for 30 h.

After IVM culture, the cumulus investments were removed from the oocytes by gentle pipetting. They were then fixed with a mixture of ethanol:acetic acid (3:1) and stained with 1% aceto-orcein solution. The nuclear status was evaluated using a phase-contrast microscope. When chromosomes were condensed and present in the equatorial view with the extrusion of the first polar body, the nuclear stage was classified as metaphase II (MII) and considered to be meiotically mature.

Experimental design

Firstly, we compared the diameter of ooplasm and maturation rate after IVM culture of oocytes derived from follicles with 400–800 (small) or ≥800 µm (large) in diameter. Secondly, effects of morphological categories and membrane integrity of cumulus cells on the diameter of ooplasm and the maturation rate were examined. Finally, the maturational competence of oocytes surrounded by several layers (category A) of intact cumulus cells (grade 1) was evaluated.

Statistical analysis

Statistical analysis was performed using JMP v8.0.2 software (SAS Institute Inc.). The difference in oocyte diameter was analysed by Student's *t*-test or by one-way analysis of variance (ANOVA) followed by Tukey-Kramer's Honestly Significant Difference test. The maturation rates of oocytes were compared by chi-squared test. Values were considered statistically significant at $p < 0.05$.

Results and Discussion

We collected 267 oocytes with normal morphology from eight cats (33.4 ± 13.9 /female, mean $\pm$ SD). Oocytes derived from the ≥800 µm (large) follicle group showed a larger diameter of ooplasm and a higher maturation rate than those from the 400–800 µm (small) group ($p < 0.05$, Table 1). These results indicate that feline oocytes grow until the follicular diameter reaches 800 µm. Luteinizing hormone binding sites were detected in the granulosa cell layer of feline

Table 1 Oocyte size and maturational ability in small (400–800 µm) vs. large (≥800 µm) follicles

Follicular diameter (µm)	Oocytes <i>n</i> (cats <i>n</i>)	Diameter of ooplasm (µm)	Nuclear maturation (%)
400–800	130 (8)	104.0 $\pm$ 6.1 ^a	60.8 ^a
≥ 800	137 (8)	107.7 $\pm$ 5.5 ^b	89.8 ^b

^{a,b}Values with different superscripts in the same column differ significantly ($p < 0.05$).

Oocyte diameter values are mean $\pm$ standard deviation (SD). Data are pooled from eight cats.

follicles that were ≥800 µm in diameter (Saint-Dizier *et al.*, 2007). The diameter of ooplasm seemed to have an effect on the maturational competence of feline oocytes, as has been shown in other species (Sato *et al.*, 1990; Fair *et al.*, 1995; Arlotto *et al.*, 1996; Otoi *et al.*, 1997; Ledda *et al.*, 1999). In a previous study, it was demonstrated that there is a strong relationship between diameter and meiotic competence of cat oocytes (Otoi *et al.*, 2001) although follicle diameter was not measured. Thus, the present results confirm earlier reports (Otoi *et al.*, 2001; Saint-Dizier *et al.*, 2007) indicating that oocyte growth occurs and maturational competence improves as follicular diameter grows up to ≥800 µm in diameter.

As shown in Table 2, there was no difference in diameter of ooplasm and maturation rate of oocytes derived from follicles with the same size, regardless of cumulus cell morphology. Bovine oocytes with a multilayered cumulus cells showed higher rates of nuclear maturation, fertilization and development after IVP than denuded oocytes and oocytes with corona radiata alone (Xu *et al.*, 1986; Shioya *et al.*, 1988; Yang & Lu, 1990; Madison *et al.*, 1992; Momozawa & Fukuda, 1995; Stojkovic *et al.*, 2001). However, we could not detect an obvious difference of maturational competence between cat oocytes with different cumulus cell morphologies. In contrast, oocytes of similar diameter from large follicles having cumulus cells with intact membrane integrity (grade 1) showed a higher maturation rate ($p < 0.05$) than oocytes having poor cumulus integrity (grade 3) (Table 3). In addition, oocytes surrounded by cumulus cells with grade 1 integrity from small follicles were larger in diameter than oocytes enclosed by cumulus cells of grades 2 and 3 ($p < 0.05$). However, the diameter of oocytes and the nuclear maturation rate was lower than those of oocytes surrounded by cumulus cells with grade 1 integrity from large follicles. Taken together, these results indicate that live and functional cumulus cells support the growth and nuclear maturation of feline oocytes.

In large follicles, maturation rate of oocytes selected by morphological category A (92.0%, $n = 75$) was similar to that of oocytes selected by grade 1 of

Table 2 Effects of cumulus morphological categories on size and maturational ability of cat oocytes from small (400–800 μm) vs. large ($\geq 800 \mu\text{m}$) follicles

Items	Follicular diameter, μm	Morphological categories of cumulus ^c (oocytes, <i>n</i>)		
		A	B	C
Diameter of ooplasm (μm)	400–800	104.4 $\pm$ 5.6 (59) ^a	104.5 $\pm$ 6.9 (41) ^a	102.6 $\pm$ 5.8 (30) ^a
	≥ 800	107.4 $\pm$ 5.2 (75) ^b	108.2 $\pm$ 6.4 (43) ^b	107.5 $\pm$ 3.9 (19) ^b
Nuclear maturation rate (%)	400–800	67.8 ^a	61.0 ^a	46.7 ^a
	≥ 800	92.0 ^b	86.0 ^b	89.5 ^b

^{a,b}Values with different superscripts in the same column differ significantly ($p < 0.05$).

^cCumulus category: A, oocytes tightly surrounded by at least three layers of cumulus cells; B, oocytes surrounded by less than three layers of cumulus cells; C, oocytes surrounded by only corona radiata cells. Oocyte diameter values are mean $\pm$ standard deviation (SD).

Table 3 Effects of membrane integrity of cumulus cells on size and maturational ability of cat oocytes from small (400–800 μm) vs. large ($\geq 800 \mu\text{m}$) follicles

Items	Follicular diameter, μm	Membrane integrity grade of cumulus cells (oocytes <i>n</i>) ^c		
		1	2	3
Diameter of ooplasm (μm)	400–800	105.7 $\pm$ 6.1 (68) ^{a,A}	102.5 $\pm$ 5.4 (31) ^{a,B}	101.8 $\pm$ 5.9 (31) ^{a,B}
	≥ 800	107.8 $\pm$ 5.6 (80) ^b	107.2 $\pm$ 4.9 (31) ^b	107.7 $\pm$ 5.8 (26) ^b
Nuclear maturation rate (%)	400–800	67.6 ^a	54.8 ^a	51.6 ^a
	≥ 800	93.8 ^{b,A}	90.3 ^{b,A,B}	76.9 ^{b,B}

^{a,b}Values with different superscripts in the same column differ significantly ($p < 0.05$).

^{A,B}Values with different superscripts in the same row differ significantly ($p < 0.05$).

^cGrade of oocytes: grade 1, one-quarter of cumulus cells stained by propidium iodide (PI); grade 2, one-half of cumulus cells stained by PI; grade 3, more than one-half of cumulus cells stained by PI. Values of oocyte diameter are means $\pm$ standard deviation (SD).

cumulus cell integrity (93.8%, $n = 80$). However, the oocytes surrounded by at least three layers of cumulus cells (category A) derived from large follicles were divided by their cumulus cell integrity into grade 1 (72%, 54/75) and grades 2 and 3 (28%, 21/75), and maturation rate of oocytes having cumulus cells with grade 1 integrity (96.3%, $n = 54$) was higher than that of oocytes having cumulus cells classified as grades 2 and 3 (81.0%, $n = 21$) ($p < 0.05$). This result also suggests that evaluating cumulus cell integrity concomitantly with cumulus morphology will improve IVP outcome in cats.

In conclusion, evaluation of membrane integrity of cumulus cells by propidium iodide staining can be used as a tool for predicting the maturational competence of oocytes.

Acknowledgements

This study was supported by a Grant-in-Aid for Scientific Research (no. 21780286) from the Ministry of Education, Science, Sports and Culture, Japan, to M. Nagano.

References

- Anderson, R.A., Sciorio, R., Kinnell, H., Bayne, R.A., Thong, K.J., de Sousa, P.A. & Pickering, S. (2009). Cumulus gene expression as a predictor of human oocyte fertilisation, embryo development and competence to establish a pregnancy. *Reproduction* **138**, 629–637.
- Arlotto, T., Schwartz, J.L., First, N.L. & Leibfried-Rutledge, M.L. (1996). Aspects of follicle and oocyte stage that affect *in vitro* maturation and development of bovine oocytes. *Theriogenology* **45**, 943–956.
- Assidi, M., Dufort, I., Ali, A., Hamel, M., Algriany, O., Dielemann, S. & Sirard, M.A. (2008). Identification of potential markers of oocyte competence expressed in bovine cumulus cells matured with follicle-stimulating hormone and/or phorbol myristate acetate *in vitro*. *Biol. Reprod.* **79**, 209–222.
- Assou, S., Haouzi, D., Mahmoud, K., Aouacheria, A., Guillemain, Y., Pantescio, V., Reme, T., Dechaud, H., De Vos, J. & Hamamah, S. (2008). A non-invasive test for assessing embryo potential by gene expression profiles of human cumulus cells: a proof of concept study. *Mol. Hum. Reprod.* **14**, 711–719.
- Assou, S., Haouzi, D., De Vos, J. & Hamamah, S. (2010). Human cumulus cells as biomarkers for embryo and pregnancy outcomes. *Mol. Hum. Reprod.* **16**, 531–538.
- Bettgowda, A., Patel, O.V., Lee, K.B., Park, K.E., Salem, M., Yao, J., Ireland, J.J. & Smith, G.W. (2008). Identification of

- novel bovine cumulus cell molecular markers predictive of oocyte competence: functional and diagnostic implications. *Biol. Reprod.* **79**, 301–309.
- Fair, T., Hyttel, P. & Greve, T. (1995). Bovine oocyte diameter in relation to maturational competence and transcriptional activity. *Mol. Reprod. Dev.* **42**, 437–442.
- Freistedt, P., Stojkovic, M. & Wolf, E. (2001). Efficient *in vitro* production of cat embryos in modified synthetic oviduct fluid medium: effects of season and ovarian status. *Biol. Reprod.* **65**, 9–13.
- Harada, M., Miyano, T., Matsumura, K., Osaki, S., Miyake, M. & Kato, S. (1997). Bovine oocytes from early antral follicles grow to meiotic competence *in vitro*: effect of FSH and hypoxanthine. *Theriogenology* **48**, 743–755.
- Johnston, L.A., O'Brien, S.J. & Wildt, D.E. (1989). *In vitro* maturation and fertilization of domestic cat follicular oocytes. *Gamete Res.* **24**, 343–356.
- Karja, N.W., Otoi, T., Murakami, M., Fahrudin, M. & Suzuki, T. (2002). *In vitro* maturation, fertilization and development of domestic cat oocytes recovered from ovaries collected at three stages of the reproductive cycle. *Theriogenology* **57**, 2289–2298.
- Katska-Ksiazkiewicz, L., Rynska, B., Kania, G., Smorag, Z., Gajda, B. & Pienkowski, M. (2003). Timing of nuclear maturation of nonstored and stored domestic cat oocytes. *Theriogenology* **59**, 1567–1574.
- Ledda, S., Bogliolo, L., Leoni, G. & Naitana, S. (1999). Follicular size affects the meiotic competence of *in vitro* matured prepubertal and adult oocytes in sheep. *Reprod. Nutr. Dev.* **39**, 503–508.
- Madison, V., Avery, B. & Greve, T. (1992). Selection of immature bovine oocytes for developmental potential *in vitro*. *Anim. Reprod. Sci.* **27**, 1–11.
- Momozawa, K. & Fukuda, Y. (1995). *In vitro* maturation and *in vitro* fertilization of bovine oocytes with heterogeneous ooplasm. *Anim. Sci. Technol.* **66**, 605–609.
- Nagano, M., Uchikura, K., Takahashi, Y. & Hishinuma, M. (2008). Effect of duration of *in vitro* maturation on nuclear maturation and fertilizability of feline oocytes. *Theriogenology* **69**, 231–236.
- Naai, H., Agung, B., Karja, N.W., Wongsrikeao, P., Shimizu, R., Taniguchi, M. & Otoi, T. (2008). Effects of the reproductive status on morphological oocyte quality and developmental competence of oocytes after *in vitro* fertilization and somatic cell nuclear transfer in cat. *Reprod. Domest. Anim.* **43**, 157–161.
- Otoi, T., Yamamoto, K., Koyama, N., Tachikawa, S. & Suzuki, T. (1997). Bovine oocyte diameter in relation to developmental competence. *Theriogenology* **48**, 769–774.
- Otoi, T., Murakami, M., Ooka, A., Karja, N.W. & Suzuki, T. (2001). Effects of size and storage temperature on meiotic competence of domestic cat oocytes. *Vet. Rec.* **148**, 116–118.
- Pope, C.E. (2000). Embryo technology in conservation efforts for endangered felids. *Theriogenology* **53**, 163–174.
- Pope, C.E., McRae, M.A., Blair, B.L., Keller, G.L. & Dresser, B.L. (1997). *In vitro* and *in vivo* development of embryos produced by *in vitro* maturation and *in vitro* fertilization of cat oocytes. *J. Reprod. Fertil. Suppl.* **51**, 69–82.
- Pope, C.E., Gomez, M.C. & Dresser, B.L. (2006). *In vitro* production and transfer of cat embryos in the 21st century. *Theriogenology* **66**, 59–71.
- Pope, C.E., Crichton, E.G., Gomez, M.C., Dumas, C. & Dresser, B.L. (2009). Birth of domestic cat kittens of predetermined sex after transfer of embryos produced by *in vitro* fertilization of oocytes with flow-sorted sperm. *Theriogenology* **71**, 864–871.
- Ruppert-Lingham, C.J., Paynter, S.J., Godfrey, J., Fuller, B.J. & Shaw, R.W. (2006). Membrane integrity and development of immature murine cumulus–oocyte complexes following slow cooling to -60°C : the effect of immediate rewarming, plunging into LN_2 and two-controlled-rate-stage cooling. *Cryobiology* **52**, 219–227.
- Saint-Dizier, M., Malandain, E., Thoumire, S., Remy, B. & Chastant-Maillard, S. (2007). Expression of follicle stimulating hormone and luteinizing hormone receptors during follicular growth in the domestic cat ovary. *Mol. Reprod. Dev.* **74**, 989–996.
- Sato, E., Matsuo, M. & Miyamoto, H. (1990). Meiotic maturation of bovine oocytes *in vitro*: improvement of meiotic competence by dibutyryl cyclic adenosine 3',5'-monophosphate. *J. Anim. Sci.* **68**, 1182–1187.
- Shioya, Y., Kuwayama, M., Fukushima, M., Iwasaki, S. & Hanada, A. (1988). *In vitro* fertilization and cleavage capability of bovine follicular oocytes classified by cumulus cells and matured *in vitro*. *Theriogenology* **30**, 489–496.
- Spinaci, M., Merlo, B., Zannoni, A., Iacono, E., De Ambrogio, M., Turba, M.E. & Zambelli, D. (2007). *In vitro* production of cat blastocysts of predetermined sex using flow cytometrically sorted semen. *Theriogenology* **67**, 872–877.
- Spindler, R.E. & Wildt, D.E. (1999). Circannual variations in intraovarian oocyte but not epididymal sperm quality in the domestic Cat. *Biol. Reprod.* **61**, 188–194.
- Stojkovic, M., Machado, S.A., Stojkovic, P., Zakhartchenko, V., Hutzler, P., Goncalves, P.B. & Wolf, E. (2001). Mitochondrial distribution and adenosine triphosphate content of bovine oocytes before and after *in vitro* maturation: correlation with morphological criteria and developmental capacity after *in vitro* fertilization and culture. *Biol. Reprod.* **64**, 904–909.
- Tanghe, S., Van Soom, A., Nauwynck, H., Coryn, M. & de Kruif, A. (2002). Minireview: Functions of the cumulus oophorus during oocyte maturation, ovulation, and fertilization. *Mol. Reprod. Dev.* **61**, 414–424.
- Uchikura, K., Nagano, M. & Hishinuma, M. (2010). Evaluation of follicular development and oocyte quality in pre-pubertal cats. *Reprod. Domest. Anim.* **45**, e405–e411.
- Wood, T.C. & Wildt, D.E. (1997). Effect of the quality of the cumulus–oocyte complex in the domestic cat on the ability of oocytes to mature, fertilize and develop into blastocysts *in vitro*. *J. Reprod. Fertil.* **110**, 355–360.
- Xu, K.P., Greve, T., Smith, S. & Hyttel, P. (1986). Chronological changes of bovine follicular oocyte maturation *in vitro*. *Acta Vet. Scand.* **27**, 505–519.
- Yang, Y.B. & Lu, K.H. (1990). The influence of bovine oocyte type on *in vitro* fertilization and subsequent development *in vitro*. *Theriogenology* **33**, 355.