

HOKKAIDO UNIVERSITY

Title	Odor Discrimination and Signal Transduction Mechanisms in Olfactory and Vomeronasal Systems	
Author(s)	Taniguchi, Mutsuo	
Citation	北海道大学. 博士(薬学) 甲第3598号	
Issue Date	1995-03-24	
DOI	10.11501/3082600	
Doc URL	http://hdl.handle.net/2115/50175	
Туре	theses (doctoral)	
File Information	000000285052.pdf	



Odor Discrimination and Signal Transduction Mechanisms in Olfactory and Vomeronasal Systems

3598

Mutsuo Taniguchi



Odor Discrimination and Signal Transduction Mechanisms in Olfactory and Vomeronasal Systems

by

Mutsuo Taniguchi

A Doctoral Dissertation Submitted

to

Faculty of Pharmaceutical Sciences Hokkaido University Sapporo, 060 Japan



CONTENTS	(page)
GENERAL INTRODUCTION	1
LIST OF ABBREVIATIONS	14
PART 1	5. 53
Odor discrimination and signal transduction	
mechanisms in olfactory system	15
could meleotides into purio remember deceptor.	
Chapter I Difference in odor intensity and quality	
between <i>l</i> - and <i>d</i> -optical isomers	15
I-1 Difference in odor intensity between <i>l</i> - and <i>d</i> -optical	
isomers	15
I-2 Difference in odor quality between <i>l</i> - and <i>d</i> -optical	
isomers	27
Chapter II Effects of treatment of the turtle	
olfactory epithelium with various	
lipids on olfactory responses	44

Signal transduction mechanisms in vomeronasal

66

system

Chapter III	Electrophysiological properties of turtle vomeronasal receptor neurons	
Chapter IV	Signal transduction mechanisms in	
	vomeronasal receptor neurons	
IV-1 The res	ponses induced by intracellular injection of	
cyclic	nucleotides into turtle vomeronasal receptor	
neuror	IS and the second are block by the certain	
IV-2 The res	ponses induced by intracellular injection of	
IP3 in	to turtle vomeronasal receptor neurons	1(
Chapter V	cAMP-dependent transduction pathway	
sceently and	does not contribute to the transduction	
	for general odorants in turtle vomerona	sal
	neurons	11
SUMMARY		12
CONCE VID Y		

Augusta en el reporter de récentification de monter aporter el dec lege

GENERAL INTRODUCTION

There are many types of internal receptors in our bodies for detecting chemical substances such as hormones, neurotransmitters or blood glucose. These internal receptors perceive only a limited number of substances such specified substances. On the other hand, olfactory receptors sensitively detect various chemical substances present in the external environment, where there are many types of substances.

During the past decade, a great deal of work in biochemistry, electrophysiology and molecular biology has enhanced our understanding of the mechanism of signal transduction in olfactory systems. As a results of these works, several concepts have been proposed for the mechanism by which olfactory neurons transduce a chemical stimulus into an electrical response (Fig. I-1). The most generally accepted one is that an olfactory response is induced by binding of an odorant to specific receptor proteins, followed by a Gprotein-mediated increase in the internal cAMP and IP3 concentration to open the second messenger-activated cation channels on plasma membranes [1-12]. Recently Buck and Axel cloned the genes of an extremely large multigene family that encodes proteins with seven transmembrane domains and is expressed in the olfactory epithelium [13]. This report attracted a wide attention, but it is not as yet verified whether or not these proteins are true receptor proteins at present. Raming et al. reported the identification of another members of the gene family encoding putative odorant receptors and demonstrated that

ana o

they were introvided in educing marging neutropy (14). They inferred a cell line derived frem Specioprers Antiperio. (30) with Stepheriotics harbouring the receptor-incoding complementery DRS These com-



upin the ollactory epithelium it may be either predominantly or

Figure I-1. A schematic drawing representing the several signal transduction pathways in olfactory neurons. See text for details. AC: adenylate cyclase. ATP: adenosine triphosphate. cAMP: cyclic adenosine monophosphate. PLC: phospholipase C. PIP₂: phosphatidyl inositol bisphosphate. IP₃: inositol trisphosphate.



they were transcribed in olfactory receptor neurons [14]. They infected a cell line derived from *Spodoptera frugiperda* (Sf9) with baculovirus harbouring the receptor-encoding complementary DNA. These nonneuronal surrogate cells infected with the recombinant virus generated IP₃ responses upon stimulation with appropriate odorants, indicating that the receptors recognize odorants and couple to G-proteins of the host cells [14]. The above findings support the idea that an olfactory response is induced by binding of an odorant to specific receptor proteins.

On the other hand, it has been known that a number of nonolfactory systems such as turtle trigeminal nerve [15], *Helix* ganglions [16], *Xenopus* melanophores [17] and neuroblastoma cells [18, 19] respond to various odorants. For example, the sensitivity of the turtle trigeminal nerve to odorants is similar to that of the turtle olfactory neuron [15]. It is demonstrated by using Northern blot analysis that the expression of the members of the gene family encoding putative olfactory receptors is restricted to the olfactory epithelium, and that within the olfactory epithelium it may be either predominantly or exclusively expressed by olfactory sensory neurons [13]. Thus these observations suggest that these specific receptor proteins are not required for odor reception in these non-olfactory systems and there is an olfactory transduction pathway which is not mediated by the receptor proteins. For instance, it is reported that the bitter tastant, quinine, directly activates transducin and Gi/Go-proteins [20]. Bitter

tastants are generally hydrophobic as similar to odorants. There is a

possibility that some odorants penetrate the lipid layer of the olfactory receptor membranes and directly stimulate G-proteins.

Recently, Kashiwayanagi *et al.* reported that odorants, which did not increase IP3 but increased cAMP in olfactory cilia, elicited large responses after application of high concentrations of forskolin, IBMX or cpt-cAMP, suggesting that there exists a pathway independent of a second messenger in olfactory transduction [21].

Putative olfactory receptor proteins are surrounded by membrane lipids, and G-proteins are located inside of the chemosensory membrane. Supposing that an odorant directly interacts with certain Gprotein, the odorant has to traverse the lipid layers of the chemosensory membrane before contact with the G-protein. Odorants are generally hydrophobic and likely to interact directly with the lipid layers of the membrane. Thus it has been pointed out that interaction of odorants with lipid layers of olfactory receptor membranes may also play an important role in generation of olfactory responses even if olfactory transduction is mediated by either pathway including receptor proteins, or direct activation of G-proteins, or alternative pathways.

In the first Chapter of this study, we examine to what extent the olfactory system discriminates various odorants. It is well known that the olfactory system is exquisitely sensitive chemodetector recognizing odorants and discriminating among thousands of distinct odors. Although a great number of studies on odor discrimination have been carried out, there are few quantitative studies on the ability of olfactory

system to discriminate odorous compounds. In order to explore the mechanism of odor discrimination, it is necessary to examine to what extent the olfactory system discriminates various odorants. We measure the turtle olfactory bulbar responses to six pairs of highly pure optical isomers and compare the differences in odor intensity obtained between the optical isomers [22, 23]. The results indicate that there is no difference in odor intensity of optical isomers for all odorants examined. We also compare odor qualities of d- and l-isomers using a cross-adaptation method for quantitative analysis on odor quality [22, 23]. The results indicate that turtle olfactory system discriminates the odor quality difference between l- and d-isomers and that extent of discrimination of optical isomers varies among species of odorants. Recently, Hanada et al. reported that odor quality of d- and l-isomer is not discriminated at 40 °C [24], while it is well discriminated at 20 °C. This result suggests that odor is not recognized by binding of a ligand to a specific protein as seen in an interaction between enzyme and substrate or between a transmitter and its receptor. It is more likely that lipid layers of the olfactory receptor membrane are involved in the reception of odorant molecules.

In order to examine the role of lipids in the *in vivo* olfactory system, we treat the turtle olfactory epithelium with various lipids and examine its effects on the olfactory responses to various odorants [25-30] (Chapter II). The results indicate that the olfactory responses to

various odorants are greatly affected by the treatment of olfactory

epithelium with various lipids, confirming that lipids play an important role in the odor reception in the olfactory system.

The vomeronasal organ is a chemoreceptive structure situated at the base of the nasal septum of most terrestrial vertebrates and is anatomically and functionally distinct from the main olfactory system. In most terrestrial vertebrates, the vomeronasal system plays important roles in the perception of chemical stimuli related to feeding, social and reproductive behavior [31, 32]. These stimuli can induce hormonal changes, affect the success of pregnancy, alter the course of puberty, modulate female cyclicity and ovulation, elicit courtship and attraction, and modulate reproductive behavior and aggression [32]. Our understanding of the functional properties of vomeronasal system in vertebrates has been mainly deepened by behavioral, biochemical and molecular biological studies. In contrast, few studies on the electrophysiological properties of vomeronasal receptor neurons in vomeronasal systems have been carried out. Chapter III describes the electrophysiological properties of the turtle vomeronasal receptor neuron with whole-cell patch clamp [33].

Up to date, no study has demonstrated the signal transduction mechanism of vomeronasal receptor neurons. In order to ascertain the existence of signal transduction pathways mediated by putative second messengers, which are considered to be involved in the main olfactory transduction, in the vomeronasal system, cAMP, cGMP (Chapter IV-2)

and IP3 (Chapter IV-3) were injected into turtle vomeronasal receptor neurons under whole-cell patch clamp. The results described in this

Chapter show that each intracellular application of cAMP, cGMP and IP3 from the patch pipette to turtle vomeronasal receptor neurons elicits the membrane current under the condition of the whole-cell patch clamp, suggesting the existence of second messenger-mediated transduction pathways in the turtle vomeronasal system [33, 34].

Turtle vomeronasal organ senses general odorants as well as chemical stimuli of a social nature [35]. For instance, Hatanaka *et al.* and Shoji *et al.* recorded turtle accessory olfactory bulb wave induced by odor stimuli and showed that the vomeronasal organ responds to many kinds of odorants [37, 38].We examine to what extent cAMPmediated pathway found in this study contributes to the generation of the vomeronasal response to general odorants in *in vivo* system by measuring the turtle accessory olfactory bulbar response (Chapter V) [36]. The results described in this Chapter indicate that a general odorant such as citralva, which does not increase IP₃ but increase cAMP in olfactory cilia of the bullfrogs and the rats, elicits a large response after application of high concentrations of forskolin. This suggests that the cAMP-mediated pathway does not greatly contribute to the generation of the vomeronasal response to a general odorant.

and Transduction in Thiss and Officianian' in. by Brand, Jeff., Trainer, J. H., Cuppe R. F. and Kare, M. R., Mortel Dekker, New



REFERENCES

- 1) Boekhoff, I., Tarelius, E., Strotmann, J. and Breer, H. Rapid activation of alternative second messenger pathways in olfactory cilia from rats by different odorants. *EMBO J.*, **9:** 2453-2458, 1990
- Frings, S. and Lindemann, B. Current recording from sensory cilia of olfactory receptor cells *in situ*. I. The neuronal response to cyclic nucleotides. J. Gen. Physiol., 97: 1-16, 1990
- 3) Nakamura, T. and Gold, G. H. A cyclic nucleotide-gated conductance in olfactory receptor cilia. *Nature*, **325**: 442-444, 1987
- Pace, U., Hasnski, E., Salomon, Y. and Lancet, D. Odorant-sensitive adenylate cyclase may mediate olfactory reception. *Nature*, 316: 255-258, 1985
- 5) Sklar, P. B., Anholt, R. H. and Snyder, S. H. The odorant-sensitive adenylate cyclase of olfactory receptor cells: differential stimulation by distinct classes of odorants. J. Biol. Chem., 261: 15538-15543, 1986
- 6) Suzuki, N. Voltage- and cyclic nucleotide-gated currents in isolated olfactory receptor cells. "Chemical Senses, Vol. 1: Receptor Events and Transduction in Taste and Olfaction" ed. by Brand, J. G., Teeter, J. H., Cagan R. H. and Kare, M. R., Marcel Dekker, New Vork (1980) and ACO 404

York (1989) pp. 469-494.

- 7) Breer, H. and Boekhoff, I. Odorants of the same odor class activate different second messenger pathways. Chem. Senses, 16: 19-29, 1991
- 8) Kahn, A. A., Steiner, J. P. and Snyder, S. H. Plasma membrane inositol 1, 4, 5-triphosphate receptor of lymhpocytes: selective enrichment in sialic acid and unique binding specificity. Proc. Nal. Acad. Sci. USA., 89: 2849-2853, 1992
- 9) Kalinoski, D. L., Aldinger, S. B., Boyle, A. G., Huque, T., Marecek, J. F., Prestwich, G. D. and Resrepo, D. Characterization of a novel inositol 1, 4, 5-triphosphate receptor in isolated olfactory cilia. Biochem. J., 281: 449-456, 1992
- 10) Restrepo, D., Miyamoto, T., Bryant, B. P. and Teeter, J. H. Odor stimuli trigger influx of calcium into olfactory neurons of the channel catfish. Science, 249: 1166-1168, 1992
- 11) Restrepo, D., Teeter, J. H., Honda, E., Boyle, A. G., Marecek, J. F., Prestwich, G. D. and Kalinoski, D. L. Evidence for an InsP3gated channel protein in isolated rat olfactory cilia, Am. J. Physiol., **63:** C667-C673, 1992
- 12) Suzuki, N., IP3-activated ion channel activities in olfactory receptor neurons from different vertebrate species. "Olfaction and Taste XI" ed. by Kurihara, K., Suzuki, N. and Ogawa, H., Springer-Verlag, Tokyo (1994) pp. 173-177

13) Buck, L. and Axel, R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. Cell, 65: 175-187, 1991

- 14) Raning, K., Krieger, J., Strotmann, J., Boekhoff, I., Kubick, S., Baumstark, C. and Breer, H. Cloning and expression of odorant receptors. Nature, 361: 353-356, 1993
- 15) Tucker, D. Nonolfactory responses from the nasal cavity: Jacobsons' organ and the trigeminal system. "Handbook of Sensory Physiology, VI-1" ed. by Beidler, L. M., Springer-Verlag, Berlin (1971) pp. 151-181
- 16) Arvanitaki, A., Takeuchi, H. and Chalazonitis, N. Specific unitary osmereceptor potentials and spiking patterns from giant nerve cells. "Olfaction and Taste II" ed. by Hayashi, T., Pergamon Press, London (1967) pp.573-598
- 17) Lerner, M. R., Reagan, J., Gyorgyi, T. and Roby, A. Olfaction by melanophores: what does it mean? Proc. Natl. Acad. Sci. U.S.A., 85: 261-264, 1988
- 18) Kashiwayanagi, M. and Kurihara, K. Neuroblastoma cell as model for olfactory cell: mechanism of depolarization in response to various odorants. Brain Res., 293: 251-258, 1984
- 19) Kashiwayanagi, M. and Kurihara, K. Evidence for non-receptor olfactory cells. Brain Res., 359: 97-103, 1985
- 20) Naim, M., Seifert, R., Nurnberg, B., Grunbaum, L and Schultz, G. Some taste substances are direct activators of G-proteins. Biochem. J., 297: 451-454, 1994
- 21) Kashiwayanagi, M., Kawahara, H. Hanada, H. and Kurihara, K. A

large contribution of a cyclic AMP-independent pathway to turtle olfactory transduction. J. Gen. Physiol., 103: 957-974, 1994

- 22) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Intensity and quality of odors of enantiomers. Chemical Senses, 16: 205, 1990
- 23) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Quantitative analysis on odor intensity and quality of optical isomers in turtle olfactory system. Am. J. Physiol., 262: R99-R104, 1992
- 24) Hanada, T., Kashiwayanagi, M. and Kurihara, K. Temperature increase abolishes ability of turtle olfactory receptors to discriminate similar odorant. Am. J. Physiol., 266: R1816-R1823, 1994
- 25) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Enhancement of turtle olfactory responses to fatty acids by treatment of olfactory epithelium with phosphatidylserine. Brain Res., 647: 10-14, 1994
- 26) Taniguchi, M., Enomoto, S. and Kurihara, K. Large enhancement of turtle olfactory responses to fatty acids by treatment of the olfactory epithelium with phosphatidylserine liposomes. Chemical Senses, 17: 105, 1992
- 27) Taniguchi, M., Enomoto, S. and Kurihara, K. Enhancement of turtle olfactory responses to fatty acids by treatment of olfactory epothelium with phosphatidylserine "Olfaction and Taste XI" ed. by Kurihara, K., Suzuki, N. and Ogawa, H., Springer-Verlag, Tokyo (1994) pp. 152-153
- 28) Kurihara, K., Shoji, T., Enomoto, S., Taniguchi, M., Hanada, T. and Kashiwayanagi, M. Transduction mechanism in vertebrate

11

olfactory systems. "Chemical Signals in Vertebrate VI" ed. by Doty, R. L., Plenum, New York (1992) pp. 89-95

- 29) Enomoto, S., Shoji, T., Taniguchi, M. and Kurihara, K. Role of lipids of receptor membranes in odor reception. "Chemical Signals in Vertebrate VI" ed. by Doty, R. L., Plenum, New York (1992) pp.55-58
- 30) Shoji, T., Enomoto, S., Taniguchi, M. and Kurihara, K.
 Transduction mechanisms in turtle olfactory and vomeronasal organs. *Brain, Behavior and Evolution*, 41: 192-197, 1993
- 31) Halpern, M. The organization and function of the vomeronasal system. Ann. Rev. Neurosci., 10: 325-362, 1987
- 32) Wysocki, C. and Meredith, M. The vomeronasal system.
 "Neurobiology of Taste and Smell" ed. by Finger, T. E. and Silver,
 W. L., John Wiley & Sons, Inc., New York (1987) pp. 125-150
- 33) Taniguchi, M. Kashiwayanagi, M. and Kurihara, K., Intracellular injection of cyclic nucleotide monophosphate increases a conductance in membranes of turtle vomeronasal eceptor neurons in the slice preparation. in preparation
- 34) Taniguchi, M. Kashiwayanagi, M. and Kurihara, K., Intracellular injection of inositol 1, 4, 5-trisphosphate increases a conductance in membranes of turtle vomeronasal receptor neurons in the slice preparation. *Neurosci. Lett.*, in press
- 35) Tucker, D. Nonolfactory responses from the nasal cavity: Jacobson's organ and the trigeminal system. "Handbool of Sensory Physiology. Vol. IV-1" ed. by Bedler, L. M., Springer-Verlag, New York

12

(1971) pp.151-181

- 36) Taniguchi, M. Kanaki, K., Kashiwayanagi, M. and Kurihara, K. in preparation
- 37) Hatanaka, T., Shibuya, T. and Inouhchi, J. Induced wave responses of the accessory olfactory bulb to odorants in two species of turtle, *Pseudemys scripta* and *Geoclemys reevesii*. Comp. Biochem. *Physiol. A Comp. Physiol.*, 91: 377-385
- 38) Shoji, T. and Kurihara, K. Sensitivity and transduction mechanisms of responses to general odorants in turtle vomeronasal system. J. Gen. Physiol., 98: 909-919



LIST OF ABBREVIATIONS

PC; phosphatidylcholine from fresh egg yolk
PS; phosphatidylserine from bovine brain
PA; phosphatidic acid from egg yolk
CL; cardiolipin from bovine heart

Structure-helivity relationships, is odor reception have attracted a wide interest and been studied psychosperically by many investigators [1.2]. Among the relationships, whather there are differences in other quality and intensity between optical isomers has been discussed as one of the ornami topics.

PART 1 CHAPTER I DIFFERENCE IN ODOR INTENSITY AND QUALITY BETWEEN *l*-AND *d*-OPTICAL ISOMERS

I-1 Difference in Odor Intensity Between *l*- and *d*-Optical Isomers INTRODUCTION

Structure-activity relationships in odor reception have attracted a wide interest and been studied psychometrically by many investigators [1,2]. Among the relationships, whether there are differences in odor quality and intensity between optical isomers has been discussed as one of the central topics.

Laiterg *et al.* reported that threshold concentration of *l*-carvone is about 50 times lower than that of *d*-carvone [3]. Recently, Polak *et al.* reported that there are 78% of subjects whose threshold concentration of *l*-carvone was 3 to 14 times lower than that of *d*-carvone and 22% of subjects whose threshold concentration of *d*-carvone was 2 to 17 times lower than that of *l*-carvone [4]. They also reported that subjects'

relative sensitivities for a-ionone optical isomers were found to diverge widely, some subjects being much more sensitive to d- α -ionone than to

CHAFTS METWER METWER

INT ROI a wide in (1.2). At quality a of the ca is about I stopjects about I ower 5 l- α -ionone and vice versa [4]. The threshold of d-nootkatone was reported to be different from that of l-nootkatone [5], whereas there was a paper reporting no difference in odor intensity between them [6]. Thus the conclusions on the differences on odor intensity based on evaluation by human sense are ambiguous [1]. In addition, the lack of availability of pure optical isomers has hampered deduction of clear conclusions on odors of optical isomers.

In the present study [7,8], odor intensities of optical isomers are compared using highly pure odorants. Odor intensity is evaluated by measuring the summated olfactory bulbar responses in the turtle. The results clearly indicate that odor intensities of optical isomers are practically identical with each other for all odorants examined.

MATERIALS AND METHODS

Recording of olfactory bulbar response

Turtles (*Geoclemys reevesii*) weighing 160-240 g were used in the present study. Olfactory bulbar responses were recorded essentially as described by Kashiwayanagi *et al.* [9]. In brief, turtles were weakly anesthetized with the necessary and minimum amount of urethan to lessen pain in the operation of the animal, immobilized by an intramuscular injection of *d*-tubocurarine chloride (about 450 μ g/100 g

hipfed to the epidedium at a flow cur of 27 million deliver theb

body wt im), and locally anesthetized with lidocaine at the wounded and head-fixation points. The olfactory bulb was exposed using a dental

drill, and the dura mater on the olfactory bulb was removed carefully. To eliminate the possible effect of the accessory olfactory bulb activities [10], the nerve from the vomeronasal organ was cut off before entry to the accessory bulb. The stimulant-induced brain waves (bulbar responses) were recorded by attaching a pair of silver bipolar electrodes to the medial part of the anterior bulb. The responses were amplified and then integrated by electric integrator (time constant 0.3 s).

Chemical stimulation

The irrigating and stimulating solutions were applied to the olfactory epithelium through a stainless steel tube. Before application of the stimulating solution, the olfactory epithelium was irrigated with turtle Ringer solution for about 10 min. Stimulating solution, which was prepared by dissolving odorants in the turtle Ringer solution, was applied to the epithelium at a flow rate of 27 ml/min. After each application of the stimulating solution on the epithelium, the epithelium was rinsed with the Ringer solution. About 10 min were interposed between successive stimulations. Composition of the turtle Ringer solution was (in mM) 116 NaCl, 4 KCl, 2 CaCl2, and 0.5 Na2HPO4-NaH₂PO₄ (pH 7.4). All the experiments were carried out at 20 ± 3 °C.

Chemicals

Highly pure optical isomers of carvone [2-methyl-5-(1methylethenyl)-2-cyclohexene-1-one], ß-citronellol (3,7-dimethyl-6-

menthol $[(1\alpha, 2\beta, 5\alpha) - 5 - methyl - 2 - (1 - 1)]$ octen-1-ol), methylethyl)cyclohexanol], hydroxycitronellal (7-hydroxy-3,7dimethyloctanal), citronellal (3,7-dimethyl-6-octenal), and limonene [1methyl-4-(1-methylethenyl)cyclohexene] were kindly supplied from Takasago International (Tokyo, Japan). The purity of the optical isomers evaluated by gas chromatography was 99% for d-carvone, 98% for *l*-carvone, >99% for *d*- β -citronellol, >99% for *l*- β -citronellol, 99% for d-menthol, >99% for l-menthol, 99% for dhydroxycitronellal, 99% for l-hydroxycitronellal, 98% for dcitronellal, 99% for l-citronellal, 97% for d-limonene, and 96% for llimonene. All chemicals used are of best grade available. Concentrations of stock solutions used were 10-2 M for hydroxycitronellal, 10^{-3} M for carvone, β -citronellol, menthol and citronellal, and 10-4 M for limonene. These stock solutions were stored at +4 °C

RESULTS

Figure 1-1 shows the summated olfactory bulbar responses to land d-carvone of varying concentrations. The responses show a peak at onset of stimulation and decline rapidly during stimulation. The peak height of the summated bulbar response is taken as the magnitude of the

response. The magnitudes of the responses to both l- and d-carvone are increased with an increase in their concentrations.

18 .

oolenti methyleti dimethylet mothylet mothylet for /-eary for /-eary bydroxyd citronella fimonene bydroxyd bydroxyd

BESUL and d-con onset of Name 1.2 plats relative respirates of the responses to 1- and contrast knows to a freedom of oderatic concentration. Here the





Figure 1-1. Turtle olfactory bulbar responses to l-carvone (A) and d-carvone (B) of varying concentrations. Bars at the bottom of each record represent duration of application of stimulants.



Figure 1-2 plots relative magnitudes of the responses to l- and d-optical isomers as a function of odorant concentration. Here the magnitude of the response to a d-isomer of appropriate concentration is taken as unity for each dose-response curve. The responses start to appear between 10^{-9} and 10^{-7} M and increase in magnitude with an increase in odorant concentrations. The data shown in Figure 1-2 indicate that the magnitude of the response to a l- isomer is practically equal to that of the corresponding d-isomer in the whole concentration range examined.



is gure 1-2, Relative magnitude of partic bulker represente to desprictly somers (0) as a function of

odement concentration. Magnitude of scaporate is calculated relative to response to 10⁻⁴ M d carroom (A), 10⁻⁴ M d-carroom (B), 10⁻⁴ M d-carroom

A 1.5 1.0 E 0.5 l-carvone 0 d-carvone 0 -8 -6 -2 -4 log[carvone] (M) B 1.5 1.0 m 0.5 I-citronellol 0 d-citronellol 0 0 -6 -8 -4 -2 log[citronellol] (M)



odorant concentration. Magnitude of response is calculated relative to response to 10^{-4} M *d*-carvone (A), 10^{-4} M *β*-*d*-citronellol (B), 10^{-4} M *d*-menthol (C), 10^{-4} M *d*-hydroxycitronellal (D), 10^{-4} M *d*-citronellal (E), or 5×10^{-5} *d*-limonene (F), respectively. Each point is mean \pm S.E.M. of data obtained from at least 3 preparations.







DISCUSSION

Whether there are differences in odor intensity and quality between optical isomers has been discussed for a long time, but no systematic electrophysiological study has been carried out. In the present study, we have used the turtle olfactory system because of its high sensitivity to odorants; the turtle olfactory system is of comparable sensitivity to that of the dog [11]. For instance, by recording the olfactory neural responses of dog and turtle to aliphatic n-acetates and aliphatic n-fatty acids, Tonosaki and Tucker [12] reported that there was no practical difference between the thresholds of the dog and turtle olfactory responses to the odorants.

The present results clearly demonstrate that there is no difference in thresholds and intensities of the odor response between the optical isomers of all odorants examined.



REFERENCES

- 1) Beets, M. G. "Structure-Activity Relationships in Human Chemoreception." Applied Science, London (1978) p.127-148
- 2) Boelens, H. Structure-activity relationships in chemoreception by human olfaction. Trends Pharmacol. Sci., 4: 421-426, 1983
- 3) Leitereg, T. J., Guadagni, D. G., Harris, J., Mon, T. R. and Teranishi. R. Chemical and sensory data supporting the difference between the odors of the enantiomeric carvones. J. Agric. Food Chem., 19: 785-787, 1971
- 4) Polak, E. H., Fombon, A. M., Tilquin, C. and Punter P. H. Sensory evidence for olfactory receptors with opposite chiral selectivity. Behav. Brain Res., 31: 199-206, 1989
- 5) Haring, H. G., Rijkens, F., Boelens, H. and van der Gen, A. Olfactory studies on enantiomeric eremophilane sesquiterpenoids. J. Agric. Food Chem., 20: 1018-1021, 1972
- 6) Stevens, K. L., Guadagni, D. G. and Stern, D. J. Odour character and threshold values of nootkatone and related compounds. J. Sci. Food Agric., 21: 590-593, 1970
- 7) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Intensity and quality of odors of enantiomers. Chemical Senses, 16: 205, 1990

8) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Quantitative analysis on odor intensity and quality of optical isomers in turtle olfactory system. Am. J. Physiol., 262: R99-R104, 1992

- 9) Kashiwayanagi, M., Shoji, T., Taniguchi, M and Kurihara, K. Turtle olfactory bulbar responses to various chemical stimuli. *Neurosci. Prot.*, in press
- 10) Hatanaka, T., Shibuya, T. and Inouchi, J. Induced wave responses of the accessory olfactory bulb to odorants in two species of turtle, *Pseudemys scripta* and *Geoclemys reevesii*. Comp. Biochem. *Physiol. A Comp. Physiol.*, **91:** 377-385, 1988
- 11) Tonosaki. K. "Nioinokagaku" ed. by Takagi, S. and Shibuya, T.,Asakura Syoten, Japan (1989) pp. 139-144
- 12) Tonosaki, K. and Tucker, D. Olfactory receptor cell responses of dog and box turtle to aliphatic *n*-acetates and aliphatic *n*-fatty acids. *Behav. neur. Biol.*, 35: 187-199

bydroxychronellel [8]. However, the conclusions on the differences in odor quitity based on evaluation by homan sense are ambiguous [1,2]. The initial event of odor teception is adsorption of odorants on the instantor sites of olfariory cells. Odorents having different odor qualifier must be adort qualifier of various odorants, a cross-mispitalizadifferences in the adort qualifier of various odorants, a cross-mispitalizamethod is useful. Usually pross-adoptation experiments have been carried an with one part of chemical stands of found concententials in the restore in the part of chemical stands of found concententials in



I-2. Difference in Odor Quality Between *l*- and *d*-Optical Isomers

INTRODUCTION

MATERIALS AND METRON

Chapter I-1 dealt with difference in odor intensity between optical isomers. In this section, difference in odor quality between optical isomers is focused on. It was reported that odor quality of dcarvone is distinctly different from that of *l*-carvone [3,4]. The odor quality of d-nootkatone was reported to be different from that of *l*nootkatone [5]. The differences in odor qualities of optical isomers were also reported with limonene [6], β -citronellol [7], and hydroxycitronellal [8]. However, the conclusions on the differences in odor quality based on evaluation by human sense are ambiguous [1,2].

The initial event of odor reception is adsorption of odorants on the receptor sites of olfactory cells. Odorants having different odor qualities must be adsorbed in different receptor sites. To analyze differences in the odor qualities of various odorants, a cross-adaptation method is useful. Usually cross-adaptation experiments have been carried out with one pair of chemical stimuli of fixed concentrations in order to observe whether a response to the stimulus applied secondarily appears. Thus the usual cross-adaptation method gives only qualitative data. In the present study, odor quality of optical isomers is compared

27

Assing Assing

using highly pure odorants. To evaluate the difference in odor quality, we apply the quantitative cross-adaptation method [9,10] to the turtle olfactory system. The results indicate that the degree of cross adaptation greatly varies with species of odorants [11,12].

MATERIALS AND METHODS

Recording of olfactory bulbar response

Olfactory bulbar response was recorded essentially as described in Chapter I-1.

Chemical stimulation

The procedure of chemical stimulation was essentially similar to that described in Chapter I-1.

Chemicals

The sources for the reagents used except for cineole and the purity of the optical isomers evaluated by gas chromatography were described in Chapter I-1. Cineole (eucalyptol; 1,3,3-trimethyl-2-oxabicyclo[2.2.2]octane) was purchased from Nakarai Chemicals (Kyoto, Japan). The concentration of stock solution used were 10⁻³ M for cineole. The concentrations of stock solutions of other odorants

used were described in Chapter I-1. The stock solutions were stored at the same temperature as described in Chapter I-1.

using toghty we apply the officient syn praify varie

Accertance Reconstitute Chemical Chemical Chemical Chemical Chemical Chemical Chemical

RESULTS

Before the cross-adaptation experiment between optical isomers, the experiment between odorants having distinctly different odors was carried out as a control experiment. Figure 1-3A illustrates typical records for the cross-adaptation experiment between *l*-limonene having a lemonlike odor and cineole having a camphorlike odor. *l*-Limonene of varying concentrations is applied first, and then cineole of a fixed concentration (10⁻⁵ M) is applied after the response to *l*-limonene declines to a spontaneous level. As seen from the records, the magnitude of the response to cineole applied secondarily is decreased only slightly with an increase in concentration of *l*-limonene applied first.

Figure 1-3B plots the magnitude of the response to 10^{-5} M cineole (R₂) applied secondarily as a function of that to primarily applied *l*-limonene of varying concentrations (R₁). The reason why R₁ (magnitude of the response) is used instead of odorant concentration is as follows. In general, two different odorants of the same concentrations induce different magnitude of the response, and the magnitude of the response is a more valuable index than odorant concentration in the cross-adaptation experiment. The magnitudes of the responses (R₁ and R₂) are represented as relative magnitudes of the response to 10^{-5} M cineole applied alone is taken as

unity. In Figure 1-3B, $R_1 = 1.0$ means that the magnitude of the response to *l*-limonene is equal to that to cineole when each odorant is applied alone. Here the value of R_2 at $R_1 = 1.0$ is defined as the heterogeneity value. Hence, the heterogeneity value is some value between 0 and 1 where 1.0 implies that the receptor sites for two odorants are completely independent of each other, suggesting that the turtle olfactory system completely discriminates the quality of these two odorants. The heterogeneity value in the cross-adaptation between *l*-limonene and cineole is evaluated as 0.93, suggesting that the receptor site for cineole is almost independent of that for *l*-limonene.

Figure 1-4A illustrates typical records of the cross-adaptation experiment between *l*- and *d*-limonene. The response to $5 \ge 10^{-5}$ M *d*limonene is decreased with an increase in concentration of *l*-limonene applied first. Figure 1-4B delineates the relationship between the magnitudes of R₁ to *l*-limonene and R₂ to *d*-limonene. As shown in Fig. 1-4, the magnitude of the response to *d*-limonene is greatly decreased with an increase in that to *l*-limonene applied first. The heterogeneity value is evaluated as 0.26.

Figure 1.2. At Typical records of respinse to 10.7 M emeloic applied: reconstantly sites application of adorates Jopon hars, Filmeniane of varying consumminant hatchest bars, 10.5 M binacces; Ba Kelation







unity, in responses heplied a heterogen between site for b site for b

magnitude of response to 10^{-5} M cineole applied secondarily as a function of magnitude of response to *l*-limonene applied first. Magnitudes of response (R₁ and R₂) are calculated relative to response to 10^{-5} M cineole. Each point is mean \pm S.E.M. of data obtained from 4 preparations.


Figure 1-4. A: Typical records of response to 5×10^{-5} M *d*-limonene applied secondarily after application of *l*-limonene of varying concentrations applied first. Bars at the bottom of records represent duration of application of odorants (open bars, *l*-limonene of varying concentrations; closed bars, 5×10^{-5} M *d*-limonene). B: Relative magnitude of response to 5×10^{-5} M *d*-limonene applied secondarily as a function of magnitude of response to *l*-limonene applied first. Magnitudes of response (R₁ and R₂) are calculated relative to response to 5×10^{-5} M *d*-limonene. Each point is mean \pm S.E.M. of data obtained from 3 preparations.

We also carried out the cross-adaptation experiments on other pairs of optical isomers. Figure 1-5 shows the results of the crossadaptation experiments between d-carvone and l-carvone (A), between d- β -citronellol and l- β -citronellol (B), between d- menthol and lmenthol (C), between d-hydroxycitronellal and l-hydroxycitronellal (D), and between d-citronellal and l-citronellal (E). As shown in Fig. 1-5, in all cases, the magnitude of the response to a d-optical isomer applied secondarily is decreased with an increase in the magnitude of the response to the corresponding *l*-optical isomer applied first, but the degree of decrement in responses to d-optical isomers are different. For example, the response to d-carvone after l-carvone application decreases only slightly with an increase in the response to l-carvone (Fig. 1-5A), suggesting that the receptor site for d-carvone is greatly different from that for l-carvone. The case was the same with citronellol (Fig. 1-5B). On the other hand, the response to d-citronellal greatly decreases with an increase in the magnitude of the response to lcitronellal (Fig. 1-5E), suggesting that d- and l-citronellal mostly stimulates the same receptor site.

Table 1-1 shows the heterogeneity value between the l- optical isomer and the corresponding d-isomer. The values are distributed from 0.78 for carvone to 0.26 for limonene. Thus the degree of discrimination for optical isomers greatly varies with species of odorants



from 0.7 distribuli



Figure 1-5. A: Relative magnitude of olfactory responses to *d*-optical isomers applied secondarily as a function of magnitudes of responses to corresponding *l*-optical isomers applied first. Magnitudes of response (R₁ and R₂) are calculated relative to response to 10^{-4} M *d*-carvone (A), 10^{-4} M β -*d*-citronellol (B), 10^{-4} M *d*-menthol (C), 10^{-3} M *d*-hydroxycitronellal (D), or 10^{-4} M *d*-citronellal (E) applied alone, respectively. Each point is mean \pm S.E.M. of data obtained from 3 preparations.





l enirgié

ssponse arvone f-hydrox cspecify

0.5 1.0 1.5 R₁ (I-menthol)





Table 1-1. Heterogeneity value between <i>l</i> -optical isomer and corresponding <i>d</i> -optical isomer	
Odorant	Heterogeneity Value
Carvone	0.78
β-Citronellol	0.75
Menthol	0.52
Hydroxycitronellal	0.42
Citronellal	0.36
Limonene	0.26

Values are means \pm SE of data obtained from 3 preparations. See text for details.

the response to cincole applied secondarily doze not producally decrease with an increase in concentration of hitmonene applied first. Thus the furthe effectory system has an advantage over the free system on the pasts that nonspecific ishibition in the cross adaptation is very small. Is



DISCUSSION

Whether there are differences in odor intensity and quality between optical isomers has been discussed for a long time, but no systematic electrophysiological study has been carried out especially with regard to odor quality. In the previous cross-adaptation experiments using the frog olfactory system, the response to an odorant applied secondarily appreciably decreased with an increase in concentration of an odorant applied first even when odor qualities of two odorants used are quite different [10]. This is probably due to nonspecific inhibition of the response to odorant applied secondarily by odorant applied first because the frog olfactory system is not so sensitive to odorants and relatively high concentrations of odorants are used for the experiments. In the present study, we have used the turtle olfactory system because the turtle olfactory system is much more sensitive than the frog olfactory system. As seen in the combination of *l*-limonene and cineole whose odors are quite different from each other, the response to cineole applied secondarily dose not practically decrease with an increase in concentration of *l*-limonene applied first. Thus the turtle olfactory system has an advantage over the frog system on the basis that nonspecific inhibition in the cross adaptation is very small. In

the present study, we have measured the olfactory bulbar response to odorants. In general, information on odor discrimination originated from the receptor cells is considered to be much more emphasized in

the central nerve system. Hence, when one pair of odorants is discriminated by the olfactory bulb, the odorants must be discriminated first by the receptor cells, and information on the discrimination may be emphasized in the bulb. Thus the olfactory bulbar response is a good index especially when odor discrimination is examined.

Many investigators have compared odor quality and intensity of optical isomers and reported that there are differences in thresholds as well as odor quality between optical isomers [3-5]. The results shown in Chapter I-1 clearly demonstrated that there is no difference in thresholds and intensities of the odor response between the optical isomers of all odorants examined. On the other hand, the results of the cross-adaptation experiments shown in this section indicate that there are differences in odor qualities between optical isomers. Carvone is known to be one of the typical odorants whose optical isomers have different odors [3,4]. This is consistent with the present results. The magnitude of the difference in odor quality between optical isomers greatly varies with species of odorants. In contrast to carvone, the differences in the odor quality are rather small in citronellal and limonene. The present study has offered first data in quantitative difference in odor quality between optical isomers.

The data in the present study were obtained at 20 ± 3 °C. Recently, Hanada *et al.* [13] found that the ability of turtle olfactory system to discriminate odor quality of odorants having similar structure such as *d*-carvone and *l*-carvone, *trans*-3-hexenol and *cis*-3-hexenol, geraniol and nerol was greatly reduced by increasing temperature up to

about 40 °C, while the ability to discriminate the odorants having different structure such as *l*-limonene and cineole, anisole and cineole was not greatly affected by increasing temperature. As shown in the present study, the difference in odor quality between *d*-carvone and *l*-carvone at about 20 °C was highest among six pairs of optical isomers. Hence it seems that the turtle olfactory system does not discriminate odor quality between a *d*-optical isomer and a corresponding *l*-optical isomer at 40 °C in general.

There are a number of possible explanations for the results described above. There is a possibility that the temperature change induces a conformational change of a specific receptor protein for an odorants, which leads to a change in the specificity of the receptor site of the protein to the odorant. In general, the specificity of a protein is not, however, unchanged by a small temperature change from 20 to 40 °C. For example, the receptor protein for *l*-amino acid does not accept *d*-amino acid even when temperature increases up to 40 °C. Hence these phenomena are not simply explained in terms of a conformational change of receptor protein. There is another possible explanation for the abolishment of odor-discriminating ability by the temperature increase. In this case, it is supposed that odorants are adsorbed on hydrophobic pockets composed of lipids and proteins in the receptor membranes. It has been pointed out that olfactory thresholds are closely

related to partition coefficients of odorants between the organic solvent

and water [14], and interaction of odorants with lipid layers mimics in vitro odor reception [15, 16]. Hanada et al. [13] reported that the

membrane fluidity of cells isolated from turtle olfactory epithelia and liposomes made of lipids extracted form the epithelia changed in a similar temperature range as for the decrease of the odordiscriminating ability, suggesting that an increase in membrane fluidity is correlated with the abolishment of the odor-discriminating ability. These studies support the above idea. According to this assumption, the mechanism for the phenomena is as follows. At room and lower temperatures, the lipid structure is rather rigid and different odorants are adsorbed on different pockets. At a higher temperature (40 °C), the fluidity of the lipid layers increases in magnitude and then the pockets for the odorants become flexible. At this temperature, the receptor pocket for a d-optical isomer accepts a corresponding l-optical isomer and hence the receptor pocket for a d-optical isomer is desensitized by application of a corresponding l-optical isomer. In any case, the fact that the turtle olfactory system does not discriminate optical isomers at 40 °C supports an idea that the lipids in the olfactory receptor membrane play an important role in odor reception.

b) Friedman, R. and Willion, J. G. Udby moongrafty and chirality.

) Rhunneker, R. vini Ohloff, G. Ophich skilles h-fuosellol inte (v Codar (-)-pinan, Asyster, Chemiz 75: 243, 1961

Wiskertunetz, W., Giger, H. mid Oldoff (D. Darstefforg von (a)- med



REFERENCES

- 1) Beets, M. G. Structure-Activity Relationships in Human Chemoreception. Applied Science, (1978) p.127-148
- 2) Bowlines, H. Structure-activity relationships in chemoreception by human olfaction. Trends Pharmacol. Sci., 4: 421-426, 1983
- 3) Leitereg, T. J., Guadagni, D. G., Harris, J., Mon, T. R. and Teranishi, R. Chemical and sensory data supporting the difference between the odors of the enantiomeric carvones. J. Agric. Food Chem., 19: 785-787, 1971
- 4) Leitereg, T. J., Guadagni, D. G., Harris, J., Mon, T. R. and Teranishi, R. Evidence for the difference between the odours of the optical isomers (+)- and (-)-carvone. Nature, 230: 455-456, 1971
- 5) Haring, H. G., Rijkens, F., Boelens, H. and van der Gen, A. Olfactory studies on enantiomeric eremophilane sesquiterpenoids. J. Agric. Food Chem., 20: 1018-1021, 1972
- 6) Friedman, L. and Miller, J. G. Odor incongruity and chirality. Science Wash. DC, 172: 1044-1046, 1971
- 7) Rienacker, R. and Ohloff, G. Optisch aktives b-citronellol aus (+)odor (-)-pinan. Angew. Chem., 73: 240, 1961
- 8) Skorianetz, W., Giger, H. and Ohloff. G. Darstellung von (+)- und (-)-7-hydroxy-dihydrocitronellal aus (+)-pulegon. Ein beitrag zur

42

kenntnis olfaktorischer eigenschaften von enantiomeren. Helv.

Chim. Acta, 54: 1797-1801, 1971

- 9) Ohno, T., Yoshii, K. and Kurihara, K. Multiple receptor types for amino acids in the carp olfactory cells revealed by quantitative cross adaptation. *Brain Res.*, **310**: 13-21, 1984
- 10) Ohno, T., Yoshii, K. and Kurihara, K. Quantitative analysis on discrimination of various odorants at receptor sites of the frog olfactory cell revealed by a cross adaptation method. *Comp. Biochem. Physiol.*, 82: 153-159, 1985
- 11) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Intensity and quality of odors of enantiomers. *Chemical Senses*, **16**: 205, 1990
- 12) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Quantitative analysis on odor intensity and quality of optical isomers in turtle olfactory system. *Am. J. Physiol.*, **262**: R99-R104, 1992
- 13) Hanada T., Kashiwayanagi, M. and Kurihara, K. Temperture increase abolishes ability of turtle olfactory receptors to discriminate similar odorant. Am. J. Physiol., 35: R1816-1823
- 14) Davies, J. T. Olfactory theories. "Handbook of Sensory Physiology. Vol. IV-1" ed. by Beidler, L. M., Springer-Verlag, Berlin (1971) pp. 322-350
- 15) Koyama, N. and Kurihara, K. Effects of odorants on lipid monolyers from bovine olfactory epithelium. *Nature*, 236: 402-404, 1972
- 16) Nomura, T. and Kurihara, K. Liposomes as a model for olfactory cells: changes in membrane potential in response to various

43

odorants. Biochemistry, 26: 6135-6140, 1987

CHAPTER II

EFFECTS OF TREATMENT OF THE TURTLE OLFACTORY EPITHELIUM WITH VARIOUS LIPIDS ON OLFACTORY RESPONSES

INTRODUCTION

It is generally considered that an olfactory response is induced by binding of an odorant to specific receptor proteins in olfactory receptor membranes. Buck and Axel cloned the genes of an extremely large multigene family that encodes seven transmembrane domain proteins whose expression is restricted to the olfactory epithelium [1]. These proteins encoded by the genes are the most probable candidates for the receptor proteins.

On the other hand, as described in Chapter I-2, odor quality of d- and l-isomer is not discriminated at least at 40 °C [2], while it is well discriminated at 20 °C. This result suggests that lipid layers of the olfactory receptor membrane are also involved in the reception of odorant molecules.

It has been known that a number of non-olfactory systems such as *Tetrahymena* [3], fly [4], and frog [5] taste cells, turtle trigeminal nerves [6], *Helix* ganglions [7] and neuroblastoma cells [8, 9] respond to

various odorants. There is a close relation between minimum concentrations of odorants to induce the responses in these systems and

those to induce a membrane potential change in liposomes [10-12]. These results suggest that there exists a non-receptor mediated pathway in odor reception.

In addition, Enomoto et al. [12] reported that the sensitivity of liposomes to odorants varied with the species of odorant. They also reported that addition of phosphatidylserine (PS) to phosphatidylcholine (PC) liposomes greatly enhances membrane potential changes, which were monitored with a fluorescence dye, in response to odorants, especially fatty acids such as n-valeric acid, isovaleric acid and nbutyric acid [13]. In order to examine whether PS enhances the response in in vivo olfactory system, the turtle olfactory epithelium is treated with lipids, including PS, cardiolipin (CL) and phosphatidic acid (PA), and its effects on the olfactory responses to various odorants are examined [14, 15]. The results obtained indicate that the effects of the lipid-treatment on the olfactory responses vary with species of lipids used for treatment. Especially the PS-treatment greatly enhances the responses to fatty acids [14-19], which corresponds to the results obtained with PS-containing liposomes system. On the basis of these results obtained, possible mechanisms of odor reception and discrimination are discussed.

45

MATERIALS AND METHODS

Recording of olfactory bulbar response

Olfactory bulbar response was recorded essentially as described in Chapter I.

In general, the magnitude of the turtle olfactory bulbar responses to an odorant gradually decreases during the experiments. The response to *n*-amyl acetate was practically unchanged before and after the lipid treatment. Hence the response to *n*-amyl acetate was often measured during the experiment as a control response and the magnitude of the olfactory responses to various odorants was calculated relative to a response to 10^{-4} M *n*-amyl acetate. The data obtained were statistically analyzed by the Student's *t*-test.

Chemical stimulation

The chemical stimulation was carried out as described in Chapter I. Because the relative magnitude of the responses to odorants such as fatty acids, lilial and lyral were relatively smaller than those to the others used in the present study, there is a possibility that responses to these odorants are obscured by background noise. To obtain the sufficient signal-to- noise resolution, we used the odorants of 10⁻⁴ M in the experiments descried in Chapter II except for that described in Fig. 2-3. At this concentration, turtle olfactory system discriminated the odorants used in this chapter and the magnitude of the response to any odorant used did not reach a plateau level.

Treatment of the olfactory epithelium with lipids

The lipid suspension was prepared as follows. Chloroform solution of lipid in a round-bottom flask was evaporated to dryness using a rotary vacuum evaporator. After chloroform was evaporated completely, glass beads were added to the flask and the dried lipid film was dispersed with turtle Ringer solution of an appropriate volume by shaking with a Vortex mixer at room temperature. The final concentration of lipid suspension was 20 mg/ml for PS-, CL- or PAsuspension. Before treatment of olfactory epithelium with lipid suspension, the epithelium was irrigated with the Ringer solution for about 10 min. Lipid suspension in the Ringer solution was applied to the olfactory epithelium through a stainless steel tube in the same way as that for application of irrigating solution. To save the lipid suspension, the lipid suspension dripped down from the turtle internal nostril was collected into a chamber and again applied to the epithelium using a reflux pump (AC-2110, ATTO Co., Ltd., Tokyo, Japan). The epithelium was incubated with the lipid suspension for 1 hour and washed out with the Ringer solution for 15 min. Then the olfactory bulbar responses to odorants were measured.

Measurement of $[^{14}C]PS$ incorporated into olfactory epithelium

Figure 3-1 shows a schematic diagram illustrating procedure of the measurement of the amount of the [14C]PS incorporated into

olfactory epithelium. Cold PS (3.0 mg) and 0.03 mg of L-phosphatidyl-L-[3-14C]serine ([14C]PS, 1.96 GBq/mmol, Amersham, Japan) were

mixed and dissolved in chloroform solution in a flask was evaporated to dryness and 1.5 ml of Ringer solution was added to the flask. The PS suspension containing [14C]PS was prepared by a similar method to that described above. The turtle internal nostril was covered with the surgical bond, the nasal cavity was filled with about 0.25 ml of the PSsuspension containing [14C]PS. After incubation for 1 hour, the turtle internal nostril was opened again and the epithelium was washed out thoroughly with Ringer solution for 15 min. The turtle head was cut off along a median plane and the olfactory epithelium treated with PSsuspension containing [14C]PS was carefully excised from the bone supporting the epithelium. The excised epithelium was collected in a scintillation vial and dissolved in the scintillation solution containing Triton X-100 and toluene (1:1, v/v). The radioactivity of the solution was then measured with a liquid scintillation counter and ratio of radioactivity incorporated into the epithelium to that of the PS suspension used was calculated. The experiments were carried out with six preparations and the mean ratio was obtained.

Pigure 2-1. A disgram illustrating processare of the measurances of the simulat of (PCJPS incorporated date obtactory epithelium. See text fig





excision of epithelia



masking of the internal nostril







[¹⁴C]PS-treatment (for 1 hour)



dissolution of epithelia in the scintillation solution containing Triton X-100 and toluene





Figure 2-1. A diagram illustrating procedure of the measurement of the amount of [¹⁴C]PS incorporated into olfactory epithelium. See text for details.



Chemicals

PS (Lot 57F1355) and CL (Lot 60H8377) were purchased form Sigma Chemical Co. (St. Louis, MO, USA). PA was kindly supplied from Kao Corporation (Tokyo, Japan). Lilial (4-(1,1-dimethylethyl)-amethylbenzenepropanal), geraniol (3,7-dimethyl-2,6-octadien-1-ol), lyral (mixture of 4-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1carboxaldehyde and 3-(4-hydroxy-4-methylpentyl)-3-cyclohexene-1carboxaldehyde), citralva (3,7-dimethyl-2,6-octadienenitrile), citral (3,7-dimethyl-2,6-octenal), l-citronellal (3,7-dimethyl-6-octenal) and lmenthol [$(1\alpha, 2\beta, 5\alpha)$ -5-methyl-2-(1-methylethyl)cyclohexanol] were kindly supplied from Takasago International (Tokyo, Japan). n-Amyl acetate, sec-amyl acetate, n-amyl chloride, tert-amyl chloride, n-amyl alcohol, isobutyric acid, n-valeric acid, isovaleric acid, anisole (methoxybenzene), and Triton X-100 were purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan). n-Butyric acid, β -ionone (4-(2,6,6-trimethyl-1-cyclohexen-1-yl)-3-buten-2-one) and cineole (eucalyptol; 1,3,3-trimethyl-2-oxabicyclo[2.2.2]octane) were purchased from Nakarai Chemicals (Kyoto, Japan). All chemicals used are of best grade available. Concentrations of stock solutions used were 10-4 M for all odorants examined.

RESULTS



Effects of the treatment of the olfactory epithelium with various lipids on the turtle olfactory bulbar responses

The turtle olfactory epithelium was treated with PS-suspension and its effects on the olfactory bulbar responses were examined. Figure 2-2 shows typical records of the turtle olfactory bulbar responses before and after PS-treatment. The response to n-amyl acetate was practically not affected. In contrast, the response to n-valeric acid before the PS-treatment was much smaller than that to n-amyl acetate, but the response to n-valeric acid after the PS-treatment became remarkably large. The responses to other fatty acids were also greatly increased by the PS-treatment as will be described later.

Figure 2-3 plots relative magnitudes of the responses to *n*-valeric acid before and after PS-treatment as a function of its concentration. Here the magnitude of the response to 10^{-3} M *n*-valeric acid before PS-treatment is taken as unity. The data shown in Fig. 2-2 indicate that the PS-treatment lowers the threshold of the response to n-valeric acid by a factor about 50 and enhances the magnitude of the responses over the concentration range examined. The enhanced responses to fatty acids by the treatment returned to the original level about 10 hour after the PS-treatment (data not shown).





Figure 2-2. Typical records of the turtle olfactoy bulber responses before and after PS-treatment to *n*-amyl acetate and *n*-valeric acid. Bars under the records represent duration of stimurants (clased bars, 10^{-4} M *n*-amyl acetate; open bars, 10^{-4} M *n*-valeric acid).





Figure 2-3. Relative magnitudes of turtle bulbar responses to *n*-valeric acid before and after PS-treatment as a function of odorant concentration. Magnitude of response is calculated relative to the response to 10^{-3} M *n*-valeric acid before PS-treatment. Each point is mean \pm S.E.M. of data obtained from three preparations.



Figure 2-4 shows the effects of PS-treatment on the responses to various odorants. Here Ro and R represent the magnitude of the olfactory responses to an odorant before and after PS-treatment, respectively. The PS-treatment enhanced the responses to fatty acids such as n-valeric acid, isovaleric acid and n-butyric acid by a factor of 4-5. The responses to other odorants were a little increased or practically unchanged by the PS-treatment.

The olfactory epithelium was also treated with other lipids and its effects on the olfactory bulbar responses were observed. Figure 2-5 shows the ratio of the magnitudes of the responses before and after CLtreatment. The responses to responses to *n*-butyric acid, *n*-amyl alcohol and lyral were reduced and the response to cineole was slightly increased by the treatment.

Figure 2-6 shows the ratio of the magnitudes of the responses before and after PA-treatment. The responses to *n*-butyric acid and isovaleric acid were decreased by the treatment, while the responses to *n*-amyl acetate, *sec*-amyl acetate and anisole were not practically affected.

The results shown in Figs. 2-4, 2-5 and 2-6 indicate that the effect on the olfactory responses greatly varies with species of lipids used for the treatment of the olfactory epithelium.





Figure 2-4. Ratio (R/Ro) of the olfactory bulbar responses to various odorants (10⁻⁴ M) before and after PS-treatment. Ro and R represent the magnitude of the olfactory bulbar response to an odorant before and after PS-treatment, respectively. The responses to *n*-valeric acid was increased by the treatment in a level of P < 0.001 (*). The responses to *n*-butyric acid and isovaleric acid were increased by the treatment in a level of P < 0.05 (**). The responses to *l*-menthol was slightly increased by the

treatment in a level of P < 0.001 (*). The responses to β -ionone, cineole, *l*-citronellal and citralva are slightly increased in a level of P < 0.05 (**). Each point is mean \pm S.E.M. obtained from at least five preparations.



Figure 2-5. Ratio (R/Ro) of the olfactory bulbar responses to various odorants (10⁻⁴ M) before and after CL-treatment. Ro and R represent the magnitude of the olfactory bulbar response to an odorant before and after CL-treatment, respectively. The responses to n-amyl alcohol and lyral were decreased by the treatment in a level of P < 0.001 (*). The response to *n*-butyric acid was decreased by the treatment in a level of P < 0.05 (**). The response to cineole was slightly increased in a level of P < 0.05

0.001 (**). Each point is mean \pm S.E.M. obtained from at least six preparations.

scorpolition of ry live alfactory applied on the

elinetary responses to the famy solds. It indentify enhanced the statistic responses to the famy solds. It indentify enhanced the feet for the maximum infiniteportune into the office ry epidetium, do with him was incolated with PS expression essentiating 1. (CPS 1) four and washed out thereughly with the kinger solution for 15 can see Fig. 3-1). Among 0.51 mg PS used, L4 1.9.5 uncar h S.E.M., is a



Figure 2-6. Ratio (R/Ro) of the olfactory bulbar responses to various odorants (10⁻⁴ M) before and after PA-treatment. Ro and R represent the magnitude of the olfactory bulbar response to an odorant before and after PA-treatment, respectively. The responses to *n*-butyric acid and isovaleric acid were decreased by the treatment in a level of P < 0.05 (**). Each point is mean \pm S.E.M. obtained from six preparations.



Incorporation of PS into olfactory epithelium

As mentioned above, the PS-treatment greatly enhanced the olfactory responses to the fatty acids. In order to examine whether PS used for the treatment is incorporated into the olfactory epithelium, the epithelium was incubated with PS-suspension containing [14C]PS for 1 hour and washed out thoroughly with the Ringer solution for 15 min (see Fig. 3-1). Among 0.51 mg PS used, 1.4 ± 0.2 (mean \pm S.E.M., n = 6) μ g PS were incorporated into one nostril. That is , 0.28 \pm 0.04 (mean \pm S.E.M., n = 6)% of PS used for the treatment were incorporated into the olfactory epithelium.

DISCUSSION

The present results showed that the olfactory responses to the fatty acids, *n*-butyric acid, *n*-valeric acid and isovaleric acid were greatly enhanced by the PS-treatment, while the responses to other odorants examined were a little enhanced or unaffected by the treatment. The experiment using $[1^{4}C]PS$ suggests that PS was incorporated into the olfactory epithelium. It is uncertain in what part of the epithelium PS was incorporated, but there is a possibility that PS was incorporated into the olfactory receptor membranes [20, 21]. It is well known that interaction of PC, one of the phospholipids, with

erythrocytes eventually results in hemolysis [22, 23]. Tanaka et al. [21] reported that hemolysis occurred through incorporation of

dilauroylglyceroPC to human erythrocytes, supporting the above possibility in regard to demonstrating that phospholipids interact with animal cells by being incorporated into the plasma membranes. Hence a simple explanation for the present results is as follows. PS is incorporated into the olfactory receptor membranes and modifies the receptor site for the fatty acids so that affinity of the receptor membranes to the fatty acids is increased, which leads to enhancement of the olfactory response.

There is another possibility that PS is incorporated into the olfactory receptor membranes and increases the affinity of the receptor proteins to the fatty acids, which leads to enhancement of the olfactory response. It is, however, noted that addition of PS to PC liposomes greatly enhanced membrane potential changes in response to the fatty aids [13]. The fact that the enhancement of the olfactory responses to the fatty acids by PS-treatment closely resemble those observed with the liposomes supports the former mechanism.

As described in general introduction, Naim *et al.* [22] reported that quinine, the hydrophobic bitter tastant, directly activated transducin and Gi/Go-proteins [24], suggesting a possibility that some odorants penetrate the lipid layer of the olfactory receptor membranes and directly stimulate G-proteins. Therefore, it is possible to explain the mechanism of enhanced responses to fatty acids by PS-treatment as follows. PS incorporated into the lipid bilayer of the olfactory receptor

membranes and modifies the receptor membranes. As a result, affinity and permeability to fatty acids to lipid layers of receptor membranes



are increased, which facilitates the direct activation of G-protein by fatty acids so that the olfactory responses to them were enhanced.

In general, phospholipids are amphiphilic substances [23]. It is, therefore, conceivable that certain proteins in the olfactory receptor membranes are solubilized by detergent effect of PS, which leads to enhancement of the olfactory response. However, the enhancement of the responses to the fatty acids was not observed by the treatment with CL or PA which has a similar detergent effect to that of PS. Therefore, the above possibility may be excluded.

It is known that PS activates adenylate cyclase [26] or protein kinase C [27]. Hence there is another possibility that PS is incorporated into the olfactory receptor membranes and activates the second messenger systems, which leads to enhancement of the olfactory responses. The olfactory response to isovaleric acid, which dose not induce an elevation of the cAMP level but induces an elevation of the IP3 level [28], was greatly increased by the PS-treatment. The olfactory responses to geraniol, β -ionone and citronellal, which increase cAMP [29], were not much enhanced by the PS-treatment. It is the same case of the olfactory responses to lilial and lyral, which do not increase cAMP but increase IP3 as well as isovaleric acid [28]. Hence it is unlikely that the PS-treatment enhanced the responses to the odorants via the cAMP [26, 29] or IP3 second messenger cascade [27, 28].

The present results suggest an importance of lipid layers in odor

reception. This notion was also demonstrated as follows. As mentioned previously in Chapter I-2, Hanada *et al.* [2] reported that an increase of

temperature of the turtle olfactory epithelium up to 40 °C has little effect on the magnitude of odor intensity, but abolishes the ability of the olfactory receptor to discriminate odorants having similar odors such as d-carvone and l-carvone, trans-3-hexenol and cis-3-hexenol and n-amyl acetate and isoamyl acetate. The decrease of odor-discriminating ability was closely related to an increase of the membrane fluidity of lipid layers of the cells isolated from turtle olfactory epithelium.

The present results support the following hypothesis for odor reception. It is assumed that odorants are adsorbed on hydrophobic pocket composed of lipids and proteins in olfactory receptor membranes. The composition of the lipids and proteins of each olfactory cell membrane is assumed to vary from cell to cell. Variation in combinations of lipids and proteins provides many different adsorption sites for odorants. The qualities of odors are recognized by firing pattern among various olfactory axons and the quality of odors is recognized in the brain which summarizes these firing one. As shown in general introduction (Fig. I-1), numerous theories regarding the molecular basis of odorant recognition have been proposed. The present results together with the above observations strongly suggest that lipid layers as well as the receptor proteins play an important role in odor reception.

Tasts IF ad by Heyrshill Te Pergeinon Press London (1967) pp.



REFERENCES

- Buck, L. and Axel, R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell*, 65: 175-187, 1991
- Hanada, T., Kashiwayanagi, M. and Kurihara, K. Temperature increase abolishes ability of turtle olfactory receptors to discriminate similar odorant. Am. J. Physiol., 266: R1816-R1823, 1994
- Ueda, T. and Kobatake, Y. Hydrophobicity of biosurface as shown by chemoreceptive thresholds in *Tetrahymena*, *Physarum* and *Nitella.*. J. Membr. Biol., 34: 351-368, 1977
- 4) Dethier, V. G. Sensitivity of the contact chemoreceptors of the blowfly to vapors. *Proc. Natl. Acad. Sci. USA*, 69: 2189-2192, 1972
- Kashiwagura, T., Kamo, N., Kurihara, K. and Kobatake, Y., Responses of the frog gustatory receptors to various odorants. *Comp. Biochem. Physiol.*, 56C: 105-108, 1977
- 6) Tucker, D. "Handbook of Sensory Physiology, VI-1" ed. by Beidler,
 L. M., Springer-Verlag, Berlin (1971) pp. 151-181
- Arvanitaki, A., Takeuchi, H. and Chalazonitis, N. "Olfaction and Taste II" ed. by Hayashi, T., Pergamon Press, London (1967) pp. 573-598



- Kashiwayanagi, M. and Kurihara, K. Neuroblastoma cell as model for olfactory cell: mechanism of depolarization in response to various odorants. *Brain Res.*, 293: 251-258, 1984
- 9) Kashiwayanagi, M. and Kurihara, K. Evidence for non-receptor olfactory cells. *Brain Res.*, **359**: 97-103, 1985
- Nomura, T. and Kurihara, K. Liposomes as a model for olfactory cells: changes in membrane potential in response to various odorants. *Biochemistry*, 26: 6135-6140, 1987
- Nomura, T. and Kurihara, K. Effects of changed lipid composition on responses of liposomes to various odorants: possible mechanism of odor discrimination. *Biochemistry*, 26: 6141-6145, 1987
- 12) Enomoto, S., Kashiwayanagi, M. and Kurihara, K. Liposomes having high sensitivity to odorants. *Biochim. Biophys. Acta*, 1062: 7-12, 1991
- 13) Enomoto, S., Kawashima, S., Yoshimura, A. and Kurihara, K. Effects of changed lipid composition and addition of proteins on odor specificity of liposomes. *Sensors and Materials*, 4: 153-164, 1992
- 14) Taniguchi, M., Kashiwayanagi, M. and Kurihara, K. Enhancement of the turtle olfactory responses to fatty aacids by treatment of olfactory epithelium with phosphatidylserine. *Brain Res.*, 647: 10-14, 1994
- 15) Taniguchi, M., Enomoto, S. and Kurihara, K. Enhancement of

63

turtle olfactory responses to fatty acids by treatment of olfactory epithelium with phosphatidylserine. "Olfaction and Taste XI", ed.

by Kurihara, K., Suzuki, N. and Ogawa, H., Springer-Verlag, Tokyo (1994) pp. 152-153

- 16) Taniguchi, M., Enomoto, S. and Kurihara, K. Large enhancement of turtle olfactory responses to fatty acids by treatment of the olfactory epithelium with phosphatidylserine liposomes. *Chemical Senses*, 17: 105, 1992
- 17) Kurihara, K., Shoji, T., Enomoto, S., Taniguchi, M., Hanada, T. and Kashiwayanagi, M. Transduction mechanism in vertebrate olfactory systems. "*Chemical Signals in Vertebrate VI*," ed. by Doty, R. L., Plenum, New York (1992) pp. 89-95
- 18) Enomoto, S., Shoji, T., Taniguchi, M. and Kurihara, K. Role of lipids of receptor membranes in odor reception. "*Chemical Signals in Vertebrate VI*," ed. by Doty, R. L., Plenum, New York (1992) pp.55-58
- 19) Shoji, T., Enomoto, S., Taniguchi, M. and Kurihara, K. Transduction mechanisms in turtle olfactory and vomeronasal organs. *Brain, Behav. Evol.*, **41:** 192-197, 1993
- Bruckdorfer, K. R., Edwards, P. A. and Green, C. Properties of aqueous dispersions of phospholipid and cholesterol. *Eur. J. Biochem.*, 4: 506-511, 1968
- 21) Tanaka, Y., Inoue, K. and Nojima, S. Interaction of dilauroylglycerophosphocholine with erythrocytes. *Biochim. Biophys. Acta*, 600: 126-139, 1980

22) Kitagawa, T., Tanaka, Y., Inoue, K. and Nojima, S. Effects of

temperature and bovine serum albumin on lysis of erythrocytes



- induced by dilauroylglycerophosphocholine and didecanoylglycerophosphocholine. *Biochim. Biophysi. Acta*, **467**: 137-145, 1977)
- 23) Tanaka, Y., Mashino, K. and Inoue, K. Mechanism of human erythrocyte hemolysis induced by short-chain phosphatidylcholines and lysophosphatidylcholine. J. Biochem., 94: 833-840, 1983
- 24) Naim, M., Seifert, R., Nurnberg, B., Grunbaum, L and Schultz, G. Some taste substances are direct activators of G-proteins. *Biochem. J.*, 297: 451-454, 1994
- 25) Alberts, B., Bray, D., Lewis, J., Raff, M., Roberts, K. and Watson,
 J. D. Membrane structure. "Molecular Biology of The Cell" 3rd ed., Garland Publishing, Inc., New York (1994) pp. 447-506
- 26) Levey, G. S., Restoration of glucagon responsiveness of solubilized myocardial adenyl cyclase by phosphatidylserine. *Biochem. Biophys. Res. Commun.*, 43: 108-113, 1971
- 27) Nishizuka, Y. The role of protein kinase C in cell surface signal transduction and tumour promotion. *Nature*, **308**: 693-699, 1984
- 28) Breer, H. and Boekhoff, I., Odorants of the same odor class activate different second messenger pathways. *Chemical Senses*, 16: 19-29, 1991
- 29) Pace, U., Hanski, E., Salomon, Y. and Lancet, D., Odorant-sensitive adenylate cyclase may mediate olfactory reception, *Nature*, 316: 255-258, 1985



PART 2

CHAPTER III ELECTROPHYSIOLOGICAL PROPERTIES OF TURTLE VOMERONASAL RECEPTOR NEURONS

or classes meninery, where the strength of

INTRODUCTION

The nasal cavities of vertebrates are innervated by several nerves that may subserve different aspects of chemical sensation and perception (Fig. 3-1). A number of behavioral, biochemical, and molecular biological studies demonstrate that the vomeronasal system plays important roles in the perception of chemical stimuli related to animal behavior in most terrestrial vertebrates [1, 2]. Interaction between chemoattractants and vomeronasal receptor neurons results in depolarizing receptor potentials which activate voltage-gated channels and generate bursts of spikes. The voltage-gated currents underlying the excitatory response form an essential part of the vomeronasal transduction process. In spite of the exponential increase of interest in the structure and biological functions of the vomeronasal system, electrophysiological properties of vomeronasal receptor neurons have been little studied. Determining their electrical properties provides a basis for further analysis of molecular transduction mechanism in this

66

system.

The electrophysiological properties of the vomeronasal receptor neurons were examined only with the frog. Trotier *et al.* reported that the resting membrane potential of the frog vomeronasal neurons was near -60 mV and overshooting repetitive action potentials were elicited by an injection of depolarizing current pulses in the range of 2 to 10 pA, which were measured using patch-clamp technique [3]. They also reported that an transient fast inward current and an outward K⁺ current were activated in these neurons. No electrophysiological property of vomeronasal receptor neurons in other species of vertebrates has been described. The vomeronasal organ is well developed in reptiles [4] In the present study, the electrophysiological features of vomeronasal receptor neurons in the turtle were studied using the whole-cell patch clamp technique.

MATERIALS AND METHODS

Slice preparation of vomeronasal epithelium

Turtles, *Geoclemys reevesii*, weighing 140-240 g, were obtained from commercial suppliers and maintained at 22 °C. Turtles were cooled to 0 °C and decapitated. The nasal cavities were opened, and the vomeronasal neuroepithelia were dissected out carefully. The epithelia were cut into slices of about 120 μ m thickness with a vibrating slicer in

normal Ringer solution at 0 °C and stored at 4 °C. One slice of the epithelium was fixed on the glass at the bottom of a recording chamber.
The preparation was viewed under upright microscope (model OPTIPHOT, Nikon, Tokyo, Japan) using a x 40 water immersion lens.

Electron microscopy

The vomeronasal neuroepithelia and its slice preparation of 400 μ m thickness were prepared as described above. After quick rinse with Ringer solution, the specimens were placed in immediately in 5% glutaraldehyde/ 4% formaldehyde, i.e. Karnovsky's fixative [5] in 100 mM sodium cacodylate buffer (pH 7.4). The fixation was carried out at room temperature for at least one night. The tissues were post-fixed in 1% osmium tetroxide aqueous solution for 2 hours and dehydrated in a graded series of ethanol. The samples, after dehydration, were critical-point dried from CO₂ and sputter-coated with gold (in model HCP-2, Hitachi Koki Co., Ltd., Ibaraki, Japan and model IB-3, Eiko engineering, Ibaraki, Japan, respectively) and examined in a Hitachi S-430 scanning electron microscope (Hitachi, Ltd., Tokyo, Japan) at 20 kV.

Data recording and analysis

Patch pipettes with resistances of 5-10 M Ω were made from borosilicate glass capillaries using a two-stage electrode puller (model PP853, Narishige Co., Tokyo, Japan) and then fire-polished. Membrane currents were recorded in the whole-cell configuration (holding

potential, -70 mV). Data were continuously recorded using an EPC-7 patch clamp amplifier (List, Darmstadt, Germany) and stored on a

video cassette recorder via a digital audio processor. Data were filtered at 10 KHz and digitized at 10 KHz. Analysis was carried out on a personal computer using pCLAMP software (Axon Instruments, Foster City, CA, USA). All averages were given as mean \pm S.E.M.

Solutions

Normal Ringer solution consisted of (in mM): 116 NaCl, 4 KCl, 2 CaCl₂, 1 MgCl₂, 15 glucose, 5 Na-pyruvate, 10 HEPES-NaOH (pH 7.4). Patch pipettes were filled with a normal internal solution (in mM): 115 KCl, 2 MgCl₂, 10 HEPES-KOH (pH 7.6).

Chemicals

All chemicals used were of best grade available.

RESULTS

Cell morphology

A transverse section of the nasal cavity showed that the olfactory mucosa occupied the dorsal region and the vomeronasal one was located in the ventral region (Fig. 3-1). Three layers of supporting cells, receptor cells and basal cells could be distinguished in transverse section of the vomeronasal mucosa [6]. We examined electrical properties of

neurons located in a receptor cell layer in the slice, having bipolar or ovoid shape (Fig. 3-2). It is reported that vomeronasal receptor neurons





Fig. 3-1. The nasal cavity of the stink turtle. A: Schematic drawing of the sagittal section of a turtle nasal cavity corresponding to scanning micrograph. B: Low-magnification scanning micrograph of a sagittal section of turtle nasal cavity. $\times 37$. There is a ridge-like structure indicated by arrow heads in the middle portion of the cavity. This ridge-like structure entirely separates the vomeronasal mucosa from the olcactory mucosa. That is, the olfactory mucosa occupied the dorsalregion and the vomeronasal mucosa was located in the ventral region. These two types of mucosa could be easily distinguished from their surface structure. EN: external nares. IN: internal nares. OM: olfactory mucosa. VM: vomeronasal mucosa.





Figure 3-2. Lateral aspect of vomeronasal epithelium (A) and detailed view of microvilli at terminals of dendrites (B). It could be seen that receptor neuron dendrite extended from round soma to the mucosal surface through the supporting cell layer, and its terminal was little swelled. The higher-magnification scanning micrograph (B) taken from the same field as that in A clealy demonstrates that the terminals of the receptor neuron dendrites possess a number of microvilli. The magnification of the micrograph are $\times 2000$ for A and $\times 8000$ for B, respectively. D: receptor neuron dendrite. M: microvilli of the receptor neuron. S: soma of the receptor neuron.



are bipolar ones and their dendrites lack cilia and possess microvilli [7, 8]. The microvilli could not be identified with the optics used for viewing the electrophysiological experiments because they were only 100 nm in diameter [8]. But several microvilli (about 100 nm in diameter) extending from the terminal ends of the dendrite of receptor neurons could be viewed with scanning electron microscope (Fig. 3-2). In whole-cell voltage-clamp [9], the vomeronasal receptor neurons were further identified by the activation of a transient inward current followed by an outward current in response to step depolarization as will be shown below.

Resting potential

With normal internal solution in the pipette, turtle vomeronasal receptor neurons maintained resting potentials ranging from -33.0 to -71.5 mV (-48.1 \pm 1.3 mV, n = 14). The input resistance was measured from the responses to injected currents of 1 sec ranging from -20 to +20 pA in 20 mV increments applied at the holding potential of -70 mV, and ranged from 0.7 to 2.8 G Ω (1.7 ± 0.1 G Ω , n = 14).

Voltage responses to injected current

In current-clamp recordings, step depolarization elicited by 7 pA stimulus current from a holding potential of about -70 mV produced an action potential (Fig. 3-3A) which had a relatively smooth

rising phase. Twenty five of 30 (80%) neurons fired one to several action potentials in response to current steps of less than 30 pA form a



conditioning potential of about -70 mV. The threshold for action potential generation in vomeronasal receptor neurons was commonly between -45 and -55 mV. A variety of spiking patterns were seen, ranging from neurons that fired only a single action potential for any suprathreshold stimulus (data not shown) to neurons that generated brief trains of action potentials (Fig. 3-3B, for instance). In the example shown in Fig. 3-3B, the action potentials were generated repetitively in response to a depolarizing current pulse of 23 pA.

Several neurons required current injection of only 3 pA to depolarize to spike threshold (data not shown), suggesting that a vomeronasal receptor neuron in turtle has high sensitivity to injected currents similar to the frog vomeronasal receptor neurons [3]. Because the membrane potentials varied from neuron to neuron, the membrane potentials of neurons examined were held at -70 mV to measure the voltage response to injected current under the same experimental condition. Thus, spike threshold was increased and spike amplitude was decreased when the membrane was equal to the resting potential (data not shown).

regure (a.). Electrical despondent of a vorticitient interprot metron. As Voltage response to current super between -5 and 7 pA for 1 pA increments). Realing least of this neuron was -70 paV and thendoold was near -55 mV. It: Voltage responde of the same rearrin to a current step of 23 pA. In: every case where the action potentials were generated repetitively, the superval between the few spikes by reased with each subsequent spike (u=3). Each bounded takes shows the concepteding current outer.







Figure 3-3. Electrical responses of a vomeronasal receptor neuron. A: Voltage response to current steps between -5 and 7 pA (in 1 pA increments). Resting level of this neuron was -70 mV and threshold was near -55 mV. B: Voltage response of the same neuron to a current step of 23 pA. In every case where the action potentials were generated repetitively, the interval between the few spikes increased with each subsequent spike (n=3). Each bottom traces shows the corresponding current pulse.



Whole-cell current

Figure 3-4A shows the two major currents elicited by depolarizing steps of voltage from a holding potential of -70 mV. A transient inward current activated between -60 and -40 mV became larger, faster, and more transient with more depolarizing steps reaching up to 1 nA at -20 mV. With further depolarization to 0 mV, this current diminished but was followed in time by a more slowly developing sustained outward current. The current-voltage curves taken at the peak of the inward current and during the sustained portion of the outward current are shown in Fig. 3-4B. Outward currents are activated near -40 mV and display slight inactivation during a 60-ms step.

Figure 3-4. Typical whole-cell currents in an concremental receptor optimon in a since preparation. At Responses to voltage steps. Step levels are shown in the bottom traces. Transient lowerd and delayed out ward currents were elicited in response to 60 ms voltage steps between -100 and 60 mV in 20 mV increments from a bolding potential of 70 mV. By Current-voltage relationsticies from a bolding potential of 70 mV. By after the onset of the voltage step during the austrined pintesa of the outward current. (6) measured from the records in V. The gipeno





Figure 3-4. Typical whole-cell currents in an vomeronasal receptor neuron in a slice preparation. A: Responses to voltage steps. Step levels are shown in the bottom traces. Transient inward and delayed out ward currents were elicited in response to 60 ms voltage steps between -100 and 60 mV in 20 mV increments from a holding potential of -70 mV. B: Current-voltage relationships of peak inward currents (O) and 60 ms after the onset of the voltage step during the sustained plateau of the outward current () measured from the records in A. The pipette contained normal internal solution and the bath contained Ringer solution.



DISCUSSION

Both vomeronasal receptor neurons and olfactory ones have functions to detect odorant and have a similar cell size. In the following, we compare the electric properties of both neurons.

Resting potential and input resistance

While the resting potential of isolated vomeronasal neurons of frog was about -61 mV [3], the mean resting level of turtle vomeronasal receptor neurons was -48 mV. In olfactory neurons of frog [10], salamander [11], newt [12] and rat [13], the resting potentials were -40 mV, -55 mV, -44 mV and -48 mV, respectively. Thus, it appears that turtle vomeronasal receptor neurons have a similar resting potential to those of sensory neurons of other species.

Input resistance of the receptor neurons which were observed in this study was near 1.7 G Ω . This value is slightly lower than that of olfactory neurons which was reported to be near 2-10 G Ω , but is not greatly different.

Voltage activated properties

Present results showed that the transient inward currents activated between -60 to -40 mV reached a peak at about -20 mV and

inactivated rapidly (Fig. 3-4). These properties are similar to those measured in isolated rat olfactory neurons [13], rat olfactory neurons in

culture [14], salamander olfactory neurons [11] and frog vomeronasal receptor neurons [3], but differ from the properties of acutely isolated rat olfactory neurons where the sodium currents were activated at more negative potentials [14]. Outward currents were activated between -50 and -40 mV and displayed only slight inactivation during at 60-ms step. These properties are similar to those measured in isolated rat olfactory neurons [12-16]. A detailed analysis of these voltage activated currents was not carried out in the present study, but the transient inward currents and the outward currents were tentatively identified as sodium and potassium currents, respectively since the properties of there currents were similar to those of sodium and potassium currents observed in many neurons [17].

Voltage response and spike activity

Because of the high input resistance of these neurons, only 3 pA of injected current was required to reach spike threshold. This is similar to the case of rat olfactory neurons [18] and frog vomeronasal neurons [3]. The olfactory system is known to be an exquisitely sensitive chemodetector recognizing odorants at concentrations as low as the level of a few nM. It is desirable for sensory neurons to generate spikes by minimum membrane depolarization. That is, the smaller membrane depolarization of sensory neurons generates action potentials, the more sensitively the neurons can detect odorants.



As the results described above, turtle vomeronasal receptor neurons appear to be similar to olfactory neurons and frog vomeronasal receptor neurons in regard to their passive electrical characteristics being electrotonically compact and possessing a low spike threshold. They are also homogeneous in regard to their spike responses and the gated currents underlying those responses.

- W. L., John Willey & Sides, Inc., New York (1997) pp. 125-150
 Trotter, D.; D'Orang, K. B. and Rosin, F.F. Village dependen currents in million feedback receptor, cells or the freq vomeronau bream En. 7. Main and Science and a multiplication of the set of the s
- Touosaki, K., Nioinollagaka, ed. by Taking, S. and Shibaya, T. Azakura System, Japan (1989) pp. 41-49
- Kemovsky, M. J. A formaldoliyde glataradehyde fizzawe of high osmolarity for use in electron microscopy. J. Cell Riol., 27: 137-138, 1965
- 6) Habsunka, T. Structure of the conformation system and induced wave in the necessory offactory built on real cased on the Proc. 20th Tep Sympox. Tasks and Smell, 183-186, 1986.
- 7) Graziatlei, P. P. C., bod Turker, D. Vomeronazal receptors in turtles, Zelsschrift für Zellferseitung, 108: 493-514, 1970
- 8) Hatanaka,T., Matsuzoki, O. and Shibuya, T. Fhis structure of vomeronasal receptor calls in the Reevers tastle, Geoclemys



REFERENCES

recording, from, calls and cell-free membership perchase P/12/19/22

- 1) Halpern, M. The organization and function of the vomeronasal system. Ann. Rev. Neurosci., 10: 325-362, 1987
- 2) Wysocki, C. and Meredith, M. The vomeronasal system.
 "Neurobiology of Taste and Smell" ed. by Finger, T. E. and Silver,
 W. L., John Wiley & Sons, Inc., New York (1987) pp. 125-150
- Trotier, D., DΦving, K. B. and Rosin, J-F. Voltage-dependent currents in microvillar receptor cells of the frog vomeronasal organ, Eur. J. Neurosci., 5: 995-1002, 1993
- Tonosaki. K. "Nioinokagaku" ed. by Takagi, S. and Shibuya, T., Asakura Syoten, Japan (1989) pp. 41-49
- Karnovsky, M. J. A formaldehyde-glutaraldehyde fixative of high osmolarity for use in electron microscopy. J. Cell Biol., 27: 137-138, 1965
- 6) Hatanaka, T. Structure of the vomeronasal system and induced wave in the accessory olfactory bulb or red eared turtle. Proc. 20th Jap. Sympos. Taste and Smell, 183-186, 1986
- 7) Graziadei, P. P. C., and Tucker, D. Vomeronasal receptors in turtles. Zeitschrift für Zellforschung, 105: 498-514, 1970
- 8) Hatanaka, T., Matsuzaki, O. and Shibuya, T. Fine structure of vomeronasal receptor cells in the Reeve's turtle, Geoclemys

80

reevesii. Zoological Magazine, 91: 190-193, 1982

- 9) Hamill, O. P., Neher, M. E. Sakmann, B. and Sigworth, F. J. Improved patch-clamp techniques for high-resolution current recording from cells and cell-free membrane patches. *Pflügers Arch.*, **391**: 85-100, 1981
- Trotier, D. A patch-clamp analysis of membrane currents in salamander olfactory receptor cells. *Pflügers Arch.*, 407: 589-595, 1986
- 11) Firestein, S. and Werblin, F. S. Gated currents in isolated olfactory receptor neurons of the larval tiger salamander. *Proc. Natl. Acad. Sci. USA*, 84: 6292-6296, 1987
- 12) Kurahashi, T. and Shibuya, T. Ca²⁺-dependent adaptive properties in the solitary olfactory receptor cell of the newt. *Brain Res.*, 515: 261-268, 1990
- 13) Okada, Y., Teeter, J. H. and Restrepo D. Inositol 1, 4, 5trisphosphate-gated conductance in isolated rat olfactory neurons. J. Neurophysiol., 71: 595-602, 1994
- 14) Trombley, P. Q. and Westbrook, G. L. Voltage-gated currents in identified rat olfactory receptor neuron. J. Neurosci., 11: 435-444, 1991
- 15) Rajendra, S. Lynch, J. W. and Barry, P. H. An analysis of Na+ currents in rat olfactory receptor neurons. *Pflügers Arch.*, 420: 342-346, 1992
- 16) Lynch, J. W. and Barry, P. H. Properties of transient K+ currents

81

and underlying single K+ channels in rat olfactory receptor neurons. J. Gen. Physiol., 97: 1043-1072, 1991

- 17) Hille B. "Ionic channels of excitable Membranes" 2nd ed., Sinauer Associates Inc., Sunderland, Massachusetts (1992) pp.59-82
- 18) Lynch, J. W. and Barry, P. H. Action potentials initiated by single channels opening in a small neuron (rat olfactory receptor). Biophys. J., 55: 755-768, 1989

that offactory responses are triggered via the G-protein-linked cAMP pathway (1-6) or IPs dependent pathway (7-12). Hence, there is the possibility that the second messenger-nordiated pathway is involved in the signal translating system in the comercises system.

Recently, in the hybridization experiment showed that some tensory neurons in the neuro-vomeroussal organ expressed offictory G-protein coupled exector nENA [11]. This observation also suggests that the vomerones d system shares a similar transduction pathway to that the officiery system in addition, has at an reported the Gproteins (Gs. Gi and Go) were tentencely identified in the vomeronesal issue membrine preparations of gener snakes using immunove activity and ADP-ribosylation techniques (14). These G proteins were suggested to be coupled with the receptor for ES20 [14] which was the formestimetian for the snakes [15].

In spite of these interesting studies, no electrophysiological study on the signal transduction mechanism of vomercumsi receptor ocurous has been carried out. The most suitable candidates for second

messengers of someronasal transduction are considered to be called

CHAPTER IV SIGNAL TRANSDUCTION MECHANISM IN VOMERONASAL RECEPTOR NEURONS

INTRODUCTION

In vertebrate olfactory transduction, it is generally considered that olfactory responses are triggered via the G-protein-linked cAMP pathway [1-6] or IP3-dependent pathway [7-12]. Hence, there is the possibility that the second messenger-mediated pathway is involved in the signal transducing system in the vomeronasal system.

Recently, *in situ* hybridization experiment showed that some sensory neurons in the mouse vomeronasal organ expressed olfactory G-protein coupled receptor mRNA [13]. This observation also suggests that the vomeronasal system shares a similar transduction pathway to that in the olfactory system. In addition, Luo *et al.* reported that Gproteins (Gs, Gi and Go) were tentatively identified in the vomeronasal tissue membrane preparations of garter snakes using immunoreactivity and ADP-ribosylation techniques [14]. These G-proteins were suggested to be coupled with the receptor for ES20 [14] which was the chemoattractant for the snakes [15].

In spite of these interesting studies, no electrophysiological study on the signal transduction mechanism of vomeronasal receptor neurons has been carried out. The most suitable candidates for second

messengers of vomeronasal transduction are considered to be cAMP and IP3. In Section IV-1 and IV-2, we describe the responses induced

by intracellular injection and cAMP or IP3 into the turtle vomeronasal receptor neurons, respectively.

IV-1 The Responses Induced by Intracellular Injection of cyclic Nucleotides into Turtle Vomeronasal Receptor Neurons

INTRODUCTION

Luo et al. [14] reported that both GTPYS and forskolin increased the cAMP level in vomeronasal receptor neurons of garter snakes. These observations suggest that there exists cAMP-dependent pathway in signal transduction in the vomeronasal receptor neurons.

In spite of this interesting observation, any study has demonstrated the existence of cAMP-activated membrane conductance in the vomeronasal system. In the present study, we injected cAMP into the turtle vomeronasal receptor neurons under whole-cell patch clamp and found that intracellular application of cAMP elicits the membrane current.

In the olfactory neurons, intracellular injection of cGMP also elicits inward current [16]. The studies using patches of ciliary membrane have clearly shown that both cGMP and cAMP directly activate a ciliary membrane conductance [3, 6, 17, 18]. In the present study [19], we found that intracellular application of cGMP from the patch pipette to turtle vomeronasal receptor neurons also elicits the

84

membrane current under the whole-cell patch clamp.

MATERIALS AND METHODS

Preparations

The slice preparations were obtained as described in Chapter III. Data recording and analysis

Membrane currents were recorded in the cell attached or wholecell configurations of the patch clamp as described in Chapter III. In the whole-cell recordings, holding potential was -70 mV. Data were analyzed as described Chapter III.

Solutions

The compositions of normal Ringer solution and normal internal solution were the same as described in Chapter III. For the stimulation with adenosine 3' : 5'-cyclic monophosphate (cAMP) and guanosine 3' : 5'-cyclic monophosphate (cGMP), each cyclic nucleotide was dissolved in the internal solution to give the desired final concentrations. The stock solutions of cAMP and cGMP at appropriate concentrations derived into 1.8 ml aliquots were stored at -80 °C and thawed just prior to use. Stocked forskolin solution was prepared by dissolving it in ethanol at 10 mM and appropriate volume was added to Ringer solution to give desired concentrations. These forskolin solutions were prepared

daily. The final concentration of ethanol never exceeded 0.5%. This

concentration of ethanol alone had no measurable effect on the electrical property of the neurons.

Gravity was used to deliver a constant stream of Ringer solution from the stimulating tube. Three electrically actuated valves were used to switch adapting Ringer solution and stimulating solutions. The stimulating tube with a lumen 160-200 μ m in diameter was placed under visual control within about 500 μ m of the neuron. The concentrations of stimuli were reported as the pipette concentration; no attempt was made to correct for dilution.

Chemicals

cAMP, forskolin and cGMP were purchased from Boehringer Mannheim GmbH (Mannheim, FRG), Wako Pure Chemical Industries Ltd. (Osaka, Japan) and Yamasa Shoyu Co. Ltd. (Choshi, Japan), respectively. All chemicals used were of best grade available.

RESULTS

As described in Chapter III in details, we examined electrical properties of neurons located in a receptor cell layer in the slice, having bipolar or ovoid shape. The vomeronasal receptor neurons were further identified by the activation of a transient inward current

followed by an outward current in response to depolarizing voltage steps from a holding potential of -70 mV (Fig. 3-4).

Transient inward current induced by cAMP

In order to examine whether cAMP-mediated pathway exists in the vomeronasal receptor neurons, the response to forskolin, a direct activator of adenylate cyclase, was recorded from an vomeronasal receptor neuron of the turtle using the cell attached configuration (Fig. 4-1). Before application of the 10 μ M forskolin, the vomeronasal receptor neurons generate spikes spontaneously. Bath application of 10 μ M forskolin caused a remarkable increase in the spike rate, suggesting the possibility that there exists adenylate cyclase and cAMP-dependent ion channels in the vomeronasal receptor neurons.

To explore the existence of cAMP-mediated pathway in vomeronasal receptor neurons directly, cAMP was introduced into a proximal part of the dendrite or a part of cell soma by whole-cell dialysis. Figure 4-2 shows the currents induced by intracellular injection of cAMP of varying concentrations into vomeronasal receptor neurons.

When the pipette was filled with an cAMP-free inner solution, the neurons held a steady baseline over the test interval of about 3-10 min after membrane rupture (left trace). On the other hand, introducing cAMP into the neurons evoked prolonged, inward currents within a few seconds after membrane rupture. The magnitudes of the responses to cAMP introduced intracellularly were increased with an

increase in their concentrations.

-	Concernent about minden and because Barris and an an and the second statements a second statement of the second statements	
. Mar is a second such	MUNICIPAL AND ADDRESS OF A DESCRIPTION OF A DESCRIPTION OF A DESCRIPTION DESCRIPTION OF A D	
	in an hill die bein eine andere werd. Die bei der die die geschen der bein ein beiner mit die der Berrichten d	
	forokolin	
	ATOrskollin	
Harry H	I that all we had a later to a second and the second stands by the second	
Long description		
	1000	

Figure 4-1. Cell-attached extracellular recordings of a single turtle vomeronasal receptor neuron *in situ*. An arrow-head represents time of 10 μ M forskolin onset.



In the present study, 30 manual ware searchafold, stimulant by mM cAMP. Thenty one neurons (70%) displayed an increme in inward current with adaptation of susrent after the peak response. The emplitude of the inward current induced by cAMP varied from 0 in 756 pA (176 z. 34 pA, mean z S; E, M, m(30), in some neurons, the cAMP-induced current was not adapted. The data obtained from these searches were excluded because b was united, whether the current



Figure 4-2. Response induced by intracellular application of cAMP from the patch pipette to a vomeronasal receptor neuron bathed in normal Ringer solution. The concentrations of cAMP contained in the pipette are shown at the top of each trace. Holding potential, -70 mV.

ingle dose to each overest, cumulative results from a group of neurone

indicated the dose-denshdence of the response magnitude. The currents

nercase is a cAMP concentration-dependent manner, which reach the

In the present study, 30 neurons were successfully stimulated by 1 mM cAMP. Twenty one neurons (70%) displayed an increase in inward current with adaptation of current after the peak response. The amplitude of the inward current induced by cAMP varied from 0 to 756 pA (176 \pm 34 pA, mean \pm S. E. M., n=30). In some neurons, the cAMP-induced current was not adapted. The data obtained from these neurons were excluded because it was unclear whether the current observed represented an inward current induced by cAMP or an artificial leak.

Kashiwayanagi *et al.* [20] applied 1 mM cAMP into the turtle olfactory neuron and observed that cAMP induced an inward current with an average peak amplitude of 252 ± 30 pA (n=31). The magnitude of the response induced by cAMP in the turtle vomeronasal receptor neurons is smaller than that in the turtle olfactory neurons. The time to peak for the response of the vomeronasal neuron ranged from 5 to 102 s (28 ± 5 s, mean \pm S. E. M., n=30). This time was slower than that for cAMP-induced response of the newt olfactory neurons (near 4 s) [16], but similar to that of turtle olfactory neurons (26 ± 4 s; mean \pm S.E.M, n=22) [20].

Figure 4-3 plots the magnitudes of the responses induced by intracellular application of cAMP as a function of cAMP concentration. Although the present method of nucleotide application allowed only a single dose to each neuron, cumulative results from a group of neurons

indicated the dose-dependence of the response magnitude. The currents

increase in a cAMP concentration-dependent manner, which reach the

maximum level at 1 mM cAMP. It should be noted that 0.1 mM cAMP can elicit the inward currents. The sensitivity of the turtle vomeronasal receptor neurons to cAMP applied intracellularly is similar to that of isolated olfactory neurons of the newt [16].

The current-voltage relationships were examined by applying a voltage ramp either from -70 to +50 mV (480 mV/s) or from -100 to +60 mV (43.7 mV/s) to voltage-clamped vomeronasal neurons before, during and after the response induced by cAMP (Fig. 4-4). There is no significant difference between I-V curves obtained by applying voltage ramp either from -70 to +50 mV (480 mV/s) or from -100 to +60 mV (43.7 mV/s). The I-V relationship measured before the response induced by intracellular injection of cAMP into neurons was similar to that measured in control cells with normal internal solution (data not shown). The slope of the I-V curve measured during the cAMP-induced response was steeper than that measured before the response, indicating that cAMP increases the membrane conductance. It returned not completely but reversibly to the basal level after the cAMP-induced response had been adapted. The reversal potential was estimated to be -14.8 ± 2.6 mV (n=12), which was more negative than the potentials observed in isolated newt olfactory neurons [6] and the patch membrane excised from the cilia of the frog [3, 18] and rat [18].











Figure 4-4. Whole-cell current-voltage relationships for the current evoked by intracellular application of 0.5 mM cAMP. The current was measured by applying a voltage ramp (480 mV/s) from -70 to +50 mV before, during and after the response induced by 0.5 mM cAMP. These traces were obtained from the same cell. The inset shows the record of the cAMP-induced response of this cell under the whole-cell voltage clamp at -70 mV. The current transients were produced by voltage ramps (480 mV/s) from -70 to +50 mV. The reversal potential of the current induced by intracellular application of 0.5 mM cAMP to this neuron was estimated to be -13.8 mV.

(43.7 mV/s) to voltage-clamped vometonesul acurnes during and alter

Transient inward current induced by cGMP

Figure 4-5 shows the currents induced by intracellular injection of cGMP of varying concentrations into vomeronasal receptor neurons. cGMP was introduced into a proximal part of the dendrite or a part of cell soma by whole-cell dialysis in the same manner as that for cAMP. When the pipette was filled with an cGMP-free inner solution, the neurons held a steady baseline over the test interval of about 3-10 min after membrane rupture (left trace). On the other hand, introducing 1 mM cGMP into the neurons evoked prolonged, inward currents within a few seconds after membrane rupture in 8 of 10 (80%) neurons which displayed an increase in inward current with adaptation of current after the peak response. The amplitude of the inward current induced by cGMP varied from 0 to 555 pA (227 \pm 61.4 pA , mean \pm S. E. M.). The data obtained from these neurons in which the cGMP-induced current was not adapted were excluded because of indistinctness whether the current observed represents an inward current induced by cGMP or an artificial leak. The time to peak for the response of the vomeronasal neuron varied from 4 to 55 s (28 ± 7 , mean \pm S. E. M., n=8). This time is similar to that for cAMP-induced response of the turtle vomeronasal receptor neuron.

Figure 4-6 shows the voltage dependence of the cGMP-induced currents examined by applying a voltage ramp from -100 to +60 mA

(43.7 mV/s) to voltage-clamped vomeronasal neurons during and after the response induced by 1 mM cGMP. Because of difficulty of the

measurement, I-V relationships before the response induced by cGMP failed to be obtained. The I-V relationship measured after the introduction of cGMP into neurons was similar to that measured in control cells with normal internal solution (data not shown). The slope of the I-V curve measured during the cGMP-induced response is steeper than that measured after the response, indicating that cGMP increases the membrane conductance. The reversal potential was estimated to be -11.5 ± 8.7 mV (n=6), which was more negative than that observed in the patch membrane excised from the cilia of the frog [3, 18] and rat [18]. The reversal potential was similar to that observed in the response to intracellular application of cAMP to turtle vomeronasal receptor neurons as shown in Fig. 4-4, suggesting that the two nucleotides act at the same site.

Figure 4-5. As Response teduces by intrincellular application of 0 mbi eGMP (normal internal solution) from the patch pipette to a concernment reponter neuron bathed in normal Klager solution (control). Its Response induced by introcellular application of 1 mbit eCMP from the patch pipette to a volumentment morphy of 1 mbit eCMP from the patch solution. Open and, cloud, him above proces indicate period, of intracellular dialysis of 0 and 1 mbit eCMP in normal internet, in indicate respectively. Holding potential, -70 mV.





Figure 4-5. A: Response induced by intracellular application of 0 mM cGMP (normal internal solution) from the patch pipette to a vomeronasal receptor neuron bathed in normal Ringer solution (control). B: Response induced by intracellular application of 1 mM cGMP from the patch pipette to a vomeronasal receptor neuron bathed in normal Ringer solution. Open and closed bars above traces indicate period of intracellular dialysis of 0 and 1 mM cGMP in normal internal solution, respectively. Holding potential, -70 mV.





Figure 4-6. Whole-cell current-voltage relationships for the current evoked by intracellular application of 1 mM cGMP. The current was measured by applying a voltage ramp (43.7 mV/s) from -100 to +60 mV before, during and after the response induced by 1 mM cGMP. These traces were obtained from the same cell. The inset shows the record of the cGMP-induced response of this cell under the whole-cell voltage clamp at -70 mV. The current transients were produced by applying voltage ramps (43.7 mV/s) from -100 to +60 mV.

induced by intracellular application of 1 mM cGMP to this neuron was estimated to be -13.2 mV.

DISCUSSION

The present study clearly demonstrated that the turtle vomeronasal receptor neurons respond to intracellular dialysis of cAMP with an increase in inward current at negative membrane potentials in a dose dependent manner. In the neurons, cAMP induced an inward current accompanied by an increase in conductance (Fig. 4-4). Intracellular dialysis of cGMP into the turtle vomeronasal receptor neurons also elicited the inward currents similar to that of cAMP regarding the peak amplitude and the reversal potential of the responses. Although we did not record single-channel activity in these experiments, some of the results presented here are pertinent to the characterization of cyclic nucleotide-gated channels found in many vertebrate olfactory neurons [3, 6, 17, 21-27]. For instance, Nakamura et al., recording from patches of ciliary membrane in toad, reported a cationic current that was directly activated by both cAMP and cGMP [3]. Therefore, it is likely that the cyclic nucleotide-gated channels in turtle vomeronasal receptor neurons are directly activated by both cAMP and cGMP

In vertebrate photo transduction, cGMP acts as a second messenger, directly activating the light-dependent membrane

conductance [28, 29]. In the turtle vomeronasal receptor neuron, intracellular application of 1 mM cGMP elicited a rather large response



than that of cAMP at the same concentration. These results suggest that cGMP, rather than cAMP, is the second messenger for the signal transduction pathway. On the other hand, the turtle vomeronasal neurons responded to forskolin (Fig. 4-1). In addition, it was reported that forskolin increased the cAMP level in vomeronasal receptor neuron of garter snake. These observations indicate that vomeronasal receptor neurons have adenylate cyclase activities enough to elicit sufficient responses. As will be shown in Chapter V, forskolin elicits a large vomeronasal response of turtle. In turtle vomeronasal system, therefore, it is more likely that cAMP acts as a second messenger for signal transduction pathway.

In the frog, injection of cAMP into vomeronasal receptor neurons failed to elicit a membrane current [30]. This disagreement may be due to the difference in the preparations; enzymatically isolated olfactory neurons and neurons in slice preparation. In enzymatically isolated olfactory neurons, transient inward currents elicited by intracellular application of cAMP are, however, commonly observed [16, 20, 31]. Hence, it is not likely that enzymatic dissociation of the neurons inflicts unintentional damage on channel activities. It is more likely that this disagreement may due to the difference in species of animals used.

In the turtle vomeronasal organ, neither a chemoattractant nor a substance which activates the cAMP cascade has been identified.

Therefore, biological roles of cAMP channels found in the present study are unknown. In general, the vomeronasal organ has been

reported to play important roles in feeding, social and reproductive behaviors [32, 33]. Hence cAMP channels in the turtle vomeronasal neurons seem to be involved in the transduction pathway for the chemicals related to the above behaviors. On the other hand, turtle vomeronasal system sensitively responds to various general odorant [34, 35]. Thus, there remains a possibility that the cyclic nucleotide-gated channels contribute to the transduction pathway for general odorants. This issue will be discussed in Chapter V.

- in ellactory receptor citin. Meteric 325: 442-444, 1987
- ademytate cyclinic may hiedicte elistency reception. Miliare 1100
- S) Skiar, P. B., Anbolt, R. R. H. and Snyder, S. H. The objective sensitive adenyiate cycluse of oldaptory receptor cells: differential sumplation by distinct cleases of odorates, *I. Biol. Chem.*, 2611 15539-15543, 1986
- 6) Suiraki, M. Vritage- and cyclic interentias-gated currents in isolated offsetury receptor cells. "Chemical Senses, Vol. 1: Receptor Events and Twatesheeting in Taste and Offsetion" ed. by Brand, J. G., Teeler, J. H., Cagan R. H. and Kate, M. R. Marcel Ochkert New York (1989) on 269-494.
- Th Dreep 14 and bushbarr I. Odorants of the trune odor class activate



REFERENCES

- 1) Boekhoff, I., Tarelius, E., Strotmann, J. and Breer, H. Rapid activation of alternative second messenger pathways in olfactory cilia from rats by different odorants. EMBO J., 9: 2453-2458, 1990
- 2) Frings, S. and Lindemann, B. Current recording from sensory cilia of olfactory receptor cells in situ. I. The neuronal response to cyclic nucleotides. J. Gen. Physiol., 97: 1-16, 1990
- 3) Nakamura, T. and Gold, G. H. A cyclic nucleotide-gated conductance in olfactory receptor cilia. Nature, 325: 442-444, 1987
- 4) Pace, U., Hanski, E., Salomon, Y. and Lancet, D. Odorant-sensitive adenylate cyclase may mediate olfactory reception. Nature, 316: 255-258, 1985
- 5) Sklar, P. B., Anholt, R. R. H. and Snyder, S. H. The odorantsensitive adenylate cyclase of olfactory receptor cells: differential stimulation by distinct classes of odorants. J. Biol. Chem., 261: 15538-15543, 1986
- 6) Suzuki, N. Voltage- and cyclic nucleotide-gated currents in isolated olfactory receptor cells. "Chemical Senses, Vol. 1: Receptor Events and Transduction in Taste and Olfaction" ed. by Brand, J. G., Teeter, J. H., Cagan R. H. and Kare, M. R., Marcel Dekker, New York (1989) pp. 469-494.
- 7) Breer, H. and boekhoff, I. Odorants of the same odor class activate

101

different second messenger pathways. Chem. Senses, 16: 19-29, 1991

- 8) Kahn, A. A., Steiner, J. P. and Snyder, S. H. Plasma membrane inositol 1, 4, 5-triphosphate receptor of lymhpocytes: selective enrichment in sialic acid and unique binding specificity. Proc. Nal. Acad. Sci. USA., 89: 2849-2853, 1992
- 9) Kalinoski, D. L., Aldinger, S. B., Boyle, A. G., Huque, T., Marecek, J. F., Prestwich, G. D. and Restrepo, D. Characterization of a novel inositol 1, 4, 5-trisphosphate receptor in isolated olfactory cilia. Biochem. J., 281: 449-456, 1992
- 10) Restrepo, D., Miyamoto, T., Bryant, B. P. and Teeter, J. H. Odor stimuli trigger influx of calcium into olfactory neurons of the channel catfish. Science, 249: 1166-1168, 1990
- 11) Restrepo, D., Teeter, J. H., Honda, E., Boyle, A. G., Marecek, J. F., Prestwich, G. D. and Kalinoski, D. L. Evidence for an InsP3gated channel protein in isolated rat olfactory cilia, Am. J. Physiol., **263:** C667-C673, 1992
- 12) Suzuki, N., IP3-activated ion channel activities in olfactory receptor neurons from different vertebrate species. "Olfaction and Taste XI" ed. by Kurihara, K., Suzuki, N. and Ogawa, H., Springer-Verlag, Tokyo (1994) pp. 173-177
- 13) Kishimoto, J., Cox, H., Keverne, E. B. and Emson, P. C., Cellular localization of putative odorant receptor mRNAs in olfactory and chemosensory neurons: a non radioactive in situ hybridization study. Mol. Brain Res., 23: 33-39, 1994





- 14) Luo, Y., Lu, S., Chen, P., Wang, D. and Halpern, M., Identification of chemoattractant receptors and G-proteins in the vomeronasal system of garter snakes, J. Biol. Chem., 269: 16867-16877, 1994.
- 15) Jiang, X. C., Inouchi, J., Wang, D. and Halpern, M., Purification and characterization of a chemoattractant from electric shockinduced earthworm secretion, its receptor binding, and signal transduction through the vomeron sal system of garter snakes, J. Biol. Chem., 265: 8736-8744, 1990
- 16) Kurahashi, T. The response induced by intracellular cyclic AMP in isolated olfactory receptor cells of the newt. J. Physiol., 430: 355-371, 1990
- 17) Trotier, D. and MacLeod, P. cAMP and cGMP open channels and depolarize olfactory receptor cells. Chemical Senses, 11: 674, 1986
- 18) Frings, S., Lynch, J. W. and Lindemann, B. Properties of cyclic Nucleotide-gated channels mediating olfactory transduction; activation, selectivity, and blockage. J. Gen. Physiol., 100: 45-67, 1992
- 19) Taniguchi, M. Kashiwayanagi, M. and Kurihara, K., Intracellular injection of cyclic nucleotide monophosphate increases a conductance in membranes of turtle vomeronasal eceptor neurons in the slice preparation. in preparation
- 20) Kashiwayanagi, M., Kawahara, H., Hanada, T. and Kurihara, K. A large contribution of a cyclic AMP-independent pathway to turtle

olfactory transduction. J. Gen. Physiol., 103: 957-974, 1994
- 21) Firestein, S. and Werblin, F. Odor-induced membrane currents in vertebrate olfactory receptor neurons. Science, 244: 79-82, 1989
- 22) Bruch, R. C. and Teeter, J. H. Cyclic AMP links amino acid chemoreceptors to ion channels in olfactory cilia. Chemical Senses, 15: 419-430, 1990
- 23) Firestein, S., Zufall, F. and Shepherd, G. M. Single odor-sensitive channels in olfactory receptorneurons are also gated by cyclic nucleotides. J. Neurosci., 11: 3565-3572, 1991
- 24) Firestein, S. Lindemann, B. Current recording from sensory cilia of olfactory receptor cells in situ. I. The neuronal response to cyclic nucleotides. J. Gen. Physiol., 97: 1-16, 1991
- 25) Lowe, G. and Gold, G. H. The spatial distributions of odorant sensitivity and odorant-induced currents in salamander olfactory receptor cells. J. Physiol., 442: 147-168, 1991
- 26) Zufall, F., Firestein, S. and Shepherd, G. M. Analysis of single nucleotide-gated channels in olfactory receptor cells. J. Neurosci., 11: 3573-3580, 1991
- 27) Zufall, F., Shepherd, G. M. and Firestein, S. Inhivition of the olfactory cycllic nucleotide gated ion channel by intracellular calcium. Proc. Royal Soc. Lond., B246: 225-230, 1991
- 28) Fesenko, E. E., Kolesnikov, S. S. and Lyubarsly, A. L. Induction by cyclic GMP of cationic conductance in plasma membrane of retinal rod outer segment. Nature, 313: 310-313, 1985



- 29) Yau, K. W. and Nakatani, K. Light-suppressible, cyclic GMPsensitive conductance in the plasma membrane of a truncated rod outer segment. *Nature*, **317**: 252-255, 1985
- 30) Trotier, D., Doving, K. B. and Rosin, J-F., Voltage-dependent currents in microvillar receptor cells of the frog vomeronasal organ, *Eur. J. Neurosci.*, **5:** 995-1002, 1993
- 31) Firestein, S., Darrow, B. and Shepherd G. M. Activation of the sensory current in salamander olfactory receptor neurons depends on a G protein-mediated cAMP second messenger system. *Neuron*, 6: 825-835, 1991
- 32) Halpern, M. The organization and function of the vomeronasal system. Ann. Rev. Neurosci., 10: 325-362, 1987
- 33) Wysocki, C. and Meredith, M. The vomeronasal system.
 "Neurobiology of Taste and Smell" ed. by Finger, T. E. and Silver,
 W. L., John Wiley & Sons, Inc., New York (1987) pp. 125-150
- 34) Hatanaka, T., Shibuya, T. and Inouhchi, J. Induced wave responses of the accessory olfactory bulb to odorants in two species of turtle, *Pseudemys scripta* and *Geoclemys reevesii*. Comp. Biochem. *Physiol.*, 91A: 377-385, 1988
- 35) Shoji, T. and Kurihara, K. Sensitivity and transduction mechanisms of responses to general odorants in turtle vomeronasal system. J. Gen. Physiol., 98: 909-919, 1991



IV-2 The Responses Induced by Intracellular Injection of Inositol 1, 4, 5-Trisphosphate into Turtle Vomeronasal Receptor Neurons

INTRODUCTION

In the garter snakes, the binding of ES20, which was the chemoattractant for the snakes [1], to its receptors resulted in an increase in the basal level of IP3 [2]. This observation supports the idea that IP3 acts as a second messenger in signal transduction in the vomeronasal receptor neurons. In the present section [3], we show that intracellular application of IP3 from the patch pipette to turtle vomeronasal receptor neurons elicits the membrane current under the whole-cell patch clamp.

MATERIALS AND METHODS

Preparations

The slice preparations were obtained as described in Chapter III. Data recording and analysis

Membrane currents were recorded in the whole-cell

In on the other of the second se

configurations of the patch clamp as described in Chapter III (holding potential, -70 mV). Data were analyzed as described in Chapter III.

Solutions

The compositions of normal Ringer solution and normal internal solution were the same as described in Chapter III. For the stimulation with IP3 from the patch pipette, IP3 was dissolved in the internal solution to provide the appropriate final concentration. The stock solution of 0.1 mM IP3 derived into 0.5 ml aliquots was stored at -80 °C in 0.5 ml aliquots and thawed just prior to use.

Chemicals

IP3 and ruthenium red were purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan). All chemicals used are of best grade available.

RESULTS

IP3 was introduced into a proximal part of the dendrite or a part of cell soma by whole-cell dialysis. When the pipette was filled with an IP3-free inner solution, the neurons held a steady baseline over the test interval of about 3-10 min after membrane rupture (Fig. 4-7A). On the other hand, introducing 0.1 mM IP3 into the neurons evoked prolonged, inward currents within a few seconds after membrane

rupture (Fig, 4-7B). In the present study, 98 neurons were successfully stimulated by 0.1 mM IP3. Fifty two neurons (53%) displayed an





Figure 4-7. A: Response induced by intracellular application of 0 mM IP3 (normal internal solution) from the patch pipette to a vomeronasal receptor neuron bathed in normal Ringer solution (control). B: Response induced by intracellular application of 0.1 mM IP3 from the patch pipette into a vomeronasal receptor neuron bathed in normal Ringer solution. C: Response induced by intracellular application of 0.1 mM IP3 from the patch pipette to a vomeronasal receptor neuron bathed in Ringer solution containing 10 μ M ruthenium red. Open and hatched bars above traces indicate period of intracellular dialysis of 0 and 0.1 mM IP3 in normal internal solution, respectively. Holding potential, -70 mV.



increase in inward current with adaptation of current after the peak response. The amplitude of the inward current induced by IP3 varied from 0 to 736 pA (90 \pm 11 pA, mean \pm S. E. M.). In some neurons, the IP3-induced current was not adapted. The data obtained from these neurons were excluded because it was unclear whether the current observed represents an inward current induced by IP3 or an artificial leak.

Kashiwayanagi and Kurihara [unpublished data] applied 0.1 mM IP3 into the turtle olfactory neuron and observed that IP3 induced an inward current in 56% of neurons examined at -70 mV with an average peak amplitude of 131 ± 7 pA (n=27). The magnitude of the response induced by IP3 in the turtle vomeronasal receptor neurons is comparable with that in the turtle olfactory neurons. The time to peak for the response of the vomeronasal neuron varied from 3 to 247 s (32 \pm 5, mean \pm S. E. M., n=52). This time is much faster than that for IP₃induced response of the rat olfactory neuron (30 to 1200 s) [4].

In the olfactory neurons of the catfish [5] and lobster [6], the IP3-induced response was inhibited by 10 μ M ruthenium red. We examined the effect of ruthenium red on IP3-induced response in the turtle vomeronasal neurons (Fig. 4-7C). Bathing the neurons in 10 μ M ruthenium red solution greatly reduced IP3-evoked inward currents. The peak amplitude of the inward current was reduced to 18.0 ± 4.6 pA (n=5) from 89.9 ± 10.9 pA (n=52).

The voltage dependence of the IP3-induced currents was examined by applying a voltage ramp from -100 to +60 mA (43.7

mV/s) to voltage-clamped vomeronasal neurons before, during and after the response induced by 0.1 mM IP3 (Fig. 4-8A). The I-V relationship measured before the introduction of IP3 into neurons was similar to that measured in control cells with normal internal solution (data not shown). At negative membrane potentials, the slope of the I-V curve measured during the IP3-induced response was steeper than that measured before the response, indicating that IP3 increases the membrane conductance. It returned reversibly to the basal level after the IP3-induced response had been adapted. The reversal potential was estimated to be -32.3 ± 1.5 mV (n=6), which was similar to that observed in rat olfactory neurons [4].

Figure 4-8B shows the IP₃-sensitive component obtained by subtracting the current measured before the response from that during the response. The IP₃-sensitive component was strongly outward rectifying at the membrane potential ranging from -30 to 0 mA, followed by large (4 of 5 cells) and small (1 of 5 cells) reduction in the current amplitude at positive potentials. A similar negative slope of the IP₃-induced current in the I-V curve was observed in the rat olfactory neurons [4]. Ca²⁺-activated K⁺ channel seems to contribute to the negative slope as Okada *et al.* [4] suggested.

1914 induced response of this bell mater the whole-call voltage charpe Bi -70 mV. The current transients ware produced by voltage campe (43.7 mV/s) from -100 to +60 mV. The revenuel potencial of the current



DISCUSSION

The present is the first and featuristic states of the present of



Figure 4-8. Whole-cell current-voltage relationships for the current evoked by intracellular application of 0.1 mM IP3. The current was measured by a voltage ramp (43.7 mV/s) from -100 to +60 mV before, during and after the response induced by 0.1 mM IP3 (A). These traces were obtained from the same cell. The inset shows the record of the IP3-induced response of this cell under the whole-cell voltage clamp at -70 mV. The current transients were produced by voltage ramps (43.7 mV/s) from -100 to +60 mV. The reversal potential of the current induced by intracellular application of 0.1 mM IP3 to this neuron was estimated to be -39.5 mV. IP3-sensitive component of this current was obtained by subtracting the current measured before the IP3-induced response from that measured during the response (B).

DISCUSSION

The present study demonstrated that turtle vomeronasal receptor neurons responded to intracellular dialysis of IP3 with an increase in inward current at negative membrane potentials. This response was induced by IP3 of mM concentration and inhibited by bath application of ruthenium red. The reversal potential was similar to that in rat olfactory neurons [4]. These results suggest that the IP3-induced depolarization in turtle vomeronasal receptor neurons is mediated by activation of IP3-gated channels.

In general, the vomeronasal organ plays important roles in feeding, social and reproductive behaviors. Hence IP₃-channels conductance in the turtle vomeronasal neurons seem to be involved in the transduction pathway for the chemicals related to the above behaviors. This is the first observation that demonstrated existence of IP₃-activated conductance in the vomeronasal receptor membranes.

Acumphysiol., 37a, 595-692, 1994

- 5) Repricipit, D., Miymmold, T., Bryant, B. P. and Tester, J. H. Udmi
- ehamel cathisti Science, 349, 1166-1168, 1990
- 6) Fudard D. A. and Ache B. W. Planne membrane inositel b. 4, 5-



REFERENCES

- 1) Jiang, X. C., Inouchi, J., Wang, D. and Halpern, M., Purification and characterization of a chemoattractant from electric shockinduced earthworm secretion, its receptor binding, and signal transduction through the vomeronasal system of garter snakes, J. Biol. Chem., 265: 8736-8744, 1990
- 2) Luo, Y., Lu, S., Chen, P., Wang, D. and Halpern, M., Identification of chemoattractant receptors and G-proteins in the vomeronasal system of garter snakes, J. Biol. Chem., 269: 16867-16877, 1994
- 3) Taniguchi, M. Kashiwayanagi, M. and Kurihara, K., Intracellular injection of inositol 1, 4, 5-trisphosphate increases a conductance in membranes of turtle vomeronasal receptor neurons in the slice preparation. submitted to Neurosci. Lett.
- 4) Okada, Y., Teeter, J. H. and Restrepo D., Inositol 1, 4, 5trisphosphate-gated conductance in isolated rat olfactory neurons, J. Neurophysiol., 71: 595-602, 1994
- 5) Restrepo, D., Miyamoto. T., Bryant, B. P. and Teeter, J. H., Odor stimuli trigger influx of calcium into olfactory neurons of the channel catfish, Science, 249: 1166-1168, 1990
- 6) Fadool D. A. and Ache B. W., Plasma membrane inositol 1, 4, 5trisphosphate-activated channels mediate signal transduction in

lobster olfactory receptor neurons, Neuron, 9: 907-918, 1992

CHAPTER V

CAMP-DEPENDENT TRANSDUCTION PATHWAY DOES NOT GREATLY CONTRIBUTE TO THE TRANSDUCTION FOR GENERAL ODORANTS IN TURTLE VOMERONASAL NEURONS

INTRODUCTION

The results described in chapter IV demonstrated the existence of cyclic nucleotide or IP3-mediated pathway in the vomeronasal receptor neurons. However, the biological role of these putative second messenger-mediated pathways in vomeronasal transduction is unknown, since neither a chemoattractant nor a substance which activates the putative second messenger cascade has yet been identified in the turtle vomeronasal organ.

On the other hand, the vomeronasal organ responds to general odorants as well as chemical stimuli of a social nature [1, 2]. It is reported that the turtle vomeronasal organ sensitively responds to many kinds of odorants [3, 4]. Recently, it was reported that the vomeronasal organ of garter snakes responded to chemical stimuli including general odorants as well as proteins in prey washes, indicating that the vomeronasal system is sensitive to a variety of odorants [5]. Hence there

is the possibility that the putative second messenger-activated

conductance may be involved in transduction mechanism for general odorants as well as natural stimuli.

Data shown in Chapter IV indicated that the population of the turtle vomeronasal receptor neurons having cyclic nucleotide-mediated pathway is larger than that of neurons having IP3-mediated pathway. Thus, present study aims to examine to what extend the cyclic nucleotide-mediated pathway contributes to *in vivo* turtle vomeronasal transduction for general odorants [6]. The results obtained show that the turtle vomeronasal response to citralva appears even after complete desensitization of cAMP-mediated pathway by applying sufficient amounts of forskolin, suggesting that the cAMP-mediated pathway does not greatly contribute to the generation of the vomeronasal response to a general odorant.

MATERIALS AND METHODS

Recordings of main and accessory olfactory bulbar responses

Accessory olfactory bulbar response (vomeronasal response) was essentially recorded similarly to the recording of main olfactory bulbar response described in Chapter I. Turtles (*Geoclemys reevesii*) weighing 140-240 g were used in the present study. Turtles were weakly anesthetized with the necessary and minimum amount of

urethane to lessen pain in the operation of the animal, immobilized by

an intramuscular injection of d-tubocurarine chloride (450 μ g/100 g

body weight), and locally anesthetized with lidocaine at the wounded and head-fixation points. The accessory olfactory bulb was exposed using a dental drill, and the dura mater on the accessory olfactory bulb was removed carefully. To eliminate the possible effect of the main olfactory bulb activities (3), the nerve from the main olfactory organ was cut off before entry to the main olfactory bulb. The stimulantinduced brain waves (bulbar responses) were recorded by attaching a pair of silver bipolar electrodes to the medial part of the anterior bulb. The responses were amplified and then integrated by electric integrator (time constant, 0.3 sec).

Activity of the main olfactory bulbar responses was recorded as described in Chapter I.

Chemical stimulation

The irrigating and stimulating solutions were applied to the vomeronasal epithelium instead of olfactory epithelium through a stainless steel tube. The other procedure of chemical stimulation was essentially similar to that described in Chapter I. The chemical stimulation of olfactory epithelium was carried out as described in Chapter I

Chemicals

Lilial (4-(1,1-dimethylethyl)- α -methylbenzenepropanal),

geraniol (3,7-dimethyl-2,6-octadien-1-ol), lyral (mixture of 4-(4hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde and 3-(4-

hydroxy-4-methylpentyl)-3-cyclohexene-1-carboxaldehyde), citralva (3,7-dimethyl-2,6-octadienenitrile), hedione (3-oxo-2pentylcyclopentaneacetic acid methyl ester), eugenol (2-methoxy-4-(2propenyl)phenol), *l*-carvone (2-methy*l*-5-(1-methylethenyl)-2cyclohexene-1-one), *d*-carvone, *d*-limonene (1-methy*l*-4-(1methylethenyl)cyclohexene), *l*-citronellal (3,7-dimethyl-6-octenal), *d*citronellal were kindly supplied from Takasago International (Tokyo, Japan). *n*-Amyl acetate, triethylamine, ethylvanillin and forskolin were purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan). Cineole (eucalyptol; 1,3,3-trimethyl-2-oxabicyclo[2.2.2]octane) was purchased from Nakarai Chemicals (Kyoto, Japan). All chemicals used were of best grade available.

RESULTS

Figure 5-1A shows the summated accessory olfactory bulbar responses to 5 x 10^{-5} M forskolin and 10^{-4} M citralva dissolved in Ringer solution. As seen from the figure, the responses show a peak at onset of stimulation and decline rapidly during stimulation. We take the peak height of the summated bulbar response as the magnitude of the response. The response to forskolin is much larger than that to citralva.

Figure 5-2 shows the relative magnitudes of the vomeronasal responses (A) and main olfactory responses (B) to various stimuli. Here

the magnitudes of the main and vomeronasal responses to 10⁻⁴ M





Figure 5-1. Typical records of the accessory olfactory bulbar responses to 50 μ M forskolin and 100 μ M citralva. Hatched bars under the records represent duration of stimulation.



A: vomeronasal response



B: olfactory response



Figure 5-2. Relative magnitude of the accessory olfactory bulbar responses (A) and main olfactory bulbar responses to various stimuli (B). Magnitude of accessory (A) and main (B) olfactory bulbar responses are calculated relative to the response to 10^{-4} M citralva, respectively. Each point is mean \pm S.E.M. 119 citralva are taken as unity, respectively. The relative magnitude of the main olfactory response to 5 x 10^{-5} M forskolin is smaller than or comparable to that to odorants. On the contrary, the relative magnitude of the vomeronasal response to 5 x 10^{-5} M forskolin is much larger than those of odorants.

To examine directly to what extend cAMP-mediated pathway contributes to *in vivo* turtle vomeronasal transduction for general odorants, we recorded the vomeronasal responses to citralva after cAMP pathway had been desensitized by forskolin. Figure 5-3A illustrates typical records of the accessory olfactory bulbar responses to 10^{-4} M citralva applied alone and after the response to 5 x 10^{-5} M forskolin was adapted to a spontaneous level. As seen from the figure, the response to citralva clearly appeared even after 10^{-5} M forskolin.

Figure 5-3B plots the magnitude of the response to forskolin applied first and that to 10^{-4} M citralva after forskolin as a function of forskolin concentrations. Here the magnitude of the responses to forskolin and citralva are calculated relative to those to 5 x 10^{-5} M forskolin and 10^{-4} M citralva applied alone, respectively. The response to forskolin was increased with an increase in forskolin concentrations and reached a plateau level at $10\text{-}20 \ \mu\text{M}$. The minimum concentration (threshold) of forskolin which elicits the vomeronasal response was 10^{-8} M. The response to a fixed concentration (10^{-4} M) of citralva applied after forskolin decreased with an increase in forskolin concentration up

to 1-10 μ M and plateaued. The fact that the response to citralva was

partially suppressed by forskolin suggests that cAMP-mediated pathway



Figure 5-3. Effects of varying concentrations of forskolin on the response to citralva. A: typical records of response to 100 μ M citralva applied alone (left trace) and 100 μ M citralva applied secondarily after adaptation to 50 μ M forskolin (right trace). Hatched bars, duration of application of stimuli. B: Relative magnitude of the accessory olfactory bulbar responses to forskolin of varying concentration applied alone (\bigcirc) and citralva of a fixed concentration (100 μ M) applied secondarily (\bigcirc) as a function of forskolin concentration. Magnitudes of the responses to forskolin and citralva were measured from a basal level. Magnitudes of the responses to 50 μ M forskolin applied alone (\bigcirc) and that to citralva applied alone (\bigcirc) and that to citralva after adaptation to forskolin (\bigcirc), respectively. Mean magnitude of the response to 50 μ M forskolin was 4.9 times as large as that to 100 μ M

somewhat contribute to the generation of the vomeronasal response to a general odorant such as citralva. On the other hand, the magnitude of the response to 10^{-4} M citralva after 5 x 10^{-5} M forskolin, where the response to forskolin itself was sufficiently saturated, was 65 % of that to citralva applied alone. This suggests that the cAMP-mediated pathway does not greatly contribute to the generation of the vomeronasal response to a general odorant such as citralva.

DISCUSSION

In the present study, we measured main and accessory olfactory bulbar responses to various stimuli. The whole-cell recordings offer direct information on olfactory transduction in a single neuron, but have some disadvantages. For example, function of isolated single neuron to respond to odorants is declined by repeated application of drugs such as forskolin. On the other hand, the accessory olfactory bulbar responses can be measured stably and reproducibly for a long time (e.g., 3 days). In addition, the recordings of the accessory olfactory bulbar responses allow us to perform repeated application of forskolin to the vomeronasal epithelium under *in vivo* conditions without any irreversible effect. Consequently, recordings from accessory olfactory bulb offer quantitative and reproducible data.

Present results obtained from measurements of the accessory olfactory bulbar responses indicated that forskolin elicits conspicuously



large responses in the turtle vomeronasal system. This is consistent with the results that forskolin elicits a large current in the turtle vomeronasal neurons as described in Chapter IV (Fig. 4-1). The present results also showed that forskolin reduces the response to citralva applied after adaptation to forskolin. This suggests the possibility that cAMPmediated pathway partly contributes to vomeronasal transduction pathway for general odorants. Another possible reason for the suppression by forskolin is that the response to citralva applied secondarily is partly suppressed by nonspecific effects of forskolin, because forskolin has various actions besides activation of adenylate cyclase [7]. Thus, we cannot conclude that all of the forskolinsuppressive components in the responses to cAMP-producing odorants are cAMP-dependent components.

Present results also indicated that the accessory olfactory bulbar response to forskolin was much larger than those to general odorants. This does not support the idea that the vomeronasal response to a general odorant is mediated dominantly by the cAMP-dependent pathway. In addition, the response to citralva remained more than 60% of that applied alone even after application of high concentrations of forskolin, suggesting that the cAMP-independent pathway greatly contributes to the generation of the vomeronasal response to general odorants such as citralva.

The cAMP-insensitive component of the response to citralva may be

generated via the pathways such as IP3-dependent pathway, cGMPdependent pathway, direct activation of G protein, or the other pathway

(see Fig. I-1). Further study will be needed to explore the transduction mechanism for general odorants and chemoattractants in the vomeronasal organ.

nisponter to odotable "Offeriow and Turk" ed. by Zoherman, Y. Morinitim Publishing Co., New York (1963) pp. 15-69

- Turker, D. Notothactory responses from the name cavery successory
 Turker, D. Notothactory responses from the name cavery Physiology.
 Vol. IV-P. ed. by Betdler, D. M. Springer-Verlag, New York
 (1971) pp 151-181
- Matanaka, T., Shibuya, T. and frombohi, J. Influited wave responses of the accessory officitory bulk to obtents in two species of territy. Pseudemys Actions and Geoclemys recesses Comp. Blockers, Physiol., 91A4 377-345, 1988.
- 4) Shoji, T. and Kutihma, K. Sensitivity and transduction mechanisms of responser to general adamats in turtle vomeroaasal system. J. Gen. Physica, 48: 909-919, 1991.
- Investig, J., Wang, D., Hang, M. C.; Kubie, J. and Halpein; M. Electrophysiclogical analysis of the nucl chemical senses in gamer snakes, Brain, Bellay, Evol., 41: 171-192, 1993
- 6) Tanigucht, M. Kamelei, K., Kashiwayanagi, M. and Kamilana, K. ar preparation (
- Th Laureenter A., Suttowski, T. M. and Scientin, K. B. Ferskolin, &



REFERENCES

- 1) Tucker, D. Olfactory, vomeronasal and trigeminal receptor responses to odorants. "Olfaction and Taste" ed. by Zotterman, Y., Macmillan Publishing Co., New York (1963) pp. 45-69
- 2) Tucker, D. Nonolfactory responses from the nasal cavity: Jacobson's organ and the trigeminal system. "Handbook of Sensory Physiology. Vol. IV-1" ed. by Beidler, L. M., Springer-Verlag, New York (1971) pp.151-181
- 3) Hatanaka, T., Shibuya, T. and Inouhchi, J. Induced wave responses of the accessory olfactory bulb to odorants in two species of turtle, Pseudemys scripta and Geoclemys reevesii. Comp. Biochem. Physiol., 91A: 377-385, 1988
- 4) Shoji, T. and Kurihara, K. Sensitivity and transduction mechanisms of responses to general odorants in turtle vomeronasal system. J. Gen. Physiol., 98: 909-919, 1991
- 5) Inouchi, J., Wang, D., Jiang, X. C., Kubie, J. and Halpern, M. Electrophysiological analysis of the nasal chemical senses in garter snakes. Brain, Behav. Evol., 41: 171-192, 1993
- 6) Taniguchi, M. Kanaki, K., Kashiwayanagi, M. and Kurihara, K. in preparation
- 7) Laurenza, A., Sutkowski, E. M. and Seamon, K. B. Forskolin: a

specific stimulator of adenylyl cyclase or a diterpene with multiple sites of action? Trend Pharm. Sci., 10: 442-447, 1989



SUMMARY

- [I] Difference in odor intensity and quality between *l* and *d*optical isomers
 - (1) The turtle olfactory bulbar responses to six pairs of highly pure optical isomers were measured and the difference in odor intensity between the optical isomers was compared. With all odorants examined, there was no difference in odor intensity between optical isomers in the whole concentration ranges examined.
 - (2) We developed the quantitative cross-adaptation method in which the concentration of one odorant was varied and that of another remained constant. The difference in odor quality between optical isomers was evaluated by the quantitative cross-adaptation method. The degree of cross-adaptation between optical isomers were greatly different among species of odorants. The rank order of the magnitude of the difference in odor quality between optical isomers was carvone > β -citronellol > menthol> hydroxycitronellal > citronellal > limonene.

[II] Effects of treatment of the turtle olfactory epithelium with various lipids on olfactory responses

The turtle olfactory epithelium was treated with suspensions of

various lipids and their effects on the olfactory responses were examined by measuring the olfactory bulbar responses.

- (1) The phosphatidylserine (PS)-treatment greatly lowered the threshold for *n*-valeric acid and enhanced its responses at all concentrations examined. The responses to isovaleric acid and *n*-butyric acid were also greatly enhanced by the PS-treatment. The responses to twelve odorants examined were a little enhanced or practically unchanged by the PS-treatment. The enhanced responses to the fatty acids returned to the original level about 10 h after the treatment.
- (2) The experiment using [¹⁴C]PS indicated that PS was incorporated into the olfactory epithelium.
- (3) In contrast to PS-treatment, the treatment of the epithelium with cardiolipin or phosphatidic acid unchanged or suppressed the olfactory responses to odorants including the fatty acids.

[III] Electrophysiological properties of turtle vomeronasal receptor neurons

The turtle vomeronasal receptor neurons in the slice preparation were studied with the whole-cell configuration of the patch clamp technique.

- (1) Typical resting potentials were near -48 mV, and typical input resistance was 1-2 G Ω .
- (2) The response to a current step consisted of either a single spike regardless of stimulus strength, or a train of spikes. Only 3 pA of

injected current was required to depolarize the neuron to spike



threshold near -55 mV, suggesting that a turtle vomeronasal receptor neuron has high sensitivity to injected currents.

- (3) Voltage-clamped vomeronasal receptor neurons displayed transient inward currents followed by sustained outward currents in response to depolarizing voltage steps. Because of their similar properties to Na⁺-channels and K⁺-channels found in many neurons, the transient inward currents were tentatively identified as Na⁺-currents and the outward currents as K⁺-currents, respectively.
- (4) These results suggest that electrophysiological properties of turtle vomeronasal receptor neurons are similar to those of olfactory neurons.
- [IV] Signal transduction mechanism in vomeronasal receptor neurons

cAMP, cGMP or IP3 were injected into turtle vomeronasal receptor neurons in the slice preparation under the whole-cell patch clamp, and the evoked current was measured.

- (1) In cell-attached recording, 10 μ M forskolin added to the bath caused a transient increase of spike rate for 4 s, suggesting that there exists cAMP-dependent pathway in the vomeronasal receptor neurons.
- (2) Intracellular application of cAMP evoked a prolonged, inward

current in a dose-dependent manner. Intracellular application of 0.1 mM cAMP was enough to elicit an inward current in these

neurons. The magnitude of the response to cAMP reached a plateau level at 1 mM with an average peak amplitude of 176 ± 34 pA (n=30). The reversal potential of the response induced by cAMP was estimated to be -14.8 ± 2.6 mV (n=12).

- (3) Intracellular application of 1 mM cGMP also evoked a prolonged, inward current with an average peak amplitude of 227 \pm 61 pA. The reversal potential of the response induced by cGMP was estimated to be $-11.5 \pm 8.7 \text{ mV}$ (n=6).
- (4) The currents elicited by either intracellular cAMP or cGMP reversed near -14 mV and had similar current-voltage relationships, suggesting that the two nucleotides act at the same site.
- (5) Application of 0.1 mM IP3 also evoked a prolonged, inward current with an average peak amplitude of 89.9 ± 10.9 pA. The reversal potential of this response was estimated to be -32.3 ± 1.5 mV (n=6).
- (6) The present results demonstrated the existence of second messenger-activated conductance in the vomeronasal receptor membranes.
- [V] cAMP-dependent transduction pathway does not greatly contribute to the transduction for general odorants in turtle vomeronasal neurons

In order to examine to what extent cAMP-mediated pathway contributes to the generation of the vomeronasal response to

general odorants in *in vivo* system, the responses of the vomeronasal organ to various stimuli in the turtle were measured by recording the accessory olfactory bulbar responses.

- (1) The vomeronasal response to forskolin was much larger than that to a general odorant.
- (2) The cAMP-mediated pathway seems to be fully activated or desensitized when a sufficient amount of forskolin is applied to the olfactory neurons from external solutions. Large vomeronasal response to citralva appeared even after the response to forskolin itself was sufficiently saturated, suggesting that the cAMP-independent pathway greatly contributes to the *in vivo* vomeronasal transduction for general odorants such as citralva.

important cole in the olfactory reception.

vomeronanal avisem indicated that such intracellular application of CAMP, sGMP and IP5 to the tasks vomeronanal receptor neurons elicits the minimum durrent. These results clearly demonstrated that the membranes of the turble vomeronanal neurons carry cyclic nucleotide and/or IP3 activated conductance. The present statilits also showed this the cyclic authentide mediated pathway does not grantly contrabute to vomeronanal transduction for peperal odorants. The present results classified for possibility first tipse second messenger-



CONCLUDING REMARKS

The present study showed that there was no difference in odor intensity between optical isomers with all odorants examined. There was certain difference in odor quality between optical isomers. Especially, the odor quality was greatly different between d- and lcarvone. However, the difference in odor quality between d- and lcarvone disappeared at 40 °C. These results suggested that the receptor mechanism of the olfactory system is different from those of hormones and neurotransmitters. In addition, we showed that the PS-treatment of the olfactory epithelium greatly enhanced the responses to the fatty acids. All these results suggest that lipids as well as proteins play an important role in the olfactory reception.

The present study on signal transduction mechanisms in turtle vomeronasal system indicated that each intracellular application of cAMP, cGMP and IP3 to the turtle vomeronasal receptor neurons elicits the membrane current. These results clearly demonstrated that the membranes of the turtle vomeronasal neurons carry cyclic nucleotide and/or IP3-activated conductance. The present results also showed that the cyclic nucleotide-mediated pathway does not greatly contribute to vomeronasal transduction for general odorants. The present results raised the possibility that these second messenger-

mediated pathways are involved in the transduction for the chemoattractant in the vomeronasal organ.





