Title	Growth and Leaf Gas Exchange in Three Birch Species Exposed to Elevated Ozone and CO ₂ in Summer
Author(s)	Hoshika, Yasutomo; Watanabe, Makoto; Inada, Naoki; Koike, Takayoshi
Citation	Water, Air, & Soil Pollution, 223(8), 5017-5025 https://doi.org/10.1007/s11270-012-1253-y
Issue Date	2012-09
Doc URL	http://hdl.handle.net/2115/50360
Rights	The final publication is available at www.springerlink.com
Type	article (author version)
File Information	WASP223-8_5017-5025.pdf

[Instructions for use](#)

**Growth and leaf gas exchange in three birch species
exposed to elevated ozone and CO₂ in summer**

Yasutomo HOSHIKA, Makoto WATANABE, Naoki INADA and
Takayoshi KOIKE

Silviculture and Forest Ecological Studies, Hokkaido University,

Sapporo 060-8689, Japan

Corresponding author: Takayoshi Koike Tel: +81-11-706-3854, Fax:
+81-11-706-2517 E-mail: tkoike@for.agr.hokudai.ac.jp

Abstract

We examined the effects of ozone and elevated CO₂ concentration in summer on the growth and photosynthetic traits of three representative birch species in Japan (mountain birch, Monarch birch and white birch). Seedlings of the three birch species were grown in 16 open-top chambers and were exposed to two levels of ozone (6 nmol mol⁻¹ and 60 nmol mol⁻¹ for 7 h per day) in combination with two levels of CO₂ (370-380 μmol mol⁻¹ and 600 μmol mol⁻¹ for daytime) from July to October. No adverse effects of ozone were found in the Monarch birch or the white birch, but elevated ozone in summer reduced branch biomass and net photosynthesis, and accelerated leaf abscission, in the mountain birch. Elevated CO₂ promoted root development and thereby reduced the ratio of shoot dry mass (stem + branch) to root dry mass (S/R ratio) in the mountain birch and white birch. In contrast, there was no difference in dry mass between ambient and elevated CO₂ for the Monarch birch, due to down-regulation of photosynthesis. Studies of the combined effect of CO₂ and ozone revealed that elevated CO₂ did not ameliorate the effect of ozone on mountain birch in late summer. In considering the ameliorating effect of CO₂ on ozone damage, it is necessary to take account of the species and the season.

Key words: compensative effects; tropospheric ozone; elevated CO₂; birch; photosynthesis

1. INTRODUCTION

Tropospheric ozone (O_3) is a widespread phytotoxic air pollutant, and also a significant greenhouse gas (Bytnerowicz et al. 2007; Serengil et al., 2011). Ground surface O_3 concentrations are increasing in East Asia because of rapidly increasing emission of the main O_3 precursors, nitrogen oxides and volatile organic compounds (Naja and Akimoto, 2004). Ohara and Sakata (2003) reported that the annual average concentrations of photochemical oxidant, mainly O_3 , increased in Japan at the large rate of $0.33 \text{ nmol mol}^{-1} \text{ year}^{-1}$ from 1985 to 1999, and are still increasing (Yamaji et al., 2008). Ohara et al. (2001) reported that the increase in O_3 concentration may also be influenced by trans-boundary air pollution. Since the 1990s, many experimental studies using open-top chambers have studied the sensitivity to O_3 of various Japanese forest tree species (Yamaguchi et al., 2011). Watanabe et al. (2010a, 2012) suggested that the present O_3 concentration in Japan may have a negative impact on the growth of forest tree species.

The effect of O_3 on tree species depends on the atmospheric conditions (e.g., Karnosky et al., 2003; Watanabe et al., 2010b). One of the topical subjects is whether elevated CO_2 concentration expected in future can ameliorate the adverse effect of O_3 on tree species (Karnosky et al., 2001). Some studies found the ameliorating effect of CO_2 on O_3 -induced growth reduction and photosynthetic impairment (e.g., Grams et al., 1999; Lütz et al., 2000; Matsumura et al., 2005), but others reported no compensatory effects of CO_2 on the impact of O_3 (e.g., Barnes et al., 1995; King et al., 2005). Elevated CO_2 is generally known to induce stomatal closure (Paoletti and Grulke,

2005). A plausible mechanism for the ameliorating effect of CO₂ is stomatal closure induced by elevated CO₂, thereby limiting O₃ uptake by stomata (e.g., Broadmeadow et al., 1999).

Birch is a pioneer species that is rapidly established following fire or other site disturbance in cool temperate region in Northeast Asia including northern Japan (Linder, 1987; Koike, 1995b). These species are commercially important because of their high growth rate, great timber quality (Koike, 1988; Mao et al., 2010), and comprise 12% of total forest stock in northern Japan (Hokkaido prefecture, 2011; Kawaguchi et al., accepted). Recently declining or die-back birch has been observed in northern Japan (Ohno et al., 2010; Noguchi et al., 2011) and near Tokyo, capital of Japan (the Oku-Nikko region) (Shimizu and Feng, 2007). Ozone might be one of the factors related to these declines. Shimizu and Feng (2007) suggested that dry mass and photosynthesis of mountain birch were reduced by exposure to O₃ at a concentration of 50 nmol mol⁻¹ (daily average) as recorded in the Oku-Nikko region of Japan, a mountainous area. The effects of elevated O₃ and elevated CO₂ on tree growth and on physiological traits were determined by OTC experiments over the entire growing season (April to October) in Japan for two kinds of birch species in central Japan (Matsumura et al., 2005). They reported compensatory effects of CO₂ on the impact of O₃ in white birch (*Betula platyphylla* var. *japonica*).

This ameliorating effect of CO₂ may also depend on seasonal influences (Grams et al., 1999; Lütz et al., 2000). Lütz et al. (2000) suggested that the amelioration of O₃-induced reduction in photosynthesis was attenuated in late summer for European beech

(*Fagus sylvatica* L.). High O₃ peak concentrations, exceeding 100 nmol mol⁻¹, were frequently recorded during summer in suburban and mountainous areas of Japan (Shimizu and Feng, 2007; Takeda and Aihara, 2007; Takigawa et al., 2007). Moreover, the peak photosynthetic rate in birch leaves in Japan occurs in late summer (Koike and Sakagami, 1985, Koike 1995b). Late-season's photosynthesis contributes large biomass increment in poplar clone in north-central U.S.A. (Nelson and Isebrands, 1983). The effect of late summer exposure to O₃ is a key to develop our knowledge of the realistic effects of O₃ on birch species native to Japan. Therefore, it is a matter of discussion if elevated CO₂ may provide no compensatory effect on O₃ impacts in late summer for birch species of Japan.

To test this prediction, we examined the effects of elevated O₃ in combination with elevated CO₂ on the growth and photosynthesis of three birch species from late July to October. Based on the data, we discuss the interaction of the effects of O₃ and CO₂ on growth.

2. MATERIALS AND METHODS

2.1 Plant materials

We used 2-year old seedlings of mountain birch (*B. ermanii*), Monarch birch (*B. maximowicziana*) and white birch (*B. platyphylla* var. *japonica*) obtained from the nursery at Naganuma town in central part of Hokkaido, i.e., main part of distribution for these birches, in order to adjust the acclimation of day length for these birches (e.g. Evans, 1963; Larcher, 2004). The seeds were collected around central Hokkaido, northern Japan. The height and stem basal diameter at the beginning of the experiment were 35.6±4.5 cm and 4.3±0.4 mm for

mountain birch, 65.1 ± 4.8 cm and 6.3 ± 0.5 mm for Monarch birch, and 53.3 ± 4.5 cm and 5.7 ± 0.6 mm for white birch. Before bud break, these bare rooted birch seedlings were planted in late May 2010 in 7-liter pots filled with 1:1 (v/v) mixture of the Kanuma pumice soil and clay soil (infertile plant culture material). After leaf initiation, diluted liquid fertilizer (Nutrient balanced fertilizer, N:P:K=6:10:5, Hyponex, Ohio, U.S.A.) was supplied to all potted soils, amounting to total nitrogen (N) application of $192 \text{ mg N pot}^{-1}$. All plants were supplied with water at 3-7 day intervals to avoid water stress.

2.2 O₃ and CO₂ treatments

The experiment was carried out in open-top chambers (OTC) in the Sapporo Experimental Forest of Hokkaido University, in northern Japan ($43^{\circ}04' \text{ N}$, $141^{\circ}20' \text{ E}$, 15 m a.s.l., annual mean temperature: 11.5° C , total precipitation: 1325 mm in 2010). All birch seedlings were exposed to the following treatments from July to October 2010: control ($\text{CO}_2 = 370\text{-}380 \text{ } \mu\text{mol mol}^{-1}$; $\text{O}_3 = 6 \text{ nmol mol}^{-1}$, charcoal filtered air), elevated O_3 ($62 \pm 17 \text{ nmol mol}^{-1}$: 7 hours, 10:00-17:00 CET), elevated CO_2 ($600 \text{ } \mu\text{mol mol}^{-1}$ during daytime) and a combination (elevated O_3 and CO_2). Ozone was generated from pure oxygen by an O_3 generator (Model PZ-1C, Kofloc, Kyoto, Japan). A proportional-integrative-differential (PID) control algorithm was applied to maintain the desired concentration of O_3 . The 16 open-top chambers ($1.2\text{m} \times 1.2\text{m} \times 1.2\text{m}$) were set in the experimental forest. Each treatment was replicated four times. The chambers were made of steel frame with polyvinyl chloride film having a transmittance of 88% of full sunlight. Four potted plants per each birch species were set in

each OTC. The plant position was changed every 2-weeks within each OTC and every month between the OTCs in each treatment because of the elimination of positional and chamber effects (Koike et al., 1995a).

2.3 Measurement of plant growth

The height, diameter and number of leaves of the seedlings were measured at the beginning and end of the experiment. Height and diameter increment was calculated as absolute difference between the final and initial diameters. The number of defoliated leaves was estimated by counting the leaf traces. In October 2010 all seedlings were harvested, and were separated into each organ (i.e., stem, branch, leaf and root). The plant organs were dried at 70°C for 1 week and weighed.

2.4 Measurement of leaf gas exchange

At the beginning of September 2010, leaf gas exchange was measured for fully expanded sun leaves, using a portable infra-red gas analyzer (Model 6400, Li-Cor instruments, Lincoln, NE, USA) at controlled values of the leaf temperature (25 °C) and the leaf-to-air vapour pressure deficit (VPD, 1.2 kPa), according to Watanabe et al. (2011a). One or two seedlings per treatment-chamber combination were selected randomly for the measurements. The intercellular CO₂ concentration (C_i) response curve of the net photosynthetic rate (A), i.e., the A/C_i curve, was obtained by measurements over 12 CO₂ steps (C_a, 380, 300, 220, 140, 60, 300, 380, 600, 800, 1100, 1400, 1700 μmol mol⁻¹ for the seedlings grown in ambient CO₂ treatments and C_a=600 380, 300, 220, 140, 60, 380, 600, 800, 1100, 1400, 1700 μmol mol⁻¹ for

the seedlings grown in elevated CO₂ treatments). The wait-time for stability of net photosynthesis at each level of CO₂ concentration was followed the procedure by Long and Bernacchi (2003). We determined A at the long-term CO₂ concentration (380 and 600 $\mu\text{mol mol}^{-1}$ in the control and elevated treatment, respectively, A_{growth}), and also at 1700 $\mu\text{mol mol}^{-1}$ (A_{max}), and the stomatal conductance at the long-term CO₂ concentrations (G_s). The maximum rate of carboxylation (V_{cmax}) and the maximum rate of electron transport (J_{max}) were calculated from the A/C_i curve (Farquhar et al., 1980; Long and Bernacchi, 2003). The values of the Rubisco (Ribulose-1,5-bisphosphate carboxylase/oxygenase) Michaelis constants for CO₂ (K_c) and O₂ (K_o), and the CO₂ compensation point in the absence of dark respiration (Γ^*), for analysis of the A/C_i curve, were derived according to the methodology of Bernacchi et al. (2001). All gas exchange measurements of A/C_i curve were carried out on days with clear sky between 9:00 and 15:00 CET.

2.5 Statistics

Statistical unit was single OTC. Data were checked for normal distribution (Kolmogorov-Smirnov D test) and homogeneity of variance (Levene's test). The effects of O₃ and CO₂ on leaf gas exchange, and on plant biomass, were tested via two-way analysis of variance (ANOVA). Result were considered significant at $p < 0.05$; a tendency toward significance was considered when $0.05 < p < 0.1$. All statistical analyses were performed with SPSS software (10.0, SPSS, Chicago, USA).

3. Results

3.1 Biomass and growth of seedlings

Elevated CO₂ significantly increased the diameter increments of the Monarch birch (Fig. 1). Height increments in the mountain birch were decreased with elevated CO₂. Attached leaves of mountain birch showed a tendency to be decreased by O₃ on Oct 6th (Fig. 2). The tendency to decrease in attached leaves was also found in monarch birch on Sep 29th. Defoliated leaves of the mountain birch tended to be larger with elevated O₃ (Fig. 3).

At the end of the experimental period, no significant difference in whole plant biomass was found between the elevated O₃ and/or CO₂ treatments for the mountain birch and Monarch birch (Table 1); for the white birch, whole plant mass tended to be larger with elevated CO₂. Interactive effect of O₃ and CO₂ on leaf biomass was recorded ($P=0.081$) in mountain birch. With elevated O₃, the branch biomass of the mountain birch tended to decrease. Elevated CO₂ increased the root biomass of the white birch. Interactive effect of O₃ and CO₂ on the ratio of shoot dry mass (stem + branch) to root dry mass (S/R ratio) of Monarch birch seedlings was found between the treatments, while elevated CO₂ reduced the S/R ratio of the mountain birch ($P=0.038$) and white birch ($P=0.058$).

3.2 Photosynthetic parameters

Leaf gas-exchange measurements showed that O₃ exposure significantly reduced A_{growth} , A_{max} , V_{cmax} and J_{max} of the mountain birch (Table 2). Elevated CO₂ significantly increased A_{growth} of the mountain birch and white birch. No significant increase in A_{growth} was

found for the Monarch birch. A decrease in V_{cmax} under elevated CO_2 was also found in the Monarch birch.

4. Discussion

Late summer exposure to O_3 reduced branch biomass (Table 1) and accelerated leaf abscission in the mountain birch (Fig. 3). In the present study, a reduction in A_{growth} was also found under elevated O_3 for the mountain birch (Table 2). Maintenance of O_3 -injured leaves may raise respiratory costs for repair of the damaged parts (Matyssek and Sandermann, 2003). As a result, leaf abscission may be accelerated. Growth of long shoots is known to result from current year assimilation (Linder, 1987; Koike, 1995b), and thus is limited by O_3 -induced early defoliation (e.g., Matyssek et al., 1993).

Late summer exposure to elevated CO_2 induced down-regulation of photosynthesis in the Monarch birch (Table 2). The present result shows that the photosynthetic down-regulation was due not to stomatal closure but to a reduction in the assimilation capacity, such as V_{cmax} (Table 2). As a result, the value of A_{growth} for Monarch birch under elevated CO_2 did not differ from that under ambient conditions. Similar findings were reported from the result of a Free-Air CO_2 Enrichment (FACE) experiment on Monarch birch (Eguchi et al., 2008; Watanabe et al., 2011b). Elevated CO_2 reduced the S/R ratio of mountain birch and white birch. This implies that elevated CO_2 stimulated root growth in the mountain birch and white birch, rather than aboveground growth. In contrast, Monarch birch did not show an enhancement of root growth in elevated CO_2 . Absence of stimulation of root development at elevated CO_2 may limit the acquisition of

nutrients by the Monarch birch. Although biomass production was initially accelerated by high CO₂, the excess of carbon gain and deficiency of nutrients may cause dilution of nutrients with rapid growth, and thus depressed Rubisco activity (Watanabe et al., 2011a). The insufficient sink capacity resulting from environmental and/or genetic factors may also cause acclimation of photosynthetic capacity under elevated CO₂ conditions (Long et al., 2004; Ainsworth and Rogers, 2007).

Concurrent exposure of elevated O₃ and CO₂ in late summer changed the allocation to each organ for Monarch birch (Table 1). Interactive effect of O₃ and CO₂ on S/R ratio of Monarch birch seedlings was found. Monarch birch appeared to allocate photosynthates to leaf production rather than other organs as a compensative response to O₃ under elevated O₃ and CO₂ conditions during early period of the experiment (Fig. 2). Previous study also reported that the compensative leaf growth in response to elevated O₃ was stimulated under enhanced CO₂ conditions for *Fagus crenata* seedlings (Watanabe et al., 2010b). Elevated CO₂ is expected to enhance A_{growth} during the early period of the experiment as a short-term response to elevated CO₂, and thus may develop compensatory leaf growth. The observed compensatory response to O₃ may depend on CO₂ supply (Kolb and Matyssek, 2001).

In late summer, exposure to elevated CO₂ did not ameliorate the adverse effects of O₃ on dry mass or photosynthetic parameters that were observed in the mountain birch (Table 1; 2). Previous studies by season-long exposure to O₃ and CO₂ showed the compensatory effect of elevated CO₂, and the ameliorating effect of CO₂ may be caused by

reducing G_s , i.e., limiting O_3 uptake by stomata (Broadmeadow et al., 1999; Matsumura et al., 2005). In the present study, however, late summer exposure to elevated CO_2 did not induce stomatal closure in the three birch species studied (Table 2). The result is supported by a phytotron study of European beech (*Fagus sylvatica*) (Grams et al., 1999), which found that a reduction of G_s was induced by elevated CO_2 in early summer, but the effects of CO_2 on G_s diminished in late summer. Stomatal aperture is also affected by leaf-to-air vapor pressure (VPD) in summer, because of high temperatures. Heath (1998) found that stomatal sensitivity to VPD was reduced under elevated CO_2 . Moreover, O_3 is known to induce sluggish stomatal response to light (e.g., Hoshika et al., 2012), VPD (Grulke et al., 2007; Uddling et al., 2009) and CO_2 (Onandia et al., 2011). A reduction in stomatal responsiveness to CO_2 under high VPD conditions may decrease the protective effects of elevated CO_2 against O_3 damage (Heath, 1998).

In conclusion, our hypothesis was supported by the results of birch responses to CO_2 and O_3 in late summer. The result presented here suggests that elevated CO_2 did not exert any ameliorating effect on the impact of late summer exposure to O_3 on birch species. Surface concentrations of O_3 are expected to increase continuously in East Asian countries (Yamaji et al., 2008). Only two reports are available on the combined effects of O_3 and elevated CO_2 on tree species native to Japan (Matsumura et al. 2005; Watanabe et al., 2010b). Our findings contribute to develop knowledge of forest dynamics of birch species in northern Japan under the atmospheric conditions expected in future, with elevated O_3 and CO_2 .

Acknowledgements

Thanks for financial support are due to the Environment Research & Technology Development Fund of Japan (B-1105) and for a Grant-in-aid from the Japanese Society for Promotion of Science (Type B 23380078, and Young Scientists B 24710027 and ditto B 24780239).

REFERENCES

- Ainsworth, E.A. & Rogers, A. (2007). The response of photosynthesis and stomatal conductance to rising [CO₂]: mechanisms and environmental interactions. *Plant, Cell and Environment*, 30, 258-270.
- Barnes, J.D., Pfirrmann, T., Steiner, K., Lütz, C., Busch, U., Küchenhoff, H. & Payer H.-D. (1995). Effects of elevated CO₂, elevated O₃ and potassium deficiency on Norway spruce [*Picea abies* (L.) Karst.]: seasonal changes in photosynthesis and non-structural carbohydrate content. *Plant, Cell and Environment*, 18, 1345-1357.
- Bernacchi, C.J., Singsaas, E.L., Pimentel, C., Portis Jr., A.R. & Long, S.P. (2001). Improved temperature response functions for models of Rubisco-limited photosynthesis. *Plant, Cell and Environment*, 24, 253-259.
- Broadmeadow, A., Heath, J. & Randle, T.J. (1999). Environmental limitations to O₃ uptake - some key results from young trees growing at elevated CO₂ concentrations. *Water, Air, and Soil Pollution*, 116, 299–310.
- Bytnerowicz, A., Omasa, K. & Paoletti, E. (2007). Integrated effects of

- air pollution and climate change on forests: A northern hemisphere perspective. *Environmental Pollution*, 147, 438–445.
- Eguchi, N., Karatsu, K., Ueda, T., Funada, R., Takagi, K., Hiura, T., Sasa, K. & Koike, T. (2008). Photosynthetic responses of birch and alder saplings grown in a free air CO₂ enrichment system in northern Japan. *Trees - Structure and Function*, 22, 437-447.
- Evans, L.T. (1963) Environmental control of plant growth, Academic press.
- Farquhar, G.D., von Caemmerer, S. & Berry, J.A. (1980). A biochemical model of photosynthetic CO₂ assimilation in leaves of C3 species. *Planta*, 149, 78–90.
- Grams, T.E.E., Anegg, S., Haberle, K-H., Langebartels, C. & Matyssek, R. (1999). Interactions of chronic exposure to elevated CO₂ and O₃ levels in the photosynthetic light and dark reactions of European beech (*Fagus crenata*). *New Phytologist*, 144, 95-107.
- Grulke, N.E., Neufeld, H.S., Davison, A.W., Roberts, M. & Chappelka, A.H. (2007). Stomatal behavior of ozone-sensitive and -insensitive coneflowers (*Rudbeckia laciniata* var. *digitata*) in Great Smoky Mountains National Park. *New Phytologist*, 173, 100-109.
- Heath, J. (1998). Stomata of trees growing in CO₂-enriched air show reduced sensitivity to vapour pressure deficit and drought. *Plant, Cell and Environment*, 21, 1077–1088.
- Hokkaido Prefecture. (2011) Forestry statistics of Hokkaido in the fiscal year 2010. <http://www.pref.hokkaido.lg.jp/sr/sum/kcs/rin-toukei/22rtk.htm>, retrieved on 10th Jan. 2012. (In Japanese)
- Hoshika, Y., Watanabe, M., Inada, N. & Koike, T. (2012).

Ozone-induced stomatal sluggishness develops progressively in Siebold beech (*Fagus crenata*). *Environmental Pollution*, 166, 152-156.

Karnosky, D.E., Oksanen, E., Dickson, R.E. & Isebrands, J.G. (2001). Impact of interacting greenhouse gases on forest ecosystems. In: Karnosky, D.F., Scarascia-Mugnozza, G., Geulemans, R. & Innes, J.L. (eds.) The impacts of carbon dioxide and other greenhouse gases on forest ecosystems. CABI, Wallingford, pp. 253-267.

Karnosky, D.F., Percy, K.E., Chappelka, A.H., Simpson, C. & Pikkarainen, J.M. (2003). Air Pollution, Global Change and Forests in the New Millennium. Elsevier, Amsterdam, 469 pp.

King, J.S., Kubiske, M.E., Pregitzer, K.S., Hendrey, G.R., McDonald, E.P., Giardina, C.P., Quinn, V.S. & Karnosky D.F. (2005). Tropospheric O₃ compromises net primary production in young stands of trembling aspen, paper birch and sugar maple in response to elevated atmospheric CO₂. *New Phytologist*, 168, 623-636.

Koike, T. (1988). Leaf structure and photosynthetic performance as related to the forest succession of deciduous broad-leaved trees. *Plant Species Biology*, 3, 77-87.

Koike, T. (1995a). Effects of CO₂ in interaction with temperature and soil fertility on the foliar phenology of alder, birch, and maple seedlings. *Canadian Journal of Botany*, 73, 149-157.

Koike, T. (1995b) Physiological ecology of the growth characteristics of Japanese mountain birch in northern Japan: a comparison with Japanese mountain white birch, In: E.O. Box et al. (eds.) Vegetation Science in Forestry: Global Perspective based on Forest Ecosystems of East & Southeast Asia. Kluwer Academic Publishers, The

Netherlands, pp.409-422.

- Koike, T. & Sakagami, Y. (1985). Comparison of the photosynthetic responses to temperature and light of *Betula maximowicziana* and *Betula platyphylla* var. *japonica*. *Canadian Journal of Forest Research*, 15, 631–635.
- Kolb, T.E. & Matyssek, R. (2001). Limitation and perspectives about scaling ozone impacts in trees. *Environmental Pollution*, 115, 373–393.
- Larcher, W. (2004). *Physiological plant ecology* 4 ed. Springer Verlag, Berlin, Heidelberg.
- Linder, S. (1987). Responses to water and nutrients in coniferous ecosystems. In: E.-D. Schulze and H.Z. Wolfer (eds.). *Potentials and Limitations of Ecosystems Analysis*. Ecological Studies, 61, Springer-Verlag, Berlin Heidelberg, pp.180-202.
- Long, S.P. & Bernacchi, C.J. (2003). Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. *Journal of Experimental Botany*, 54, 2393-2401.
- Long, S.P., Ainsworth, E.A., Rogers, A. & Ort, D.R. (2004). Rising atmospheric carbon dioxide: plants FACE the future. *Annual Review of Plant Biology*, 55, 591-628.
- Lütz, C., Anegg, S., Gerant, D., Alaoui-Sosse, B., Gerard, J. & Dizengremel, P. (2000). Beech trees exposed to high CO₂ and to simulated summer ozone levels: Effects on photosynthesis, chloroplast components and leaf enzyme activity. *Physiologia Plantarum*, 109, 252-259.
- Mao, Q., Watanabe, M. & Koike, T. (2010). Growth characteristics of

- two promising tree species for afforestation, birch and larch in the northeastern part of Asia. *Eurasian Journal of Forest Research*, 13, 69-76.
- Matsumura, H., Mikami, C., Sakai, Y., Murayama, K., Izuta, T., Yonekura, T., Miwa, M. & Kohno, Y. (2005). Impacts of elevated O₃ and/or CO₂ on growth of *Betula platyphylla*, *Betula ermanii*, *Fagus crenata*, *Pinus densiflora* and *Cryptomeria japonica* seedlings. *Journal of Agricultural Meteorology*, 60, 1121-1124.
- Matyssek, R., Keller, T. & Koike, T. (1993). Branch growth and leaf gas exchange of *Populus tremula* exposed to low ozone concentrations throughout two growing seasons. *Environmental Pollution*, 79, 1-7.
- Matyssek, R., Günthardt-Georg, M.S., Maurer, S. & Keller, T. (1995). Nighttime exposure to ozone reduces whole-plant production in *Betula pendula*. *Tree Physiology*, 15, 159-165.
- Matyssek, R. & Sandermann, H. (2003). Impact of ozone on trees: an ecophysiological perspective. In: Esser, K., Lüttge, U., Beyschlag, W. & Hellwig F (eds). *An Ecophysiological Perspective. Progress in Botany vol.64*. Springer-Verlag, Berlin Heidelberg, pp. 349-404.
- Naja, M. & Akimoto, H. (2004). Contribution of regional pollution and long-range transport to the Asia-Pacific region: Analysis of long-term ozonesonde data over Japan. *Journal of Geophysical Research*, 109, D21306, doi: 10.1029/2004JD004687.
- Nelson, N. D. & Isebrands, J. G. (1983). Late-season photosynthesis and photosynthate distribution in an intensively-cultured *Populus nigra* x *P. laurifolia* clone. *Photosynthetica* 17, 537-549
- Noguchi, I. (2011). Atmospheric condition and forest declining of

mountain birch in the somma of Lake Mashu. *Northern Forestry* 63, 29 (in Japanese).

- Ohara, T., Uno, I., Wakamatsu, S. & Murano, K. (2001). Numerical simulation of the springtime trans-boundary air pollution in East Asia. *Water, Air, and Soil Pollution*, 130, 295-300.
- Ohno, Y., Umeki, K., Yasaka, M., Watanabe, I. & Takiya, M. (2010). Competition as a predisposing factor of crown dieback in a secondary forest of *Betula maximowicziana* in Hokkaido, northern Japan. *Journal of Forest Research*, 15, 161-168.
- Onandia, G., Olsson, A.-K., Barth, S., King, J.S. & Uddling, J. (2011). Exposure to moderate concentrations of tropospheric ozone impairs tree stomatal response to carbon dioxide. *Environmental Pollution*, 159, 2350-2354.
- Paoletti, E. & Grulke, N.E. (2005). Does living in elevated CO₂ ameliorate tree response to ozone? A review of stomatal responses. *Environmental Pollution*, 137, 483-493.
- Serengil, Y., Augustaitis, A., Bytnerowicz, A., Grulke, N., Kozovitz, A.R., Matyssek, R., Müller-Starck, G., Schaub, M., Wieser, G., Coskun, A.A. & Paoletti, E. (2011). Adaptation of forest ecosystems to air pollution and climate change: a global assessment on research priorities. *iForest – Biogeosciences and Forestry*, 4, 44-48.
- Shimizu, H. & Feng, Y.W. (2007). Ozone and/or water stress could have influenced the *Betula ermanii* Cham. forest decline observed at Oku-Nikko, Japan. *Environmental Monitoring and Assessment*, 128, 109-119.
- Takeda, M. & Aihara, K. (2007). Effects of ambient ozone concentrations on beech (*Fagus crenata*) seedlings in the Tanzawa

- mountains, Kanagawa prefecture, Japan. *Journal of Japan Society of Atmospheric Environment*, 42, 107-117 (in Japanese with English summary).
- Takigawa, M., Niwano, M., Akimoto, H. & Takahashi, M. (2007). Development of a one-way nested global-regional air quality forecasting model. *Sola*, 3, 81-84.
- Uddling, J., Teclaw, R.M., Pregitzer, K.S. & Ellsworth, D.S. (2009). Leaf and canopy conductance in aspen and aspen-birch forests under free-air enrichment of carbon dioxide and ozone. *Tree Physiology*, 29, 1367–1380.
- Watanabe, M., Umemoto-Yamaguchi, M., Koike, T. & Izuta, T. (2010b). Growth and photosynthetic response of *Fagus crenata* seedlings to ozone and/or elevated carbon dioxide. *Landscape Ecological Engineering*, 6, 181-190.
- Watanabe, M., Matsuo, N., Yamaguchi, M., Matsumura, H., Kohno, Y. & Izuta, T. (2010a). Risk assessment of ozone impact on the carbon absorption of Japanese representative conifers. *European Journal of Forest Research*, 129, 421-430.
- Watanabe, M., Watanabe, Y., Kitaoka, S., Utsugi, H., Kita, K. & Koike, T. (2011a). Growth and photosynthetic traits of hybrid larch F₁ (*Larix gmelinii* var. *japonica* × *L. kaempferi*) under elevated CO₂ concentration with low nutrient availability. *Tree physiology*, 31, 965-975.
- Watanabe, M., Mao, Q., Novriyanti, E., Ito, H., Ueda, T., Takagi, K., Sasa, K. & Koike, T. (2011b). Photosynthetic activity of 3 birches grown under free air CO₂ enrichment system. *Boreal Forest Research*, 59, 39-40 (in Japanese).

- Watanabe, M., Yamaguchi, M., Matsumura, H., Kohno, Y. & Izuta, T. (2012). Risk assessment of ozone impact on *Fagus crenata* in Japan: consideration of atmospheric nitrogen deposition. *European Journal of Forest Research*, 131, 475-484.
- Yamaguchi, M., Watanabe, M., Matsumura, H., Kohno, Y. & Izuta, T. (2011). Experimental studies on the effects of ozone on growth and photosynthetic activity of Japanese forest species. *Asian Journal of Atmospheric Environment*, 5, 65-78.
- Yamaji, K., Ohara, T., Uno, I., Kurokawa, J., Pochanart, P. & Akimoto, H. (2008). Future prediction of surface ozone over east Asia using models-3 community multiscale air quality modeling system and regional emission inventory in Asia. *Journal of Geophysical Research*, 113, D08306, doi: 10.1029/2007JD008663.

Figure captions

Fig.1 Height increments (A) and stem basal diameter increments (B) of three birch species treated with elevated O_3 (60 nmol mol^{-1} during daytime) and/or CO_2 ($600 \text{ } \mu\text{mol mol}^{-1}$ during daytime) during the experimental period from July to October 2010. Each value is the mean of four replications. The vertical bar indicates standard deviation.

Fig.2 Seasonal variation of number of attached leaves of three birch species (A, mountain birch; B, Monarch birch; C, white birch) treated with elevated O_3 (60 nmol mol^{-1} during daytime) and/or CO_2 ($600 \text{ } \mu\text{mol mol}^{-1}$ during daytime) during the experimental period from July to October 2010. Each value is the mean of four replications. The vertical bar indicates standard deviation.

Fig.3 Number of defoliated leaves of three birch species treated with elevated O_3 (60 nmol mol^{-1} during daytime) and/or CO_2 ($600 \text{ } \mu\text{mol mol}^{-1}$ during daytime) during the experimental period from July to October 2010. Each value is the mean of four replications. The vertical bar indicates standard deviation.

Fig.1

	<i>P</i> value for Two-way ANOVA		
	O ₃	CO ₂	CO ₂ +O ₃
Height increment			
Mountain birch	0.274	0.011	0.136
Monarch Birch	0.848	0.901	0.260
White Birch	0.995	0.578	0.582
Diameter increment			
Mountain birch	0.191	0.123	0.223
Monarch Birch	0.369	0.001	0.328
White Birch	0.741	0.452	0.747

Fig.2

Fig.3

Tables

Table 1 Dry mass of plant organs per plant and the ratio of shoot dry mass (stem + branch) to root dry mass (S/R ratio) of three birch species treated with O₃ (60 nmol mol⁻¹ during daytime) and/or CO₂ (600 µmol mol⁻¹ during daytime) at the end of the experimental period

	Control	O ₃	CO ₂	CO ₂ +O ₃	<i>P</i> value for Two-way ANOVA				
					O ₃	CO ₂	CO ₂ +O ₃		
Mountain Birch									
Stem biomass (g)	5.38 (0.64)	5.25 (0.54)	4.87 (0.44)	4.91 (0.27)	0.857	0.111	0.738		
Branch biomass (g)	1.39 (0.11)	1.25 (0.20)	1.47 (0.20)	1.29 (0.20)	0.095	0.519	0.792		
Leaf biomass (g)	1.80 (0.85)	1.14 (0.14)	1.05 (0.03)	1.29 (0.40)	0.388	0.239	0.081		
Root biomass (g)	13.09 (2.45)	13.18 (1.09)	14.57 (1.52)	14.10 (1.84)	0.837	0.205	0.761		
Whole plant biomass (g)	19.86 (3.12)	19.68 (1.54)	20.92 (1.33)	20.36 (2.17)	0.739	0.436	0.862		
S/R ratio	0.52 (0.04)	0.49 (0.04)	0.45 (0.07)	0.44 (0.05)	0.464	0.038	0.713		
Monarch Birch									
Stem biomass (g)	11.47 (0.36)	11.64 (1.37)	12.62 (0.88)	11.66 (1.43)	0.485	0.307	0.329		
Branch biomass (g)	2.78 (0.62)	2.95 (0.30)	2.83 (0.33)	3.07 (0.28)	0.346	0.674	0.874		
Leaf biomass (g)	2.71 (0.88)	2.70 (0.79)	3.39 (0.66)	3.13 (0.98)	0.754	0.208	0.778		
Root biomass (g)	20.07 (3.64)	17.13 (3.00)	20.69 (4.35)	21.06 (2.93)	0.481	0.220	0.367		
Whole plant biomass (g)	32.90 (6.04)	31.71 (3.89)	36.15 (5.02)	35.80 (4.55)	0.761	0.163	0.868		
S/R ratio	0.72 (0.09)	0.88 (0.12)	0.80 (0.11)	0.71 (0.04)	0.460	0.355	0.025		
White Birch									
Stem biomass (g)	9.31 (0.32)	9.11 (1.05)	9.62 (0.79)	9.80 (1.13)	0.987	0.282	0.672		
Branch biomass (g)	1.88 (0.47)	1.88 (0.20)	1.93 (0.17)	1.78 (0.17)	0.596	0.870	0.600		
Leaf biomass (g)	4.83 (0.61)	5.06 (0.31)	4.94 (0.65)	4.38 (1.10)	0.661	0.456	0.302		
Root biomass (g)	17.39 (3.74)	17.01 (1.20)	21.50 (3.66)	20.81 (4.27)	0.757	0.040	0.929		
Whole plant biomass (g)	28.59 (4.44)	28.00 (2.23)	31.37 (0.97)	32.39 (5.01)	0.905	0.067	0.660		
S/R ratio	0.67 (0.10)	0.67 (0.04)	0.57 (0.06)	0.60 (0.10)	0.725	0.058	0.710		

Each value is the mean (±SD) of 4 replicates.

Table 2 Photosynthetic parameters of three birch species treated with elevated O₃ (60 nmol mol⁻¹ during daytime) and/or CO₂ (600 μmol mol⁻¹ during daytime), measured at the beginning of September 2010.

	Control	O ₃	CO ₂	CO ₂ +O ₃	<i>P</i> value for Two-way ANOVA					
					O ₃	CO ₂	CO ₂ +O ₃			
Mountain Birch										
A _{growth} (μmol m ⁻² s ⁻¹)	12.9 (2.5)	10.9 (1.7)	17.7 (2.3)	15.5 (1.8)	0.069	0.001	0.973			
A _{max} (μmol m ⁻² s ⁻¹)	25.5 (3.5)	21.7 (2.4)	24.4 (3.5)	22.7 (1.5)	0.083	0.973	0.466			
G _s (mmol m ⁻² s ⁻¹)	373.6 (88.5)	298.3 (85.4)	364.8 (65.0)	323.4 (23.1)	0.110	0.763	0.587			
V _{cmax} (μmol m ⁻² s ⁻¹)	54.0 (11.4)	47.0 (5.1)	54.2 (3.9)	45.9 (6.1)	0.047	0.977	0.793			
J _{max} (μmol m ⁻² s ⁻¹)	114.2 (15.1)	99.9 (12.0)	112.7 (11.8)	104.9 (5.8)	0.044	0.523	0.856			
Monarch Birch										
A _{growth} (μmol m ⁻² s ⁻¹)	7.6 (1.1)	7.7 (2.3)	7.1 (1.1)	7.3 (3.1)	0.888	0.694	0.941			
A _{max} (μmol m ⁻² s ⁻¹)	13.3 (1.9)	12.5 (5.4)	9.9 (1.2)	10.2 (2.7)	0.870	0.102	0.745			
G _s (mmol m ⁻² s ⁻¹)	233.3 (45.5)	187.1 (42.7)	223.2 (44.4)	188.8 (102.3)	0.231	0.897	0.856			
V _{cmax} (μmol m ⁻² s ⁻¹)	27.8 (3.9)	26.4 (10.5)	18.8 (2.7)	19.9 (7.6)	0.975	0.045	0.721			
J _{max} (μmol m ⁻² s ⁻¹)	60.6 (9.2)	51.8 (19.8)	45.7 (5.0)	48.7 (12.6)	0.661	0.186	0.376			
White Birch										
A _{growth} (μmol m ⁻² s ⁻¹)	10.3 (1.4)	10.5 (1.7)	12.7 (1.8)	13.7 (1.9)	0.588	0.019	0.696			
A _{max} (μmol m ⁻² s ⁻¹)	17.6 (2.6)	17.5 (2.3)	16.0 (2.5)	18.3 (3.2)	0.368	0.759	0.351			
G _s (mmol m ⁻² s ⁻¹)	232.5 (13.3)	219.6 (23.8)	201.6 (29.4)	233.1 (49.3)	0.568	0.593	0.188			
V _{cmax} (μmol m ⁻² s ⁻¹)	46.3 (9.6)	43.8 (11.5)	41.0 (11.6)	39.6 (8.7)	0.633	0.259	0.893			
J _{max} (μmol m ⁻² s ⁻¹)	82.2 (11.7)	78.1 (13.6)	77.8 (15.7)	82.6 (15.7)	0.955	0.997	0.471			

A_{growth}, net photosynthetic rate at the chosen long-term CO₂ concentration;

A_{max}, net photosynthetic rate at CO₂ saturation; G_s, stomatal conductance for water vapor;

V_{cmax}, maximum rate of carboxylation; J_{max}, maximum rate of electron transport.

Each value is the mean (±SD) of 4 replicates.