

# HOKKAIDO UNIVERSITY

Title	Aceticlastic and NaCI-Requiring Methanogen "Methanosaeta pelagica" sp. nov., Isolated from Marine Tidal Flat Sediment				
Author(s)	Mori, Koji; Iino, Takao; Suzuki, Ken-Ichiro; Yamaguchi, Kaoru; Kamagata, Yoichi				
Citation	Applied and Environmental Microbiology, 78(9), 3416-3423 https://doi.org/10.1128/AEM.07484-11				
Issue Date	2012-05				
Doc URL	http://hdl.handle.net/2115/50384				
Rights	© 2012 by the American Society for Microbiology				
Туре	article (author version)				
File Information	AEM78-9_3416-3423.pdf				



Hokkaido University Collection of Scholarly and Academic Papers : HUSCAP

## Title

2 Aceticlastic and NaCl-requiring methanogen "*Methanosaeta pelagica*" sp. nov., isolated from marine tidal flat sediment.

4

# Authors

- Koji Mori,<sup>1</sup> Takao Iino,<sup>2</sup> Ken-ichiro Suzuki,<sup>1</sup> Kaoru Yamaguchi<sup>1</sup> & Yoichi Kamagata<sup>3</sup>
   <sup>1</sup>NITE Biological Resource Center (NBRC), National Institute of Technology and Evaluation (NITE),
- 2-5-8 Kazusakamatari, Kisarazu, Chiba 292-0818, Japan
   <sup>2</sup>Japan Collection of Microorganisms, RIKEN BioResource Center, 2-1 Hirosawa, Wako, Saitama
- 10 351-0198, Japan

<sup>3</sup>Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology
 (AIST), Central 6, 1-1-1 Higashi, Tsukuba, Ibaraki 400-8511, Japan

# 14 **Corresponding author**

Koji Mori

- 16 NITE Biological Resource Center (NBRC), National Institute of Technology and Evaluation (NITE),
   2-5-8 Kazusakamatari, Kisarazu, Chiba 292-0818, Japan
- 18 Tel, +81-438-20-5763; Fax, +81-438-52-2329; E-mail, mori-koji@nite.go.jp

# 20 Running title

Marine aceticlastic methanogen, Methanosaeta pelagica

### 2 ABSTRACT

Acetate is a key compound for anaerobic organic matter degradation, and so far, two genera,

- 4 *Methanosaeta* and *Methanosarcina*, are only contributors for acetate degradation among methanogens. An aceticlastic methanogen, designated strain 03d30q<sup>T</sup>, was isolated from a tidal flat sediment in
- 6 Futtsu, Japan. The phylogenetic analyses based on 16S rRNA and *mcrA* genes revealed that the isolate belonged to the genus *Methanosaeta*, but the optimal Na<sup>+</sup> concentration for growth shifted to
- 8 marine environments unlike the other known *Methanosaeta* species. The quantitative estimation by using a real-time PCR indicated that the 16S rRNA gene of the genus *Methanosaeta* was detected in
- 10 the sediments and the relative abundance ranged from 3.9% to 11.8% of the total archaeal 16S rRNA genes. Also, the amount of the genus *Methanosaeta* increased with increasing depth and was much
- 12 higher than that of the genus *Methanosarcina*. This is the first report of marine *Methanosaeta* species, and on the basis of phylogenetic and characteristic studies, the novel species is proposed,

14 *"Methanosaeta pelagica"* sp. nov., with types strain 03d30q<sup>T</sup>.

#### 2 INTRODUCTION

The global budget of atmospheric  $CH_4$  is on the order of 500-600 Tg  $CH_4$  per year (26), and 4 approximately 74% of the emitted  $CH_4$  is derived from biological methanogenesis (5, 12, 17). Acetate is a key compound for anaerobic organic matter degradation, and it accounts for two-thirds of

6 total CH<sub>4</sub> generated (18). However, so far, only two genera, *Methanosarcina* and *Methanosaeta* are known as aceticlastic methanogens. *Methanosarcina* is a relative generalist that prefers methanol and

8 methylamine to acetate (2), while *Methanosaeta* is a specialist that uses only acetate (3). Analyses of the complete genome sequences of *Methanosarcina acetivorans*, *Methanosarcina mazei* and

10 *Methanosaeta thrmophila* reveled that two genera employ different enzymes to catalyze the first step of acetoclastic methanogenesis while the core steps of methanogenesis are similar in the two genera

12 (10, 14, 51). In fact, as phenotype, the distinction of affinities for acetate between two genera has been known among the scientists working in this field: *Methanosaeta* shows a much lower minimum

14 threshold for acetate utilization (7-70 μM) than *Methanosarcina* (0.2-1.2 mM) (17). Accordingly, *Methanosaeta* adapts to the low acetate environments, and it was often detected more dominantly than

16 Methanosarcina in the environments such as rice fields (15), landfill sites (24, 34) and anaerobic digestors (25, 49). Methanosaeta is widely distributed in nature and probably the predominant CH<sub>4</sub>

- producer on earth.Methanogens are abundant in habitats where electron acceptors such as oxygen, nitrate and sulfate are
- 20 limiting. In anaerobic marine habitats, owing to the existence of sulfate in seawater, organic matter degradation is usually performed by sulfate-reducing bacteria, and anaerobic oxidation of CH<sub>4</sub> that is
- 22 produced by methanogenesis beneath the sulfate-reducing layers is also performed by ANME clusters (1, 22, 28, 42). Meanwhile, many studies in marine sediments have demonstrated that the
- 24 methanogenesis also occur in the high sulfate concentration layer and the anaerobic methane oxidation layer as well as the deep sulfate-depleted layer (6, 8, 45). Methylotrophic methanogens containing
- 26 *Methanosarcina* are regarded as major contributors to the CH<sub>4</sub> production here because methylated compounds are not utilized efficiently by sulfate-reducing bacteria and are termed noncompetitive
- substrates (40). In marine sediments, particularly deep sulfate-depleted layer,  $CH_4$  has been considered to originate from  $H_2/CO_2$  rather than acetate (60). To date, methylotrophic, hydrogen and

acetate-utilizing methanogens have been isolated from the marine sediments (13, 20, 52).
 As acetoclastic methanogens, *Methanosarcina* have been isolated from the marine sediments while

- 32 *Methanosaeta* have not yet. However, because acetate concentration of pore water in marine sediments are usually below 20  $\mu$ M (19, 44), it would appear that the conditions are suitable for
- 34 *Methanosaeta* rather than *Methanosarcina*. In the present study, we report the isolation of an aceticlastic methanogen belonging to *Methanosaeta* from tidal flat sediment, and based on the results
- 36 of the phylogenetic relationship and ecological investigation, we will discuss the contribution to  $CH_4$  production from acetate in marine sediments.

#### MATERIALS AND METHODS

- 2 **Microorganisms and growth conditions.** Strain  $03d30q^{T}$  (= NBRC  $105920^{T}$  = DSM  $24271^{T}$ ) was isolated as follows in this study. *Escherichia coli* NBRC 3301 (K-12), *Methanosarcina barkeri*
- 4 NBRC 100474<sup>T</sup>, *Methanosaeta thermophila* NBRC 101360<sup>T</sup>, *Methanosaeta concilii* NBRC 103675<sup>T</sup> and *Methanosaeta harundinacea* NBRC 104789<sup>T</sup> were used for analyses. The cultivating media
- 6 were NBRC media No. 802, 837, 897, 994 and 925 (38), respectively, and the cultivating temperatures were 37°C except for *M. thermophila* (55°C).
- 8 **Study area, sample collection, and measurements.** Futtsu tidal flat is a foreshore sandy flat situated from the estuary of Koito River to sea in Tokyo Bay, Chiba Prefecture, Japan. Sediment
- 10 samples were collected on 24 June 2005 by using a peat sampler (model DIK-105A; Daiki Rika Kogyo Co., Ltd.). For the cultivation, the collected sample was transported to our laboratory in a
- 12 sealed nylon bag with an O<sub>2</sub>-absorbing and a CO<sub>2</sub>-generating agent (Anaero-Pack; Mitsubishi Gas Chemical) and inoculated to a medium as soon as possible. For the molecular analysis, the collected
- samples in 2-ml tubes were stored at -80°C until extraction of the microbial DNA.Temperature of the sediment was measured on site by a thermometer. Salinity and pH of pore water
- 16 were measured on site by a refractometer (ATC-S/Mill-E; ATAGO) and a pH meter (D-13; Horiba), respectively.
- 18 **Enrichment and isolation of acetoclastic methanogen.** In order to enrich and isolate aceticlastic methanogens, NBRC medium No. 1108 was used. The medium was composed of the following salts
- and solutions (liter<sup>-1</sup>): 1.19 g KH<sub>2</sub>PO<sub>4</sub>, 0.21 g K<sub>2</sub>HPO<sub>4</sub>, 3.05 g MgCl<sub>2</sub>·6H<sub>2</sub>O, 0.15 g CaCl<sub>2</sub>·2H<sub>2</sub>O, 0.54 g
   NH<sub>4</sub>Cl, 20 g NaCl, 6.56 g sodium acetate, 1.5 g Bacto Yeast Extract (Difco), 0.4 g Bacto Tryptone
- (Difco), 0.14 g Coenzyme M (2-Mercaptoethanesulfonic acid sodium), 2.5 g NaHCO<sub>3</sub>, 1 mg resazurin,
  2 ml trace element solution (36), 10 ml vitamin solution (36), and 0.36 g Na<sub>2</sub>S·9H<sub>2</sub>O. The medium
- was prepared in culture vessel with butyl-rubber stoppers and aluminum caps under  $N_2/CO_2$  (80/20 [vol/vol]). For the enrichments and routine cultivations, 50-ml culture vessel containing 20 ml of the
- 26 liquid medium were used.

Study of phenotypic and chemotaxonomic futures. The optimum temperature, pH, and  $Na^+$  concentration for growth were determined by examining the time course of  $CH_4$  production. The pH

- concentration for growth were determined by examining the time course of  $CH_4$  production. The pH of medium was adjusted by adding  $Na_2CO_3$  or HCl at room temperature.  $Na^+$  concentration range for
- 30 growth of each *Methanosaeta* species was examined by using each medium adding various concentration of NaCl. The gas phase of the cultures was analyzed by gas chromatography using a
- thermal conductivity detector and a Molecular Sieve 60/80 column (both of Shimadzu).
   The G + C content of extracted genomic DNA was analyzed using high performance liquid
- chromatography (HPLC) with a reverse-phase column (31).
  Observation. An Olympus AX70 microscope was used for routine observation. For the
  observation of ultrathin section of cells, cells were prepared by using rapid freezing and the
  freeze-substitution method (63). After the procedures, cells were embedded, stained (63), and
- 38 observed using a transmission electron microscope (H-7000; Hitachi) operating at 100 kV.

DNA extraction, PCR, sequencing, and quantitative PCR. The extraction and purification of the genomic DNA of microorganisms were performed as previously described (37). The 16S rRNA gene of microorganisms was amplified and sequenced as previously described (35). The partial *mcrA* 

- 4 (alpha subunit of methyl-coenzyme M reductase) gene was amplified using primers MR1mod (5'-GAC CTS CAC TWC GTV AAC AAC-3') and ME2mod (5'-TCA TBG CRT AGT TNG GRT
- 6 AGT-3') that were slightly modified primers MR1 (50) and ME2 (16) based on the genome sequence of *M. thermophila* (CP000477). The PCR amplification and sequencing of the products were
- 8 performed as previously described conditions (33).The genomic DNA of the microbial cells in the sediments was extracted using Power Soil DNA
- 10 Isolation Kit (MoBio). An analysis of a quantitative PCR was performed as previously described (35) except for primer sets and standards. The following 16S rRNA gene-targeted PCR primer sets
- 12 were used: Bac349F and Bac806R (55) for the domain *Bacteria*, A344F (4) and ARC915 (54) for the domain *Archaea*, MX825cm (5'- GCT AGG TGT CRG YCA CGG TGC GA-3') modified MX825c
- 14 (7) and ARC915 for the genus *Methanosaeta*, and MS821c (5'- GCT CGC TAG GTG TCA GGC ATG GCG-3') (47) and ARC915 for the genus *Methanosarcina*. For the construction of the template
- 16 standards for each primer set, the dilution series of the 16S rRNA gene PCR products of *E. coli* (for *Bacteria*), *M. thermophila* (for *Archaea* and *Methanosaeta*) and *M. barkeri* (for *Methanosarcina*) were
- 18 used. These PCR products were used in each real-time PCR analysis to calculate the copy number of the 16S rRNA genes in the sediment samples. Two tests were performed to confirm the specificity of
- 20 the real-time PCR assay. First, a melting-curve analysis was performed after the amplification. Second, the sizes of the PCR products were confirmed by gel electrophoresis. In addition to them, as
- for the assays of the genera *Methanosaeta* and *Methanosarcina*, the real-time PCR products were cloned and sequenced using the TOPO TA Cloning Kit (Invitrogen) and M13 primer set to confirm the
- 24 specificity.

Phylogenetic analysis. Phylogenetic analyses were carried out using the16S rRNA gene sequence and the deduced amino acid sequence of the *mcrA* gene. The 16S rRNA gene sequences were aligned against an ARB data set using the ARB program (29), and the resulting alignment was

- 28 manually refined based on primary and secondary structural considerations. The data set of the *mcrA* gene was aligned using program Clustal X (56). Phylogenetic trees were inferred by
- 30 neighbor-joining (NJ) with the software packages of Clustal X, and 1,000 replicate data sets were used for bootstrap analysis (48, 56).
- 32 **Nucleotide sequence accession number.** The GenBank/EMBL/DDBJ accession numbers for the sequences of 16S rRNA and *mcrA* genes are XXXXXX-XXXXXX.
- 34

# RESULTS

- 36 **Site description.** Core samples of sandy sediments were collected from 2 points (samples 031 and 032) at low tide. Because the sample 031 was near a sea grass and the sediment was blackish in
- 38 whole, the conditions were inferred to be totally anaerobic. As for the 032 sample, the sediment

gradually came to black from gray with increasing depth, and the deeper sediment probably maintained anaerobic conditions. Each sediment sample at 3 depths (10, 35 and 60 cm) was selected for the molecular analysis, and the 032 samples at 35 and 60 cm in depths were used for the enrichment of aceticlastic methanogens. The temperature, pH and salinity of the sediments were 21°C, 7.3 and 32‰ on average, respectively. Those of the surrounding sea water were 25°C, 7.8 and

6 31‰, respectively.

**Enrichment and isolation.** For the primary enrichment of aceticlastic methanogens, 1 g of each sediment sample was inoculated into the medium and incubated at 20 and 30°C. After a 1-month cultivation, microbial growth and CH<sub>4</sub> formation were observed in all cultures, and then enriched

10 cultures were transferred to the fresh medium. The procedure was repeated several times. After that, microscopic observation indicated that *Methanosaeta*-like cells were enriched in the culture of

12 the sample 032 (60 cm in depth) at 30°C. Therefore, we focused on the enrichment and tried to obtain a pure culture. Because growth of enriched microorganisms could not be achieved

- 14 successfully on medium solidified with 2% agar, an attempt was made to isolate microorganisms by the serial dilution method. We found that the *Methanosaeta*-like cells were dominant and the
- 16 transferred culture at higher dilution (> $10^7$ ) continuously grew in the medium removing yeast extract and tryptone although more than 3 months were necessary for confirming its growth. After repeating
- 18 the maximum dilution several times, an aceticlastic methanogen, designated strain 03d30q<sup>T</sup>, was obtained (Fig. 1A). The purity of the isolate was verified by determination of the 16S rRNA gene
- 20 sequence amplified from the extracted DNA using the various primer sets (36) and inoculation into following media: the medium under H<sub>2</sub>/CO<sub>2</sub> (80/20 [vol/vol], 150 kPa) as substitute for N<sub>2</sub>/CO<sub>2</sub>; the
- 22 medium supplemented with 10 mM sodium lactate, 10 mM sodium sulfate and 10 mM sodium thiosulfate; NBRC medium No. 325 (38) for aerobic heterotrophs. The results showed that a trace of
- contaminant was not observed and culture of strain 03d30q<sup>T</sup> was pure.
   Phylogenetic analysis. Almost the full-length 16S rRNA gene sequences of strain 03d30q<sup>T</sup> (1,383)
- 26 bp; XXXXXXX) and *M. concilii* NBRC 104675<sup>T</sup> (1,407 bp; XXXXXXX) were determined. Phylogenetic analysis based on the 16S rRNA gene sequence (Fig. 2) indicated that strain  $03d30q^{T}$
- 28 belonged to the genus *Methanosaeta* and sequence similarities between the isolate and the valid species, *M. concilii*, *M. thermophila* and *M. harundinacea* were 92.8, 92.5 and 97.0%, respectively.
- 30 In the genus *Methanosaeta*, strain 03d30q<sup>T</sup> formed a cluster with environmental clone sequences retrieved from saline environments such as hydrothermal sediments, cold-seep sediments, oil
- 32 reservoirs, marine methane hydrates and ancient seawater (11, 20, 32, 39, 43, 46) although it included the sequences from terrestrial environments and *M. harundinacea*, which were isolated from a UASB
- 34 reactor treating beer-manufacture wastewater (30). The cluster was clearly separated from the ANME clusters that were often retrieved from marine sediments. The sequence divergence for the

36 cluster was 9.4%.

In addition to the 16S rRNA gene sequence, the *mcrA* gene sequences of strain  $03d30q^{T}$  (1,160 bp; 38 XXXXXX), *M concilii* NBRC  $103675^{T}$  (1,160 bp; XXXXXX) and *M. harundinacea* NBRC

104789<sup>T</sup> (1,160 bp; XXXXXXX) were also determined. The neighbor-joining tree for McrA (Fig. 3)

- demonstrated that strain 03d30q<sup>T</sup> was also associated with the genus *Methanosaeta* and the similarities between the isolate and *M. concilii*, *M. thermophila* and *M. harundinacea* were 82.2, 85.6 and 90.9%,
   remeatively.
- 4 respectively.
- Physiology of the isolate and Na<sup>+</sup> requirement. Morphologically, the cells of strain 03d30q<sup>T</sup> were
  weakly auto-fluorescent rods, and many multicellular filaments were microscopically observed (Fig. 1A). There was a tubular sheath enclosing some cells and the individual cells within the sheath had
  cytoplasmic membrane (Fig. 1B). The G+C content of the genomic DNA of strain 03d30q<sup>T</sup> was 45.4 mol%. Strain 03d30q<sup>T</sup> utilized acetate as the sole energy source, and yeast extract stimulated its
- 10 growth. Strain  $03d30q^{T}$  only used acetate as energy and carbon sources. Growth and methane formation were not observed on H<sub>2</sub>/O<sub>2</sub> (80/20; 125 kPa), formate (40 mM), methanol (20 mM),
- 12 ethanol (20 mM), methylamine (20 mM), dimethylamine (20 mM), trimethylamine (20 mM) or dimethyl sulfide (4 mM). Strain 03d30q<sup>T</sup> was able to grow between 20°C and 35°C, and pH 6.0 and
- 14 7.8: its optimum growth conditions were 30°C and pH 7.5. Effect of Na<sup>+</sup> concentration in the medium on growth as  $CH_4$  production of strain 03d30q<sup>T</sup> and *Methanosaeta* species were shown in Fig.
- 16 4. The Na<sup>+</sup> concentrations for growth of strain  $03d30q^{T}$  ranged from 0.20 to 0.80 M, with an optimum of 0.28 M. The other *Methanosaeta* species optimally grew in each medium containing the
- 18 lowest Na<sup>+</sup> concentration. The upper Na<sup>+</sup> concentration for growth of *M. concilii*, *M. thermophila* and *M. harundinacea* were 0.40, 0.56 and 0.34 M, respectively. Strain  $03d30q^{T}$  had a doubling time
- of 12.4 day under the optimum growth conditions (30°C, pH 7.5 and in 0.28M Na<sup>+</sup>).
   Environmental quantification. For the field estimation of aceticlastic methanogens in the tidal flat
- 22 sediment, quantitative real-time PCR assays using the 16S rRNA gene copy number were performed (Fig. 5). The relative abundance of the genera *Methanosaeta* and *Methanosarcina* were calculated as
- the percentages of the total archaeal 16S rRNA gene copy number in the DNA extracts (Table 1). There was a trend that *Bacteria* decreased and *Archaea* increased with increasing depth. Like the
- tendency of *Archaea*, the amount of the genus *Methanosaeta* increased with increasing depth although their proportions in the total *Archaea* ranged from 3.9 to 11.8% and did not correlate with depth. The
- amount of the genus *Methanosarcina* was less than one-tenth of that of the genus *Methanosaeta*, and at a maximum, the proportion was 0.9% of the total *Archaea*.
- 30

## DISCUSSION

- 32 **Ecology** In the present study, we firstly succeeded in the isolation of aceticlastic methanogen belonging to the genus *Methanosaeta* from marine environments. The phylogenetic analyses based
- 34 on the 16S rRNA gene have indicated that *Methanosaeta* species inhabit saline environments such as marine sediments, gas hydrate sediments, hydrothermal sediments, oil reservoir and natural gas filed
- 36 (11, 20, 32, 39, 43, 46), although it is difficult to distinguish clearly between saline and terrestrial *Methanosaeta* species by the phylogenetic analysis (Fig. 2). These analyses has also suggested that
- 38 *Methanosaeta* species detected as clonal genes in saline environments are probably the saline-adapted

species. Our isolated strain is obviously halophilic and adapts to saline environments unlike the other

- 2 *Methanosaeta* species isolated so far (Fig. 4). We firstly demonstrated that *Methanosaeta* species can grow in saline environments and saline-adapted *Methanosaeta* species are ubiquitous in saline
- 4 environments. Accordingly, *Methanosaeta* species seems to contribute to anaerobic acetate degradation in saline environments as well as terrestrial environments.
- 6 There are much larger areas of non-seep marine sediments, but these have been much less studied. Some reports revealed that the constituent members in the sediments were similar to those in some
- 8 seep sediment (21, 23, 44, 57, 61, 62). Global  $CH_4$  production in marine sediments is very significant but the large fraction is converted to  $CO_2$  by ANME in seep and non-seep sediments and by
- 10 aerobic methanotrophs in the water column (1, 41). These CH<sub>4</sub> is mainly produced from methanogens living under the ANME layer and methylotrophic methanogens although hydrogen and
- 12 acetate are mainly consumed by sulfate-reducing bacteria in surface layer (27, 40). However, generally sulfate is depleted at a depth of tens of centimeters to several meters, and potentially
- 14 methanogenesis becomes the dominant terminal oxidation process in the sulfate-depleted layer. In addition, CH<sub>4</sub> production has been detected and methanogen has been retrieved from the sediments
- 16 with high sulfate concentration (6, 8, 19, 20, 45). In the present study, by the real-time PCR analysis using the *Methanosaeta*-specific primer set, the 16S rRNA gene fragments of the genus were detected
- 18 in the tidal flat sediments and the amount increased with increasing depth (Fig. 5). Interestingly, the amount of *Methanosaeta* was ten times higher than that of *Methanosarcina*. The result may reflect
- 20 that the CH<sub>4</sub> production from acetate originates from *Methanosaeta* rather than *Methanosaeta* and *Methanosaeta* contributes to anaerobic organic matter degradation in tidal flat sediments. Acetate
- 22 concentration of pore water in marine sediments are usually below 20 μM (20, 44). Therefore, in marine sediments, *Methanosaeta* has an advantage in acetate utilization than *Methanosarcina* from the
- aspect of affinity for acetate (17).

Although several strains of aceticlastic methanogens belonging to Methanosarcina have been isolated

from marine sediments (13, 20, 52, 58), isotopic evidence suggests that acetate is a minor precursor to methanogenesis and the rate of acetate oxidation to  $CO_2$  is higher than that of  $CH_4$  production in the

- surface layer (44, 60). On the other hand, there is a case that growth of *Methanosaeta* could not be observed although it was detected by the molecular approach (24). Certainly, the enrichment and
- 30 isolation of *Methanosaeta* are difficult tasks among methanogens because of its slow growth. In addition, Wellsbury *et al.* reported that it may be difficult to distinguish between abiological
- thermogenic and biological aceticlastic methanogenesis in marine sediments (9, 59). These facts may suggest that aceticlastic CH<sub>4</sub> generation in marine sediments has previously been underestimated.
- <sup>34</sup> Parkes *et al.* showed that trace of *Methanosaeta* was detected and the acetate concentration was kept at a low level in marine sediments although the rate of CH<sub>4</sub> production from acetate was equivalent to
- 36 one-quarter of that from CO<sub>2</sub> (44), indicating that there are sufficient acetate-utilizing microorganisms to maintain the ecology in the site. Aceticlastic methanogens may thinly and broadly contribute
- 38 acetate degradation in marine sediments, particularly tidal flat sediments where a lot of organic matter

is often supplied to.

- 2 In conclusion, we firstly obtained the marine *Methanosaeta* species from the tidal flat sediment and suggested a possibility for the contributor for acetate degrading in saline environments. Further
- 4 analyses for acetate degradation in tidal flat sediments such as amounts of acetate, sulfate, methanogens and sulfate-reducing bacteria are needed.
- 6 **Taxonomy** Characteristics of strain  $03d30q^{T}$  and *Methanosaeta* species are summarized in Table 2. The phylogenetic tree based on 16S rRNA gene (Fig. 2) revealed that strain  $03d30q^{T}$  belonged to the
- 8 genus *Methanosaeta*, with *M. harundinaceae* as the closest relative (sequence similarity, 97.0%). Similar trend was also found for phylogenetic analysis based on McrA (Fig. 3). These results
- 10 suggested that a new species should be created for the isolate (53). In addition, the G + C content of genomic DNA, the sensitivities to Na<sup>+</sup> concentration and temperature for growth differentiated strain
- 12 03d30q<sup>T</sup> from the other *Methanosaeta* species. In particular, the difference of Na<sup>+</sup> sensitivity was a criterion for discriminating strain 03d30q<sup>T</sup> from the other *Methanosaeta* species: strain 03d30q<sup>T</sup> was a
- 14 sole halophilic species in the genus *Methanosaeta*. For the reasons given above, strain 03d30q<sup>T</sup> represents a novel species within the genus *Methanosaeta*, for which the name *Methanosaeta pelagica*
- 16 sp. nov. is proposed.

## Description of Methanosaeta pelagica sp. nov.

- 18 Methanosaeta pelagica (pe.la'gi.ca. L. fem. adj. pelagica, pertaining to the sea). Single cells are 0.5 μm wide and 2.5-11.0 μm long, and long sheathed chains of cells form. It shows
- 20 weakly blue-green autofluorescence under epifluorescence microscopy. Strictly anaerobic. Acetate is the only substrate for growth and  $CH_4$  production;  $H_2/CO_2$ , formate, methanol, ethanol,
- 22 methylamine, dimethylamine, trimethylamine or dimethyl sulfide do not support growth. Yeast extract is essential for growth. The optimum growth conditions are 30°C, pH 7.5 and in 0.28M Na<sup>+</sup>
- 24 concentration (1.6% NaCl). The DNA G + C content of the type strain is 45.4 mol%.
- The type strain,  $03d30q^{T}$  (= NBRC  $105920^{T}$  = DSM  $24271^{T}$ ), was isolated from a tidal flat sediment in Tokyo Bay, Chiba Prefecture, Japan.

## 28 ACKNOWLEDGMENTS

We thank the staff of Futtsu fishermen's union for collecting the sediment samples. We also thank 30 Kuniko Shimamura and Shinobu Iwasaki for technical support.

## 32 **REFERENCES**

- 1. **Boetius, A., et al.** 2000. A marine microbial consortium apparently mediating anaerobic oxidation of methane. Nature **407:**623-6.
- Boone, D. R., and R. A. Mah. 2001. Genus I. Methanosarcina Kluyver and van Niel 1936,
   400,<sup>AL</sup> emend. Mah and Kuhn 1984a, 266 (Nom. Cos., Opinion 63 of the Jud. Comm. 1986b,
   492), p. 269-276. *In* D. R. Boon and R. W. Castenholz (ed.), Bergey's manual of systematic
   bacteriology second edition, The *Archaea* and the deeply branching and phototrophic *Bacteria*,

vol. 1. Springer.

- Boone, D. R., W. B. Whitman, and Y. Koga. 2001. Family II. Methanosaetaceae fam. nov., p. 289-294. *In* D. R. Boon and R. W. Castenholz (ed.), Bergey's manual of systematic
   bacteriology second edition, The *Archaea* and the deeply branching and phototrophic *Bacteria*, vol. 1. Springer.
- 6 4. Casamayor, E. O., et al. 2002. Changes in archaeal, bacterial and eukaryal assemblages along a salinity gradient by comparison of genetic fingerprinting methods in a multipond solar saltern. Environ. Microbiol. 4:338-48.
- 5. Conrad, R. 2009. The global methane cycle: recent advances in understanding the microbial
   processes involved. Environmental Microbiology Reports 1:285-292.
- 6. Cragg, B. A., et al. 1996. Bacterial populations and processes in sediments containing gas
  hydrates (ODP Leg 146: Cascadia Margin). Earth and Planetary Science Letters 139:497-507.
- 7. Crocetti, G., M. Murto, and L. Bjönsson. 2006. An update and optimisation of
   oligonucleotide probes targeting methanogenic Archaea for use in fluorescence in situ
   hybridisation (FISH). Journal of Microbiological Methods 65:194-201.
- D'Hondt, S., et al. 2004. Distributions of microbial activities in deep subseafloor sediments. Science 306:2216-21.
- de Graaf, W., P. Wellsbury, R. J. Parkes, and T. E. Cappenberg. 1996. Comparison of acetate turnover in methanogenic and sulfate-reducing sediments by radiolabeling and stable isotope labeling and by use of specific inhibitors: evidence for isotopic exchange. Applied and Environmental Microbiology 62:772-7.
- Deppenmeier, U., et al. 2002. The genome of *Methanosarcina mazei*: evidence for lateral gene transfer between bacteria and archaea. J Mol Microbiol Biotechnol 4:453-61.
- Dhillon, A., et al. 2005. Methanogen diversity evidenced by molecular characterization of methyl coenzyme M reductase A (mcrA) genes in hydrothermal sediments of the Guaymas
   Basin. Applied and Environmental Microbiology 71:4592-601.
- Dworkin, M., et al. 2006. The Methanogenic Bacteria, p. 165-207. *In* M. Dworkin, S. Falkow,
   E. Rosenberg, K.-H. Schleifer, and E. Stackebrandt (ed.), The Prokaryotes. Springer New York.
- 30 13. Elberson, M. A., and K. R. Sowers. 1997. Isolation of an aceticlastic strain of Methanosarcina siciliae from marine canyon sediments and emendation of the species
   32 description for Methanosarcina siciliae. Int J Syst Bacteriol 47:1258-61.
- 14. Galagan, J. E., et al. 2002. The genome of *M. acetivorans* reveals extensive metabolic and
   physiological diversity. Genome Res 12:532-42.
- 15. Grosskopf, R., P. H. Janssen, and W. Liesack. 1998. Diversity and structure of the
   methanogenic community in anoxic rice paddy soil microcosms as examined by cultivation
   and direct 16S rRNA gene sequence retrieval. Appl Environ Microbiol 64:960-9.
- 38 16. Hales, B. A., et al. 1996. Isolation and identification of methanogen-specific DNA from

blanket bog peat by PCR amplification and sequence analysis. Appl Environ Microbiol **62:**668-75.

Jetten, M. S. M., A. J. M. Stams, and A. J. B. Zehnder. 1992. Methanogenesis from acetate:
 a comparison of the acetate metabolism in *Methanothrix soehngenii* and *Methanosarcina* spp. FEMS Microbiology Reviews 88:181-198.

 $\mathbf{2}$ 

- Is. Jones, W. J. 1991. Diversity and physiology of methanogens. *In* J. E. Rogers and W. B. Whitman (ed.), Microbial production and consumption of greenhouse gased: methane, nitrogen, oxides, and halomethanes. ASM Press.
- 19. Kendall, M. M., and D. R. Boone. 2006. Cultivation of methanogens from shallow marine
   sediments at Hydrate Ridge, Oregon. Archaea 2:31-8.
- 20. Kendall, M. M., et al. 2007. Diversity of *Archaea* in marine sediments from Skan Bay,
   12 Alaska, including cultivated methanogens, and description of I sp. nov. Appl Environ Microbiol 73:407-14.
- 14 21. Kim, B. S., H. M. Oh, H. Kang, and J. Chun. 2005. Archaeal diversity in tidal flat sediment as revealed by 16S rDNA analysis. Journal of microbiology **43**:144-51.
- 16 22. Knittel, K., T. Losekann, A. Boetius, R. Kort, and R. Amann. 2005. Diversity and distribution of methanotrophic archaea at cold seeps. Appl Environ Microbiol **71**:467-79.
- 18 23. Kopke, B., R. Wilms, B. Engelen, H. Cypionka, and H. Sass. 2005. Microbial diversity in coastal subsurface sediments: a cultivation approach using various electron acceptors and substrate gradients. Applied and Environmental Microbiology 71:7819-30.
- 24. Laloui-Carpentier, W., T. Li, V. Vigneron, L. Mazeas, and T. Bouchez. 2006.
   22 Methanogenic diversity and activity in municipal solid waste landfill leachates. Antonie Van Leeuwenhoek 89:423-34.
- 24 25. Leclerc, M., J. P. Delgenes, and J. J. Godon. 2004. Diversity of the archaeal community in
   44 anaerobic digesters as determined by single strand conformation polymorphism analysis
   and 16S rDNA sequencing. Environmental Microbiology 6:809-19.
- 26. Lelieveld, J., P. J. Crutzen, and F. J. Dentener. 1998. Changing concentration, lifetime and
   climate forcing of atmospheric methane. Tellus B 50:128-150.
- Liu, Y., and W. B. Whitman. 2008. Metabolic, phylogenetic, and ecological diversity of the
   methanogenic archaea. Annals of the New York Academy of Sciences 1125:171-89.
- 28. Losekann, T., et al. 2007. Diversity and abundance of aerobic and anaerobic methane
   oxidizers at the Haakon Mosby Mud Volcano, Barents Sea. Appl Environ Microbiol
   73:3348-62.
- 34 29. Ludwig, W., et al. 2004. ARB: a software environment for sequence data. Nucleic Acids Res.32:1363-71.
- 36 30. Ma, K., X. Liu, and X. Dong. 2006. *Methanosaeta harundinacea* sp. nov., a novel acetate-scavenging methanogen isolated from a UASB reactor. Int. J. Syst. Evol. Microbiol.
   38 56:127-31.

- 31. Mesbah, M., U. Premachandran, and W. B. Whitman. 1989. Precise mesurement of the
   2 G+C content of deoxyribonucleic acid by high-performance liquid chromatography. International Journal of Systematic Bacteriology 39:159-167.
- 4 32. **Mochimaru, H., et al.** 2007. Methanogen diversity in deep subsurface gas-associated water at the Minami-Kanto gas field in Japan. Geomicrobiol. J. **24:**93-100.
- Mori, K., and S. Harayama. in press. *Methanobacterium petrolearium* sp. nov. and *Methanobacterium ferruginis* sp. nov., novel mesophilic methanogens isolated from salty
   environments. Int. J. Syst. Evol. Microbiol.
- 34. Mori, K., R. Sparling, M. Hatsu, and K. Takamizawa. 2003. Quantification and diversity of
   the archaeal community in a landfill site. Can J Microbiol 49:28-36.
- 35. Mori, K., et al. 2008. First cultivation and ecological investigation of a bacterium affiliated
  with the candidate phylum OP5 from hot springs. Appl Environ Microbiol 74:6223-9.
- 36. Mori, K., and K. Suzuki. 2008. *Thiofaba tepidiphila* gen. nov., sp. nov., a novel obligately
   14 chemolithoautotrophic, sulfur-oxidizing bacterium of the *Gammaproteobacteria* isolated from a hot spring. Int. J. Syst. Evol. Microbiol. 58:1885-91.
- Mori, K., H. Yamamoto, Y. Kamagata, M. Hatsu, and K. Takamizawa. 2000. *Methanocalculus pumilus* sp. nov., a heavy-metal-tolerant methanogen isolated from a waste-disposal site. Int. J. Syst. Evol. Microbiol. 50 Pt 5:1723-9.
- 38. NBRC. 2010. NBRC Catalogue of Biological Resources: Microorganisms,
   microorganism-related DNA resources, and human-related DNA resources., 2nd ed. National Institute of Technology and Evaluation (NITE), Chiba, Japan.
- 39. Orcutt, B., et al. 2010. Impact of natural oil and higher hydrocarbons on microbial diversity, distribution, and activity in Gulf of Mexico cold-seep sediments. Deep-sea Res. II, Top. Stud.
   Oceanogr. 57:2008-2021.
- 40. Oremland, R. S., and S. Polcin. 1982. Methanogenesis and sulfate reduction: competitive
   and noncompetitive substrates in estuarine sediments. Appl Environ Microbiol 44:1270-6.
- 41. Orphan, V. J., C. H. House, K. U. Hinrichs, K. D. McKeegan, and E. F. DeLong. 2001.
   Methane-consuming archaea revealed by directly coupled isotopic and phylogenetic analysis. Science 293:484-7.
- 30 42. Orphan, V. J., C. H. House, K. U. Hinrichs, K. D. McKeegan, and E. F. DeLong. 2002. Multiple archaeal groups mediate methane oxidation in anoxic cold seep sediments. Proc Natl
   32 Acad Sci U S A 99:7663-8.
- 43. Pachiadaki, M. G., V. Lykousis, E. G. Stefanou, and K. A. Kormas. 2010. Prokaryotic
   34 community structure and diversity in the sediments of an active submarine mud volcano (Kazan mud volcano, East Mediterranean Sea). FEMS microbiology ecology 72:429-44.
- 44. Parkes, R. J., et al. 2007. Biogeochemistry and biodiversity of methane cycling in subsurface marine sediments (Skagerrak, Denmark). Environmental Microbiology 9:1146-61.
- 45. Parkes, R. J., B. A. Cragg, and P. Wellsbury. 2000. Recent studies on bacterial populations

and processes in subseafloor sediments: A review. Hydrogeology Journal 8:11-28.

- 2 46. Pham, V. D., et al. 2009. Characterizing microbial diversity in production water from an Alaskan mesothermic petroleum reservoir with two independent molecular methods.
   4 Environmental Microbiology 11:176-87.
- 47. Raskin, L., J. M. Stromley, B. E. Rittmann, and D. A. Stahl. 1994. Group-specific 16S
  6 rRNA hybridization probes to describe natural communities of methanogens. Appl Environ Microbiol 60:1232-40.
- 8 48. Saitou, N., and M. Nei. 1987. The neighbor-joining method a new method for reconstructing phylogenetic trees. Mol. Biol. Evol. 4:406-425.
- 10 49. **Sekiguchi, Y., et al.** 1998. Phylogenetic diversity of mesophilic and thermophilic granular sludges determined by 16S rRNA gene analysis. Microbiology **144 ( Pt 9):**2655-65.
- 12 50. **Simankova, M. V., et al.** 2003. Isolation and characterization of new strains of methanogens from cold terrestrial habitats. Syst Appl Microbiol **26:**312-8.
- 14 51. Smith, K. S., and C. Ingram-Smith. 2007. *Methanosaeta*, the forgotten methanogen? Trends in Microbiology 15:150-155.
- Sowers, K. R., S. F. Baron, and J. G. Ferry. 1984. Methanosarcina acetivorans sp. nov., an Acetotrophic Methane-Producing Bacterium Isolated from Marine Sediments. Appl Environ Microbiol 47:971-8.
- 53. Stackebrandt, E., and B. M. Goebel. 1994. Taxonomic note: a place for DNA-DNA
   reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. International Journal of Systematic Bacteriology 44:846-849.
- Stahl, D. A., and R. Amann. 1991. Development and application of nucleic acid probes., p. 205-248. *In* E. Stackebrandt and M. Goodfellow (ed.), Nucleic Acid Technicques in Bacterial
   Systematics. Jon Wiley & Sons, Chichester.
- 55. Takai, K., and K. Horikoshi. 2000. Rapid detection and quantification of members of the
   archaeal community by quantitative PCR using fluorogenic probes. Appl Environ Microbiol
   66:5066-72.
- 56. Thompson, J. D., T. J. Gibson, F. Plewniak, F. Jeanmougin, and D. G. Higgins. 1997. The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by
   quality analysis tools. Nucleic Acids Res. 25:4876-82.
- 57. Treude, T., M. Krüger, A. Boetius, and B. B. Jørgensen. 2005. Environmental control on
   anaerobic oxidation of methane in the gassy sediments of Eckernförde Bay (German Baltic).
   Limnol Oceanogr 50:1771-1786.
- 34 58. von Klein, D., H. Arab, H. Völker, and M. Thomm. 2002. *Methanosarcina baltica*, sp. nov., a novel methanogen isolated from the Gotland Deep of the Baltic Sea. Extremophiles
  36 6:103-110.
- 59. Wellsbury, P., et al. 1997. Deep marine biosphere fuelled by increasing organic matter
  availability during burial and heating. Nature 388:573-576.

- 60. Whiticar, M. J., E. Faber, and M. Schoell. 1986. Biogenic methane formation in marine and
   freshwater environments: CO<sub>2</sub> reduction vs. acetate fermentation--Isotope evidence.
   Geochimica et Cosmochimica Acta 50:693-709.
- 4 61. Wilms, R., H. Sass, B. Kopke, H. Cypionka, and B. Engelen. 2007. Methane and sulfate profiles within the subsurface of a tidal flat are reflected by the distribution of sulfate-reducing
  6 bacteria and methanogenic archaea. FEMS microbiology ecology 59:611-21.
- Wilms, R., et al. 2006. Specific bacterial, archaeal, and eukaryotic communities in tidal-flat
   sediments along a vertical profile of several meters. Applied and Environmental Microbiology
   72:2756-64.
- 10 63. Yamaguchi, K., K.-i. Suzuki, and K. Tanaka. 2010. Examination of electron stains as a substitute for uranyl acetate for the ultrathin sections of bacterial cells. Journal of Electron
   12 Microscopy 59:113-118.

TABLE 1. Relative abundances of Archaea, and the genera Methanosaeta and Methanosarcinacalculated by real-time PCR analyses.

	Sample	Depth	Abundance of the 16S rRNA gene number (%)			
18		( <b>cm</b> )	Archaea	Methanosaeta	Methanosarcina	
			in the total Prokaryotes	in the total Archaea	in the total Archaea	
20	031	10	0.5	6.6	0.2	
		35	1.3	4.6	0.4	
22		60	2.4	4.9	0.7	
	032	10	0.1	11.8	0.9	
24		35	0.5	3.9	0.4	
		60	3.0	7.8	0.2	
26						

- $\mathbf{2}$

**TABLE 2.** Comparison of properties among the species of the genus *Methanosaeta*.

	Species	"M. pelagica"	M. concilii	M. thermophila	M. harundinacea					
16	Type strain	03d30q <sup>T</sup>	GP6 <sup>T</sup>	$\mathbf{P}_{\mathrm{T}}^{\mathrm{T}}$	$8Ac^{T}$					
	Habitat	Tidal-flat sediment	Anaerobic digestor	Anaerobic digestor,	Anaerobic digestor					
18				mud in thermal lake						
	Optimum growth conditions:									
20	Temp. (°C	C) 30	35-40	55-60	34-37					
	pН	7.5	7.1-7.5	6.5-6.7	7.2-7.6					
22	Na <sup>+</sup> conc.	( <b>M</b> ) 0.28	< 0.06	< 0.13	< 0.02					
Effect on yeast extract for growth										
24		Require	Inhibit	Inhibit*	Require					
	G+C conten	t (mol%)								
26		45.4 (HPLC)	50.3 (Tm)	52.7 (HPLC)	55.7 (Tm)					

\*Yeast extract inhibits growth of some strains of *M. thermophila*.

### 2 **FIGURE LEGENDS**

**Fig. 1.** Phase-contrast micrograph of strain  $03d30q^{T}$  (A; bar, 10 µm). Ultrathin section of strain  $03d30q^{T}$  observed with a transmission electron microscope (B; bar, 2 µm).

Fig. 2. Neighbor-joining tree based on 16S rRNA gene sequence of strain 03d30qT and relatives. The probability scores at branch points obtained are indicated by solid circles (above 95%) and open
 circles (above 90%). The sequence obtained from isolates are indicated in bold. Environmental

10 GenBank/EMBL/DDBJ accession numbers are shown in parentheses. Bar, 0.02 substitutions per compared nucleotide site.

clonal sequences retrieved from saline environments are indicated by gravish squares.

12

Fig. 3. Neighbor-joining tree showing the relationships between strain 03d30q<sup>T</sup> and relatives based
on deduced McrA sequences. The probability scores at branch points obtained are indicated by solid
circles (above 95%) and open circles (above 90%). GenBank/EMBL/DDBJ accession numbers are
shown in parentheses. Bar, 0.02 substitutions per compared positions.

- **Fig. 4.** Effect of Na<sup>+</sup> concentration on CH<sub>4</sub> production of strain  $03d30q^{T}$  (solid circles), *M. concilii* (open triangles), *M. thermophila* (open squares) and *M. harundinaceae* (open circles).
- 20

Fig. 5. Copy numbers of 16S rRNA genes of *Bacteria* (triangles), *Archaea* (squares), *Methanosaeta*(circles) and *Methanosarcina* (diamonds) detected by real-time PCR analyses. Open and solid symbols are indicated the sediment samples collected from 031 and 032 points, respectively.



Figure 1, Mori et al.







Figure 4, Mori et al.

