Title	Molecular basis of the dopaminergic system in the cricket <i>Gryllus bimaculatus</i>
Author(s)	Watanabe, Takayuki; Sadamoto, Hisayo; Aonuma, Hitoshi
Citation	Invertebrate neuroscience, 13, 107-123 https://doi.org/10.1007/s10158-013-0153-1
Issue Date	2013
Doc URL	http://hdl.handle.net/2115/52729
Type	article (author version)
File Information	Accepted_2013May_InvNeurosci.pdf

[Instructions for use](#)

IVNS-D-12-00151

Molecular basis of the dopaminergic system in the cricket *Gryllus bimaculatus*

Takayuki Watanabe¹, Hisayo Sadamoto² and Hitoshi Aonuma^{1*}

¹Research Institute for Electronic Science, Hokkaido University, Sapporo, Japan

²Laboratory of Functional Biology, Kagawa School of Pharmaceutical Sciences, Tokushima
Bunri University, Sanuki, Japan

Running title: dopamine related genes in the cricket

*Corresponding author. Hitoshi Aonuma, D.Sc., Research Institute for Electronic Science,
Hokkaido University, Nishi 7 Chome, Kita 12 Jo, Kitaku, Sapporo, Hokkaido 060-0812,
Japan. Tel/Fax: +81-11-706-3832. Email: aon@es.hokudai.ac.jp

Pages: 45 pages

Figure and tables: 10 figures and two tables

ABSTRACT

In insects, dopamine modulates various aspects of behavior such as learning and memory, arousal and locomotion, and is also a precursor of melanin. To elucidate the molecular basis of the dopaminergic system in the field cricket *Gryllus bimaculatus* DeGeer, we identified genes involved in dopamine biosynthesis, signal transduction and dopamine re-uptake in the cricket. Complementary DNA of two isoforms of tyrosine hydroxylase (TH), which convert tyrosine into L-3,4-dihydroxyphenylalanine (L-DOPA), were isolated from the cricket brain cDNA library. In addition, four dopamine receptor genes (*Dop1*, *Dop2*, *Dop3* and *DopEcR*) and a high-affinity dopamine transporter (*DAT*) gene were identified. The two TH isoforms contained isoform-specific regions in the regulatory ACT domain, and showed differential expression patterns in different tissues. In addition, the dopamine receptor genes had a receptor subtype-specific distribution: the *Dop1*, *Dop2* and *DopEcR* genes were broadly expressed in various tissues at differential expression levels, and the *Dop3* gene was restrictedly expressed in neuronal tissues and the testicles. Our findings provide a fundamental basis for understanding the dopaminergic regulation of diverse physiological processes in the cricket.

Keywords: dopamine, tyrosine hydroxylase, dopamine receptors, high-affinity dopamine transporter, *Gryllus bimaculatus*

INTRODUCTION

Biogenic amines function as intracellular messenger molecules that play essential roles in regulating physiological processes and in controlling various behaviors in insects (Evans, 1980; Roeder, 2005). Pharmacological and behavioral studies have been carried out to investigate roles of the biogenic amine system in several insects including the field cricket *Gryllus bimaculatus* DeGeer. For example, pharmacological manipulation of the octopaminergic and dopaminergic systems revealed that these two biogenic amines are involved in appetitive and aversive learning in the cricket, respectively (Unoki et al., 2005; Unoki et al., 2006; Mizunami et al., 2009). Application of octopamine receptor antagonists and dopamine/octopamine depletion induced by blocking their biosynthesis with α -methyl-p-tyrosine lead to a decrease in aggression in male crickets (Stevenson et al., 2005; Rillich and Stevenson, 2011; Rillich et al., 2011). Recently, transgenic techniques became applicable in *G. bimaculatus* (Nakamura et al., 2008), which allow that physiological roles of the biogenic amine system in the cricket nervous system can be analyzed using a neurogenetic approach. However, although physiological and pharmacological studies have been extensively carried out to elucidate the functional roles of the cricket biogenic amine system, little is known about its molecular basis. In a previous report, we identified genes involved in the cricket serotonergic system and examined their expression patterns (Watanabe et al., 2011). Here, we focus on the dopaminergic system and investigate the molecular basis of the dopaminergic system in *G. bimaculatus*.

In insects, dopamine functions as a neurotransmitter/neurohormone that is synthesized in and released from dopaminergic neurons in the nervous system, and also serves as a precursor of melanin that plays important roles in cuticular sclerotization/pigmentation and in the innate immune system. Dopamine biosynthesis requires two enzymes, tyrosine hydroxylase (TH) and aromatic L-amino acid decarboxylase

(AADC, also known as dopa decarboxylase (DDC)). As the initial and rate-limiting step of dopamine biosynthesis, TH converts L-tyrosine into L-3,4-dihydroxyphenylalanine (L-DOPA). Then, AADC converts L-DOPA into dopamine. AADC is involved not only in dopamine biosynthesis but also in serotonin biosynthesis, and is expressed ubiquitously in cricket tissues (Watanabe et al., 2011). Once dopamine is released extracellularly as a neurotransmitter or neurohormone, it is received by dopamine receptors on the plasma membrane of target cells, or undergoes re-uptake via the high-affinity dopamine transporter. Insect dopamine receptors belong to the seven-transmembrane segments (7TM)-containing G protein-coupled receptor (GPCR) superfamily, and are classified into four subtypes: the D₁-like dopamine receptors (Dop1), invertebrate dopamine receptors (INDRs, also known as Dop2), the D₂-like dopamine receptors (Dop3) and the dopamine/ecdysteroid receptors (DopEcR) (Mustard et al., 2005; Srivastava et al., 2005). Once dopamine receptors are activated by dopamine, downstream signaling systems, such as the cyclic adenosine monophosphate (cAMP) second messenger pathway and the inositol trisphosphate (IP₃) second messenger pathway, are activated. The high-affinity dopamine transporter (DAT) is a 12-TM spanning Na⁺/Cl⁻-dependent sodium: neurotransmitter symporter that belongs to the solute carrier family of membrane transport proteins.

In the present study, we identified six genes involved in biosynthesis, transduction and transport of dopamine expressed in the brain of the field cricket *G. bimaculatus*. We obtained a partial open reading frame (ORF) clone of two isoforms of the *TH* gene, full-length ORF clones of three dopamine receptor genes (*Dop1*, *Dop2* and *DopEcR*), a partial ORF clone of a D₂-like dopamine receptor gene (*Dop3*) and a full-length ORF clone of a *Gryllus* *DAT* gene. Expression analysis of the genes revealed an isoform-specific distribution of the two *TH* isoforms and a receptor-subtype specific distribution of the dopamine receptor genes. Our data demonstrate that the dopaminergic system is

1 widespread in the CNS and in peripheral tissues of the cricket.

1 MATERIALS AND METHODS

2 Animals

3 Crickets *Gryllus bimaculatus* DeGeer were reared on a 14-h:10-h light/dark cycle at
4 27°C. They were fed a diet of insect food pellet (Oriental Yeast Co., Tokyo, Japan), chopped
5 carrot and water *ad libitum*.

7 Complementary DNA Cloning of *TH*, *Dop1*, *Dop2*, *Dop3*, *DopEcR* and *DAT* genes

8 First, we cloned partial cDNAs of target genes by reverse transcriptase-polymerase
9 chain reaction (RT-PCR). Complementary DNA synthesis was carried out according to the
10 experimental procedure described previously (Watanabe et al., 2011). In order to design gene
11 specific primers to clone a partial cDNA of the *Gryllus TH* gene, we searched EST
12 (expressed sequence tag) clones corresponding to *Gryllus TH* on the GenBank database, and
13 obtained an EST sequence encoding the N-terminus of TH protein (GenBank accession
14 number: AK278042). To obtain a cDNA fragment encoding the C-terminal region of TH
15 protein, we designed a forward gene specific primer (GSP) and a reverse degenerate primer
16 on the basis of the nucleotide sequence of the *Gryllus TH* EST clone and a conserved amino
17 acid sequence (MSRPFEV) among the insect TH proteins, respectively. The partial cDNA
18 fragments of *Gryllus Dop1* and *Dop2* genes were cloned by Hamada et al (2009). A partial
19 cDNA fragment of the *Gryllus Dop3* gene was amplified as a side product in our previous
20 study (Watanabe et al., 2011). Partial cDNA fragments of the *Gryllus DopEcR* and *DAT*
21 genes were amplified using degenerate primers designed on the basis of conserved amino
22 acid sequences among the insect DopEcR proteins (SVYTFMWI and CQCWMV) and DAT
23 proteins (GIPLFYM and YVDFYYNVII), respectively.

24 Next, to obtain the 5' and 3' region of target mRNAs, we performed 5' and 3' rapid
25 amplification of cDNA ends (RACE) using the FirstChoice RLM-RACE kit (Ambion,

Austin, TX, USA). Finally, we performed RT-PCR to amplify the cDNA fragments containing the full-length or partial open reading frame (ORF) of the genes. We determined nucleotide sequences of at least three independent cDNA clones, and registered their consensus sequences to GenBank. All PCRs were carried out according to the experimental procedure described previously (Watanabe et al., 2011). The primers used to amplify the cDNAs containing the full-length or partial ORF are listed in Table 1.

Sequence comparison, prediction of transmembrane segments, PEST domain, N-glycosylation sites and phosphorylation sites

The deduced full-length amino acid sequences of *Gryllus* Dop1, Dop2, DopEcR and DAT, and the deduced partial amino acid sequences of *Gryllus* TH and Dop3 were compared with those of the corresponding parts of the known homologous genes of other species by using the MAFFT or MUSCLE algorithms on the Geneious 5.6 program (Drummond et al., 2011). The GenBank accession numbers of the proteins used for the comparison are listed in Table 2. The transmembrane regions of the dopamine receptors and of DAT were predicted by the TMHMM v. 2.0 program (Sonnhammer et al., 1998; <http://www.cbs.dtu.dk/services/TMHMM/>). The PEST domain was searched in insect TH proteins using a web-based algorithm, PESTFind (<http://emboss.bioinformatics.nl/cgi-bin/emboss/pepfind>). Potential N-glycosylation and phosphorylation sites were predicted using the NetNGlyc 1.0 program (<http://www.cbs.dtu.dk/services/NetNGlyc>) and the NetPhos 2.0 program (Blom et al., 1999; <http://www.cbs.dtu.dk/services/NetPhos/>), respectively.

Molecular phylogenetic analyses

Molecular phylogenetic trees of the *Gryllus TH*, *Dop1*, *Dop2*, *DopEcR* and *DAT* genes were calculated using their deduced amino acid sequences. Because the deduced amino acid sequence of the partial cDNA of *Gryllus Dop3* is too short to construct phylogenetic tree, we constructed the molecular phylogenetic tree of the *Gryllus* D₂-like dopamine receptor (*Dop3*) gene using their nucleotide sequences corresponding to the protein-coding region. The protein sequences were aligned with the corresponding parts of the known homologous genes of other species by using the MAFFT or MUSCLE algorithm on the Geneious 5.6 program (Drummond et al., 2011). The cDNA sequence of *Gryllus Dop3* encoding the C-terminus of the protein was aligned with the corresponding parts of the known homologous genes of other species by using the ClustalW algorithm on the Geneious 5.6 program. Phylogenetic trees were constructed from the aligned sequences by the bootstrap neighbor-joining algorithms on the MEGA 5 program (Tamura et al., 2011) and visualized with the Geneious 5.5 program. The GenBank accession numbers of the proteins/nucleotides used for the comparison are listed in Table 2.

Tissue-specific expression analysis of *TH*, *Dop1*, *Dop2*, *Dop3*, *DopEcR* and *DAT* genes

RT-PCR analysis was performed to assess the tissue-specific expression of the *Gryllus TH*, *Dop1*, *Dop2*, *Dop3*, *DopEcR* and *DAT* genes. The following tissues were subjected to expression analysis: (1) the central brain, (2) suboesophageal ganglion (SOG), (3) optic lobe (lamina + medulla) and retina, (4) corpus cardiacum-corpora allata complex (CC + CA), (5) thoracic muscles, (6) salivary glands, (7) midgut, (8) Malpighian tubules, (9) testes and (10) ovaries. RNA extraction and reverse transcription were carried out according to the experimental procedure described previously (Watanabe et al., 2011). Briefly, tissue samples were dissected from 1-week-old adult crickets, and total RNA was extracted with TRIzol reagent (Invitrogen). After DNase treatment, 1 µg of each total RNA was reverse-

transcribed in a 20 µl reaction using the Transcriptor First Strand cDNA Synthesis Kit (Roche Applied Science, Tokyo, Japan). A random hexamer and an anchored oligo(dT)₁₈ primer were used as primers. PCR was carried out using *Ex taq* polymerase (TaKaRa, Shiga, Japan). 0.2 µl of cDNA solution was added to a 10 µl PCR reaction. PCR amplification of target genes was performed for 35 cycles at 96°C for 15 s, 58 °C for 15 s, and 72°C for 60 s, followed by a final extension at 72°C for 5 min. PCR products were run through a 1.5% agarose gel and visualized by etidium brimide. The *Eflalpha* gene, which is ubiquitously and stably expressed in all examined tissue (Watanabe et al., 2011), was amplified as an internal control gene. To confirm that genomic DNA was not present in cDNA solutions, we amplified the *Eflalpha* gene and checked the absence of an amplification product in the RT negative controls (RT(-)). Primers used for the RT-PCR analysis are listed in Table 1.

RESULTS

Identification and expression analyses of *Gryllus* tyrosine hydroxylase gene

Nucleotide sequence and structural features of *Gryllus* TH isoforms: We obtained partial cDNA fragments of two isoforms of tyrosine hydroxylase gene (*TH*) expressed in the cricket. We performed RT-PCR with the gene-specific primers (GSPs) and obtained a 1408-bp cDNA fragment of the *TH-A* isoform and a 1606-bp cDNA fragment of the *TH-B* isoform (GenBank accession numbers: AB720738 and AB720737, respectively). These genes have the following properties: *TH-A* isoform, partial ORF = 29–1408 bp, protein product = 460 amino acids; *TH-B* isoform, partial ORF = 29–1606 bp, protein product = 526 amino acids (Figure 1A and B). Several functional domains and residues, such as the ACT domain, catalytic domain and several catalytic residues are conserved in both *Gryllus* TH proteins (Figure 1A). In addition, two conserved arginine residues (Arg²⁹ and Arg³⁰), important for catecholamine-mediated inhibition of enzyme activity (Nakashima et al., 1999; Nakashima et al., 2000), were conserved in the N-terminal region of both *Gryllus* TH proteins (Figure 1A). We failed to obtain a cDNA clone of the 3' region of the *TH* ORF corresponding to the C-terminal coiled-coil domain important for dimerization. The *Gryllus* TH-B isoform contains a 66-amino-acid insertion in the ACT domain. The position of this insertion was conserved in other insect TH-B isoforms such as those found in *Mythimna separata* and *Drosophila melanogaster* (Birman et al., 1994; Ninomiya and Hayakawa, 2007) (Figure 1B). The inserted sequences of the three insect TH-B isoforms are predicted as the PEST domain that is found in many rapidly degraded proteins (Rogers et al., 1986).

A comparison of the deduced amino acid sequences of the *Gryllus* TH isoforms with those of other insect amino acid hydroxylases indicates that the *Gryllus* TH isoforms are closely related to other known insect THs (*Gryllus* TH-A is 73.1% identical to *Drosophila* pale-A (NP_476897), 74.1% identical to *Mythimna* TH (type 2, brain form; BAF32574) and

75.1% identical to *Apis* TH (NP_001011633). The molecular phylogenetic analysis of aromatic amino acid hydroxylase genes also indicates that *Gryllus* TH is closely related to the insect TH protein (Figure 2).

Tissue-specific expression of *Gryllus* TH isoforms: We examined the tissue-specific expression pattern of total *TH* transcript and each *TH* isoform in the cricket by using RT-PCR (Figure 3). Our results indicate an isoform-specific expression pattern of the *Gryllus* *TH* gene: the *Gryllus* *TH-A* isoform is predominantly expressed in neural tissues, whereas the *Gryllus* *TH-B* isoform is ubiquitously expressed in all examined tissues. An intensely stained band of PCR product of the *Gryllus* *TH* common region (cDNA fragments corresponding to the catalytic domain) was detected in the lanes of the central brain, SOG and the ovaries. A weaker stained band of PCR product of the *Gryllus* *TH* common region was detected in the lanes of the optic lobe + retina, CC + CA, thoracic muscles, salivary gland and testes. An intensely stained band of PCR product of the *Gryllus* *TH-A* isoform was detected in the lanes of the central brain, SOG, optic lobe + retina and CC + CA. A weaker stained band of PCR product of the *Gryllus* *TH-A* isoform was detected in the lanes of the thoracic muscles and testicle. An intensely stained band of PCR product of the *Gryllus* *TH-B* isoform was detected in the lanes of the thoracic muscles, testicle and ovary. A weaker stained band of PCR product of the *Gryllus* *TH-B* isoform was detected in the lanes of the other tissues except for the midgut and Malpighian tubulus.

Identification and expression analysis of three *Gryllus* dopamine receptor genes

D₁-like and invertebrate dopamine receptors: We identified the D₁-like dopamine receptor gene (*Dop1*) and the invertebrate dopamine receptor gene (*Dop2*) in the cricket. We performed RT-PCR with the GSPs designed at the 5' and 3' UTRs of the genes, and obtained a 1478-bp cDNA fragment of *Gryllus* *Dop1*, and a 1458-bp cDNA fragment of *Gryllus* *Dop2*

(GenBank accession numbers: AB720739 and AB720740, respectively). These genes have the following properties: *Dop1*, full-length ORF = 9–1463 bp, protein product = 485 amino acids; *Dop2*, full-length ORF = 50–1447 bp, protein product = 465 amino acids (Figure 4A and B).

The cDNA fragment of *Gryllus Dop1* encoded the seven transmembrane (TM) segments (TM1–TM7 in Figure 4A). The N-terminal extracellular region contained a putative N-glycosylation site. The first, second, and third cytoplasmic loops (the region between TM1-TM2, TM3-TM4 and TM5-TM6, respectively) and the C-terminal intracellular region contained several potential phosphorylation sites. The second cytoplasmic loop contained a conserved DRY motif. The cDNA fragment of *Gryllus Dop2* encoded the seven transmembrane (TM) segments (TM1-TM7 in Figure 4B). The N-terminal extracellular region contained three putative N-glycosylation sites. The first and third cytoplasmic loops (the region between TM1-TM2 and TM5-TM6, respectively) and the C-terminal intracellular region contained several potential phosphorylation sites. The second cytoplasmic loop contained a conserved DRY motif.

A comparison of the deduced amino acid sequences of *Gryllus Dop1* and *Dop2* with those of other insect G-protein coupled receptors indicates that *Gryllus Dop1* is closely related to other known insect Dop1 (53.2% identical to *Bombyx Dop1* (NP_001108459), 53.2% identical to *Apis Dop1* (NP_001011595) and 53.0% identical to *Drosophila Dop1* (NP_477007)), and that *Gryllus Dop2* is closely related to other known insect Dop2 (68.7% identical to *Bombyx Dop2* (NP_001108338), 66.4% identical to *Apis Dop2* (NP_001011567) and 61.2% identical to *Drosophila Dop2* (NP_733299)). The molecular phylogenetic analysis of the D₁-like dopamine receptors and the INDRs also indicates that *Gryllus Dop1* and *Dop2* are closely related to insect Dop1 and Dop2 proteins, respectively (Figure 5).

D₂-like dopamine receptor: We identified an insect D₂-like dopamine receptor gene, *Dop3*, expressed in the cricket brain. We performed RT-PCR with the GSPs and obtained a 484-bp cDNA fragment of the *Gryllus Dop3* gene (GenBank accession number: AB720741). This clone contains an ORF spanning 2–201 bp, resulting in a protein product of 63 amino acids. The cDNA fragment of *Gryllus Dop3* encoded the two transmembrane (TM) segments (TM6 and TM7 in Figure 6A). In addition, we isolated another cDNA clone that differed by an alternatively spliced intron within the 3' UTR (*Gryllus Dop3* long 3' UTR variant; GenBank accession number: AB720740).

Comparison of the deduced amino acid sequence of *Gryllus Dop3* with those of the corresponding part of other insect G-protein coupled receptors indicates that *Gryllus Dop3* is closely related to other known insect D₂-like dopamine receptors (78.5% identical to *Apis Dop3* (NP_001014983), 85.1% identical to *Tribolium Dop3* (XP_969037), and 74.3% identical to *Drosophila DDR2* (NP_001014760)). The molecular phylogenetic analysis of the D₂-like dopamine receptors also indicates that *Gryllus Dop3* is closely related to insect D₂-like receptors (Figure 6B).

Dopamine/ecdysteroid receptor: We identified an insect dopamine/ecdysteroid receptor gene, *DopEcR*, in the cricket. We performed RT-PCR with the GSPs designed on the 5' and 3' UTRs of the gene, and obtained a 1046-bp cDNA fragment of *Gryllus DopEcR* (GenBank accession number: AB720743). This clone contains an ORF spanning 23–1003 bp, resulting in a protein product of 326 amino acids (Figure 7A). The cDNA fragment of *Gryllus DopEcR* encoded the seven transmembrane (TM) segments (TM1–TM7 in Figure 7A). The second cytoplasmic loop contained a DRY motif.

Comparison of the deduced amino acid sequences of *Gryllus DopEcR* with those of other insect G-protein coupled receptors indicates that the *Gryllus DopEcR* is closely related to other known insect DopEcR (72.4% identical to *Apis DopEcR* (XP_396491), 66.3%

identical to *Anopheles* DopEcR (XP_315694) and 66.9% identical to *Drosophila* DopEcR (NP_647897). The molecular phylogenetic analysis of the DopEcR indicates that the *Gryllus* DopEcR is closely related to other insect DopEcR proteins (Figure 7B).

Tissue-specific expression of four dopamine receptor genes: We examined the tissue-specific expression pattern of the four identified dopamine receptor genes by using RT-PCR. Our results indicate that the *Gryllus* dopamine receptor genes exhibit receptor subtype-specific distributions (Figure 8): *Gryllus Dop1* and *Dop2* genes were ubiquitously expressed in all examined tissues, whereas the *Gryllus Dop3* and *DopEcR* genes showed restricted expression in specific tissues. An intensely stained band of PCR product of *Gryllus Dop1* gene was detected in the lanes of the central brain, SOG, CC + CA, salivary gland, midgut, testicle and ovaries. A weaker stained band of PCR product of *Gryllus Dop1* gene was detected in the lanes of the optic lobe + retina, thoracic muscles and Malpighian tubules. An intensely stained band of PCR product of *Gryllus Dop2* gene was detected in the lanes of the central brain, SOG, CC + CA, thoracic muscles, midgut, testicle and ovaries. A weaker stained band of PCR product of *Gryllus Dop2* gene was detected in the lanes of the optic lobe + retina, salivary gland and Malpighian tubules. An intensely stained band of PCR product of *Gryllus Dop3* gene was detected in the lanes of the central brain, SOG and testicle. An intensely stained band of PCR product of the *Gryllus DopEcR* gene was detected in the lanes of the central brain, SOG, CC + CA, thoracic muscles, midgut, testicle and ovaries. A weaker stained band of PCR product of *Gryllus DopEcR* gene was detected in the lanes of the salivary gland. Expression on *Gryllus DopEcR* gene was not detected in the lane of the Malpighian tubules.

Identification and expression analysis of a *Gryllus* high-affinity dopamine transporter gene

We identified a high-affinity dopamine transporter gene, *DAT*, expressed in the cricket brain. We performed RT-PCR with the GSPs designed on the 5' and 3' UTRs of the *Gryllus DAT* gene and obtained an 1880-bp cDNA fragment (GenBank accession number: AB720744). This clone contains an ORF spanning 17–1855 bp, resulting in a protein product of 612 amino acids. The cDNA fragment of *Gryllus DAT* encoded the twelve TMs (TM1–TM12 in Figure 9A). The second and third extracellular loop (the region between TM3-TM4 and TM5-TM6) contained potential N-glycosylation sites. The N-terminal intracellular region and the second intracellular loop (the region between TM4-TM5) contained potential phosphorylation sites. In addition, two cysteine residues that are required for *DAT* biosynthesis and/or its delivery to the cell surface (Chen et al., 2007) were conserved in the second extracellular loop (“S-S bond” in Figure 9A).

A comparison of the deduced amino acid sequence of *Gryllus DAT* with those of other insect monoamine transporters indicates that *Gryllus DAT* is closely related to other known insect *DAT* (75.7% identical to *Apis DAT* (NP_001139210), 74.3% identical to *Bombyx DAT* (NP_001037362) and 72.0% identical to *Drosophila DAT* (NP_523763)). The molecular phylogenetic analysis of biogenic amine transporter proteins, including the serotonin transporter (SERT), octopamine transporter (OAT) and noradrenaline transporter (NAT), also indicates that the *Gryllus DAT* is closely related to the insect *DAT* proteins (Figure 9B).

Next, we performed RT-PCR analyses to investigate the tissue-specific expression of the *Gryllus DAT* gene. The PCR product of *Gryllus DAT* was detected in the lanes of all examined tissues (Figure 10).

DISCUSSION

Multiple TH protein isoforms were found in vertebrates (four isoforms in human, two isoforms in monkey) (Nagatsu, 1989; Lewis et al., 1993; Lewis et al., 1994; Haycock, 2002), which are produced by alternative splicing at the N-terminal portion of the regulatory ACT domain. In the present study, we isolated partial cDNA fragments of two *TH* isoforms (the short variant, *TH-A*; the long variant, *TH-B*) generated by alternative splicing in the middle of the ACT domain. Similar alternatively spliced forms of the *TH* gene were identified in *D. melanogaster* and the armyworm *M. separata* (Birman et al., 1994; Ninomiya and Hayakawa, 2007). In the cricket, the *TH-A* isoform is predominantly expressed in the CNS, whereas the *TH-B* isoform is ubiquitously expressed. In *D. melanogaster* and *M. separata*, the corresponding *TH* isoforms show similar expression patterns (Friggi-Grelín et al., 2003; Ninomiya and Hayakawa, 2007). These data suggest that the existence of two TH isoforms and the isoform-specific expression pattern of the *TH* gene are evolutionary conserved in insects. Evolutionary conserved tissue-specific distribution of two *TH* isoforms suggests that the *TH-A* isoform is predominantly responsible for synthesis of the neurotransmitter dopamine, while the *TH-B* isoform is also responsible for synthesis of dopamine as a precursor of melanin, which is required for cuticular tanning/sclerotization and the innate immune system.

The inserted sequence of the *Gryllus* *TH-B* isoform and its homologous sequences in *D. melanogaster* and *M. Separata* are predicted as the PEST domain (Figure 1B). The PEST domain is found in many rapidly degraded proteins (Rogers et al., 1986). The degradation of a PEST-containing protein is mediated via the ubiquitin/proteasome system or calpain (Shumway et al., 1999; Reverte et al., 2001; Spencer et al., 2004). Moreover, recombinant *Drosophila* TH isoforms exhibited distinct enzymatic properties *in vitro* (i.e. cofactor

binding) (Vié et al., 1999). Further biochemical studies are necessary to reveal the functional roles of the inserted region of the insect TH-B isoforms.

The change of the dopamine level in the nervous systems has been reported in several insects. In the brain of the wood ant *Formica japonica*, dopamine levels decrease in response to starvation stress and aging (Wada-Katsumata et al., 2011; Aonuma and Watanabe, 2012). In the honeybee, dopamine level in the optic lobes changes with the light-dark cycle (Carrington et al., 2007). **In social insects, changes in brain dopamine levels are associated with division of labor, reproductive status and caste (Brandes et al., 1990; Taylor et al., 1992; Schulz and Robinson, 1999; Bloch et al., 2000; Harano et al., 2005).** Since TH catalyzes the rate-limiting step of dopamine biosynthesis, regulation of the TH enzymatic activity is one of the key mechanisms underlying the change of the dopamine level in the CNS. The N-terminal region of the TH protein is important for the regulation of its enzyme activity (Nakashima et al., 2009; Nakashima et al., 1999; Nakashima et al., 2000) and protein stability (Nakashima et al., 2011). The N-terminal region of the *Gryllus* TH isoforms contains several putative phosphorylation sites (Ser³², Ser⁵⁰, Tyr⁶⁵ and Ser⁷⁸) which are conserved in *Drosophila* TH (pale, see Figure 1). The Ser³² is conserved in TH proteins of both vertebrates and invertebrates, and enzymatic activity of *Drosophila* TH is increased by phosphorylation at Ser³² *in vitro* (Vié et al., 1999). The Ser⁴⁰ residue of vertebrate TH proteins, which is homologous to the Ser³² of insect TH, is phosphorylated by various kinases including PKA, PKG and PKC (Dunkley et al., 2004). Further studies will reveal the cellular mechanisms underlying the changes in dopamine levels in the insect CNS.

Except for a dopamine-gated chloride channel found in nematodes [LGC-53 in *Caenorhabditis elegans* (Ringstad et al., 2009) and HcGGR3 in *Haemonchus contortus* (Vijayaraghava et al., 2009)], all dopamine receptors belong to the GPCR superfamily, and are classified into three subtypes: the D₁-like receptors, INDRs and D₂-like receptors.

1 Recently another GPCR-type receptor, DopEcR, was identified in *D. melanogaster*, which
2 shows high affinity for dopamine as well as ecdysteroids (Srivastava et al., 2005). D₁-like
3 receptors are coupled to G_{as} protein, which subsequently activates adenylyl cyclase to
4 produce cAMP, and the INDRs are coupled to Ca²⁺ signaling as well as cAMP (Beggs et al.,
5 2011). On the other hand, D₂-like dopamine receptors are coupled to G_{ai} protein that inhibits
6 adenylyl cyclase. D₂-like dopamine receptors can also activate the mitogen-activated protein
7 kinase (MAPK) pathway (Yan et al., 1999). Activation of DopEcR by dopamine leads to an
8 increase in intracellular cAMP levels as well as activation of the phosphoinositol 3-phosphate
9 pathway. Ecdysteroids, in contrast, inhibit the effects of dopamine and induce the activation
10 of the MAPK pathway (Srivastava et al., 2005). In the present study, we searched GPCRs
11 structurally associated with *Gryllus* DopEcR in order to construct a molecular phylogenetic
12 tree of DopEcR, and found that the vertebrate GPR21 and GPR52 genes are most similar to
13 insect DopEcR. GPR21 and GPR52 were first identified from the human expressed sequence
14 tags database and genome database (O'Dowd et al., 1997; Sawzdargo et al., 1999), and their
15 homologues were then found in other vertebrate genomes (see Table 2). Although the natural
16 agonists, downstream signal transduction pathways and physiological roles of the vertebrate
17 GPR21 and GPR52 are still unknown, the mRNA of human GPR21 is expressed in several
18 brain areas (e.g. the frontal cortex, caudate nucleus, putamen, thalamus) (O'Dowd et al.,
19 1997). Gene function analyses of the DopEcR/GPR21/GPR52 family are necessary to reveal
20 the physiological roles of the novel GPCR family.

21 Dopamine is widely distributed in the CNS and in peripheral tissues of both
22 vertebrates and invertebrates. Application of agonists/antagonists of specific dopamine
23 receptors has revealed physiological functions of dopamine in various insect tissues. To
24 elucidate the sites of dopamine action in the cricket, we investigated the tissue-specific
25 expression of four dopamine receptor genes in the tissue of adult crickets.

Central nervous system: Hamada et al. (2009) cloned partial cDNA fragments of the *Gryllus Dop1* and *Dop2* genes, and examined their mRNA distributions in the brain. They reported that mRNAs of both dopamine receptors were expressed predominantly in the Kenyon cells of the mushroom bodies in the cricket brain. In *Drosophila*, the DAMB (a homologue of *Gryllus Dop2*) is exclusively expressed in the mushroom bodies (Han et al., 1996), in which the dopaminergic system plays essential roles in learning and memory (Waddell, 2010). In the brain of the honeybee and silkworm, mRNAs of dopamine receptors are also detected in the Kenyon cells of the mushroom bodies (Kurshan et al., 2003; Kyle et al., 2005; Mitumasu et al., 2008). In the cricket, pharmacological inhibition of D₁-like dopamine receptor impaired aversive olfactory and visual conditioning and memory recall (Unoki et al., 2005; Unoki et al., 2006; Mizunami et al., 2009). In *Drosophila*, the D₁-like dopamine receptor DopR (a homologue of *Gryllus Dop1*) regulates caffeine-induced arousal in the mushroom bodies (Andretic et al., 2008), as well as stress-induced arousal in the central complex and sleep-wake arousal in the lateral-ventral neurons (Lebestky et al., 2009). Moreover, the *DopEcR* gene is strongly expressed in the cricket CNS although its physiological functions in the nervous system are still unknown in insects. Contrary to the *Gryllus Dop1*, *Dop2* and *DopEcR* genes, which are expressed in all neuronal tissues examined in this study, the *Gryllus* D₂-like dopamine receptor gene, *Dop3*, showed restricted expression in the brain and SOG, and its expression was not detected in the optic lobes and CC + CA. Draper et al. (2007) examined the distribution of the *Drosophila* D₂-like receptor (DD2R) in the CNS using anti-DD2R antibodies, and found that a small number of neurons including the Ap-let cohort of peptidergic neurons was immunoreactive in the larval and adult CNS. They also generated DD2R RNA-interference (RNAi) lines and demonstrated that the RNAi-mediated knock-down of the *DD2R* gene resulted in the reduction of locomotor activity in the adult flies. On the other hand, in the honeybee, RNAi-mediated knockdown of

1 *AmDOP2* receptor gene affects locomotion (Mustard et al., 2010). Functional analysis of
2 each dopamine receptor in the cricket CNS will reveal the differential involvement of specific
3 dopamine receptor subtypes in behavior.

4 **Corpus cardiacum-corpora allata (CC-CA) complex:** The CC-CA complex functions as
5 an endocrine center in insects. Woodring and Hoffmann (1994) reported that dopamine
6 application has no effect on juvenile hormone biosynthesis in the isolated CA of adult
7 crickets. On the other hand, dopamine receptor blockers decreased release of adipokinetic
8 hormone (AKH) from the corpus cardiacum in the locust *Schistocerca gregaria*
9 (Samaranayaka, 1976). Dopamine stimulates cAMP accumulation in the CC of the cockroach
10 *Periplaneta americana* (Gole et al., 1987). In the CC of the locust *Locusta migratoria*,
11 dopamine potentiates cAMP-induced release of AKH, and is abundantly contained in the
12 storage part of the CC (Passier et al., 1995). Our study demonstrates that mRNAs of *Gryllus*
13 *Dop1* and *Dop2*, which are positively coupled to adenylyl cyclase, are expressed in the CC-
14 CA complex of adult crickets. Therefore, these receptors might be involved in the dopamine-
15 mediated stimulation of AKH release in the CC. In addition, the expression analysis revealed
16 that the *Gryllus DopEcR* gene, which mediates rapid response to ecdysteroids, was expressed
17 in the CC-CA complex of adult crickets. Our data suggest the presence of a feedback
18 regulation of the ecdysone system mediated by *DopEcR* in the CC-CA complex of the cricket.

19 **Salivary gland:** In insects, the aminergic and peptidergic systems control salivation (Ali,
20 1997; Walz et al. 2006). In *G. bimaculatus*, dopaminergic innervations in the salivary gland
21 originate from the SOG (SN1 neurons) (Helle et al., 1995; Hörner et al., 1995; Ali, 1997). In
22 the cockroach salivary gland, dopamine induces production of cAMP that acts as a second
23 messenger in the acinar cells to cause the secretory response (Grewe and Kebabian, 1982;
24 Gray et al., 1984; Marg et al. 2004; Rietdorf et al. 2005). Our study demonstrates that
25 mRNAs of *Gryllus Dop1*, *Dop2* and *DopEcR*, which are positively coupled with adenylyl

cyclase, are expressed in the cricket salivary gland. Therefore, these receptors might be involved in the dopamine-mediated stimulation of salivation in the salivary gland.

Testicle and ovaries: Gonadotropic effect of dopamine has been reported in several insects. In the cockroach *Blattella germanica*, dopamine stimulates oocyte growth just before vitellogenesis, whereas it has an inhibitory effect at the end of vitellogenesis (Pastor et al., 1991). In *Drosophila*, depletion of dopamine in newly eclosed female flies resulted in abnormal development of the ovaries (Neckameyer, 1996). Gonadotropic effect of dopamine was also demonstrated in eusocial hymenoptera such as *Polistes chinensis* (Sasaki et al., 2009) and *Apis mellifera* (Dombroski et al., 2003). In *A. mellifera*, *AmDop1* and *AmDop3* are expressed in the worker ovaries, and their expression is associated with the reproductive status of workers (Vergoz et al., 2012). In contrast, in the cricket, *Gryllus Dop1*, *Dop2* and *DopEcR* genes are expressed in the ovaries, while the expression of the D₂-like dopamine receptor gene (*Dop3*) was not detected. We collected ovary samples from female crickets two weeks after adult molt. At this time the ovaries are fully activated. Like in *A. mellifera*, the expression of the dopamine receptor genes might be affected by the reproductive status in the cricket. To our knowledge, our study is the first report on the expression of dopamine receptor genes in the insect testicle. Our data suggest that the activity of both the female ovaries and the male testicle is regulated by dopamine in insects. Further histological and pharmacological studies are needed to understand the actions of dopamine on the testicle in the cricket.

Other non-neuronal tissues: *Gryllus Dop1* and *Dop2* genes were expressed in the midgut and Malpighian tubules. The catecholamine-containing nerve endings are present in the visceral muscle in insects (Klemm, 1972; Klemm, 1979). In *Locusta migratoria*, TH-like immunoreactive neurons are present in the stomatogastric nerve system, and dopamine inhibits phasic contraction of the foregut muscle (Lange and Chan, 2008). In *Drosophila*, the

D₂-like receptor is expressed in a small number of cells in the ventriculus and in the Malpighian tubules (Draper et al., 2007). In the cricket midgut and Malpighian tubules, we did not detect the expression of the D₂-like receptor gene by RT-PCR, but the expression of the *DopEcR* gene was detected in the midgut of adult crickets. Further studies are necessary to elucidate functional roles of dopamine receptor subtypes expressed in the cricket visceral organs. Contrary to the visceral muscle, little is known about the dopaminergic modulation of skeletal muscle activity in insect. In *Drosophila* larval neuromuscular junctions, dopamine reduces presynaptic activity but does not affect the postsynaptic receptiveness to glutamate (Cooper and Neckameyer, 1999). In the thoracic muscles of the cricket, the two D₁-like dopamine receptor genes and the *DopEcR* gene are expressed; therefore, dopamine can affect muscular activity *in vivo*. Further investigation is necessary to elucidate the dopaminergic control of skeletal muscle activity in the cricket.

In the present study, we determined the full-length coding sequence of the *Gryllus* *DAT* gene and examined tissue-specific distribution of the gene. Interestingly, our expression analysis revealed that the *Gryllus* *DAT* gene is ubiquitously expressed in the all cricket tissues examined in this study. In the other insects such as *Drosophila* and moths, it is reported that the *DAT* gene is predominantly expressed in the nervous tissues. In *Drosophila*, the *DAT* gene is detected in the brain and thoracic-abdominal ganglion but not in the other tissues (FlyAtlas Anatomical Expression Data; Chintapalli et al., 2007). In the larva of the cabbage looper moth *Trichoplusia ni*, Northern blot analysis revealed that mRNA of *DAT* gene was contained in the head (including the brain and SOG) but not in the fat body and epidermal tissues (Gallant et al., 2003). Further investigation is necessary to elucidate the physiological function of the *DAT* gene expressed in the non-neuronal tissues in the cricket.

1 In summary, we identified six genes involved in the biosynthesis, transduction and re-
2 uptake of dopamine in the cricket *G. bimaculatus*. Two TH isoforms showed isoform-specific
3 distribution in cricket tissues. Tissue-specific expression analysis of dopamine receptor genes
4 showed that the dopaminergic system is widely distributed in the cricket, and that the
5 dopaminergic system might regulate various aspects of physiological phenomena via distinct
6 dopamine receptor pathways.

7 **ACKNOWLEDGEMENTS**

8 We thank Dr. Miriam Henze (Lunds Universitet, Sweden) for reviewing. This research was
9 partly supported by Grant-in-Aid for JSPS Fellows to T. Watanabe, grants-in-aid for
10 Scientific Research (KAKENHI) from the MEXT, Scientific Research on Priority Areas
11 (Area No. 454) to H. Aonuma (No. 17075001) and from the JSPS to H. Aonuma (No.
12 23300113).

REFERENCES

- Ali, D. (1997). The aminergic and peptidergic innervation of insect salivary glands. *J Exp Biol*, 200:941-949.
- Andretic, R., Kim, Y. C., Jones, F. S., Han, K. A. & Greenspan, R. J. (2008). *Drosophila* D1 dopamine receptor mediates caffeine-induced arousal. *Proc Natl Acad Sci U S A*, 105:20392-20397.
- Aonuma, H. & Watanabe, T. (2012). Changes in the content of brain biogenic amine associated with early colony establishment in the queen of the ant, *Formica japonica*. *PLoS ONE*, 7: e43377.
- Beggs, K. T., Hamilton, I. S., Kurshan, P. T., Mustard, J. A. & Mercer, A. R. (2005). Characterization of a D2-like dopamine receptor (AmDOP3) in honey bee, *Apis mellifera*. *Insect Biochem Mol Biol*. 35:873-882.
- Beggs, K. T., Tyndall, J. D. & Mercer, A. R. (2011). Honey bee dopamine and octopamine receptors linked to intracellular calcium signaling have a close phylogenetic and pharmacological relationship. *PLoS One*. 6:e26809.
- Bendtsen, J. D., Nielsen, H., von Heijne, G. & Brunak, S. (2004). Improved prediction of signal peptides: SignalP 3.0. *J Mol Biol*, 340:783-795.

Birman, S., Morgan, B., Anzivino, M. & Hirsh, J. (1994). A novel and major isoform of tyrosine hydroxylase in *Drosophila* is generated by alternative RNA processing. *J Biol Chem* 269:26559-26567.

Bloch, G., Simon, T., Robinson, G. E. & Hefetz, A. (2000). Brain biogenic amines and reproductive dominance in bumble bees (*Bombus terrestris*). *J Comp Physiol A*. 186: 261-268.

Blom, N., Gammeltoft, S. & Brunak, S. (1999). Sequence and structure-based prediction of eukaryotic protein phosphorylation sites. *J Mol Biol* 294:1351-1362.

Brandes, C., Sugawa, M. & Menzel, R. (1990). High-performance liquid chromatography (HPLC) measurement of catecholamines in single honeybee brains reveals caste-specific differences between worker bees and queens in *Apis mellifera*. *Comp Biochem Physiol* 97: 53-57.

Carrington, E., Kokay, I. C., Duthie, J., Lewis, R. & Mercer, A. R. (2007). Manipulating the light/dark cycle: effects on dopamine levels in optic lobes of the honey bee (*Apis mellifera*) brain. *J Comp Physiol A Neuroethol Sens Neural Behav Physiol*, 193:167-180.

Chen, R., Wei, H., Hill, E. R., Chen, L., Jiang, L., Han, D. D. & Gu, H. H. (2007). Direct evidence that two cysteines in the dopamine transporter form a disulfide bond. *Mol Cell Biochem*, 298:41-48.

Chintapalli, V. R., Wang, J., Dow, J. A. (2007). Using FlyAtlas to identify better *Drosophila melanogaster* models of human disease. *Nat Genet*. 39:715-720.

- 1
- 2 Cooper, R. L. & Neckameyer, W. S. (1999). Dopaminergic modulation of motor neuron
- 3 activity and neuromuscular function in *Drosophila melanogaster*. *Comp Biochem Physiol B*
- 4 *Biochem Mol Biol*, 122:199-210.
- 5
- 6 Dombroski, T. C. D., Simoes, Z. L. P. & Bitondi, M. M. G. (2003). Dietary dopamine causes
- 7 ovary activation in queenless *Apis mellifera* workers. *Apidologie*, 34:281-289
- 8
- 9 Donly, B. C. & Caveney, S. (2005). A transporter for phenolamine uptake in the arthropod
- 10 CNS. *Arch Insect Biochem Physiol*, 59:172-183.
- 11
- 12 Draper, I., Kurshan, P. T., McBride, E., Jackson, F. R. & Kopin, A. S. (2007). Locomotor
- 13 activity is regulated by D2-like receptors in *Drosophila*: an anatomic and functional analysis.
- 14 *Dev Neurobiol*, 67:378-393.
- 15
- 16 Drummond, A. J., Ashton, B., Buxton, S., Cheung, M., Cooper, A., Duran, C., Field, M.,
- 17 Heled, J., Kearse, M., Markowitz, S., Moir, R., Stones-Havas, S., Sturrock, S., Thierer, T. &
- 18 Wilson, A. (2011). Geneious v5.4, Available from <http://www.geneious.com/>
- 19
- 20 Dunkley, P. R., Bobrovskaya, L., Graham, M. E., von Nagy-Felsobuki, E. I. & Dickson, P. W.
- 21 (2004). Tyrosine hydroxylase phosphorylation: regulation and consequences. *J Neurochem*,
- 22 91:1025-1043.
- 23
- 24 Evans, P. D. (1980). Biogenic amines in the insect nervous system. *Adv Insect Physiol*,
- 25 15:317-473.

- 1
- 2 Evans, P. D. & Maqueira, B. (2005). Insect octopamine receptors: a new classification
3 scheme based on studies of cloned *Drosophila* G-protein coupled receptors. *Invert Neurosci*,
4 5:111-118.
- 5
- 6 Friggi-Grelín, F., Iché, M. & Birman, S. (2003). Tissue-specific developmental requirements
7 of *Drosophila* tyrosine hydroxylase isoforms. *Genesis*, 35:260-269.
- 8
- 9 Gallant, P., Malutan, T., McLean, H., Verellen, L., Caveney, S., Donly, C. (2003).
10 Functionally distinct dopamine and octopamine transporters in the CNS of the cabbage
11 looper moth. *Eur J Biochem*. 270:664-674.
- 12
- 13 Gole, J. W. D., Orr, G. L. & Downer, R. G. H. (1987). Pharmacology of octopamine-,
14 dopamine-, and 5-hydroxytryptamine-stimulated cyclic AMP accumulation in the corpus
15 cardiacum of the American cockroach, *Periplaneta americana* L. *Arch Insect Biochem*
16 *Physiol*, 5:119-128.
- 17
- 18 Gray, D. C., Ginsborg, B. L. & House, C. R. (1984). Cyclic AMP as a possible mediator of
19 dopamine stimulation of cockroach gland cells. *Q J Exp Physiol*, 69:171-186.
- 20
- 21 Grewe, C. W. & Kebabian, J. W. (1982). Dopamine stimulates production of cyclic AMP by
22 the salivary gland of the cockroach, *Nauphoeta cinerea*. *Cell Mol Neurobiol*, 2:65-69.
- 23

- 1 Hamada, A., Miyawaki, K., Honda-sumi, E., Tomioka, K., Mito, T., Ohuchi, H. & Noji, S.
2 (2009). Loss-of-function analyses of the fragile X-related and dopamine receptor genes by
3 RNA interference in the cricket *Gryllus bimaculatus*. *Dev Dyn*, 238:2025-2033.
- 4
- 5 Han, K. A., Millar, N. S., Grotewiel, M. S. & Davis, R.L. (1996). DAMB, a novel dopamine
6 receptor expressed specifically in *Drosophila* mushroom bodies. *Neuron*, 16:1127-1135.
- 7
- 8 Harano, K., Sasaki, K. & Nagao, T. (2005). Depression of brain dopamine and its metabolite
9 after mating in the European honeybee (*Apis mellifera*) queens. *Naturwiss.* 92:310-313.
- 10
- 11 Haycock, J. W. (2002). Species differences in the expression of multiple tyrosine
12 hydroxylase protein isoforms. *J Neurochem*, 81:947-953.
- 13
- 14 Helle, J., Dirksen, H., Eckert, M., Nässel, D.R., Spörhase-Eichmann, U. & Schürmann, F.W.
15 (1995). Putative neurohemal areas in the peripheral nervous system of an insect, *Gryllus*
16 *bimaculatus*, revealed by immunocytochemistry. *Cell Tissue Res*, 281:43-61.
- 17
- 18 Hörner, M., Spörhaseeichmann, U., Helle, J., Venus, B. & Schurmann, F.W. (1995). The
19 distribution of neurons immunoreactive for β -tyrosine hydroxylase, dopamine and serotonin
20 in the ventral nerve cord of the cricket, *Gryllus bimaculatus*. *Cell Tissue Res*, 280:583-604.
- 21
- 22 Klemm, N. (1972). Monoamine-containing nervous fibres in foregut and salivary gland of the
23 desert locust, *Schistocerca gregaria* Forskal (Orthoptera, Acrididae). *Comp Biochem Physiol*
24 *A Comp Physiol*, 43:207-211.
- 25

- 1 Klemm, N. (1979). Biogenic amines in the stomatogastric nervous system of several insect
2 orders. *Entomol Gen*, 5:113-121.
- 3
- 4 Lange, A. B. & Chan, K. (2008). Dopaminergic control of foregut contractions in *Locusta*
5 *migratoria*. *J Insect Physiol*, 54:222-230.
- 6
- 7 Lebestky, T., Chang, J. S., Dankert, H., Zelnik, L., Kim, Y. C., Han, K. A., Wolf, F. W.,
8 Perona, P. & Anderson, D.J. (2009). Two different forms of arousal in *Drosophila* are
9 oppositely regulated by the dopamine D1 receptor ortholog DopR via distinct neural circuits.
10 *Neuron*, 64:522-536.
- 11
- 12 Lewis, D. A., Melchitzky, D. S. & Haycock, J. W. (1993). Four isoforms of tyrosine
13 hydroxylase are expressed in human brain. *Neuroscience*, 54:477-492.
- 14
- 15 Lewis, D. A., Melchitzky, D. S. & Haycock, J. W. (1994). Expression and distribution of two
16 isoforms of tyrosine hydroxylase in macaque monkey brain. *Brain Res*, 656:1-13.
- 17
- 18 Marg, S., Walz, B. & Blenau, W. (2004). The effects of dopamine receptor agonists and
19 antagonists on the secretory rate of cockroach (*Periplaneta americana*) salivary glands. *J*
20 *Insect Physiol*, 50:821-830.
- 21
- 22 Mizunami, M., Unoki, S., Mori, Y., Hirashima, D., Hatano, A. & Matsumoto, Y. (2009).
23 Roles of octopaminergic and dopaminergic neurons in appetitive and aversive memory recall
24 in an insect. *BMC Biol*, 7:46.
- 25

- 1 Monastirioti, M., Linn, C. E. Jr. & White, K. (1996). Characterization of *Drosophila*
2 tyramine beta-hydroxylase gene and isolation of mutant flies lacking octopamine. *J Neurosci*,
3 16:3900-3911.
- 4
- 5 Mustard, J. A., Beggs, K. T. & Mercer, A. R. (2005). Molecular biology of the invertebrate
6 dopamine receptors. *Arch Insect Biochem Physiol*, 59:103-117.
- 7
- 8 Mustard, J. A., Pham, P. M. & Smith, B. H. (2010). Modulation of motor behavior by
9 dopamine and the D1-like dopamine receptor *AmDOP2* in the honey bee. *J Insect Physiol*.
10 56:422-430.
- 11
- 12 Nagatsu, T. (1989). The human tyrosine hydroxylase gene. *Cell Mol Neurobiol*, 9:313-321.
- 13
- 14 Nakamura, T., Yoshizaki, M., Ogawa, S., Okamoto, H., Shinmyo, Y., Bando, T., Ohuchi, H.,
15 Noji, S. & Mito, T. (2010). Imaging of transgenic cricket embryos reveals cell movements
16 consistent with a syncytial patterning mechanism. *Curr Biol*, 20:1641-1647.
- 17
- 18 Nakashima, A., Mori, K., Suzuki, T., Kurita, H., Otani, M., Nagatsu, T. & Ota, A. (1999).
19 Dopamine inhibition of human tyrosine hydroxylase type 1 is controlled by the specific
20 portion in the N-terminus of the enzyme. *J Neurochem*, 72:2145-2153.
- 21
- 22 Nakashima, A., Hayashi, N., Mori, K., Kaneko, Y. S., Nagatsu, T. & Ota, A. (2000). Positive
23 charge intrinsic to Arg³⁷-Arg³⁸ is critical for dopamine inhibition of the catalytic activity of
24 human tyrosine hydroxylase type 1. *FEBS Lett*, 465:59-63.
- 25

- 1 Nakashima, A., Hayashi, N., Kaneko, Y. S., Mori, K., Sabban, E. L., Nagatsu, T. & Ota, A.
2 (2009). Role of N-terminus of tyrosine hydroxylase in the biosynthesis of catecholamines. *J*
3 *Neural Transm*, 116:1355-1362.
- 4
- 5 Nakashima, A., Mori, K., Kaneko, Y. S., Hayashi, N., Nagatsu, T. & Ota, A. (2011).
6 Phosphorylation of the N-terminal portion of tyrosine hydroxylase triggers proteasomal
7 digestion of the enzyme. *Biochem Biophys Res Commun*, 407:343-347.
- 8
- 9 Neckameyer, W. S. (1996). Multiple roles for dopamine in *Drosophila* development. *Dev*
10 *Biol*, 176:209-219.
- 11
- 12 Ninomiya, Y. & Hayakawa, Y. (2007). Insect cytokine, growth-blocking peptide, is a primary
13 regulator of melanin-synthesis enzymes in armyworm larval cuticle. *FEBS J*, 274:1768-1777.
- 14
- 15 O'Dowd, B. F., Nguyen, T., Jung, B. P., Marchese, A., Cheng, R., Heng, H. H., Kolakowski,
16 L. F. Jr., Lynch, K. R. & George, S. R. (1997). Cloning and chromosomal mapping of four
17 putative novel human G-protein-coupled receptor genes. *Gene*, 187:75-81.
- 18
- 19 Passier, P. C., Vullings, H. G., Diederer, J. H. & Van der Horst, D. J. (1995). Modulatory
20 effects of biogenic amines on adipokinetic hormone secretion from locust corpora cardiaca *in*
21 *vitro*. *Gen Comp Endocrinol*, 97 :231-238.
- 22
- 23 Pastor, D., Piulachs, M. D., Cassier, P., André, M. & Bellés, X. (1991). Etude *in vivo* et *in*
24 *vitro* de l'action de la dopamine sur la croissance des ovocytes et la production d'hormone

juvénile chez *Blattella germanica* (L.) (Dictyoptera; Blattellidae). *C R Acad Sci III*, 313:207-212.

Rao, V. T., Siddiqui, S. Z., Prichard, R. K. & Forrester, S. G. (2009). A dopamine-gated ion channel (HcGGR3*) from *Haemonchus contortus* is expressed in the cervical papillae and is associated with macrocyclic lactone resistance. *Mol Biochem Parasitol*, 166:54-61.

Reverte, C. G., Ahearn, M. D. & Hake, L. E. (2001). CPEB degradation during *Xenopus* oocyte maturation requires a PEST domain and the 26S proteasome. *Dev Biol*, 231:447-458.

Rietdorf, K., Blenau, W. & Walz, B. (2005). Protein secretion in cockroach salivary glands requires an increase in intracellular cAMP and Ca^{2+} concentrations. *J Insect Physiol*, 51:1083-1091.

Rillich, J. & Stevenson, P. A. (2011). Winning fights induces hyperaggression via the action of the biogenic amine octopamine in crickets. *PLoS One*, 6:e28891.

Rillich, J., Schildberger, K. & Stevenson, P. A. (2011). Octopamine and occupancy: an aminergic mechanism for intruder-resident aggression in crickets. *Proc Biol Sci*, 278:1873-1880.

Ringstad, N., Abe, N. & Horvitz, H.R. (2009). Ligand-gated chloride channels are receptors for biogenic amines in *C. elegans*. *Science*, 325:96-100.

Roeder, T. (1999). Octopamine in invertebrates. *Prog Neurobiol*, 59:533-561.

- 1
- 2 Roeder, T. (2005). Tyramine and octopamine: ruling behavior and metabolism. *Annu Rev*
3 *Entomol*, 50:447-477.
- 4
- 5 Rogers, S., Wells, R. & Rechsteiner, M. (1986). Amino acid sequences common to rapidly
6 degraded proteins: the PEST hypothesis. *Science*, 234:364-368.
- 7
- 8 Samaranayaka, M. (1976). Possible involvement of monoamines in the release of
9 adipokinetic hormone in the locust *Schistocerca gregaria*. *J Exp Biol*, 65:415-425.
- 10
- 11 Sasaki, K., Yamasaki, K., Tsuchida, K. & Nagao, T. (2009). Gonadotropic effects of
12 dopamine in isolated workers of the primitively eusocial wasp, *Polistes chinensis*.
13 *Naturwissenschaften*, 96:625-629.
- 14
- 15 Sawzdargo, M., Nguyen, T., Lee, D. K., Lynch, K. R., Cheng, R., Heng, H. H., George, S. R.
16 & O'Dowd, B. F. (1999). Identification and cloning of three novel human G protein-coupled
17 receptor genes GPR52, Ψ GPR53 and GPR55: GPR55 is extensively expressed in human
18 brain. *Brain Res Mol Brain Res*, 64:193-198.
- 19
- 20 Schulz, D. J. & Robinson, G. E. (1999). Biogenic amines and division of labor in honey bee
21 colonies: behaviorally related changes in the antennal lobes and age-related changes in the
22 mushroom bodies. *J Comp Physiol A*. 184:481-488.
- 23
- 24 Shumway, S. D., Maki, M. & Miyamoto, S. (1999). The PEST domain of I κ B α is necessary
25 and sufficient for *in vitro* degradation by μ -calpain. *J Biol Chem*, 274:30874-30881.

1

2 Srivastava, D. P., Yu, E. J., Kennedy, K., Chatwin, H., Reale, V., Hamon, M., Smith, T., &

3 Evans, P. D. (2005). Rapid, nongenomic responses to ecdysteroids and catecholamines

4 mediated by a novel *Drosophila* G-protein-coupled receptor. *J Neurosci*, 25:6145-6155.

5

6 Stevenson, P. A., Dyakonova, V., Rillich, J. & Schildberger, K. (2005). Octopamine and

7 experience-dependent modulation of aggression in crickets. *J Neurosci*, 25:1431-41.

8

9 Spencer, M. L., Theodosiou, M. & Noonan, D. J. (2004). NPDC-1, a novel regulator of

10 neuronal proliferation, is degraded by the ubiquitin/proteasome system through a PEST

11 degradation motif. *J Biol Chem*, 279:37069-37078.

12

13 Stevenson, P. A., Hofmann, H. A., Schoch, K. & Schildberger, K. (2000). The fight and

14 flight responses of crickets depleted of biogenic amines. *J Neurobiol*, 43, 107-120.

15

16 Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. & Kumar S. (2011). MEGA5:

17 Molecular Evolutionary Genetics Analysis using Maximum Likelihood, Evolutionary

18 Distance, and Maximum Parsimony Methods. *Mol Biol Evol*, 28:2731-2739.

19

20 Taylor, D. J., Robinson, G. E., Logan, B. J., Lavery, R. & Mercer, A. R. (1992). Changes in

21 brain amine levels associated with the morphological and behavioural development of the

22 worker honeybee. *J Comp Physiol A*. 170:715-721.

23

1 Unoki, S., Matsumoto, Y. & Mizunami, M. (2005). Participation of octopaminergic reward
2 system and dopaminergic punishment system in insect olfactory learning revealed by
3 pharmacological study. *Eur J Neurosci*, 22:1409-1416.

4
5 Unoki, S., Matsumoto, Y. & Mizunami, M. (2006). Roles of octopaminergic and
6 dopaminergic neurons in mediating reward and punishment signals in insect visual learning.
7 *Eur J Neurosci*, 24:2031-2038.

8
9 Vergoz, V., Lim, J. & Oldroyd, B. P. (2012). Biogenic amine receptor gene expression in the
10 ovarian tissue of the honey bee *Apis mellifera*. *Insect Mol Biol*, 21:21-29.

11
12 Vié, A., Cigna, M., Toci, R. & Birman, S. (1999). Differential regulation of *Drosophila*
13 tyrosine hydroxylase isoforms by dopamine binding and cAMP-dependent phosphorylation. *J*
14 *Biol Chem*, 274:16788-16795.

15
16 Wada-Katsumata, A., Yamaoka, R. & Aonuma, H. (2011). Social interactions influence
17 dopamine and octopamine homeostasis in the brain of the ant *Formica japonica*. *J Exp Biol*,
18 214:1707-1713.

19
20 Waddell, S. (2010). Dopamine reveals neural circuit mechanisms of fly memory. *Trends*
21 *Neurosci*, 33:457-464.

22
23 Wallace, B. G. (1976). The biosynthesis of octopamine-characterization of lobster tyramine
24 beta-hydroxylase. *J Neurochem*, 26:761-770.

- 1 Watanabe, T., Sadamoto & H., Aonuma, H. (2011). Identification and expression analysis of
2 the genes involved in serotonin biosynthesis and transduction in the field cricket *Gryllus*
3 *bimaculatus*. *Insect Mol Biol*, 20:619-635.
- 4
- 5 Walz, B., Baumann, O., Krach, C., Baumann, A. & Blenau, W. (2006). The aminergic
6 control of cockroach salivary glands. *Arch Insect Biochem Physiol*, 62:141-152.
- 7
- 8 Woodring, J. & Hoffmann, K. H. (1994). The effects of octopamine, dopamine and serotonin
9 on juvenile hormone synthesis, *in vitro*, in the cricket, *Gryllus bimaculatus*. *J Insect Physiol*,
10 40: 797-802.
- 11
- 12 Yan, Z., Feng, J., Fienberg, A. A. & Greengard, P. (1999). D₂ dopamine receptors induce
13 mitogen-activated protein kinase and cAMP response element-binding protein
14 phosphorylation in neurons. *Proc Natl Acad Sci U S A*, 96:11607-11612.
- 15

FIGURES AND TABLES

Figure 1

Comparison of the amino acid sequences of *Gryllus* TH with other TH proteins. (A) Alignment of the deduced amino acid sequence of *Gryllus* TH-A isoform with that of *Drosophila* pale-A isoform (NP_476897), *Mythimna* TH-A (type 2, brain form; BAF32574) and *Apis* TH (NP_001011633). Identical amino acids are printed in white letters on a black background. The catalytic domain is surrounded by a gray box. Black lines indicate the N-terminal regulatory (ACT) domain and the C-terminal coiled-coil region. The tetrahydrobiopterin (BH₄)-binding sites and iron binding sites are indicated by white and black arrowheads, respectively. The putative phosphorylation sites conserved between *Gryllus* TH and *Drosophila* pale proteins are indicated by white circles above the alignments. Two arginine residues important for catecholamine-mediated inhibition of enzyme activity (Nakashima et al., 1999; Nakashima et al., 2000) are indicated by black circles above the alignments. (B) Alignment of the deduced amino acid sequence of the ACT domain of insect TH-A and -B isoforms. GenBank accession numbers of aligned insect TH isoforms are as follows: *Drosophila* pale-A (NP_476897), *Drosophila* pale-B (NP_476898) *Mythimna* TH-B (BAF32573).

Figure 2

Molecular phylogenetic tree of *Gryllus* TH and other aromatic amino acid hydroxylases. The scale bar indicates 0.08 substitutions per site. The species name and the GenBank accession numbers of aromatic amino acid hydroxylases are as follows: *Drosophila* pale-A (*Drosophila melanogaster*, NP_476897), *Mythimna* TH (*Mythimna separata*, BAF32574), *Apis* TH (*Apis mellifera*, NP_001011633), *Tribolium* TH (*Tribolium castaneum*, NP_001092299), *Caenorhabditis* CAT-2 (*Caenorhabditis elegans*, ADZ54165), *Mus* TH

(*Mus musculus*, NP_033403), *Danio* TH (*Danio rerio*, NP_571224), *Danio* TH2 (*Danio rerio*, NP_001001829), *Drosophila* dTRH (*Drosophila melanogaster*, NP_612080), *Apis* TRH (*Apis mellifera*, XP_394674), *Tribolium* TRH (*Tribolium castaneum*, XP_967413), *Gryllus* TRH (*Gryllus bimaculatus*, BAJ83476), *Caenorhabditis* tph-1 (*Caenorhabditis elegans*, NP_495584), *Aplysia* TPH (*Aplysia californica*, ABF18968), *Mus* TpH1 (*Mus musculus*, NP_033440), *Danio* TpH1 (*Danio rerio*, NP_840091), *Mus* TpH2 (*Mus musculus*, NP_775567), *Danio* TpH2 (*Danio rerio* NP_999960), *Danio* tphd2 (*Danio rerio*, AAT38217), *Drosophila* Henna (*Drosophila melanogaster*, NP_523963), *Apis* TPH (*Apis mellifera*, XP_623300), *Tribolium* TPH (*Tribolium castaneum*, XP_967025), *Gryllus* TPH (*Gryllus bimaculatus*, BAJ83477), *Caenorhabditis* pah-1 (*Caenorhabditis elegans*, NP_495863), *Mus* TPH (*Mus musculus*, NP_032803), *Danio* TPH (*Danio rerio*, NP_956845).

Figure 3

Tissue-specific expression patterns of TH isoforms. The cDNA fragments corresponding to the catalytic domain (*TH* common region) and the isoform-specific ACT domains (*TH-A* isoform and *TH-B* isoform) were amplified by RT-PCR from ten cricket tissues. *Gryllus Eflalpha* gene was amplified as an internal control gene. PCR products were run on a 1.5% agarose gel and stained with ethidium bromide. SOG = suboesophageal ganglion; CC + CA = corpus cardiacum-corpora allata complex.

Figure 4

Comparison of the amino acid sequences of *Gryllus* D₁-like dopamine receptor (Dop1) and INDR (Dop2) with *Apis* homologues. (A) Alignment of the deduced amino acid sequence of *Gryllus* Dop1 with that of *Drosophila* Dop1 (AAA85716), *Bombyx* Dop1 (NP_001108459) and *Apis* Dop1 (NP_001011595). (B) Alignment of the deduced amino acid

sequence of *Gryllus* Dop2 with that of *Drosophila* Dop2 (NP_733299), *Bombyx* Dop2 (NP_001108338) and *Apis* Dop2 (NP_001011567). Identical amino acids are printed in white letters on a black background. Black lines indicate the transmembrane segments (TM). The conserved DRY motif is indicated by white circles above the alignments. Putative N-glycosylation sites of *Gryllus* dopamine receptors are indicated by white arrowheads above the alignments. Putative phosphorylation sites conserved between dopamine receptors are indicated by black arrowheads under the alignments.

Figure 5

Molecular phylogenetic tree of *Gryllus* Dop1, Dop2 and other dopamine receptors. Vertebrate D₂-like dopamine receptors are used as an outgroup. The scale bar indicates 0.08 substitution per site. The species name and the GenBank accession numbers of the proteins are as follows: *Drosophila* Dop1 (*Drosophila melanogaster*, AAA85716), *Apis* Dop1 (*Apis mellifera*, NP_001011595), *Nasonia* Dop1 (*Nasonia vitripennis*, XP_001606438), *Acyrtosiphon* Dop1 (*Acyrtosiphon pisum*, XP_001947683), *Drosophila* Dop2 (*Drosophila melanogaster*, NP_733299), *Aedes* Dop2 (*Aedes aegypti*, XP_001651499), *Bombyx* Dop2 (*Bombyx mori*, NP_001108338), *Apis* Dop2 (*Apis mellifera*, NP_001011567), *Nasonia* Dop2 (*Nasonia vitripennis*, NP_001155849), *Aplysia* D₁-like (*Aplysia californica*, NP_001191631), *Danio* D₁ (*Danio rerio*, NP_001129448), *Danio* D₅-like 1 (*Danio rerio*, XP_001341592), *Danio* D₅-like 2 (*Danio rerio*, XP_692025), *Mus* D₁ (*Mus musculus*, NP_034206), *Mus* D₂ (*Mus musculus*, NP_034207), *Mus* D₃ (*Mus musculus*, NP_031903), *Mus* D₄ (*Mus musculus*, NP_031904), *Mus* D₅ (*Mus musculus*, NP_038531), *Mus* α_{1A} adrenergic receptor (*Mus musculus*, NP_038489), *Mus* α_{1B} adrenergic receptor (*Mus musculus*, NP_031442), *Danio* α_{1A} adrenergic receptor-like (*Danio rerio*, XP_001338938), *Danio* α_{1B} adrenergic receptor

(*Danio rerio*, NP_001007359), *Danio* α_{1B} adrenergic receptor-like (*Danio rerio*, XP_001922013), *Danio* α_{1D} adrenergic receptor (*Danio rerio*, XP_697043).

Figure 6

Comparison of the amino acid sequences of the *Gryllus* D₂-like dopamine receptor (Dop3) with other D₂-like dopamine receptors. (A) Alignment of the deduced amino acid sequence of *Gryllus* Dop3 with that of *Drosophila* DD2R (NP_001014760), *Apis* Dop3 (NP_001014983) and *Tribolium* Dop3 (XP_969037). Identical amino acids are printed in white letters on a black background. Black lines indicate the transmembrane segments (TM). (B) Molecular phylogenetic tree of *Gryllus* Dop3 and other dopamine receptors. Vertebrate D₁-like dopamine receptors are used as an outgroup. The scale bar indicates 0.07 substitutions per site. The species name and the GenBank accession numbers of the cDNAs are as follows: *Drosophila* DDR2 (*Drosophila melanogaster*, NM_001014760), *Apis* Dop3 (*Apis mellifera*, NM_001014983), *Nasonia* Dop3 (*Nasonia vitripennis*, XM_001602460), *Tribolium* Dop3 (*Tribolium castaneum*, XM_963944), *Pediculus* Dop3 (*Pediculus humanus corporis*, XM_002426878), *Panulirus* Dop3 (*Panulirus interruptus*, DQ900655), *Caenorhabditis* Dop-2 (*Caenorhabditis elegans*, NM_001028876), *Brugia* Dop-2 (*Brugia malayi*, XM_001901847), *Mus* D₁ (*Mus musculus*, NP_034206), *Mus* D₂ (*Mus musculus*, NM_010077), *Mus* D₃ (*Mus musculus*, NM_007877), *Mus* D₄ (*Mus musculus*, NM_007878), *Mus* D₅ (*Mus musculus*, NP_038531), *Danio* D_{2a} (*Danio rerio*, NM_183068), *Danio* D_{2b} (*Danio rerio*, NM_197936), *Danio* D_{2-like} (*Danio rerio*, NM_197935), *Danio* D₃ (*Danio rerio*, NM_183067), *Danio* D_{4a} (*Danio rerio*, NM_001012616), *Danio* D_{4b} (*Danio rerio*, NM_001012618), *Danio* D_{4-related} (*Danio rerio*, NM_001012620).

Figure 7

Comparison of the amino acid sequences of *Gryllus* DopEcR with other G-protein coupled receptors. (A) Alignment of the deduced amino acid sequence of *Gryllus* DopEcR with that of *Drosophila* DopEcR (NP_647897), *Apis* DopEcR (XP_396491) and *Tribolium* DopEcR (XP_968380). Identical amino acids are printed in white letters on a black background. Black lines indicate the transmembrane segments (TM). The DRY motif is indicated by white circles above the alignments. (B) Molecular phylogenetic tree of *Gryllus* DopEcR and other G-protein coupled receptors. *Mus musculus* histamine H₂ receptor is used as an outgroup. The scale bar indicates 0.15 substitutions per site. The species name and the GenBank accession numbers of the proteins are as follows: *Drosophila* DopEcR (*Drosophila melanogaster*, NP_647897), *Aedes* DopEcR (*Aedes aegypti*, XP_001654794), *Apis* DopEcR (*Apis mellifera*, XP_396491), *Tribolium* DopEcR (*Tribolium castaneum*, XP_968380), *Mus* GPR21 (*Mus musculus*, NP_001139802), *Gallus* GPR21 (*Gallus gallus*, XP_001233342), *Xenopus* GPR21 (*Xenopus tropicalis*, XP_002931480), *Mus* GPR52 (*Mus musculus*, NP_796357), *Gallus* GPR52 (*Gallus gallus*, XP_001234532), *Danio* GPR52 (*Danio rerio*, CAK04352), *Drosophila* OA β_1 (*Drosophila melanogaster*, AJ880687), *Drosophila* OA β_2 (*Drosophila melanogaster*, NP_001034049), *Mus* β_1 adrenergic receptor (*Mus musculus*, NP_031445), *Mus* β_2 adrenergic receptor (*Mus musculus*, NP_031446), *Mus* β_3 adrenergic receptor (*Mus musculus*, NP_038490), *Mus* histamine H₂ receptor (*Mus musculus*, NP_032312).

Figure 8

Tissue specific expression patterns of mRNA of *Gryllus* dopamine receptor genes. The cDNA fragments of *Gryllus* *Dop1*, *Dop2*, *Dop3* and *DopEcR* genes were amplified by RT-PCR from ten cricket tissues. The *Gryllus* *Eflalpha* gene was amplified as an internal control

gene. PCR products were run on a 1.5% agarose gel and stained with ethidium bromide. SOG = suboesophageal ganglion; CC + CA = corpus cardiacum-corpora allata complex.

Figure 9

Comparison of the amino acid sequences of *Gryllus* DAT with other monoamine transporters. (A) Alignment of the deduced amino acid sequence of *Gryllus* DAT with that of *Drosophila* DAT (NP_523763), *Bombyx* DAT (NP_001037362) and *Apis* DAT (NP_001139210). Identical amino acids are printed in white letters on a black background. Black lines indicate the transmembrane segments (TM). Putative N-glycosylation sites of *Gryllus* DAT are indicated by white arrowheads above the alignments. Putative phosphorylation sites conserved between *Gryllus* and *Drosophila* DAT are indicated by black arrowheads under the alignments. ‘S-S bond’ indicates two conserved cystine residues that form a disulfide bond (Chen et al., 2007). (B) Molecular phylogenetic tree of *Gryllus* DAT and other monoamine transporters. The scale bar indicates 0.06 substitutions per site. The species name and the GenBank accession numbers of the biogenic amine transporters are as follows: *Drosophila* DAT (*Drosophila melanogaster*, NP_523763), *Aedes* DAT (*Aedes aegypti*, XP_001654246), *Bombyx* DAT (*Bombyx mori*, NP_001037362), *Trichoplusia* DAT (*Trichoplusia ni*, AAN52844), *Apis* DAT (*Apis mellifera*, NP_001139210), *Caenorhabditis* dat-1 (*Caenorhabditis elegans*, NP_499043), *Mus* DAT (*Mus musculus*, NP_034150), *Danio* DAT (*Danio rerio*, NP_571830), *Mus* NAT (*Mus musculus*, NP_033235), *Danio* NAT (*Danio rerio*, XP_694138), *Trichoplusia* OAT (*Trichoplusia ni*, AF388173), *Tribolium* OAT (*Tribolium castaneum*, XP_975356), *Pediculus* OAT (*Pediculus humanus corporis*, XP_002425932), *Acyrtosiphon* OAT (*Acyrtosiphon pisum*, XP_001949303), *Drosophila* SERT (*Drosophila melanogaster*, NP_523846), *Bombyx* SERT (*Bombyx mori*, NP_001037436), *Manduca* SERT (*Manduca sexta*, AAN59781), *Tribolium* SERT (*Tribolium*

castaneum, XP_968717), *Acyrtosiphon* SERT (*Acyrtosiphon pisum*, XP_001944311), *Mus* SERT (*Mus musculus*, NP_034614).

Figure 10

Tissue-specific expression pattern of *Gryllus* *DAT* gene. The cDNA fragments of *Gryllus* *DAT* gene were amplified by RT-PCR from ten cricket tissues. *Gryllus* *Eflalpha* gene was amplified as an internal control gene. PCR products were run on a 1.5% agarose gel and stained with ethidium bromide. SOG = suboesophageal ganglion; CC + CA = corpus cardiacum-corpora allata complex.

1 **Table 1. List of primers used in this study**

Degenerate primers

	Forward primer	Reverse primer
<i>TH</i> (C-terminal - region)	5'-	5'-CACYTCTGAANGGYCGSGACAT-3'
<i>DopEcR</i>	GGTNTAYACNTTYATGTGGAT -3'	5'- GRCCATCCARCAYTGRCA -3'
<i>DAT</i>	5'- GGCATHCCYYTNTTCTACATG G-3'	5'- ATDATGACGTTRTAGTAGAARTCNACG TA-3'

Primers for full-length/partial ORF amplification

	Forward primer	Reverse primer
<i>TH</i>	5'-CCGCGACAACAACCTCTCAG-3'	5'-CATGCCGGTGGACACCCAC-3'
<i>Dop1</i>	5'- GCGCCGCCATGGAGGACGACG -3'	5'-GCGGCCGTCTCTTCAGATGG-3'
<i>Dop2</i>	5'- GCCGACGAAACAACCTCGAG-3'	5'-CCGGCTGTTTGTTCAGGTGG-3'

<i>Dop3</i>	5'-GCCCTTCTTCACGTGCAAC- 3'	5'-CCAGCGCGTGTTGTTACG-3'
	5'-	
<i>DopEcR</i>	CCCTGCACCCCTCACCGCGAC	5'-CCAGCGCGTGTTGTTACG-3'
	TATG-3'	
		5'-
<i>DAT</i>	5'-CGCTGAGGAGCTGCAGATG- 3'	GAGCACCTTTTGGCAAATAGAGAAATC ACAC-3'

Primers for tissue-specific expression

analysis

	Forward primer	Reverse primer
<i>TH</i>		
common region	5'-GGCCTTCCGGATCTTCC-3'	5'-GGGTGTCCACCGGCATG-3'
<i>TH-A</i> isoform	5'-CCGCGACAACAACCTCTCAG-3'	5'-CCGTCAGCTCAGCGTCGTTG-3'
<i>TH-B</i> isoform	5'-CCGCGACAACAACCTCTCAG-3'	5'-CCGTCAGCTCAGCGTCATC-3'
<i>Dop1</i>	5'-CGCCGTACCACGTGTCTG-3'	5'-CGCCGAGTTGGAGTAGC-3'
<i>Dop2</i>	5'-CCAACTCCAGCATGAATCC-3'	5'-GAGTGCGCCAGCATCATGG-3'

Dop3 5'-GCCATGTGCACCAAGCTG-3' 5'-CCAGCGCGTGTTGTTACG-3'

5'-

DopEcR CGTACAGCGGTGGGTGTATGG 5'-CGTCTGCACCGTCTCGTAGC-3'

-3'

5'-

5'-

DAT GTGGCGAGATCATCAAATAGC GAGCACCTTTTGGCAAATAGAGAAATC

TGC-3'

ACAC-3'

5'-

Eflalpha GTGTTCTGAAGCCAGGTATGG-5'-CTCCAGCAACATAACCACGAC-3'

3'

A

Gryllus TH-A 1 MMVAAQAQKNREMPAIIKSSYSIENGYPARRRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Drosophila pale-A 1 MMVAAQAQKNREMPAIIKSSYSIENGYPARRRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Mythimna TH-A 1 MMVAAQAQKNREMPAIIKSSYSIENGYPARRRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Apis TH 1 MMVAAQAQKNREMPAIIKSSYSIENGYPARRRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL

Regulatory (ACT) domain

96 DGVGALARILKTIENFKGSVTHVEHRAAGTGARADALVRVDISROQLLLLRALRQSAALSAVALLADTHVSIKDPWFFRHASLDNCNHLMTK
Gryllus TH-A 96 DGVGALARILKTIENFKGSVTHVEHRAAGTGARADALVRVDISROQLLLLRALRQSAALSAVALLADTHVSIKDPWFFRHASLDNCNHLMTK
Drosophila pale-A 96 DGVGALARILKTIENFKGSVTHVEHRAAGTGARADALVRVDISROQLLLLRALRQSAALSAVALLADTHVSIKDPWFFRHASLDNCNHLMTK
Mythimna TH-A 96 DGVGALARILKTIENFKGSVTHVEHRAAGTGARADALVRVDISROQLLLLRALRQSAALSAVALLADTHVSIKDPWFFRHASLDNCNHLMTK
Apis TH 96 DGVGALARILKTIENFKGSVTHVEHRAAGTGARADALVRVDISROQLLLLRALRQSAALSAVALLADTHVSIKDPWFFRHASLDNCNHLMTK

191 YEPFLDMNHHPGFADKVVYRERRKMTADIAFAKRGDPIPTTGYAAATATWVVFNNVDFLPHACAEYRRVGMGSAANIFGPRIPQDEMS
Gryllus TH-A 191 YEPFLDMNHHPGFADKVVYRERRKMTADIAFAKRGDPIPTTGYAAATATWVVFNNVDFLPHACAEYRRVGMGSAANIFGPRIPQDEMS
Drosophila pale-A 191 YEPFLDMNHHPGFADKVVYRERRKMTADIAFAKRGDPIPTTGYAAATATWVVFNNVDFLPHACAEYRRVGMGSAANIFGPRIPQDEMS
Mythimna TH-A 191 YEPFLDMNHHPGFADKVVYRERRKMTADIAFAKRGDPIPTTGYAAATATWVVFNNVDFLPHACAEYRRVGMGSAANIFGPRIPQDEMS
Apis TH 191 YEPFLDMNHHPGFADKVVYRERRKMTADIAFAKRGDPIPTTGYAAATATWVVFNNVDFLPHACAEYRRVGMGSAANIFGPRIPQDEMS

Catalytic domain

286 FLNRHGTGFLRPAAGLLTARDFLASLAFRNFQSTQYVHRHSSSPHPTPEPCDIEHMGHMLLADPFKAQFSQIEGLASLGASDAIEKLSTVYWF
Gryllus TH-A 286 FLNRHGTGFLRPAAGLLTARDFLASLAFRNFQSTQYVHRHSSSPHPTPEPCDIEHMGHMLLADPFKAQFSQIEGLASLGASDAIEKLSTVYWF
Drosophila pale-A 286 FLNRHGTGFLRPAAGLLTARDFLASLAFRNFQSTQYVHRHSSSPHPTPEPCDIEHMGHMLLADPFKAQFSQIEGLASLGASDAIEKLSTVYWF
Mythimna TH-A 286 FLNRHGTGFLRPAAGLLTARDFLASLAFRNFQSTQYVHRHSSSPHPTPEPCDIEHMGHMLLADPFKAQFSQIEGLASLGASDAIEKLSTVYWF
Apis TH 286 FLNRHGTGFLRPAAGLLTARDFLASLAFRNFQSTQYVHRHSSSPHPTPEPCDIEHMGHMLLADPFKAQFSQIEGLASLGASDAIEKLSTVYWF

381 RVEFGLCCKEHLKAYGAGLLSSVYGLLHALSDCKEHRPPEPAATGVPYODQYQVPIYVAESFEDAKDKFRWVSTGM-----
Gryllus TH-A 381 RVEFGLCCKEHLKAYGAGLLSSVYGLLHALSDCKEHRPPEPAATGVPYODQYQVPIYVAESFEDAKDKFRWVSTGM-----
Drosophila pale-A 381 RVEFGLCCKEHLKAYGAGLLSSVYGLLHALSDCKEHRPPEPAATGVPYODQYQVPIYVAESFEDAKDKFRWVSTGM-----
Mythimna TH-A 381 RVEFGLCCKEHLKAYGAGLLSSVYGLLHALSDCKEHRPPEPAATGVPYODQYQVPIYVAESFEDAKDKFRWVSTGM-----
Apis TH 381 RVEFGLCCKEHLKAYGAGLLSSVYGLLHALSDCKEHRPPEPAATGVPYODQYQVPIYVAESFEDAKDKFRWVSTGM-----

475 EVLDSVDKLETLVHOMNTEHLHLTNALSKRRP-
Gryllus TH-A 475 EVLDSVDKLETLVHOMNTEHLHLTNALSKRRP-
Drosophila pale-A 474 EVLDSVDKLETLVHOMNTEHLHLTNALSKRRP-
Mythimna TH-A 474 EVLDSVDKLETLVHOMNTEHLHLTNALSKRRP-
Apis TH 476 EVLDSVDKLETLVHOMNTEHLHLTNALSKRRP-

Coiled coil region**B**

29 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Gryllus TH-A 29 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Mythimna TH-A 30 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Drosophila pale-A 30 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
 30 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Gryllus TH-B 30 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Mythimna TH-B 29 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL
Drosophila pale-B 30 RRSVVDDARFETLVVKOTKQSVLEEARQNSDADLTTEEVLHANAASSSSDAPAAQSAALVLRRL

B isoform-specific insertion in the ACT domain

61 -----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Gryllus TH-A 61 -----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Mythimna TH-A 60 -----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Drosophila pale-A 61 -----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
 107 PEPFPDPTPEQL-----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Gryllus TH-B 107 PEPFPDPTPEQL-----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Mythimna TH-B 98 -----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD
Drosophila pale-B 108 PEPFPDPTPEQL-----DALLTEPEMLTATAAASAEAAQVQVAAALVLRRLDVGAGARIILKINENFKGCVTVVRAAAGTGVD

130 ADAIVRVVD
Gryllus TH-A 130 ADAIVRVVD
Mythimna TH-A 129 ADAIVRVVD
Drosophila pale-A 130 ADAIVRVVD
 196 ADAIVRVVD
Gryllus TH-B 196 ADAIVRVVD
Mythimna TH-B 181 ADAIVRVVD
Drosophila pale-B 201 ADAIVRVVD

Central brain
SOG
Lamina + Medulla + Retina
CC + CA
Thoracic muscles
Salivary gland
Midgut
Malpighian tubules
Testicle
Ovary

0000 0000 0000 0000 0000 0000 0000 0000 0000 0000

[illegible]

RT(+)
RT(-)

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	105	106	107	108	109	110	111	112	113	114	115	116	117	118	119	120	121	122	123	124	125	126	127	128	129	130	131	132	133	134	135	136	137	138	139	140	141	142	143	144	145	146	147	148	149	150	151	152	153	154	155	156	157	158	159	160	161	162	163	164	165	166	167	168	169	170	171	172	173	174	175	176	177	178	179	180	181	182	183	184	185	186	187	188	189	190	191	192	193	194	195	196	197	198	199	200	201	202	203	204	205	206	207	208	209	210	211	212	213	214	215	216	217	218	219	220	221	222	223	224	225	226	227	228	229	230	231	232	233	234	235	236	237	238	239	240	241	242	243	244	245	246	247	248	249	250	251	252	253	254	255	256	257	258	259	260	261	262	263	264	265	266	267	268	269	270	271	272	273	274	275	276	277	278	279	280	281	282	283	284	285	286	287	288	289	290	291	292	293	294	295	296	297	298	299	300	301	302	303	304	305	306	307	308	309	310	311	312	313	314	315	316	317	318	319	320	321	322	323	324	325	326	327	328	329	330	331	332	333	334	335	336	337	338	339	340	341	342	343	344	345	346	347	348	349	350	351	352	353	354	355	356	357	358	359	360	361	362	363	364	365	366	367	368	369	370	371	372	373	374	375	376	377	378	379	380	381	382	383	384	385	386	387	388	389	390	391	392	393	394	395	396	397	398	399	400	401	402	403	404	405	406	407	408	409	410	411	412	413	414	415	416	417	418	419	420	421	422	423	424	425	426	427	428	429	430	431	432	433	434	435	436	437	438	439	440	441	442	443	444	445	446	447	448	449	450	451	452	453	454	455	456	457	458	459	460	461	462	463	464	465	466	467	468	469	470	471	472	473	474	475	476	477	478	479	480	481	482	483	484	485	486	487	488	489	490	491	492	493	494	495	496	497	498	499	500	501	502	503	504	505	506	507	508	509	510	511	512	513	514	515	516	517	518	519	520	521	522	523	52
--	---	---	---	---	---	---	---	---	---	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	----

A

Gryllus Dopl1 1 MTNMAIRAIAAAGVGSAATVATSTTSSISSSTTIINTSSATTIGGNHTSGSTGFSTNSTLLDADHLPQLTTAKVDLDIEIDQLLTNGYDGT ME-
Drosophila Dopl1 1 MTNMAIRAIAAAGVGSAATVATSTTSSISSSTTIINTSSATTIGGNHTSGSTGFSTNSTLLDADHLPQLTTAKVDLDIEIDQLLTNGYDGT ME-FAST
Bombyx Dopl1 1
Apis Dopl1 1

Gryllus Dopl1 8 VNG---SSASAGPSGGAGGASADSNDDELLSPSPVGVLCALLSLIFLSHAGNGLVCVAIYTRGLRRRIGNLFLASLAADLFVAALVMTFAV
Drosophila Dopl1 96 TILTSFYNESSMTNASEMTITGGEPEPLSPVGVLCALLSLIFLSHAGNGLVCVAIYTRGLRRRIGNLFLASLAADLFVAALVMTFAV
Bombyx Dopl1 7 ---LAE---NVEPLIDAEVVDGDAVLGLVGLLILFLFSHAGNGLVCVAIYTRGLRRRIGNLFLASLAADLFVAALVMTFAV
Apis Dopl1 1 -----MILSONNITDQKEDDPENTFSLSPVGVLCALLSLIFLSHAGNGLVCVAIYTRGLRRRIGNLFLASLAADLFVGGVMTFAV

oco TM1 TM2

Gryllus Dopl1 100 NDLLGHVDFGEALCDTWAFADVCMSTASILNLCAISLDRIYHIKDPRLRYGRWVRRVAVGSIATVWLLAALVSFVPISLGLHRPDEPPPAARG
Drosophila Dopl1 188 NDLLGVHDFGEALCDTWAFADVCMSTASILNLCAISLDRIYHIKDPRLRYGRWVRRVAVGSIATVWLLAALVSFVPISLGLHRPDEPPPAARG
Bombyx Dopl1 95 NDLLGVHVFGEFCDTWAFADVCMSTASILNLCAISLDRIYHIKDPRLRYGRWVRRVAVGSIATVWLLAALVSFVPISLGLHRPDEPPPAARG
Apis Dopl1 85 NDLLGVHVFGEFCDTWAFADVCMSTASILNLCAISLDRIYHIKDPRLRYGRWVRRVAVGSIATVWLLAALVSFVPISLGLHRPDEPPPAARG

TM3 TM4

Gryllus Dopl1 195 PRGEBELTCADLLAPFYAVSSSVQSTPCLVHVIYCYRILYIAQKHKVSRATVTFWQVDEG-----AGDEPSHVRVDEPSQSSPH
Drosophila Dopl1 279 DKGKVEBELTCADLLAPFYAVSSSVQSTPCLVHVIYCYRILYIAQKHKVSRATVTFWQVDEG-----AGDEPSHVRVDEPSQSSPH
Bombyx Dopl1 186 ODPPKPYETCALDLPYTYAVSSVCSIFLPCVHVIYCYRILYIAQKHKVSRATVTFWQVDEG-----AGDEPSHVRVDEPSQSSPH
Apis Dopl1 176 DSKVEBELTCADLLAPFYAVSSSVQSTPCLVHVIYCYRILYIAQKHKVSRATVTFWQVDEG-----AGDEPSHVRVDEPSQSSPH

TM5

Gryllus Dopl1 279 VSDHKAATVGVINGVFLVCVWVFFCVNIHAAPCKTICPGAFRVLTWLGYSNSAFNPYIYSIFNKEFRDAPFRILANAAVDEGCGSCCPACC
Drosophila Dopl1 371 VSDHKAATVGVINGVFLVCVWVFFCVNIHAAPCKTICPGAFRVLTWLGYSNSAFNPYIYSIFNKEFRDAPFRILANAAVDEGCGSCCPACC
Bombyx Dopl1 268 VSDHKAATVGVINGVFLVCVWVFFCVNIHAAPCKTICPGAFRVLTWLGYSNSAFNPYIYSIFNKEFRDAPFRILANAAVDEGCGSCCPACC
Apis Dopl1 265 VSDHKAATVGVINGVFLVCVWVFFCVNIHAAPCKTICPGAFRVLTWLGYSNSAFNPYIYSIFNKEFRDAPFRILANAAVDEGCGSCCPACC

TM6 TM7

Gryllus Dopl1 372 GEFVSSRSLSLAAGQGRS-KASVDCGLGGMGAGAGCGGNGACAADAV---EPRSSAGAAAAVPTAVHASKPPPPQSPAQPPPPSPF
Drosophila Dopl1 456 -QDVGNHTR-RNSDRFIDYGA---RNVVVMI-SG---RSSA
Bombyx Dopl1 352 GCGVNRTP---RNDFFVDYGA---RNVVVMI-SG---RSSA
Apis Dopl1 344 ---N-RARGNP---STSSCG---FRSVVVKI---GSM-TECNI---SPRSSACQVGM

Gryllus Dopl1 463 QPRLAETRVSDTSDP-EEEVSAI
Drosophila Dopl1 489 -----LEQ---VSAI
Bombyx Dopl1 402 -----YH
Apis Dopl1 391 -----AQRHDEQ---VSAI

B

[illegible]

A

<i>Gryllus</i> Dop3	1	PFFTCNIMDAMCTKLSLDCS	PGVTAFILTTWLGYNMNSFVNPIVYITIN	NPEFRKAFKKLMALG	T
<i>Drosophila</i> DDR2	545	PPFSCNIMDAMCTKFKKDCRPG	LTAYMTTWLGYNNSFVNPIVYITIN	NPEFRKAFKKLMHMC	-
<i>Apis</i> Dop3	634	PPFSCNIMDAMCTKLTAD	COPGVTAFTVTSWLGYNMNSFVNPIVYT	FNPEFRKAFHKLVSF	--
<i>Tribolium</i> Dop3	391	PFFTCNIMDAMCSKLN	PCOPGVAAFTLTWLGYNMNSFVNPIVYITIN	NPEFRKAFKKLMFLGP	

TM6 TM7

B

0.07

A

<i>Gryllus</i>	DopEcR	1	-----MIVLSFDHNNNTREFTLELTOAGHFVPGVAIVVNVLIATIANLFRGSPSEVNVLLSLAVADLLCGLLVLPVLSVYPAVQVRW		
<i>Drosophila</i>	DopEcR	1	-----MQMSXYLDNS-----KVEALRAVLLSGVAIVLNSLLIATANFGPGEVENVYLLSLADLLCGLLVVPVSVYPAVGEV		
<i>Apis</i>	DopEcR	1	-----MEEP-----SLEALQAGHFVPGVAIVLNSLLIATIANLFRGSPSEVNVYLLSLADLLCGLLVLPVSVYPAVVRW		
<i>Tribolium</i>	DopEcR	1	MFRTRDSTSAHTIFSDNSNDGIVGTFLELQTOAGHFVPGVAIVVNVLIATIANLFRGSPSEVNVLLSLAVADLLCGLLVLPVLSVYPAVVRW		
				TM1	TM2
<i>Gryllus</i>	DopEcR	86	VYGDVVMCRIVGYLEVLTWAVSVVFPMFNISVSDRYLAVRKPLRYETVQTKRCQCWMVFTWISAAMLCCPPLLGFNFHFD-RDAVICMLDWGMMAA		
<i>Drosophila</i>	DopEcR	82	VGGDIVCRTEGYLEVLTWAVSVVFPMFNISVDRYLAVRKPLRYETVQTKRCQCWMVFTWISAAMLCCPPLHGYMPTFENNTHICMLDWGMMAA		
<i>Apis</i>	DopEcR	75	VGGDIVCRIVGYLEVLTWAVSVVFPMFNISVDRYLAVRKPLRYETVQTKRCQCWMVFTWISAAMCCPPLGLFKRPIFD-REAAICMLDWGMMAA		
<i>Tribolium</i>	DopEcR	96	VYGDIVCRIVGYLEVLTWAVSVVFPMFNISVDRYLAVRKPLRYETVQTKRCQCWMVFTWISAAMLCCPPLLGFNFHFD-RDAVICMLDWGMMAA		
			TM3	TM4	
<i>Gryllus</i>	DopEcR	180	KSVTAVLGLGPSLITITVYTYIFNMHMKLRSGVAFHDKKYATALSENLSNPSHMSVFLVLFVWSTWPTATVRYKYVYFGVQVQVFLHFI		
<i>Drosophila</i>	DopEcR	176	TSAPLALGLGPSLITVGNVGYTFVMMKRKSGEPHDKKYATALSENLSNPSHMSVFLVAFVWWSVPTILLRLRYKYVYFGVQVQVFLHFI		
<i>Apis</i>	DopEcR	169	TVSLTSLVGLGPSITITVITVYCYIFNMHMKLRSGVAFHDKKYATALSENLSNPSHMSVFLVAFVWWSVPTAGGRIRYKYVYFGVQVQVFLHFI		
<i>Tribolium</i>	DopEcR	190	TVSLTSLVGLGPSLITITVYTYIFNMHMKLRSGVAFHDKKYATALSENLSNPSHMSVFLVAFVWWSVPTIGVRYKYVYFGVQVQVFLHFI		
			TM5	TM6	
<i>Gryllus</i>	DopEcR	275	VWLGILNSFWKMLLVA-LSPQFLRLAFVFCMTCCKRYKGRLOAEELIGHEADD		
<i>Drosophila</i>	DopEcR	271	VWLGILNSFWKMLITMGS-SLPQFLRLAFVFCMTCCKRYKGRLOAEELIGHEADD		
<i>Apis</i>	DopEcR	263	VWLGIVNSFWKAVVLGA-LSPQFLRLAFVLCMTCCKRHRLPDELIGHEADD		
<i>Tribolium</i>	DopEcR	285	VWLGILNSFWKSMLLTA-SLPQFLRLAFVFCMTCCKRYKGRLOAEELIGHEADD		
			TM7		

B

Ef1alpha

DopEcR

Dop3

Dop2

Dop1

Central brain
SOG

Lamina + Medulla + Retina
CC + CA

Thoracic muscles
Salivary gland
Midgut

Malpighian tubules
Testicle
Ovary

RT(+)

RT(-)

A

<i>Gryllus</i> DAT	1	MAP--DGAARRASPTTPVGAALVRGAGGGGGGGRGGGDDVVERETWGWKVDVFLLSVIGFAVDLANVVRFPYLYCKNKGGAFLVPYICIMLVVGGIPLPYM		
<i>Drosophila</i> DAT	1	MSPTGHIISKST-----PTTHDNDNNISDERETWGWKVDVFLLSVIGFAVDLANVVRFPYLYCKNKGGAFLVPYICIMLVVGGIPLPYM		
<i>Bombyx</i> DAT	1	-----MALK--TPT-----GVVGERETWGWKVDVFLLSVIGFAVDLANVVRFPYLYCKNKGGAFLVPYICIMLVVGGIPLPYM		
<i>Apis</i> DAT	1	MS-----SR-----VVK-----NAPKL--NCSVDGRETWGWKVDVFLLSVIGFAVDLANVVRFPYLYCKNKGGAFLVPYICIMLVVGGIPLPYM		
			▲	
<i>Gryllus</i> DAT	94	ELALGGQFRRKGAITCWGRIVPLFKGIGYAVVLIAFVDFYFNVIIAWALRRFFASFTSSLPWTSCKGNHWNTPQCRDLVD-----	TM1	TM2
<i>Drosophila</i> DAT	84	ELALGGQFRRKGAITCWGRIVPLFKGIGYAVVLIAFVDFYFNVIIAWALRRFFASFTNSLPWTSCKNHNWNTNCRPFESQNSARVPVIGNISDQ		
<i>Bombyx</i> DAT	71	ELALGGQFRRKGAITCWGRIVPLFKGIGYAVVLIAFVDFYFNVIIAWALRRFFASFTMLPWTNCDNWNTPACRPFES-----I		
<i>Apis</i> DAT	76	ELALGGQFRRKGAITCWGRIVPLFKGIGYAVVLIAFVDFYFNVIIAWALRRFFASFTSLPWTNCDNWNTPACRPFES-----I		
			TM3	
<i>Gryllus</i> DAT	175	---GCTVSAAMADRAASDNGSH-----VTSAASEYFNRAILELHESAGLHDLGIVKWDMALCLLAVYLYCYFSLWKIGISTSQKVVNFTA		
<i>Drosophila</i> DAT	178	AMCNQSLIYNEFYMGSSST---DSAVGHVEGFQSAASEYFNRAILELNRSEGHDGAIKWDMALCLLIVYLYCYFSLWKIGISTSQKVVNFTA		
<i>Bombyx</i> DAT	151	NDV--NRTRIRNTSIA--SLGIAPATP-----VTSAASEYFNRAILELQCEGLHDLGSKVWDMALCLLAVYLYCYFSLWKIGISTSQKVVNFTA		
<i>Apis</i> DAT	169	NP--GNSSIVVEDRIDRN--SNQGYN--ST-----WYTSAAQCEYFNRAILELHESEGLHDLGIKWDALCLLIVYLYCYFSLWKIGISTSQKVVNFTA		AA
			TM4	
<i>Gryllus</i> DAT	256	LFPIYVLLILLVRGVTLPGSAGITRYILSPNFSAISRAEVVWVDAATQVFFSLGPGFGVLLAYASYNKYHNNVYKDALLTSVINSATSFVAGFVIF		
<i>Drosophila</i> DAT	270	LFPIYVLLILLVRGVTLPGSAGITRYILSPNFSAISRAEVVWVDAATQVFFSLGPGFGVLLAYASYNKYHNNVYKDALLTSVINSATSFVAGFVIF		
<i>Bombyx</i> DAT	236	LFPIYVLLILLVRGVTLPGSAGITRYILSPNFSAISRAEVVWVDAATQVFFSLGPGFGVLLAYASYNKYHNNVYKDALLTSVINSATSFVAGFVIF		
<i>Apis</i> DAT	255	LFPIYVLLILLVRGVTLPGSAGITRYILSPNFSAISRAEVVWVDAATQVFFSLGPGFGVLLAYASYNKYHNNVYKDALLTSVINSATSFVAGFVIF		
			TM5	TM6
<i>Gryllus</i> DAT	351	SVLGYNMAHSG--KPIQEVATEGPGLVFVYVPAAIATMPGISTFWALIFFMMLLTGLDSSFGGSEAITALSDEFP--IICNNREIFVACLFSLYFL		TM7
<i>Drosophila</i> DAT	365	SVLGYNMAHSG--KPIQEVATEGPGLVFVYVPAAIATMPGISTFWALIFFMMLLTGLDSSFGGSEAITALSDEFP--IICNNREIFVACLFSLYFL		
<i>Bombyx</i> DAT	331	SVLGYNMAHSG--RDVQDVATEGPGLVFVYVPAAIATMPGISTFWALIFFMMLLTGLDSSFGGSEAITALSDEFP--IICNNREIFVACLFSLYFL		
<i>Apis</i> DAT	350	SVLGYNMAHSG--KSIQDVATEGPGLVFVYVPAAIATMPGISTFWALIFFMMLLTGLDSSFGGSEAITALSDEFP--IICNNREIFVACLFSLYFL		
			TM8	TM9
<i>Gryllus</i> DAT	444	VGLASCTGGGYFFHLLDRIAAGYSMLFAVLFEAIAVSWIYGRFCADIIDMIGFSPGYNWRVCWRFVAPFIFLFIIVYGLIGYEPLTYEDVYV		
<i>Drosophila</i> DAT	458	VGLASCTGGGYFFHLLDRIAAGYSMLFAVLFEAIAVSWIYGRFCADIIDMIGFSPGYNWRVCWRFVAPFIFLFIIVYGLIGYEPLTYEDVYV		
<i>Bombyx</i> DAT	424	VGLASCTGGGYFFHLLDRIAAGYSMLFAVLFEAIAVSWIYGRFCADIIDMIGFSPGYNWRVCWRFVAPFIFLFIIVYGLIGYEPLTYEDVYV		
<i>Apis</i> DAT	443	VGLASCTGGGYFFHLLDRIAAGYSMLFAVLFEAIAVSWIYGRFCADIIDMIGFSPGYNWRVCWRFVAPFIFLFIIVYGLIGYEPLTYEDVYV		
			TM10	TM11
<i>Gryllus</i> DAT	539	PGWANLIGWLIAAGSSIMIPGMAAVKLLTTPGCFORLKYLTTPWRDQ--IFA--INGVQ--TDSQORLRLA--SSPQA--DDV-----		
<i>Drosophila</i> DAT	553	PGWANALGWGIAGSSVVMIPAVAILKLSPPGSLRORFILTTPWRD--QOSMAMVLNGVT--TEVVVRL--TDTEFAKSPVDV-----		
<i>Bombyx</i> DAT	519	PGWANALGWGIAGSSVMQIPVAIYKLLTTPGSLRORLRLTTPVDSER--NGTVHNGMIVSISGGGVRLMSAVQ--PTTPQPPGVANIPAAAPAL--		
<i>Apis</i> DAT	538	PGWANLIGWLIAAGSSIAMIPGMAYKLLTTPGCFORLKYLTTPWRD--QORNA--DFSSVANGAVRRSFTA-----DQDLNLTKEQQLNLTKEQGE		
			TM12	
<i>Gryllus</i> DAT		-----		
<i>Drosophila</i> DAT		-----		
<i>Bombyx</i> DAT	613	ASSPALV-----		
<i>Apis</i> DAT	626	VMIQSRNVNGDPPPEPV		

B

Ef1 alpha

DAT

RT(+)

RT(-)

Central brain
SOG

SOG

Lamina + Medulla + Retina

CC-0

Thoracic muscles

Salivary glands

Salivary gland
Midgut

Midgut

Malpighian tubules
Testicle
Ovary

Testicle

Ovary