



Title	Oxidized DJ-1 Inhibits p53 by Sequestering p53 from Promoters in a DNA-Binding Affinity-Dependent Manner
Author(s)	Kato, Izumi; Maita, Hiroshi; Takahashi-Niki, Kazuko; Saito, Yoshiro; Noguchi, Noriko; Iguchi-Ariga, Sanae M. M.; Ariga, Hiroyoshi
Citation	Molecular and Cellular Biology, 33(2), 340-359 https://doi.org/10.1128/MCB.01350-12
Issue Date	2013-01
Doc URL	http://hdl.handle.net/2115/52746
Type	article (author version)
File Information	35118_1_merged_1351241601.pdf



[Instructions for use](#)

1 **Oxidized DJ-1 inhibits p53 by sequestering p53 from promoters in a**
2 **DNA-binding affinity-dependent manner**

3

4 **Izumi Kato,^{a,b} Hiroshi Maita,^a Kazuko Takahashi-Niki,^a Yoshiro Saito,^c Noriko**
5 **Noguchi,^c Sanae M.M. Iguchi-Ariga,^d and Hiroyoshi Ariga^a**

6

7 Graduate School of Pharmaceutical Sciences, Hokkaido University, Kita-ku, Sapporo 060-0812,
8 Japan,^a Graduate School of Life Science, Hokkaido University, Kita-ku, Sapporo 060-0812, Japan,^b
9 Faculty of Medical and Life Sciences, Doshisha University, Kyotanabe, Kyoto 610-0394, Japan,^c
10 Graduate School of Agriculture, Hokkaido University, Kita-ku, Sapporo 060-8589, Japan,^d

11

12 Running title: Oxidative Stress-dependent Repression of p53 by DJ-1

13

14 Address correspondence to Hiroyoshi Ariga, Graduate School of Pharmaceutical Sciences, Hokkaido
15 University, Kita-ku, Sapporo 060-0812, Japan. Tel.: +81 11 706 3745; Fax: 81 11 706 4988; E-mail:
16 hiro@pharm.hokudai.ac.jp

17

18

19 **ABSTRACT**

20 *DJ-1* is an oncogene and causative gene for familial Parkinson's disease. Although the oxidative
21 status of DJ-1 at cysteine at 106 (C106) is thought to affect all of the activities of DJ-1 and excess
22 oxidation leads to the onset of various diseases, the precise molecular mechanisms underlying the
23 effects of oxidation of DJ-1 on protein-protein interaction of DJ-1 remain unclear. In this study, we
24 found that DJ-1 bound to the DNA-binding region of p53 in an oxidation of C106-dependent manner.
25 Of the p53-target genes, the expression level and promoter activity of the DUSP1 gene, but not those
26 of the p21 gene, were increased in H₂O₂-treated DJ-1 (-/-) cells and were decreased in wild-type
27 DJ-1- but not C106S DJ-1-transfected H1299 cells through sequestration of p53 from the DUSP1
28 promoter by DJ-1. DUSP1 down-regulated by oxidized DJ-1 activated ERK and decreased apoptosis.
29 DUSP1 and p21 promoters harbor non-consensus and consensus p53-recognition sequences,
30 respectively, which have low affinity and high affinity to p53. However, DJ-1 inhibited p21 promoter
31 activity brought by p53 mutants harboring low DNA-binding affinity but not by wild-type p53.
32 These results indicate that DJ-1 inhibits the expression of p53-target genes in p53 DNA-binding
33 affinity- and oxidation of DJ-1 C106-dependent manners.

34

35 **INTRODUCTION**

36 *DJ-1* was identified by us as a novel oncogene that induces anchorage-independent growth of
37 fibroblasts cooperatively with activated *ras* (31) and was later found to be a causative gene for a
38 familial form of Parkinsons disease, *Park7* (2). DJ-1 has 3 cysteines located at amino acid numbers
39 46, 53 and 106 (C46, C53 and C106, respectively). Of the 3 cysteines, C106 is first oxidized as SOH,
40 SO₂H and SO₃H forms, and excessive oxidation then causes oxidation of C46 and C53 (21, 45). The
41 C106S mutant of DJ-1, which is a substitution mutant of DJ-1 at amino acid number 106 from
42 cysteine to serine, possesses no or little protective activity against neuronal cell death induced by
43 oxidative stress (5, 19, 27, 44, 45), and abnormally oxidized forms of DJ-1 were observed in patients

44 with sporadic forms of Parkinsons disease (1). From these points, C106 is the most important
45 cysteine to maintain DJ-1's function. Although oxidative status of DJ-1 affects DJ-1's activity toward
46 cells and disease, the precise molecular mechanisms remain unclear.

47 DJ-1 binds to various factors, including transcriptional factors such as androgen receptor (32, 46),
48 p53 (11, 41), polypyrimidine tract-binding protein-associated splicing factor (PSF) (51) and Keap1,
49 an inhibitor for nuclear factor erythroid-2 related factor2 (Nrf2) (8). However, it is not known how
50 DJ-1 chooses its suitable binding protein(s) during the course of oxidative stress.

51 p53 is a tumor suppressor protein that activates transcriptional programs under various types of
52 cellular stress, including oxidative stress. It is, however, not clear how p53 determines a point
53 leading to cell cycle arrest and to apoptosis. Recent reports suggest that activation of specific
54 promoters by p53 is achieved through its interaction with heterologous transcription factors such as
55 Hzf, human cellular apoptosis susceptibility (hCAS)/CSE1L and ankyrin-repeat, SH3-domain and
56 proline-rich-region containing protein (ASPP) family proteins (9, 36, 47). DJ-1 directly binds to p53
57 to restore p53 transcriptional activity by inhibiting sumoylation of p53 through interaction of DJ-1
58 with Topors/p53BP3, a SUMO-1 ligase for p53 (50). Sumoylation of DJ-1 itself is necessary for
59 DJ-1 to localize from the cytoplasm to nucleus (40) and DJ-1 is a negative regulator for sumoylation
60 (10). Moreover, DJ-1 decreases Bax expression through repressing p53 transcriptional activity by an
61 unknown mechanism (11). Although DJ-1 regulates p53 transcriptional activity through interaction
62 with a SUMO-1 ligase of p53 and regulates its location, it is still unclear whether oxidative status of
63 DJ-1 affects p53 activity. We hypothesized that oxidative status of DJ-1 contributes to its binding
64 activity to various proteins to regulate their functions.

65 In this study, we found that DJ-1 bound to the DNA-binding region of p53 in an oxidative status of
66 DJ-1-dependent manner and that the oxidation of C106 was essential for DJ-1 binding to p53,
67 resulting in altering DNA-binding affinity of p53. Furthermore, DJ-1 repressed transcriptional
68 activity of p53 in a p53 DNA-binding affinity-dependent manner.

69

70 MATERIALS AND METHODS

71 **Cell culture and mice.** HEK293T, A549, H1299 and mouse embryonic fibroblast cells were cultured
72 at 37°C in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% calf serum.
73 DJ-1-heterozygous knockout mice were kindly provided by J. Shen (13), and DJ-1-homozygous
74 knockout mice (DJ-1(-/-)) and wild-type mice with the same background (DJ-1(+/+)) were obtained.
75 Newborn mice with genotypes of DJ-1(-/-) and DJ-1 (+/+) at 1 day after birth were cut with scissors,
76 digested with trypsin, and seeded on a 10-cm dish in DMEM with 10% calf serum. These cells were
77 used as mouse DJ-1 (-/-) and DJ-1 (+/+) cells. All animal experiments were carried out in accordance
78 with the National Institutes of Health Guide for the Care and Use of Laboratory Animals, and the
79 protocols were approved by the Committee for Animal Research at Hokkaido University (the permit
80 number 08-0468).

81

82 **RT-PCR and real-time PCR.** Nucleotide sequences of primers used for RT-PCR were as follows:

83 β -actin sense: 5'-TCCTCCCTGGAGAAGAGCTA-3', β -actin as:
84 5'-CCAGACAGCACTGTGTTGGC-3', mouse p21 sense: 5'-CCGTGGACAGTGAGCAGTTG-3',
85 mouse p21 as2: 5'-GAAGACCAATCTGCGCTTGG-3', mouse NOXA sense:
86 5'-GAACGCGCCAGTGAACCCAA-3', mouse NOXA as:5'-CTTTGTCTCCAATCCTCCGG-3',
87 mouse PUMA sense: 5'-TCCTCAGCCCTCCCTGTAC-3', mouse PUMA as:
88 5'-CCATTTCTGGGGCTCCAGGA-3', mouse DUSP1 sense2:
89 5'-CAGCTCCTGGTTCAACGAGG-3', and mouse DUSP1 as:
90 5'-GCAGCTTGAGAGGTGGTGAT-3'. Nucleotide sequences of primers used for real-time PCR
91 were as follows: mACTB 192-211F: 5'-CCTAGGCACCAGGGTGTGAT-3', mACTB 734-753R:
92 5'-GCTCGAAGTCTAGAGCAACA-3', mACTB F-real-time:
93 5'-CCCTAAGGCCAACCGTGAAA-3', mACTB R-real-time:

94 5'-ACGACCAGAGGCATACAGGGA-3', mDUSP1 F-real-time:
 95 5'-CCTGGTTCAACGAGGCTATTG-3', mDUSP1 R-real-time:
 96 5'-CCAGCTTTACCCGGTTAGTCC-3', human DUSP1 sense3:
 97 5'-GTATCACGCTTCCCGCAAGG-3', human DUSP1 as5:
 98 5'-CAAACACCCTTCCTCCAGCATTC-3', human Actin sense3:
 99 5'-CGGCTGAGGAGTGGCTGG-3', and human Actin as4: 5'-CCAGCCGAGACACGGCAT-3'.

100 After mouse primary or H1299 cells had been treated with 300 μ M H₂O₂ for 0.25-4 hrs or 0.5 hrs,
 101 total RNAs were prepared and subjected to semi-quantitative RT-PCR and quantitative RT-PCR
 102 (real-time PCR) analyses. After reactions, PCR products were extracted, separated on 2% agarose
 103 gels, and stained with ethidium bromide. PCR conditions for RT-PCR were as follows: 1 min at
 104 94°C, 30 sec at 94°C, 30 sec at 58°C and 22 cycles of 1 min at 72°C for β -actin; 1 min at 94°C, 30
 105 sec at 94°C, 30 sec at 58°C and 25 cycles of 1 min at 72°C for p21; 1 min at 94°C, 30 sec at 94°C, 30
 106 sec at 58°C and 28 cycles of 1 min at 72°C for NOXA; 1 min at 94°C, 30 sec at 94°C, 30 sec at 58°C
 107 and 34 cycles of 1 min at 72°C for PUMA; and 1 min at 94°C, 30 sec at 94°C, 30 sec at 58°C and 28
 108 cycles of 1 min at 72°C for DUSP1. PCR conditions for real-time PCR were as follows: 10 sec at
 109 95°C, 5 sec at 95°C and 44 cycles of 20 sec at 60°C for β -actin; and 10 sec at 95°C, 5 sec at 95°C
 110 and 44 cycles of 20 sec at 60°C for DUSP1.

111

112 **Luciferase assay.** pGL4.10-hDUSP1 and pGL3-hp21 were digested with KpnI and HindIII and each
 113 resultant fragment was inserted into KpnI and HindIII sites of pGL4.12[luc2CP] (Promega, Madison,
 114 WI). H1299 cells in 6-well dishes were transfected with pGL4.12-hDUSP1 or pGL4.12-hp21
 115 together with pcDNA3-FLAG-p53 and pEF-DJ-1-HA or pEF-DJ-1 C106S-HA by Lipofectamine
 116 2000 (Invitrogen, Carlsbad, CA) as described in the manufacturer's protocol. pActin- β -galactosidase
 117 was also cotransfected with plasmids. At 24 hrs after transfection, cells were treated with or without
 118 300 μ M H₂O₂ or with 10 μ g/mL cycloheximide for 30 min. Whole cell extracts were then prepared

119 by addition of a Triton X-100-containing solution from the Pica gene kit (Wako Pure Chemicals,
120 Osaka, Japan) to cells. About a one-fifth volume of the extract was used for the β -galactosidase assay
121 to normalize transfection efficiencies as described previously (14), and the luciferase activity due to
122 the reporter plasmid was determined using a luminometer (Luminocounter Lumat LB 9507, EG & G
123 Berthold, Bad Wildbad, Germany). Proteins in aliquots of the cell extract were analyzed by Western
124 blotting with an anti-FLAG antibody (M2, Sigma, St. Louis, MO, USA) and anti-HA antibody
125 (1:2000, MBL, Nagoya Japan) and visualized as described in the “Western blotting and antibody”
126 section. The same experiments were repeated at least three times.

127

128 **Chromatin immunoprecipitation (ChIP) assay.** ChIP assays using cultured A549 cells were
129 performed according to the protocol of the ChIP assay kit (Millipore, Billerica, MA). Briefly, after
130 proteins had been cross-linked with DNA, cell pellets were resuspended in an SDS-lysis buffer and
131 sonicated on ice using a sonicator (UR-20P, Tomy, Tokyo, Japan) 4 times for 15 sec each time.
132 Genomic DNA was sheared to 300 to 1200 base pairs in length. Chromatin solution was
133 preincubated with salmon sperm DNA and protein A-agarose and incubated with species-matched
134 IgG or with specific antibodies overnight at 4°C. DNA fragments immunoprecipitated were then
135 used as templates for PCR with Ex taq (TaKaRa Bio, Kyoto, Japan) and reacted for 60 sec at 94°C,
136 60 sec at 94°C, 30 sec at 58°C and 35 cycles of 30 sec at 72°C. Nucleotide sequences of
137 oligonucleotides used for PCR primers were as follows: hDUSP1 sense:
138 5'-AAGAGCAGGCCGGACAGC-3', hDUSP1 as: 5'-GAGCGCGTTTATATGCGGC-3', hDUSP1
139 sense 3: 5'-CCCAATCCCTCTCCCACTAG-3', hDUSP1 as3:
140 5'-CAGAGCCGCTAAAATGGGCA-3', hp21 sense: 5'-TGCTGCTCCACCGCACTC-3', hp21 as:
141 5'-GAAAACAGGCAGCCCAAGGAC-3', hp21 sense3: 5'-CTATCAGCTGCCTCGGGG-3' and
142 hp21 as3: 5'-GGCGCCCAAGTTCCTAAC. PCR products were separated on a 2% agarose gel and
143 stained with ethidium bromide. Reverse images of black and white staining are shown.

144

145 **Western blotting and antibodies.** Proteins were extracted from cells with a buffer containing 150
146 mM NaCl, 5 mM EDTA, 50 mM Tris (pH 7.5) and 0.5% NP-40, loaded on 12% sodium dodecyl
147 sulfate (SDS)-polyacrylamide gels, and subjected to Western blotting. Antibodies used in this study
148 were as follows: anti-HA (1:2000, MBL, Nagoya Japan), anti-FLAG F7425 (1:1000, Sigma), anti-T7
149 (1:1000, Novagen, Madison, WI), anti-p53 (1:1000, DO-1, Santa Cruz Biotechnology, California,
150 CA), anti-phospho p53(serine6) (1:1000, Cell Signaling, Danvers, MA), anti-phospho p53(serine9)
151 (1:1000, Cell Signaling), anti-phospho p53(serine15) (1:1000, Cell Signaling), anti-phospho
152 p53(serine20) (1:1000, Cell Signaling), anti-phospho p53(serine37) (1:1000, Cell Signaling),
153 anti-phospho p53(serine46) (1:1000, Cell Signaling), anti-phospho p53(serine392) (1:1000, Cell
154 Signaling), anti-actin (1:4000, Chemicon, Temecula, CA), anti-phospho ERK1/2 (1:1000, Cell
155 Signaling), anti-ERK1/2 (1:1000, Santa Cruz Biotechnology, California, CA), anti-p53 (1:1000,
156 Pab240, Santa Cruz Biotechnology), anti-MKP1(1:500, C-19 , Santa Cruz Biotechnology), rat anti-
157 DJ-1 monoclonal (1:1000), anti-DJ-1 polyclonal (1:4000), mouse anti-DJ-1 monoclonal (1: 4000,
158 3E8, MBL) and anti-oxidized DJ-1 (1:1000) antibodies. Rabbit anti-DJ-1, rat anti-DJ-1 and
159 anti-oxidized DJ-1 antibodies were established by us as described previously (31, 35). After
160 membranes had been reacted with primary antibodies, they were reacted with Alexa Fluor
161 680-conjugated anti-mouse (Molecular Probes, Eugene, OR), Alexa Fluor 680-conjugated anti-rabbit
162 (Molecular Probes), IRDye 800-conjugated anti-mouse antibody (Rockland, Philadelphia, PA) or
163 IRDye 800-conjugated anti-rabbit antibody (Rockland) and visualized by using an infrared imaging
164 system (Odyssey, LI-COR, Lincoln, NE).

165

166 **Pull-down assay.** ³⁵S-labeled p53 was synthesized in vitro using reticulocyte lysate of the TNT
167 transcription-translation coupled system (Promega). Labeled proteins were reacted with GST or
168 GST-wild-type DJ-1 or GST-C106S DJ-1 expressed in and prepared from *Escherichia coli* in a

169 G-buffer containing 150 mM NaCl, 5 mM EDTA and 50 mM Tris (pH 7.5), 0.05% bovine serum
170 albumin and 0.1% Nonidet P-40 for 2 hrs at 4°C, mixed with glutathione sepharose beads, and
171 centrifuged. After washing pellets with the wash buffer containing 150 mM NaCl, 10 mM Tris-HCl
172 (pH 7.5), 0.1% Nonidet P-40, Laemmli buffer was added to pellets. Pellets were then heated at 97°C
173 for 5 min, separated on a 12% polyacrylamide gel containing SDS, and visualized by CBB staining
174 and by fluorography.

175 HEK293T or H1299 cells were transfected with expression vectors for wild-type DJ-1-HA or
176 C106S DJ-1-HA and One-STrEP-p53 and treated with H₂O₂. Proteins in cell extracts were then
177 subjected to pull-down assays with a Strep-Tactin sepharose beads according to the supplier's
178 protocol (IBA, Göttingen, Germany), and co-precipitated DJ-1 was detected by Western blotting
179 with anti-HA and anti-oxidized-DJ-1 antibodies.

180

181 **Co-immunoprecipitation assay.** Proteins were extracted from cultured cells with or without 300
182 μM H₂O₂ for 30 min by the procedure described previously (16). Proteins were immunoprecipitated
183 with a rabbit anti-DJ-1 antibody (1:500, MBL) or normal rabbit IgG, and precipitates were analyzed
184 by Western blotting with anti-p53 (1:1000, Santa Cruz Biotechnology) or mouse anti-DJ-1 antibody
185 (1:1000, 3E8, MBL). Proteins on membranes were visualized as described above.

186 HEK293T cells were transiently transfected with expression vectors for FLAG-p53 and DJ-1-HA
187 by the calcium phosphate method and were lysed by treatment of cells with or without 1 mM H₂O₂
188 for 30 min. Proteins were then immunoprecipitated by an anti-FLAG antibody and precipitates were
189 detected by anti-HA antibody or anti-DJ-1 antibody.

190

191 **Isoelectric focusing.** Cells were treated with 1 mM H₂O₂ for 30 min, and cell extracts were prepared
192 by the procedure described previously (16). Proteins in the extracts were then separated on pH 5–8
193 ranges of an isoelectric focusing gel or on 12.5% polyacrylamide gel containing SDS, transferred

194 onto nitrocellulose membranes, and reacted with an anti-HA antibody.

195

196 **ELISA assay.** GST or GST-p53 purified from E coli was loaded on ELISA plates (BD) at 4°C
197 overnight. After plates had been blocked with 0.25 x block A and washed with 0.1% Tween 20-PBS,
198 wild-type or C106S DJ-1 was added to plates and incubated for 1 hr at 4°C. Plates were then reacted
199 with an anti-rat DJ-1 antibody (1:1000) for 1 hr at 37°C and reacted with an ABTS solution, and
200 absorbance of each well was measured using a plate reader (BIO-RAD).

201

202 **Immunofluorescence.** A549 cells were treated with or without 1 mM H₂O₂ for 30 min. Cells were
203 then fixed with acetone/methanol and reacted with an anti-p53 or anti-DJ-1 antibody, and
204 immunofluorescence images of proteins were detected by a fluorescein isothiocyanate- or
205 rhodamine-conjugated secondary antibody, respectively. Nuclei were stained with DAPI.

206

207 **Cell cycle FACS analysis.** Mouse DJ-1 (+/+) and DJ-1 (-/-) cells were transfected with 50 nM
208 AllStars negative control siRNA (QIAGEN), DUSP1 siRNA-1 (Scramble siRNA cocktail that
209 contains 4 sets of siRNA, Thermo scientific, 040753-01-0005, San Jose, CA) and DUSP1 siRNA-2
210 using Lipofectamine 2000 (Invitrogen, Carlsbad, CA). Nucleotide sequences of DUSP1 siRNA-2 are
211 followed: DUSP1-2 sense: 5'-GGAUGCAGCUCCUGUAGUATT-3' and DUSP1-2 antisense:
212 5'-UACUACAGGAGCUGCAUCCTT-3'. At 48 hrs after transfection, the cells were starved for 6
213 hrs and then treated with or without H₂O₂. Cells were harvested at 40 hrs after the second H₂O₂
214 treatment, washed once with PBS, and fixed with 90% ethanol overnight. The cells were then treated
215 with 1 mg/ml RNase for 30 min at 37°C, washed once with PBS, and suspended in 300 µl of PBS
216 containing 50 µg/ml of propidium iodide (Sigma-Aldrich). After leaving for 1 hour at room
217 temperature, the cells were subjected to FACS analysis using a calibur flow cytometer
218 (Becton-Dickinson), and data obtained were analyzed using CellQuest software and ModFit

219 software.

220

221 **Statistical analyses.** Data are expressed as means \pm S.E. Statistical analyses were performed using
222 analysis of variance (one-way ANOVA) followed by unpaired Student's *t*-test. For comparison of
223 multiple samples, the Tukey-Kramer test was used.

224

225 **RESULTS**

226 **Oxidative stress enhances DJ-1-binding to p53.**

227 To address the effects of DJ-1 oxidation on interaction of DJ-1 with its target proteins, including p53,
228 FLAG-tagged proteins were co-transfected with DJ-1-HA into HEK293T cells, and the cells were
229 treated with 1 mM H₂O₂ for 30 min. Co-immunoprecipitation followed by Western blotting analyses
230 showed that p53 and Daxx were strongly bound to DJ-1 in H₂O₂-treated cells (Fig. 1 and data not
231 shown). We therefore focused on interaction of DJ-1 with p53 under the condition of oxidative stress
232 in this study.

233 The same results as those shown in Fig. 1A were obtained when FLAG-DJ-1 was cotransfected
234 with T7-p53 into HEK293T cells (Fig. 1B). In this case, both phosphorylated and unphosphorylated
235 T7-p53 was bound to DJ-1. Proteins from H₂O₂-treated A549 cells and H1299 cells in which p53 is
236 not expressed were immunoprecipitated with an anti-DJ-1 antibody and precipitates were analyzed
237 by Western blotting with an anti-p53 antibody. The results showed that endogenous p53 bound to
238 endogenous DJ-1 under normal culture conditions and that the amount of precipitated p53 was
239 enhanced after treatment of A549 cells, but not H1299 cells, with H₂O₂ (Figs. 1C and D,
240 respectively), suggesting that oxidative status of DJ-1 modulates p53-binding activity of DJ-1.

241 DJ-1 is localized in the cytoplasm, nucleus and mitochondria in cells and its localization is
242 changed by oxidative stress (24). Oxidative stress might therefore alter localization of p53 and
243 interaction of p53 with DJ-1. When A549 cells were treated with or without 300 μ M H₂O₂ for 30 min

244 and localization of proteins were analyzed by an immunofluorescence technique, DJ-1 and p53 were
245 located both in the cytoplasm and nucleus and in the nucleus, respectively, and both proteins were
246 co-localized in the nucleus in cells before and after H₂O₂ treatment (Figs.1E and 1F), indicating that
247 oxidative stress enhances binding activity of DJ-1 to p53 without affecting localization of DJ-1.

248

249 **Cysteine 106 of DJ-1 is essential for binding of DJ-1 to p53 under an oxidative stress condition.**

250 To examine whether translational modification both of p53 and DJ-1 induced by oxidative stress
251 contributes to H₂O₂-dependent increase of p53-DJ-1 complex formation, HEK293T cells were
252 exposed to 1 mM H₂O₂ for 30 min, and phosphorylation of p53 was analyzed by Western blotting
253 with several anti-phosphorylated p53 antibodies. As shown in Fig. 2A, enhanced phosphorylation of
254 p53 at serine residues 15 and 20, but not at serine residues 6, 9, 37, 46 and 392, was observed. When
255 HEK293T cells were transfected with serine mutants of p53 in which serine was changed to alanine
256 and exposed to 1 mM H₂O₂ for 30 min, expected phosphorylation of serines 15 and 20 of FLAG-p53
257 mutants was not observed by single and double substitution (Fig. 2B). HEK293T cells were then
258 co-transfected with wild-type and serine mutants of FLAG-p53 and DJ-1-HA and exposed to 1 mM
259 H₂O₂ for 30 min at 48 hrs after transfection. Co-immunoprecipitation experiments showed that of the
260 three serine mutants and wild-type p53, no or only a slight difference in binding activity of
261 FLAG-p53 to DJ-1-HA was observed under the oxidative stress condition (Fig. 2C), indicating that
262 phosphorylation of p53 does not contribute to H₂O₂-dependent enhancement of DJ-1-p53 interaction.

263 Oxidative status of cysteine106 (C106) of DJ-1 is critical for all of the functions of DJ-1.
264 HEK293T cells were co-transfected with wild-type or C106S mutant of DJ-1-HA and FLAG-p53
265 and exposed to H₂O₂ under the same conditions as those described above. The oxidation levels of
266 wild-type and C106S DJ-1-HA were first examined by using electron focusing gels. While C106S
267 DJ-1-HA had no shifted band, approximately 50% of wild-type DJ-1-HA was shifted to a more
268 acidic point in H₂O₂-treated cells (Fig. 3A). Furthermore, when HEK293T cells were treated with

269 H₂O₂ and with UV or doxorubicin, other stressors activating p53, and proteins were analyzed with
270 anti-DJ-1 and anti-C106 oxidated-DJ-1 antibodies, H₂O₂ exposure clearly enhanced C106 oxidation
271 of DJ-1 (Fig. 3B), suggesting that a shifted band observed in a sample of wild-type DJ-1-HA was
272 derived from C106. The DJ-1-p53 complex was then analyzed by immunoprecipitation using an
273 anti-FLAG antibody followed by Western blotting with an anti-DJ-1 antibody. The results showed
274 that while wild-type DJ-1 was strongly bound to p53 in H₂O₂-treated cells, the C106S mutant of
275 DJ-1 had lost p53-binding activity (Fig. 3C). Pull-down experiments were then carried out using
276 GST-wild-type DJ-1, GST-C106S DJ-1 and GST with ³⁵S-labeled p53 in the presence or absence of
277 H₂O₂ and reducing agent DTT. Recombinant DJ-1 purified from *E. coli* is comprised of a mixture
278 containing reduced and oxidized forms of C106, and the amounts of SOH and SO₂H forms of C106
279 are about 10~20% of total forms of C106 (data not shown). Doublet bands of p53 in gels must be
280 unphosphorylated and phosphorylated p53. As in the case of DJ-1 binding to p53 in 293T cells, H₂O₂
281 -treated wild-type DJ-1 was bound to p53 more strongly than did H₂O₂ -non-treated wild-type DJ-1,
282 and C106S mutant of DJ-1 had lost the enhancement of p53-binding activity (Fig. 3D). The
283 enhancement of binding activity of DJ-1 to p53 was diminished by addition of DTT, indicating that
284 DJ-1 directly interacts with p53 and that oxidation of C106 is essential for this interaction.

285 To directly examine whether C106-oxidized DJ-1 binds to p53, two experiments were carried out.
286 First, HEK293T cells were co-transfected with FLAG-wild-type or C106S mutant of DJ-1 and
287 T7-p53 and exposed to H₂O₂. Proteins were subjected to co-immunoprecipitation and Western
288 blotting analyses with an anti-C106-oxidized DJ-1 antibody (Fig. 3E). Second, HEK293T and H1299
289 cells were transfected with wild-type or C106S DJ-1-HA and One-Strep-p53 and treated with H₂O₂.
290 Proteins were pulled-down using Strep-Tactin sepharose and precipitates were analyzed by Western
291 blotting with the anti-oxidized DJ-1 antibody (Figs. 3F and 3G, respectively). The results showed that
292 oxidized wild-type DJ-1, but not C106S DJ-1, bound to p53 under the oxidative stress condition.

293

294 **DJ-1 down-regulates DUSP1 expression under an oxidative stress condition.**

295 To address the role of increased formation of DJ-1-p53 complex after treatment of cells with H₂O₂,
296 the expression levels of p53-target genes in mouse primary cells were examined by semi-quantitative
297 RT-PCR. As shown in Fig. 4A, the expression level of DUSP1 mRNA was increased at a peak of 30
298 min after H₂O₂ treatment, while transcription of other p53-target genes such as p21, NOXA and
299 PUMA was induced 1-2 hrs. The other stresses such as UV exposure and doxorubicin treatment, on
300 the other hand, did not alter the DUSP1 gene expression level, while the expression levels of Noxa
301 and p21 were increased at 0.5-6 hrs after UV exposure and doxorubicin treatment (Figs. 5A-5D),
302 suggesting that the DUSP1 gene is a primary target of p53 against oxidative stress.

303 DUSP1, a mitogen-activated protein kinase phosphatase, regulates the apoptosis signaling
304 pathway through dephosphorylating ERK and is known to be only the p53-target gene that
305 specifically responds to H₂O₂ treatment (25, 43, and Figs. 4A, 5A and 5B). Since the amount of
306 p53-DJ-1 complex was also increased at the peak of 30 min after H₂O₂ treatment in HEK293T cells
307 transfected with DJ-1-HA and FLAG-p53 and in A549 cells (Figs. 4B and 4C) and p53-DJ-1
308 complex was not detected at 1-4 hrs after H₂O₂ treatment (Fig. 4B right), we focused on the effect of
309 DJ-1 on p53-dependent DUSP1 expression.

310 The expression levels of DUSP1 and p21 mRNAs were examined using primary cells derived
311 from DJ-1-knock out (DJ-1 (-/-)) and DJ-1 (+/+) mice. Semi-quantitative RT-PCR and real-time PCR
312 analyses showed that the expression level of DUSP1 mRNA in DJ-1 (+/+) cells was increased at 30
313 min and then decreased at 2 hrs after H₂O₂ treatment and that the level at 30 min was further
314 increased in DJ-1 (-/-) cells (Figs. 4D and 4F). The expression levels of p21 mRNA were, on the
315 other hand, increased at 2 hrs and there was no difference between the expression levels in DJ-1 (-/-)
316 and DJ-1 (+/+) cells (Figs. 4E and 4G). The expression levels of DUSP1 protein were increased at 2
317 hrs in DJ-1 (+/+) cells and further increased in DJ-1 (-/-) cells (Fig. 4H). On the other hand, there
318 was little difference of expression levels of p21 protein at 2 hrs between DJ-1 (-/-) and DJ-1 (+/+)

319 cells (Fig. 4H). When DJ-1 (-/-) and DJ-1 (+/+) cells were transfected with siRNA targeting p53, the
320 expression levels of DUSP1 and p21 were decreased (Fig. 4I), suggesting that transcriptional activity
321 of p53 is still active even in DJ-1 (-/-) cells. Furthermore, no significant differences in expression
322 level and subcellular localization of p53 were observed in DJ-1 (-/-) and DJ-1 (+/+) cells (Figs. 4J
323 and 4K), indicating that up-regulation of DUSP1 expression in DJ-1 (-/-) cells was not due to change
324 of p53 localization.

325

326 **Cysteine 106 of DJ-1 is essential for repression of p53-dependent DUSP1 transcription under**
327 **an oxidative stress condition.**

328 p53 binds to the promoter region of the DUSP1 gene (25). Since luciferase generally used in reporter
329 assays (call conventional luciferase) has a relatively long half-life, it is not suitable for detecting
330 prompt or immediate change of promoter activity. In the case of prompt decrease in promoter activity
331 toward stress response, for instance, luciferase activity obtained is not parallel to promoter activity
332 due to accumulation of already synthesized conventional luciferase. To overcome this problem, we
333 set up an experimental condition using pGL4.12-Luciferase that harbors a modified version of
334 luciferase with a short half-life of ~20 min and using cycloheximide (CHX), an inhibitor for
335 translation. Even when pGL4.12-Luciferase was used, the short half-life of the modified version of
336 luciferase was still not sufficient to decrease the background caused by accumulation of modified
337 luciferase. Since CHX blocks further translation of luciferase, prompt promoter activity is parallel to
338 promoter activity. Twenty-four hrs after cells had been transfected with reporter plasmids, the cells
339 were divided to three sets. One set was a “control” set to measure the background signal and the
340 other two sets were “test” sets either for an H₂O₂-treated or non-treated sample. The “control” set
341 was first treated with CHX for 30 min just before addition of H₂O₂ to cells and then luciferase
342 activity obtained in the “control” set was subtracted as the background signal from that in the “test”
343 set. Furthermore, to measure transcriptional activity of transfected p53 without an effect of

344 endogenous p53, p53-nul H1299 cells were used.

345 Fig. 6A shows histograms of raw data of luciferase activities obtained in the three sets after
346 transfection of the luciferase reporter construct linked to the DUSP1 promoter: p53 strongly
347 activated DUSP1 promoter activity, and treatment of cells with H₂O₂ reduced its activation activity,
348 which is consistent with results reported previously (25). When wild-type or C106S DJ-1 was
349 co-transfected with p53, luciferase activities tended to increase compared to those without p53 both
350 in wild-type DJ-1- and C106S DJ-1-transfected cells, but luciferase activity was lower in wild-type
351 DJ-1-transfected cells than in C106S DJ-1-transfected cells in the presence of CHX (Fig. 6A).
352 Similar expression levels of introduced proteins were confirmed by Western blotting (Fig. 6C). To
353 compare accurate changes among these samples, ratios of luciferase activities in H₂O₂-treated cells to
354 those in non-treated cells were calculated by subtracting the background signal obtained as the
355 “control” set (Fig. 7A). The results showed that in H₂O₂-treated cells, p53 transcriptional activity
356 toward the DUSP1 promoter was significantly reduced by wild-type DJ-1 to 60% of that with p53
357 alone or with C106S DJ-1. No significant change of promoter-less luciferase activity was observed
358 both in wild-type and C106S DJ-1-transfected cells (Fig. 7C). Furthermore, to confirm that DUSP1
359 promoter activity detected by this assay is parallel with the expression level of DUSP1 mRNA,
360 H1299 cells were transfected with FLAG-p53 and wild-type DJ-1-HA or C106S DJ-1-HA.
361 Twenty-four hrs after transfection, cells were treated with H₂O₂ in the absence of CHX. The
362 expression levels of DUSP1 mRNA examined by quantitative RT-PCR (real-time PCR) were
363 decreased by wild-type DJ-1 but not C106S DJ-1 in H₂O₂-treated cells (Fig. 7D). These results
364 suggest that DJ-1 downregulates p53-dependent DUSP1 expression in a C106-dependent manner in
365 H₂O₂-treated cells and that oxidation of C106 is necessary for this activity. When the same assay
366 using the p21 promoter linked to the modified version of the luciferase gene was carried out, there
367 was no significant change in promoter activity after transfection of cells with wild-type and C106S
368 DJ-1 (Figs. 6B, 6D and Fig. 7B), indicating that the H₂O₂-dependent inhibitory effect of DJ-1 on p53

369 transactivation activity is specific to the DUSP1 gene.

370

371 **DJ-1 directly binds to the p53 DNA-binding region.**

372 p53 is comprised of five domains; an N-terminal transactivation domain (TAD) followed by a
373 proline-rich region (PRR), central DNA-binding domain (DBD), tetramerization domain (TET) and
374 extreme C-terminus (CT) (see recent review, 48). To determine DJ-1-binding region of p53,
375 GST-wild-type DJ-1 and GST were reacted with ³⁵S-labeled full-length p53 and three p53-deletion
376 mutants depicted in Fig. 8A. Pull-down experiments showed that DJ-1 bound to full-length p53,
377 p53 Δ CT and p53DBD, but not to p53 Δ DBD and that H₂O₂-treated DJ-1 more strongly bound to
378 full-length p53 and p53 Δ CT than did H₂O₂-non-treated DJ-1 (Fig. 8B), indicating that DJ-1 binds to
379 p53-DBD (Fig. 8B). Furthermore, pull-down (Fig. 8C) and ELISA (Fig. 8D) assays showed that
380 DJ-1 bound to p53-DBD in a C106-dependent manner.

381

382 **DJ-1-p53 complex stably binds to the p21 promoter but not to the DUSP1 promoter.**

383 A previous study showed that p53 binds to a 10-bp perfect palindromic site in the DUSP1 promoter
384 to activate its transcription, though the p53-binding sequence in the DUSP1 promoter is not a
385 sequence similar to the consensus sequence found in the p21 promoter (25). Chromatin
386 immunoprecipitation (ChIP) assays were carried out using DUSP1 and p21 genes. First, H1299 cells
387 were co-transfected with FLAG-p53 and wild-type or C106S DJ-1-HA, treated with H₂O₂ for 30 min
388 and subjected to ChIP assays with an anti-p53 antibody targeting two regions containing the
389 promoter and other regions (downstream region) were amplified by PCR. As shown in Fig. 9A, the
390 anti-p53 antibody precipitated the DUSP1 promoter region but not the downstream region under a
391 non-oxidative stress condition. In H₂O₂ treated-cells, precipitation of the DUSP1 promoter region by
392 the anti-p53 antibody was cancelled by transfection of wild-type DJ-1-HA but not by that of C106S
393 DJ-1-HA. Although the p21 promoter region contains the consensus p53-binding sequence,

394 precipitated p21 promoter region was not cancelled by transfection of wild-type DJ-1-HA and C106S
395 DJ-1-HA both under oxidative and non-oxidative stress conditions (Fig. 9B). The p53 levels
396 precipitated in reactions were at the similar range by Western blotting (Fig. 9C). Furthermore, ChIP
397 assays were carried out using A549 cells. While the anti-p53 antibody precipitated the DUSP1
398 promoter region but not the downstream region in H₂O₂-treated cells, an anti-DJ-1 antibody did not
399 precipitate regions even after more than 50 cycles of amplification by PCR (Fig. 9D). Both anti-p53
400 and anti-DJ-1 antibodies precipitated the p21 promoter region before and after H₂O₂ treatment and
401 the precipitated levels were increased after H₂O₂ treatment (Fig. 9E). Equal levels of precipitated p53
402 and DJ-1 were confirmed (Fig. 9F). These results indicate that binding activity of p53 to the DUSP1
403 promoter is weaker than that to the p21 promoter and that the p53-DJ-1 complex endogenously binds
404 to the p21 promoter but not to the palindromic site in the DUSP1 promoter. It is thought that
405 DJ-1-free p53 binds to the DUSP1 promoter and that DJ-1-p53 complex binds to the p21 promoter
406 (Fig. 9D). Time course analysis of ChIP assays were then carried out using A549 cells. p53 bound to
407 the DUSP1 promoter in H₂O₂-treated cells at peak of 15 min exposure to H₂O₂ and then the level of
408 promoter-bound p53 decreased (Fig. 10A). No binding of DJ-1 to the DUSP1 promoter was observed.
409 The amount of promoter-bound p53 in DJ-1(-/-) cells was larger than that in DJ-1(+/-) (Fig. 10C).
410 The precipitated levels of p53 and DJ-1 were at the similar levels during the course of H₂O₂ exposure
411 (Figs. 10B and 10D). Taken together, these results suggest that p53 binding to the DUSP1 promoter
412 is sequestered by DJ-1 through its binding to p53DBD in a C106-dependent manner under an
413 oxidative stress condition, resulting in suppression of transactivation activity of p53 toward the
414 DUSP1 gene.

415

416 **DJ-1 increases phosphorylation of ERK and reduces apoptosis under an oxidative stress**
417 **condition.**

418 Since DUSP1 dephosphorylates ERK as MAPK phosphatase, we examined whether the

419 dephosphorylation level of ERK is up-regulated in DJ-1 (-/-) cells by the increased level of DUSP1.
420 DJ-1 (-/-) and DJ-1 (+/+) cells were exposed to H₂O₂ twice and the phosphorylation level of ERK1/2
421 was analyzed by Western blotting. Since culture medium was changed before second H₂O₂ treatment,
422 cells were treated with the same concentration of H₂O₂ at respective stimulation. As shown in Figs.
423 11A and 11C, the expression level of DUSP1 was increased 2 hrs after the first H₂O₂ treatment in
424 DJ-1 (-/-) cells but not in DJ-1 (+/+) cells, and the phosphorylation level of ERK1/2 was lower in
425 DJ-1 (-/-) cells than that in DJ-1 (+/+) cells after the second H₂O₂ treatment. The similar decreased
426 curve of ERK1/2 phosphorylation level after second H₂O₂ treatment was obtained both in DJ-1 (-/-)
427 and DJ-1 (+/+) cells, indicating that DJ-1 did not prolong the ERK activation. There were no or little
428 difference in the expression levels of p53 and total ERK1/2 in DJ-1 (-/-) and DJ-1 (+/+) cells
429 regardless of H₂O₂ treatment (Fig. 11C). Furthermore, knockdown of DUSP1 using two different
430 siRNA (siDUSP1-1 and siDUSP1-2) increased ERK phosphorylation in DJ-1 (-/-) cells but not in
431 DJ-1 (+/+) cells (Figs. 11B and 11D), indicating a reverse correlation between the expression levels
432 of DUSP1 and phosphorylated ERK in DJ-1 (-/-) cells under an oxidative stress condition. Since
433 DUSP1 activates the apoptosis pathway in cells after oxidative stress, DJ-1 (-/-) and DJ-1 (+/+) cells
434 were treated twice with H₂O₂ and then treated with propidium iodide, and their apoptosis levels were
435 analyzed by using FACS in which apoptotic cells are observed in the sub-G1 phase. As shown in
436 Figs. 12A and 12B, H₂O₂ treatment induced apoptosis in both DJ-1 (-/-) and DJ-1 (+/+) cells, but the
437 level of apoptosis in DJ-1 (-/-) cells was about 10-times higher than that in DJ-1 (+/+) cells.
438 Furthermore, knockdown of DUSP1 expression in DJ-1 (-/-) cells reduced apoptosis to 60% of that
439 in control siRNA-treated cells (Fig. 12B). There was no difference between the expression levels of
440 p53 in DJ-1 (-/-) and DJ-1 (+/+) cells (Fig. 11D). These results suggest that DJ-1 decreases apoptosis
441 in cells under intermittent oxidative stress condition through downregulation of DUSP1 expression.

442

443 **DJ-1 inhibits p53 activity in a DNA-binding affinity-dependent manner.**

444 We showed that DJ-1 inhibited transcriptional activity of p53 targeting the DUSP1 gene, but not that
445 of p53 targeting the p21 gene, by inhibiting DNA-binding activity of p53 in the DUSP1 promoter but
446 not in the p21 promoter. Since DUSP1 and p21 promoters contain non-consensus and consensus
447 p53-binding sequences and since non-consensus and consensus p53-binding sequences possess low
448 and high binding affinities to p53, respectively, we considered two possible mechanisms by which
449 DJ-1 inhibited p53-binding activity only to the DUSP1 promoter: One is the difference in the
450 p53-binding region between the consensus and non-consensus sequences and the other is the
451 difference in binding affinity of p53 to DNA. To address this point, we used substitution mutants of
452 p53 at amino acid number 181 from arginine to leucine (R181L), to cysteine (R181C) and to proline
453 (R181P), which are linked to tumor development in families with the hereditary Li-Fraumeni or
454 Li-Fraumeni-like cancer susceptibility syndrome (37). DNA-binding activity of these p53 mutants
455 changed in the order of wild-type (WT) > R181L > R181C > R181P (38). Pull-down assays using
456 GST-wild-type DJ-1, GST and ³⁵S-labeled p53 mutants showed that DJ-1 bound to R181L, R181C,
457 R181P and WT p53 with the same or similar affinity (Fig. 13A). Assays using the luciferase reporter
458 gene conjugated to the p21 promoter showed that transcriptional activity of p53 mutants toward the
459 p21 promoter was reduced compared to that of WT p53 in a DNA-binding activity-dependent
460 manner and that the R181P mutant had no transcriptional activity (Fig. 13B). We therefore used
461 R181L and R181C mutants of p53 for further study. Pull-down assays showed that H₂O₂ treatment
462 enhanced DJ-1-binding activity of R181L and R181C p53 (Fig. 13C). H1299 cells were then
463 co-transfected with wild-type DJ-1 or C106S DJ-1 together with R181L or R181C p53 and treated
464 with 300 μM H₂O₂ or with 10 μg/mL cycloheximide for 30 min at 20 hrs after transfection, and their
465 luciferase activities toward the p21 promoter were examined. Luciferase activities and similar
466 expression levels of introduced proteins in transfected cells are shown in Figs.14B-14E, and ratios of
467 luciferase activity in cells treated with or without H₂O₂ were calculated. The results showed that
468 wild-type DJ-1 significantly inhibited transcriptional activities of R181L p53 and R181C p53 toward

469 the p21 promoter in H₂O₂-treated cells to 50% and 40%, respectively, of that in cells without DJ-1 or
470 in cells transfected with C106S DJ-1 (Fig. 14A). Wild-type DJ-1 suppressed p21 promoter activity
471 brought by R181C p53 more effectively than that by R181L p53, suggesting that DJ-1 inhibits p53
472 transcriptional activity in a p53 DNA-binding affinity-depending manner. Furthermore, H1299 cells
473 were co-transfected with FLAG-R181L p53 or -R181C p53 together with wild-type DJ-1-HA or
474 C106S DJ-1-HA, then treated with or without H₂O₂ and subjected to ChIP assays targeting the p21
475 promoter. As shown in Figs. 15A and 15B, the anti-p53 antibody precipitated the p21 promoter
476 region but not the downstream region under a non-oxidative stress condition, and precipitated p21
477 promoter region was inhibited by transfection of wild-type DJ-1-HA but not by that of C106S
478 DJ-1-HA under an oxidative stress condition. The precipitated p53 levels were at the similar range by
479 Western blotting (Figs. 15C and 15D). These results suggest that DJ-1 inhibits p53 activity in a
480 DNA-binding affinity-dependent manner.

481

482 **DISCUSSION**

483 In this study, we showed that oxidized DJ-1 at C106 induced by oxidative stress strongly binds to
484 p53 and that the enhanced interaction of DJ-1 with p53 under the oxidative stress condition is
485 required to suppress p53-dependent transcriptional activation of the DUSP1 gene through preventing
486 promoter recognition of p53 (Fig. 16A). C106 of DJ-1 is essential for increase of complex formation
487 with p53 (Fig. 3C). Although several studies, including studies by us, have shown direct interaction
488 of DJ-1 with p53 (11, 41), this is the first evidence that oxidative status of DJ-1 regulates its
489 interaction.

490 C106 of DJ-1 is oxidized from forms of SH (reduced), SOH, SO₂H and SO₃H, and excess
491 oxidation of C106, probably as an SO₃H form, has been found in brains of patients with Parkinsons
492 disease and Alzheimer's disease (1, 6). What type of oxidation form of C106 is the active form is still
493 under debate. Zhou et al. reported that DJ-1 at C106 with SO₂H is an active form in terms of

494 chaperone activity toward α -synuclein (53). We have, on the other hand, reported that stimulating
495 activity of DJ-1 toward tyrosine hydroxylase and L-DOPA decarboxylase requires the presence of
496 reduced and SOH forms of C106, which account for more than 50% of total forms (16). In this study,
497 cells were treated with 1 mM H₂O₂ for 30 min and pI of DJ-1 spots was shifted from pI 6.2 to 5.8.
498 (Fig. 3B). Since a spot of pI 5.8 contained more than 50% of the SO₂H form of C106 to total forms
499 of C106 (data not shown), it is possible that DJ-1 with SO₂H of C106 preferentially binds to p53 to
500 inhibit p53-dependent DUSP1 expression. Since other forms of C106 were also involved in the pI
501 5.8 spot, however, we cannot rule out the possibility that C106 forms other than SO₂H much more
502 effectively work toward p53. Establishment of methods to purify DJ-1 possessing respective forms
503 of C106 is necessary for analysis of DJ-1.

504 p53 bound to the DUSP1 promoter at 15 min after exposure of cells to H₂O₂ (Fig. 10A) and the
505 expression level of DUSP1 mRNA then peaked at 30 min (Fig. 4A). The DJ-1-binding level with p53
506 was also peaked at 30 min after H₂O₂ exposure and then decreased (Fig. 4B) concomitant with
507 decreased binding of p53 to the DUSP1 promoter (Fig. 10A). When the same experiments were
508 carried out using DJ-1(-/-) cells, the expression level of DUSP1 mRNA was increased at 30 min
509 after H₂O₂ exposure (Figs. 4D and 4E). If the p53/DJ-1 interaction is a negative feedback that shuts
510 down DUSP1 after its initial activation, binding of DJ-1 to p53 must be correlated with repression of
511 DUSP1 expression at later time points after H₂O₂. As described above, this is not a case. As shown in
512 Figs. 11A and 11C, DJ-1 increased the phosphorylation level of ERK through down-regulation of
513 DUSP1 expression at 15 min after second exposure of H₂O₂, but thereafter there was no difference of
514 decreasing curve of ERK phosphorylation between DJ-1(+/+) and DJ-1(-/-) cells, suggesting that
515 DJ-1 determines the maximal level of ERK phosphorylation. From these results, the mechanism of
516 DJ-1 action toward DUSP1 expression may be followings: DJ-1 receives oxidative stress as an
517 oxidative stress sensor and its C106 is oxidized. Oxidized DJ-1 binds to p53 at the limited time point,
518 resulting in suppression of DUSP1 expression, thereby regulating the activation level of ERK.

519 p53 is post-translationally modified by various stresses, and tetramerization of p53 occurs to
520 function. We did not find any evidence that phosphorylation of p53 affects its interaction with DJ-1.
521 p53, however, receives other post-translational modifications such as lysine-acetylation. Acetylation
522 of p53 enhances DNA-binding activity of p53 to specific DNA regions (15) and it is therefore
523 possible that acetylation of p53 affects DUSP1 expression. Furthermore, it has been reported that p53
524 induces phosphorylation of DJ-1 (34). Thus, it is thought that in addition to oxidation, DJ-1 chooses
525 its binding proteins dependently on various modifications of DJ-1.

526 Thirty-min exposure of cells to H₂O₂ induced phosphorylation of serines 15 and 20, but not
527 serine 46, of p53 and the p53-DJ-1 complex level was increased at the peak of 30 min after H₂O₂
528 treatment (Figs. 2 and 4B). Since severe DNA damage leads to phosphorylation of p53 serine 46 (33),
529 it is thought that weak DNA damage occurred under the oxidative stress condition used in this study
530 and that DJ-1 contributes to suppression of p53 under the condition of such weak damage.

531 Of the p53 DNA-binding sequences, p21 and DUSP1 promoters contain consensus and
532 non-consensus p53-binding sequences, respectively, which have high affinity and low affinity to p53
533 (12, 25). The protein level of Bax, the promoter of which contains the consensus p53-binding
534 sequence, is down-regulated by DJ-1 before and after H₂O₂ treatment of cells (11, 29), suggesting
535 that suppressive activity of DJ-1 toward Bax does not depend on the oxidative status of DJ-1. It is of
536 interest that DJ-1 inhibited p53 transcriptional activity toward the DUSP1 promoter but not that
537 toward the p21 promoter (Fig. 7) and that these phenomena were obtained through interference of the
538 DNA-binding activity of p53 toward the DUSP1 promoter by DJ-1 (Fig.9). When p53 mutants with
539 low DNA-binding affinity to the p21 promoter were used, however, DJ-1 inhibited their
540 transcriptional activity even to the p21 promoter (Fig. 14A). Binding activity of DJ-1 to p53 mutants
541 was the same as that to wild-type p53 (Fig. 13A). It is therefore thought that DJ-1 interferes with
542 p53-binding activity to the DNA region that possesses weak association with p53 and that DJ-1 does
543 not affect p53-binding activity to the DNA region that strongly interacts with p53 under an oxidative

544 stress condition. In other words, the binding affinity of p53 to the respective DNA-binding region
545 determines whether or not DJ-1 inhibits DNA-binding and transcriptional activities of p53 (Fig. 16B).
546 The present findings suggest that DJ-1 plays a novel role in suppressing the expression of p53-target
547 genes through preventing promoter recognition of p53 in a DNA-binding affinity-dependent manner.
548 Furthermore, DJ-1 and p53 were co-localized in the nucleus before and after H₂O₂ treatment (Figs.
549 1E and F) and p53 was localized to the nucleus both in DJ-1 (-/-) and DJ-1 (+/+) cells with or
550 without H₂O₂ treatment (Figs. 4J and 4K), suggesting that DJ-1 inhibits binding of p53 to specific
551 DNA regions without affecting localization of p53.

552 Exposure of cells to H₂O₂ quickly induces phosphorylation of p53 and ERK. Activated p53 then
553 up-regulates the expression of pro-apoptotic genes (4, 42). Activation of ERK via the Ras/Raf/MEK
554 pathway, on the other hand, supports cell survival (3, 49). It has been reported that inhibition of p53
555 activates ERK after H₂O₂ treatment of cells, suggesting the existence of a cross-talk of the negative
556 signaling pathway between p53 and ERK (22). Oxidized DJ-1 inhibited p53-dependent DUSP1
557 expression under an oxidative stress condition (Figs. 4D, 4F and 7A), and it increased ERK
558 phosphorylation and cell survival (Figs. 11 and 12). Suppression of DUSP1 expression also results in
559 a decrease of oxidative stress-induced cell death in SH-SY5Y cells (18), and activation of ERK
560 regulates tyrosine hydroxylase transcription through activating the orphan nuclear receptor Nurr1 (17,
561 26). In addition to ERK, DJ-1 inhibits p38, MKK3 and MKK6 activity through inactivating ASK1
562 under an oxidative stress condition (30). It is therefore thought that DJ-1 contributes to activation of
563 survival pathways by suppressing the apoptosis pathway through regulating expression of MAPK
564 phosphatase under an oxidative condition.

565 It has been reported that DJ-1 is overexpressed in many types of cancers, especially in cancer
566 cells with poor prognosis, and that almost half of the cancer cells possess p53 mutations (7, 23, 28,
567 52, 54). In this study, DJ-1 inhibited transcriptional activity of p53 mutants more effectively than that
568 of wild-type p53 after cells had been exposed to H₂O₂ (Fig. 14A): p21 promoter activity brought by

569 R181L p53 was decreased to 82% of that brought by wild-type p53 and was inhibited by wild-type
570 DJ-1 to 57.2% of that without DJ-1, indicating that the final promoter activity by R181L p53 became
571 47% of that by wild-type p53. In the case of R181C p53, the final activity was reduced only to 8% of
572 that obtained by wild-type p53 (Figs. 13B and 14A). DJ-1 activates ERK (Fig. 11A) and Akt
573 pathways by binding to PTEN (20) and has a cooperative transforming activity with H-Ras, which is
574 located upstream of both pathways (31). These results suggest that in addition to stimulation of cell
575 proliferation pathways, overexpressed DJ-1 contributes to the poor prognosis in cancer cells by
576 suppressing cell death pathways.

577

578

579 REFERENCES

- 580 1. **Bandopadhyay R, Kingsbury AE, Cookson MR, Reid AR, Evans IM, Hope AD, Pittman**
581 **AM, Lashley T, Canet-Aviles R, Miller DW, McLendon C, Strand C, Leonard AJ,**
582 **Abou-Sleiman PM, Healy DG, Ariga H, Wood NW, de Silva R, Revesz T, Hardy JA, Lees**
583 **AJ.** 2004. The expression of DJ-1 (PARK7) in normal human CNS and idiopathic Parkinson's
584 disease. *Brain* **127**:420-430.
- 585 2. **Bonifati V, Rizzu P, van Baren MJ, Schaap O, Breedveld GJ, Krieger E, Dekker MC,**
586 **Squitieri F, Ibanez P, Joosse M, van Dongen JW, Vanacore N, van Swieten JC, Brice A,**
587 **Meco G, van Duijn CM, Oostra BA, Heutink P.** 2003. Mutations in the DJ-1 gene associated
588 with autosomal recessive early-onset parkinsonism. *Science* **299**:256-259.
- 589 3. **Bonni A, Brunet A, West AE, Datta SR, Takasu MA, Greenberg ME.** 1999. Cell survival
590 promoted by the Ras-MAPK signaling pathway by transcription-dependent and -independent
591 mechanisms. *Science* **286**:1358-1362.
- 592 4. **Bulavin DV, Saito S, Hollander MC, Sakaguchi K, Anderson CW, Appella E, Fornace AJ,**
593 **Jr.** 1999. Phosphorylation of human p53 by p38 kinase coordinates N-terminal phosphorylation
594 and apoptosis in response to UV radiation. *EMBO J.* **18**:6845-6854.

- 595 5. **Canet-Aviles RM, Wilson MA, Miller DW, Ahmad R, McLendon C, Bandyopadhyay S,**
596 **Baptista MJ, Ringe D, Petsko GA, Cookson MR.** 2004. The Parkinson's disease protein DJ-1
597 is neuroprotective due to cysteine-sulfinic acid-driven mitochondrial localization. *Proc. Natl.*
598 *Acad. Sci. of the U. S. A.* **101**:9103-9108.
- 599 6. **Choi J, Sullards MC, Olzmann JA, Rees HD, Weintraub ST, Bostwick DE, Gearing M,**
600 **Levey AI, Chin LS, Li L.** 2006. Oxidative damage of DJ-1 is linked to sporadic Parkinson and
601 Alzheimer diseases. *J. Biol. Chem.* **281**:10816-10824.
- 602 7. **Chow SN, Chen RJ, Chen CH, Chang TC, Chen LC, Lee WJ, Shen J, Chow LP.** 2010.
603 Analysis of protein profiles in human epithelial ovarian cancer tissues by proteomic technology.
604 *Eur. J. Gynaecol. Oncol.* **31**:55-62.
- 605 8. **Clements CM, McNally RS, Conti BJ, Mak TW, Ting JP.** 2006. DJ-1, a cancer- and
606 Parkinson's disease-associated protein, stabilizes the antioxidant transcriptional master regulator
607 Nrf2. *Proc. Natl. Acad. Sci. of the U. S. A.* **103**:15091-15096.
- 608 9. **Das S, Raj L, Zhao B, Kimura Y, Bernstein A, Aaronson SA, Lee SW.** 2007. Hzf Determines
609 cell survival upon genotoxic stress by modulating p53 transactivation. *Cell* **130**:624-637.
- 610 10. **Fan J, Ren H, Fei E, Jia N, Ying Z, Jiang P, Wu M, Wang G.** 2008. Sumoylation is critical for
611 DJ-1 to repress p53 transcriptional activity. *FEBS Lett.* **582**:1151-1156.
- 612 11. **Fan J, Ren H, Jia N, Fei E, Zhou T, Jiang P, Wu M, Wang G.** 2008. DJ-1 decreases Bax
613 expression through repressing p53 transcriptional activity. *J. Biol. Chem.* **283**:4022-4030.
- 614 12. **Freeman J, Schmidt S, Scharer E, Iggo R.** 1994. Mutation of conserved domain II alters the
615 sequence specificity of DNA binding by the p53 protein. *EMBO J.* **13**:5393-5400.
- 616 13. **Goldberg MS, Pisani A, Haburcak M, Vortherms TA, Kitada T, Costa C, Tong Y, Martella**
617 **G, Tschertter A, Martins A, Bernardi G, Roth BL, Pothos EN, Calabresi P, Shen J.** 2005.
618 Nigrostriatal dopaminergic deficits and hypokinesia caused by inactivation of the familial
619 Parkinsonism-linked gene DJ-1. *Neuron* **45**:489-496.

- 620 14. **Graham FL, van der Eb AJ.** 1973. Transformation of rat cells by DNA of human adenovirus 5.
621 *Virology* **54**:536-539.
- 622 15. **Gu W, Roeder RG.** 1997. Activation of p53 sequence-specific DNA binding by acetylation of
623 the p53 C-terminal domain. *Cell* **90**:595-606.
- 624 16. **Ishikawa S, Taira T, Niki T, Takahashi-Niki K, Maita C, Maita H, Ariga H, Iguchi-Ariga**
625 **SM.** 2009. Oxidative status of DJ-1-dependent activation of dopamine synthesis through
626 interaction of tyrosine hydroxylase and 4-dihydroxy-L-phenylalanine (L-DOPA) decarboxylase
627 with DJ-1. *J. Biol. Chem.* **284**:28832-28844.
- 628 17. **Jacobsen KX, MacDonald H, Lemonde S, Daigle M, Grimes DA, Bulman DE, Albert PR.**
629 2008. A Nurr1 point mutant, implicated in Parkinson's disease, uncouples ERK1/2-dependent
630 regulation of tyrosine hydroxylase transcription. *Neurobiol. Dis.* **29**:117-122.
- 631 18. **Kim GS, Choi YK, Song SS, Kim WK, Han BH.** 2005. MKP-1 contributes to oxidative
632 stress-induced apoptosis via inactivation of ERK1/2 in SH-SY5Y cells. *Biochem. Biophys. Res.*
633 *Commun.* **338**:1732-1738.
- 634 19. **Kim RH, Peters M, Jang Y, Shi W, Pintilie M, Fletcher GC, DeLuca C, Liepa J, Zhou L,**
635 **Snow B, Binari RC, Manoukian AS, Bray MR, Liu FF, Tsao MS, Mak TW.** 2005. DJ-1, a
636 novel regulator of the tumor suppressor PTEN. *Cancer Cell* **7**:263-273.
- 637 20. **Kim YC, Kitaura H, Taira T, Iguchi-Ariga SM, Ariga H.** 2009. Oxidation of DJ-1-dependent
638 cell transformation through direct binding of DJ-1 to PTEN. *Int. J. Oncol.* **35**:1331-1341.
- 639 21. **Kinumi T, Kimata J, Taira T, Ariga H, Niki E.** 2004. Cysteine-106 of DJ-1 is the most
640 sensitive cysteine residue to hydrogen peroxide-mediated oxidation in vivo in human umbilical
641 vein endothelial cells. *Biochem. Biophys. Res. Commun.* **317**:722-728.
- 642 22. **Lange-Carter CA, Pleiman CM, Gardner AM, Blumer KJ, Johnson GL.** 1993. A divergence
643 in the MAP kinase regulatory network defined by MEK kinase and Raf. *Science* **260**:315-319.
- 644 23. **Le Naour F, Misek DE, Krause MC, Deneux L, Giordano TJ, Scholl S, Hanash SM.** 2001.

- 645 Proteomics-based identification of RS/DJ-1 as a novel circulating tumor antigen in breast cancer.
646 Clin. Cancer Res. **7**:3328-3335.
- 647 24. **Li HM, Niki T, Taira T, Iguchi-Arigo SM, Ariga H.** 2005. Association of DJ-1 with chaperones
648 and enhanced association and colocalization with mitochondrial Hsp70 by oxidative stress. Free
649 Radic. Res. **39**:1091-1099.
- 650 25. **Liu YX, Wang J, Guo J, Wu J, Lieberman HB, Yin Y.** 2008. DUSP1 is controlled by p53
651 during the cellular response to oxidative stress. Mol. Cancer Res. **6**:624-633.
- 652 26. **Lu L, Sun X, Liu Y, Zhao H, Zhao S, Yang H.** 2012. DJ-1 upregulates tyrosine hydroxylase
653 gene expression by activating its transcriptional factor Nurr1 via the ERK1/2 pathway. Int. J.
654 Biochem. Cell. Biol. **44**:65-71.
- 655 27. **Martinat C, Shendelman S, Jonason A, Leete T, Beal MF, Yang L, Floss T, Abeliovich A.**
656 2004. Sensitivity to oxidative stress in DJ-1-deficient dopamine neurons: an ES- derived cell
657 model of primary Parkinsonism. PLoS Biol. **2**:e327.
- 658 28. **Melle C, Ernst G, Escher N, Hartmann D, Schimmel B, Bleul A, Thieme H, Kaufmann R,**
659 **Felix K, Friess HM, Settmacher U, Hommann M, Richter KK, Daffner W, Taubig H,**
660 **Manger T, Claussen U, von Eggeling F.** 2007. Protein profiling of microdissected pancreas
661 carcinoma and identification of HSP27 as a potential serum marker. Clin Chem. **53**:629-635.
- 662 29. **Miyashita T, Reed JC.** 1995. Tumor suppressor p53 is a direct transcriptional activator of the
663 human bax gene. Cell **80**:293-299.
- 664 30. **Mo JS, Jung J, Yoon JH, Hong JA, Kim MY, Ann EJ, Seo MS, Choi YH, Park HS.** 2010.
665 DJ-1 modulates the p38 mitogen-activated protein kinase pathway through physical interaction
666 with apoptosis signal-regulating kinase 1. J. Cell. Biochem. **110**:229-237.
- 667 31. **Nagakubo D, Taira T, Kitaura H, Ikeda M, Tamai K, Iguchi-Arigo SM, Ariga H.** 1997. DJ-1,
668 a novel oncogene which transforms mouse NIH3T3 cells in cooperation with ras. Biochem.
669 Biophys. Res. Commun. **231**:509-513.

- 670 32. **Niki T, Takahashi-Niki K, Taira T, Iguchi-Ariga SM, Ariga H.** 2003. DJBP: a novel
671 DJ-1-binding protein, negatively regulates the androgen receptor by recruiting histone
672 deacetylase complex, and DJ-1 antagonizes this inhibition by abrogation of this complex. *Mol.*
673 *Cancer Res.* **1**:247-261.
- 674 33. **Oda K, Arakawa H, Tanaka T, Matsuda K, Tanikawa C, Mori T, Nishimori H, Tamai K,**
675 **Tokino T, Nakamura Y, Taya Y.** 2000. p53AIP1, a potential mediator of p53-dependent
676 apoptosis, and its regulation by Ser-46-phosphorylated p53. *Cell* **102**:849-862.
- 677 34. **Rahman-Roblick R, Hellman U, Becker S, Bader FG, Auer G, Wiman KG, Roblick UJ.** 2008.
678 Proteomic identification of p53-dependent protein phosphorylation. *Oncogene* **27**:4854-4859.
- 679 35. **Saito Y, Hamakubo T, Yoshida Y, Ogawa Y, Hara Y, Fujimura H, Imai Y, Iwanari H,**
680 **Mochizuki Y, Shichiri M, Nishio K, Kinumi T, Noguchi N, Kodama T, Niki E.** 2009.
681 Preparation and application of monoclonal antibodies against oxidized DJ-1. Significant
682 elevation of oxidized DJ-1 in erythrocytes of early-stage Parkinson disease patients. *Neurosci*
683 *Lett.* **465**:1-5.
- 684 36. **Samuels-Lev Y, O'Connor DJ, Bergamaschi D, Trigiante G, Hsieh JK, Zhong S, Campargue**
685 **I, Naumovski L, Crook T, Lu X.** 2001. ASPP proteins specifically stimulate the apoptotic
686 function of p53. *Mol. Cell* **8**:781-794.
- 687 37. **Santibanez-Koref MF, Birch JM, Hartley AL, Jones PH, Craft AW, Eden T, Crowther D,**
688 **Kelsey AM, Harris M.** 1991. p53 germline mutations in Li-Fraumeni syndrome. *Lancet*
689 **338**:1490-1491.
- 690 38. **Schlereth K, Beinoraviciute-Kellner R, Zeitlinger MK, Bretz AC, Sauer M, Charles JP,**
691 **Vogiatzi F, Leich E, Samans B, Eilers M, Kisker C, Rosenwald A, Stiewe T.** 2010. DNA
692 binding cooperativity of p53 modulates the decision between cell-cycle arrest and apoptosis.
693 *Mol. Cell* **38**:356-368.

- 694 39. **Shaw P, Bovey R, Tardy S, Sahli R, Sordat B, Costa J.** 1992. Induction of apoptosis by
695 wild-type p53 in a human colon tumor-derived cell line. Proc. Natl. Acad. Sci. of the U. S. A.
696 **89:4495-4499.**
- 697 40. **Shinbo Y, Niki T, Taira T, Ooe H, Takahashi-Niki K, Maita C, Seino C, Iguchi-Arigo SM,**
698 **Ariga H.** 2006. Proper SUMO-1 conjugation is essential to DJ-1 to exert its full activities. Cell
699 Death Diff. **13:96-108.**
- 700 41. **Shinbo Y, Taira T, Niki T, Iguchi-Arigo SM, Ariga H.** 2005. DJ-1 restores p53 transcription
701 activity inhibited by Topors/p53BP3. Int. J. Oncol. **26:641-648.**
- 702 42. **Shono T, Tofilon PJ, Schaefer TS, Parikh D, Liu TJ, Lang FF.** 2002. Apoptosis induced by
703 adenovirus-mediated p53 gene transfer in human glioma correlates with site-specific
704 phosphorylation. Cancer Res. **62:1069-1076.**
- 705 43. **Sun H, Charles CH, Lau LF, Tonks NK.** 1993. MKP-1 (3CH134), an immediate early gene
706 product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo. Cell
707 **75:487-493.**
- 708 44. **Taira T, Saito Y, Niki T, Iguchi-Arigo SM, Takahashi K, Ariga H.** 2004. DJ-1 has a role in
709 antioxidative stress to prevent cell death. EMBO Rep. **5:213-218.**
- 710 45. **Takahashi-Niki K, Niki T, Taira T, Iguchi-Arigo SM, Ariga H.** 2004. Reduced anti-oxidative
711 stress activities of DJ-1 mutants found in Parkinson's disease patients. Biochem. Biophys. Res.
712 Commun. **320:389-397.**
- 713 46. **Takahashi K, Taira T, Niki T, Seino C, Iguchi-Arigo SM, Ariga H.** 2001. DJ-1 positively
714 regulates the androgen receptor by impairing the binding of PIASx alpha to the receptor. J. Biol.
715 Chem. **276:37556-37563.**
- 716 47. **Tanaka T, Ohkubo S, Tatsuno I, Prives C.** 2007. hCAS/CSE1L associates with chromatin and
717 regulates expression of select p53 target genes. Cell **130:638-650.**
- 718 48. **WA, F-P, Prives C.** 2012. Mutant p53: one name, many proteins. Genes Dev. **26:1268-1286.**

- 719 49. **Wang JK, Gao G, Goldfarb M.** 1994. Fibroblast growth factor receptors have different
720 signaling and mitogenic potentials. *Mol. Cell. Biol.* **14**:181-188.
- 721 50. **Weger S, Hammer E, Engstler M.** 2003. The DNA topoisomerase I binding protein topors as a
722 novel cellular target for SUMO-1 modification: characterization of domains necessary for
723 subcellular localization and sumolation. *Exp. Cell Res.* **290**:13-27.
- 724 51. **Xu J, Zhong N, Wang H, Elias JE, Kim CY, Woldman I, Pifl C, Gygi SP, Geula C, Yankner**
725 **BA.** 2005. The Parkinson's disease-associated DJ-1 protein is a transcriptional co-activator that
726 protects against neuronal apoptosis. *Hum. Mol. Genet.* **14**:1231-1241.
- 727 52. **Yuen HF, Chan YP, Law S, Srivastava G, El-Tanani M, Mak TW, Chan KW.** 2008. DJ-1
728 could predict worse prognosis in esophageal squamous cell carcinoma. *Cancer Epidemiol.*
729 *Biomarkers Prev.* **17**:3593-3602.
- 730 53. **Zhou W, Zhu M, Wilson MA, Petsko GA, Fink AL.** 2006. The oxidation state of DJ-1
731 regulates its chaperone activity toward alpha-synuclein. *J. Mol. Biol.* **356**:1036-1048.
- 732 54. **Zhu XL, Wang ZF, Lei WB, Zhuang HW, Jiang HY, Wen WP.** 2010. DJ-1: a novel
733 independent prognostic marker for survival in glottic squamous cell carcinoma. *Cancer Sci.*
734 **101**:1320-1325.

735

736 **ACKNOWLEDGEMENTS**

737 This work was supported by grants-in-aid from the Ministry of Education, Science, Culture and
738 Sports and by the Program for Promotion of Fundamental Studies in Health Sciences of the National
739 Institute of Biomedical Innovation (NIBIO) in Japan. We thank Kiyomi Takaya for her technical
740 assistance.

741

742 **LEGEND OF FIGURES**

743 **FIG. 1.** Oxidative stress enhances DJ-1-binding to p53.

744 **A.** HEK293T cells were transfected with FLAG-p53 and DJ-1-HA and treated with 1 mM H₂O₂ for

745 30 min at 48 hrs after transfection. Proteins were analyzed by immunoprecipitation followed by
746 Western blotting.

747 **B.** HEK293T cells were transfected with FLAG-DJ-1 and T7-p53 and analyzed as described in a
748 legend for Fig. 1A.

749 **C, D.** A549 cells (p53+/+) (C) and H1299 cells (p53-/-) (D) were treated with 300 μ M H₂O₂ for 30
750 min, and proteins analyzed by immunoprecipitation followed by Western blotting. The intensity of
751 precipitated p53 bands in lanes 5 and 6 was quantified and their relative level is shown under the
752 figure.

753 **E, F.** A549 cells were treated with (F) or without (E) 300 μ M H₂O₂ for 30 min. Immunofluorescence
754 analyses were carried out using anti-p53 and anti-DJ-1 antibodies as described in Materials and
755 methods.

756

757 **FIG. 2.** Phosphorylation levels of serines in p53 under an oxidative stress condition.

758 **A.** HEK293T cells with or without 1 mM H₂O₂ for 30 min, and proteins were analyzed by Western
759 blotting with anti-p53 and respective anti-p-p53 antibodies.

760 **B.** HEK293T cells transfected with FLAG-wild-type and -mutants of p53 were treated with 1 mM
761 H₂O₂ for 30 min at 48 hrs after transfection. Proteins were detected by Western blotting.

762 **C.** HEK293T cells transfected with DJ-1-HA, FLAG-wild-type and mutants of p53 were treated with
763 1 mM H₂O₂ for 30 min at 48 hrs after transfection. Proteins were analyzed by immunoprecipitation
764 followed by Western blotting. A precipitated p53 band in lane 2 is overflow from a band in lane 3.

765

766 **FIG. 3.** Cysteine 106 of DJ-1 is essential for binding to p53 under an oxidative stress condition.

767 **A.** HEK293T cells were transfected with wild-type or C106S DJ-1-HA and FLAG-p53. Forty-eight
768 hrs after transfection, cells were treated with 1 mM H₂O₂ for 30 min, and proteins were analyzed by
769 an isoelectric focusing gel and subjected to Western blotting with an anti-HA antibody.

770 B. HEK293T cells were treated with 20 J/m² UV, 1 μM doxorubicin or 1 mM H₂O₂ for 0.5-4 hrs and
771 proteins were analyzed by Western blotting.

772 C. HEK293T cells were transfected with wild-type DJ-1-HA or C106S DJ-1-HA and FLAG-p53 and
773 treated with H₂O₂ as described in the legend of Fig. 2A. Proteins were analyzed by
774 immunoprecipitation followed by Western blotting.

775 D. GST, GST-wild-type DJ-1 and GST-C106S DJ-1 were incubated with ³⁵S-labeled p53 in the
776 presence or absence of H₂O₂ and DTT, and subjected to pull-down assays. CBB: Coomassie brilliant
777 blue.

778 E. HEK293T cells were transfected with FLAG-wild-type DJ-1 or FLAG -C106S DJ-1 and T7-p53
779 and treated with H₂O₂. Proteins were then analyzed by immunoprecipitation followed by Western
780 blotting.

781 F, G. HEK293T cells (F) or H1299 cells (G) were transfected with wild-type or C106S DJ-1-HA and
782 One-STrEP-p53 and treated with H₂O₂. Proteins were then subjected to pull-down assays using
783 Strep-Tactin sepharose beads.

784

785 **FIG. 4.** DJ-1 down-regulates DUSP1 expression under an oxidative stress condition.

786 A. Mouse primary cells were treated with H₂O₂ for 0.25-6 hrs, and the expression levels of
787 respective mRNA were examined by semi-quantitative RT-PCR and their relative expression to that
788 of β-actin is shown.

789 B. HEK293T cells were transfected with FLAG-p53 and DJ-1-HA and treated with 1 mM H₂O₂ for
790 15-45 min (left panel) and for 0.5-4 hrs (right panel) at 48 hrs after transfection. Proteins were
791 analyzed by immunoprecipitation followed by Western blotting.

792 C. A549 cells were treated with 300 μM H₂O₂ for 15-120 min. Proteins were analyzed by
793 immunoprecipitation with an anti-DJ-1 antibody followed by Western blotting.

794 D, E. DJ-1 (+/+) and DJ-1 (-/-) mouse cells were treated with H₂O₂ for 0.5 hrs (C) or 2 hrs (D). The

795 expression levels of DUSP1 (C) and p21 mRNA (D) were examined by semi-quantitative RT-PCR
796 and their relative expression to that of β -actin is shown.

797 **F, G.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were treated with H₂O₂ for 0.5 and 2 hrs. The expression
798 levels of DUSP1 (F) and p21 mRNA (G) were examined by quantitative RT-PCR.

799 H. DJ-1 (+/+) and DJ-1 (-/-) mouse cells were treated with H₂O₂ for 0.5 and 2 hrs. The expression
800 levels of respective protein were analyzed by Western blotting.

801 I. DJ-1 (+/+) and DJ-1 (-/-) mouse cells were transfected with control siRNA or p53 siRNA and
802 treated with 300 μ M H₂O₂ for 30 min or 2 hrs at 72 hrs after transfection. The expression levels of
803 DUSP1 and p21 mRNAs were examined by semi-quantitative RT-PCR. Nucleotide sequences of p53
804 siRNA are as follows: mp53 sense: 5'- CCAGAAGAUAUCCUGCCAUTT-3' and mp53 antisense:
805 5'- AUGGCAGGAUAUCUUCUGGTT-3'.

806 **J, K.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were treated with H₂O₂ for 0.5 and 2 hrs. p53 was
807 visualized as described in Materials and methods.

808 Values in Figures D-G are means \pm S.E. n=3 experiments. Significance: ** p < 0.01 and *** p < 0.001.
809 N.S. represents no significance.

810

811 **FIG. 5.** Expression of DUSP1, p21 and NOXA in UV- and doxorubicin-treated cells.

812 Mouse DJ-1 (+/+) cells were treated with 20 J/m² UV (A, C) or 1 μ M doxorubicin (B, D) for 0.5-6
813 hrs. The expression levels of mRNA and protein of DUSP1, p21 and NOXA were examined by
814 semi-quantitative RT-PCR (A, B) and by Western blotting (C, D), respectively.

815

816 **FIG. 6.** Raw data in three sets of luciferase assays.

817 **A, B.** H1299 cells were co-transfected with pGL4.12-DUSP1-luciferase (A) or
818 pGL4.12-p21-luciferase (B) and expression vectors for FLAG-p53 and wild-type or C106S DJ-1-HA.
819 Twenty-four hrs after transfection, cells were treated with 300 μ M H₂O₂ or with 10 μ g/mL

820 cycloheximide (CHX) for 30 min. Luciferase activities were then calculated.

821 **C, D.** Proteins were prepared from H1299 cells transfected with FLAG-p53 and wild-type or C106S
822 DJ-1-HA and analyzed by Western blotting.

823

824 **FIG. 7.** Cysteine 106 of DJ-1 is essential for repression of p53-dependent DUSP1 transcription under
825 an oxidative stress condition.

826 **A, B, C.** H1299 cells were co-transfected with pGL4.12-DUSP1-luciferase (A),
827 pGL4.12-p21-luciferase (B) or pGL4.12-luciferase (C) and FLAG-p53 and wild-type or C106S
828 DJ-1-HA. Cells were treated as described in the legend for Fig. 6. Fold repression of luciferase
829 activity was calculated as described in the text.

830 **D.** H1299 cells were transfected with FLAG-p53 and DJ-1-HA and treated with H₂O₂ for 30 min in
831 the absence of cycloheximide as described in the legend for Fig. 6. The expression levels of DUSP1
832 mRNA were examined by quantitative RT-PCR.

833 Values are means ± S.E. n=3 experiments. Significance: ***p* < 0.01, ****p* < 0.001. N.S. represents no
834 significance.

835

836

837 **FIG. 8.** DJ-1 directly binds to the p53 DNA-binding region.

838 **A.** Schematic diagram of p53 deletion mutants. FL: full-length; CT: C-terminus; DBD: DNA-binding
839 domain.

840 **B.** GST and GST-DJ-1 were incubated with ³⁵S-labeled p53 in the presence or absence of H₂O₂, and
841 then subjected to pull-down assays. I: Input; G: GST; D: GST-DJ-1; CBB: Coomassie brilliant blue.

842 **C.** GST, GST-DJ-1 and GST-C106S DJ-1 were incubated with ³⁵S-labeled p53DBD and subjected to
843 pull-down assays. CBB: Coomassie brilliant blue.

844 **D.** ELISA assays were carried out using GST-p53, GST, DJ-1 and C106S DJ-1.

845

846 **FIG. 9.** DJ-1-p53 complex stably binds to the p21 promoter but not to the DUSP1 promoter.

847 **A, B.** H1299 cells were transfected with FLAG-p53 and wild-type or C106S DJ-1-HA. Forty-eight
848 hrs after transfection, cells were treated with 300 μ M H₂O₂ for 30 min, and ChIP assays were carried
849 out targeting DUSP1 (A) and p21 (B) genes.

850 **C.** Proteins prepared from transfected H1299 cells were analyzed by Western blotting.

851 **D, E.** A549 cells were treated with 300 μ M H₂O₂ for 30 min, and ChIP assays were carried out
852 targeting DUSP1 (D) and p21 (E) genes. The intensity of precipitated DNA bands in lanes 4 and 6 of
853 Figure E was quantified and their relative level is shown.

854 **F.** Proteins prepared from H₂O₂-treated A549 cells were analyzed by Western blotting.

855

856 **FIG. 10.** A time course of endogenous DJ-1/p53 interactions and endogenous p53 ChIP on the DUSP
857 promoter in cells exposed to oxidative stress.

858 **A.** A549 cells were treated with 300 μ M H₂O₂ for 0.25-2 hrs, and ChIP assays targeting the DUSP1
859 gene were carried out.

860 **B.** Proteins prepared from H₂O₂-treated A549 cells were analyzed by Western blotting.

861 **C.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were treated with 300 μ M H₂O₂ for 0.5 hrs, and ChIP assays
862 targeting the DUSP1 gene were carried out.

863 **D.** Proteins prepared from H₂O₂-treated mouse DJ-1 (+/+) and DJ-1 (-/-) cells were analyzed by
864 Western blotting.

865

866 **FIG. 11.** Phosphorylation levels of ERK in cells under an oxidative stress condition.

867 **A, C.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were starved for 6 hrs and then treated with or without
868 two pulses of H₂O₂. Proteins were analyzed by Western blotting (C) and relative phosphorylation
869 levels of ERK were quantified (A).

870 **B, D.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were transfected with control siRNA or DUSP1 siRNA-1
871 or DUSP1 siRNA-2, starved for 6 hrs at 48 hrs after transfection, and then treated with two pulses of
872 H₂O₂. Proteins were analyzed by Western blotting (B) and relative phosphorylation levels of ERK
873 were quantified (D). Values are means ± S.E. n=3 experiments. Significance: **p* < 0.05.

874

875 **FIG. 12.** DJ-1 regulates cell survival under an oxidative stress condition.

876 **A.** DJ-1 (+/+) and DJ-1 (-/-) mouse cells were transfected with control siRNA or DUSP1 siRNA-1
877 and treated as described in the legend for Fig. 11. Forty hrs after H₂O₂ addition, cell-cycle profiles
878 were examined by using flow cytometry.

879 **B.** The sub-G1 phase of the cell cycle obtained in FIG. 12A was quantified, and relative number of
880 cells in the sub-G1 fraction compared to that in DJ-1 (+/+) cells with control siRNA and H₂O₂ was
881 calculated. Values are means ± S.E. n=3 experiments. Significance: ****p* < 0.001.

882

883 **FIG. 13.** Binding activity of p53 mutants to DJ-1

884 **A and C.** GST and GST-DJ-1 were incubated with ³⁵S-labeled wild-type and mutant p53 in the
885 presence or absence of H₂O₂, and subjected to pull-down assays. I: Input; G: GST; D: GST-wild-type
886 DJ-1; C: GST-C106S DJ-1; CBB: Coomassie brilliant blue.

887 **B.** H1299 cells were co-transfected with pGL4.12-p21-luciferase and expression vectors for
888 wild-type, R181L, R181C and R181P of FLAG-p53. Twenty-four hrs after transfection, luciferase
889 assays were carried out. Values are means ± S.E. n=3 experiments. Significance: ***p* < 0.01 . N.S.
890 represents no significance.

891

892 **FIG. 14.** DJ-1 inhibits transcriptional activity of p53 to the p21 promoter under an oxidative stress
893 condition in a DNA-affinity-dependent manner.

894 **A.** H1299 cells were co-transfected with pGL4.12-p21-luciferase and expression vectors for
895 FLAG-p53 mutants and for wild-type or C106S DJ-1-HA. Twenty-four hrs after transfection, their

896 luciferase activities were examined. Values are means \pm S.E. n=3 experiments. Significance: * p <
897 0.05 and ** p < 0.01 .

898 **B, C.** H1299 cells were co-transfected with pGL4.12-p21-luciferase and an expression vector for
899 wild-type or C106S DJ-1-HA together with a vector for FLAG-p53 R181L (B) or FLAG-p53 R181C
900 (C). Twenty-four hrs after transfection, the cells were treated with H₂O₂ and cycloheximide (CHX)
901 as described in the legend for Fig. 6 and their luciferase activities were examined.

902 **D, E.** Proteins were prepared from H1299 cells transfected with FLAG-p53 mutants and wild-type or
903 C106S DJ-1-HA and analyzed by Western blotting.

904

905 **Fig. 15.** DJ-1 inhibits DNA-binding activity of p53 mutants.

906 **A, B.** H1299 cells were transfected with FLAG-p53 R181L (A) or R181C (B) and wild-type or
907 C106S DJ-1-HA. Forty-eight hrs after transfection, cells were treated with 300 μ M H₂O₂ for 30 min,
908 and ChIP assays were carried out targeting the p21 gene.

909 **C, D.** Proteins were prepared from H1299 cells transfected with FLAG-p53 R181L (C) or R181C
910 (D) and wild-type or C106S DJ-1-HA and analyzed by Western blotting.

911

912 **FIG. 16.** Schematic models of the role of DJ-1 in inhibition of p53 transactivation

913 **A.** DJ-1 activates cell survival pathways under an oxidative stress condition through down-regulation
914 of DUSP1 expression.

915 **B.** DJ-1 suppresses transactivation activity of p53 depending on p53-DNA binding-affinity.

916

917

918




Fig. 1






Fig. 2




Fig. 3











Fig.4-1


F Real-time PCR




G Real-time PCR




H



I



J DJ-1(+/+)



K DJ-1 (-/-)




Fig. 4-2







Fig. 5








Fig. 6



A



	+	+	+
FLAG-p53	+	+	+
Empty vector	+	-	-
DJ-1 WT-HA	-	+	-
DJ-1 C106S-HA	-	-	+

pGL4.12-DUSP1-LUC



B



	+	+	+
FLAG-p53	+	+	+
Empty vector	+	-	-
DJ-1 WT-HA	-	+	-
DJ-1 C106S-HA	-	-	+

pGL4.12-p21-LUC

C




	+	+	+
FLAG-p53	+	+	+
Empty vector	+	-	-
DJ-1 WT-HA	-	+	-
DJ-1 C106S-HA	-	-	+

pGL4.12-LUC

D

DUSP1 mRNA



	+	+	+
FLAG-p53	+	+	+
Empty vector	+	-	-
DJ-1 WT-HA	-	+	-
DJ-1 C106S-HA	-	-	+

Fig. 7







Fig. 8




Fig. 9




Fig. 10




Fig. 11





Fig. 12




Fig. 13




Fig. 14






Fig. 15

A



B




Fig. 16