



Title	Nonmuscle myosin II folds into a 10S form via two portions of tail for dynamic subcellular localization
Author(s)	Kiboku, Takayuki; Katoh, Tsuyoshi; Nakamura, Akio; Kitamura, Akira; Kinjo, Masataka; Murakami, Yota; Takahashi, Masayuki
Citation	Genes to Cells, 18(2), 90-109 https://doi.org/10.1111/gtc.12021
Issue Date	2013-02
Doc URL	http://hdl.handle.net/2115/53054
Rights	The definitive version is available at www3.interscience.wiley.com
Type	article (author version)
Additional Information	There are other files related to this item in HUSCAP. Check the above URL.
File Information	Supporting_Information.pdf (Supporting Information)



[Instructions for use](#)

Legends for Supplemental Videos

Video S1. FRAP reveals that IIB-SK1·2 is less dynamic than IIB-WT. MRC-5 SV1 TG1 cells were cotransfected with pEGFP-IIB-IIB-SK1·2 (shown on the right) and pmCherry-IIB-WT (shown on the left), and photobleaching was performed. Images were captured using a LSM 510 META microscope at 1 frame/10 sec. The playback speed is 5 frames/sec. Still images are displayed in Fig. 4A.

Video S2. Photoconversion reveals that IIB-SK1·2 is less dynamic than IIB-WT. MRC-5 SV1 TG1 cells were transfected with phmKikGR-IIB-WT (left) or phmKikGR-IIB-SK1·2 (right) and photoconversion was performed. Images were captured using an inverted microscope (Ti-E) and confocal laser microscope system (A1R) at 1 frame/3 min. The playback speed is 5 frames/sec. Bar, 5 μm .

Video S3. Photoconversion reveals that IIB-SK1·2 is less dynamic than IIB-WT. Enlarged movie of **Video S2**. Bar, 2 μm . Still images are displayed in Fig. 4D.

Video S4. IIB-SK1·2 accumulates in the posterior region of migrating cells. MEF/3T3 Tet-Off cells were cotransfected with pEGFP-IIB-SK1·2 and pmCherry-IIB-WT. Images were captured using an inverted microscope (HS All-in-One Fluorescence Microscope BZ-9000; KEYENCE) equipped with an objective lens (CFI Plan Apo λ 20 \times /0.75 NA; KEYENCE). The cells were maintained in DMEM/F12 (1:1) (GIBCO) supplemented with 10% FBS, and were warmed at 37°C in a Stage Top Incubator (INU-KI-F1; Tokai Hit) during observation. The images were captured at 1 frame/3 min and analyzed using Multipoint Time-lapse BZ-H2TL software (KEYENCE). The playback speed is 10 frames/sec. Still images are displayed in Fig. S2.

Video S5. IIB-SK1·2 remains in the posterior of the daughter cells during postmitotic spreading. MEF/3T3 Tet-Off cells were cotransfected with pEGFP-IIB-SK1·2 and pmCherry-IIB-WT. Images were captured using an inverted microscope (Ti-E; Nikon), equipped with a cooled CCD camera (ORCA-R2; Hamamatsu Photonics) and an objective lens (Plan Flour 40 \times /0.75 NA; Nikon). Cells were maintained in DMEM/F12 (1:1) (GIBCO) supplemented with 10% FBS, and warmed at 37°C in a chamber (INUBG2TF-WSKM; Tokai Hit) during observation. The images were captured at 1 frame/5 min and analyzed using MetaMorph software (Molecular Devices). The playback speed is 10 frames/sec. Bar, 20 μm . Still images are displayed in Fig. S3.

Supplemental Table S1

Table S1. Primers used for mutagenic PCR.

Use	Primers*
NMHC-IIB-SK1 Forward Reverse	5'-TTTTGAGAAGCTTCAAGTTCAGCATGAGTTTCTTACACTTCTGCTTCTTCTGCTAACAGCTGG-3' 5'-GGAATCCCGCTCACTCAGCACAGAACTATTTAAGATTAAGAATGCCCTCGAGGAAGCCCTGGAGG-3'
NMHC-IIB-SK2 Forward Reverse	5'-GCAGGATCTTTCCCTCTTCATGTTCAAGAGATGCCTCTGCCTCCTCCAGCTGGGTCTCATTCC-3' 5'-GCATCCAGCTTGAGTTGAACCAAGTCAAGTCTGAGGTTGATAGAGACCTGCAAACCAGGGATGAGC-3'
NMHC-IIB-SK3 Forward Reverse	5'-TCTGCATGGACTCCACGATTCTAATGTGGTTTCTTTCATCTGGTCATTCTGCTCATCCCTGGTTTGC-3' 5'-GCACACTGGATGCTGAGATCAGGAGCAGGAATGATGCCATTAGGTCAAAGAAAAGATGGAGATAGACC-3'
NMHC-IIA-SK1 Forward Reverse	5'-TTTTGAGAAGCTTCAAGTTCAGCATGAGTTTCTTACACTTCTGCTTCTCCTCCGCCAGGAGCTGG-3' 5'-GGAATCCCGCTCACTCAGCACAGAACTATTTAAGATTAAGAATGCCCTGGAGGAAGCCATGGAGC-3'
NMHC-IIA-SK2 Forward Reverse	5'-GCAGGATCTTTCCCTCTTCATGTTCAAGAGATGCCTCTGCCTCTTCCAGCTGCGTCTTCATCTCC-3' 5'-GCATCCAGCTTGAGTTGAACCAAGTCAAGTCTGAGGTTGATCGGGACCTGCAGGGCCGGGACGAGC-3'
nmRLC Forward Reverse	5'-CTCGAGCCACCATGTGCGAGCAAAAAGGCA-3' 5'-GGTACCTCAGTCATCTTTGTCTTTGG-3'
HA tagged nmELC Forward Reverse	5'-GAATCCCACCATGTGTGACTTCACCGAAG-3' 5'-AAGCTTTCATGCATAGTCCGGGACGTCATACGGATAGCCATTCAGCACCATGCGGACGAGC-3'
L21 insertion Forward Reverse	5'-GGATCTCGGTCCGAAACCAACTCCTAAAAACCGCCACCATGTGCTACTACCATCAC-3' 5'-GTGATGGTAGTACGACATGGTGGCGGTTTTTTAGGAGTTGGTTTCGGACCGAGATCC-3'
mCherry Forward Reverse	5'-GCTACCGGTCGCCACCATGG-3' 5'-TGAGTACTTGTACAGCTCGTCCATGC-3'

*Underscores indicate primer sequences annealing to the template.

GenBank: *MYH10* (NMHC-IIB; NP_005955); *MYH9* (NMHC-IIA; NP_002464); *MYH14* (NMHC-IIC; NP_079005); *MYH11* (smooth muscle MHC; NP_001035202); *MYH1* (fast skeletal muscle MHC; NP_005954); *MYH6* (cardiac muscle MHC α ; NP_002462); *MYH7* (cardiac muscle MHC β ; NP_000248). A blue bar and red bars indicate the RLC-interacting region proposed by the in vitro crosslink experiments of Olney et al. (Olney et al. 1996) and Salzameda et al. (Salzameda et al. 2006), respectively. An orange arrow indicates the second skip residue from heptad repeat in the myosin tail, which is a putative second bending position. A green arrow also indicates the putative second bending position predicted by the single molecule negative stain electron microscopy with an image processing technique (Burgess et al., 2007).

Supplemental Figure S2




Figure S2. Still images from Video S4 focused on a direction change of migrating MEF/3T3 Tet-Off cells coexpressing EGFP-IIB-SK1-2 and mCherry-IIB-WT. Bar, 20 μm .

Supplemental Figure S3




Figure S3. Still images from Video S5 focused on cytokinesis and postmitotic spreading of MEF/3T3 Tet-Off cells coexpressing EGFP-IIB-SK1.2 and mCherry-IIB-WT. Bar, 20 μ m.

Supplemental Figure S4




Figure S4. Localization of endogenous myosin IIA in HeLa Tet-Off cells expressing EGFP-IIB-WT (A), or EGFP-IIB-SK1·2 (B). Second column is images of endogenous myosin IIA staining. Third column is merged images of EGFP-IIBs and endogenous myosin IIA. Fourth column is images of F-actin stained by coumarin-phalloidin. The other representative images corresponding to Fig. 8D were shown. Bar, 5 μ m.