



Title	Role of acid-tolerant arbuscular mycorrhizal fungi in plant acid tolerance and their distribution along soil pH gradients [an abstract of entire text]
Author(s)	河原, 愛
Citation	北海道大学. 博士(農学) 甲第11104号
Issue Date	2013-09-25
Doc URL	http://hdl.handle.net/2115/53965
Type	theses (doctoral - abstract of entire text)
Note	この博士論文全文の閲覧方法については、以下のサイトをご参照ください。
Note(URL)	https://www.lib.hokudai.ac.jp/dissertations/copy-guides/
File Information	Ai_Kawahara_summary.pdf



[Instructions for use](#)

学 位 論 文 要 約

博士の専攻分野の名称 博 士 (農学)

氏名 河原 愛

学 位 論 文 題 名

Role of acid-tolerant arbuscular mycorrhizal fungi in plant acid tolerance and their
distribution along soil pH gradients

(耐酸性アーバスキュラー菌根菌の植物耐酸性に果たす役割と土壌 pH 勾配に
沿った分布)

This thesis consists of 14 figures, 9 tables, 105 references, General introduction, two Chapters, and General discussion in a total of 89 pages with five accompanying publications.

In acidic soil high levels of aluminum ion inhibit plant root growth and reduce phosphorus (P) availability in the soil, and thus plants suffer serious P deficiency. *Miscanthus sinensis* is a common pioneer grass in early primary succession in acidic soil, but association with arbuscular mycorrhizal (AM) fungi is essential for the establishment in strongly acidic soil. AM fungi associate with most land plants and supply P to their host plants, but the physiological and ecological roles of the fungi in acidic soil have not been well understand. In the present study, the mechanism underlying the improved plant acid-tolerance by mycorrhizal formation was investigated with respect to acid-tolerance/sensitivity of AM fungi in Chapter 1, and the distribution pattern of acid-tolerant/sensitive AM fungi along soil pH gradients was investigated at the landscape level in conjunction with a model experiment in Chapter 2.

1. Role of acid-tolerant arbuscular mycorrhizal fungi in plant acid tolerance

In this chapter, a hypothesis that AM fungi confer the extra acid-tolerance on the host beyond its adaptability via providing an alternative pathway for nutrient uptake instead of the roots damaged by aluminum ion was addressed. *M. sinensis* seedlings were grown at pH 3.2 and 5.2 in the presence of either P-fertilizer, an acid-sensitive fungus *Claroideoglossum etunicatum* H1-1, or an acid-tolerant fungus *Rhizophagus* sp. RF1. At pH 5.2, the growth and shoot P concentration of the plants were significantly increased both by P-fertilizer and the mycorrhizal fungi irrespective of their sensitivity to soil acidity. Whereas, at pH 3.2, only the plants associated with the acid-tolerant fungus could survive. The plants grown at pH 3.2 showed more aluminum-deposition on the root tips, more damaged root tips, and shorter root length than those grown at pH 5.2, and no difference in the root damages among the treatments was observed. These observations suggested that the acid-tolerant AM fungus could confer the extra acid-tolerance on the host plants beyond the adaptability of the plant to acidic soil via providing an alternative pathway for nutrient uptake, and not via alleviating the root damages caused by aluminum.

2. Community structure of arbuscular mycorrhizal fungi along soil pH gradients

In this chapter, a question whether AM fungi in strongly acidic soils are specialists that preferentially occur in acidic soil or generalists that occur over wide ranges of soil pH was addressed. Six field sites that covered a soil pH range of 3.0–7.4 were chosen across Japan, and rhizosphere soils of the pioneer grass *M. sinensis* grown in the sites were collected. *M. sinensis* seedlings were grown on the soils in a glasshouse, and the community compositions of AM fungi in the roots were determined based on the sequences of the fungal large subunit ribosomal RNA gene. Overall, about 2,700 clones

were sequenced, and 52 AM fungal phylotypes (sequence types) were defined. AM fungal richness decreased significantly with increases in soil acidity. Ordination analysis showed that soil pH was a major driver of the communities, but geographic isolation and climatic factors were not. Nestedness analysis indicated a significant nested pattern along the soil pH gradients: the communities in lower-pH soils were subsets of those in higher-pH soils, implying that the fungi that occurred in acidic soil occurred over wide ranges of soil pH. For further confirmation, the AM fungal communities collected from neutral and acidic soils were grown at three different pH levels in association with *M. sinensis* seedlings. In the community from the neutral soil, fungal richness decreased with increases in soil acidity, and the communities grown at lower pH were nested within that grown at highest pH. In contrast, no significant nested pattern was observed in the community from the acidic soil. All these results suggest that soil acidity is a major constraint on AM fungal richness and acid-tolerant fungi are generalists that occur over wide ranges of pH. Given the significance of AM fungi in plant colonization in acidic soil, the community nestedness along soil pH gradient may contribute to rapid recovery and succession of plant community in acidic soil.

The present study provides new insight of the functional and ecological significance of arbuscular mycorrhizal fungi in acidic soil and is expected to contribute to restoration of acidic soil.