Title	Interannual changes in the zooplankton community structure on the southeastern Bering Sea shelf during summers of 1994-2009
Author(s)	Ohashi, Rie; Yamaguchi, Atsushi; Matsuno, Kohei; Saito, Rui; Yamada, Nao; Iijima, Anai; Shiga, Naonobu; Imai, Ichiro
Citation	Deep Sea Research Part II: Topical Studies in Oceanography, 94, 44-56 https://doi.org/10.1016/j.dsr2.2013.03.018
Issue Date	2013-10-01
Doc URL	http://hdl.handle.net/2115/54032
Type	article (author version)
File Information	Ohashi_MS.pdf

[Instructions for use](#)

Interannual changes in the zooplankton community structure on the southeastern Bering
Sea shelf during summers of 1994–2009

Rie Ohashi^a, Atsushi Yamaguchi^{a,*}, Kohei Matsuno^a, Rui Saito^a, Nao Yamada^b, Anai
Iijima^c, Naonobu Shiga^d, Ichiro Imai^a

^a Graduate School of Fisheries Sciences, Hokkaido University, 3-1-1 Minato, Hakodate,
Hokkaido 041-8611, Japan

^b Hokkaido Hakodate Fisheries High School, 2-15-3 Nanaehama, Hokuto, Hokkaido
049-0111, Japan

^c Donan Research Branch, Salmon and Freshwater Fisheries Research Institute,
Hokkaido Research Organization, 189-43 Kumaishi Ayukawacho, Yakumocho, Futami,
Hokkaido 043-0402, Japan

^d Hakodate Junior College, 52-1 Takaokacho, Hakodate, Hokkaido 042-0955, Japan

*Corresponding author.

Tel: +81-138-40-5543

Fax: +81-138-40-5542

E-mail: a-yama@fish.hokudai.ac.jp

Abstract

On the southeastern Bering Sea shelf, mesozooplankton play an important role in material transfer between primary producers and fisheries resources. The biomass of mesozooplankton in this region is known to vary annually, but little is known about annual changes in community structure and species composition. In the present study, regional and long-term changes in abundance, biomass and community structure of copepods and chaetognaths on the shelf were evaluated based on NORPAC net samples collected during summers of 1994–2009. During the study period, regime shifts occurred from high interannual variability regime (1994-1999) to low interannual variability regime with high temperature (2000-2005), then to a low interannual variability regime with low temperature (2007-2009). A total of 24 calanoid copepod species belonging to 21 genera were identified from samples. Copepod abundance ranged from 150 to 834,486 inds. m^{-2} , was greatest on the Middle shelf, and was higher in cold years, than in warm years. Copepod biomass ranged from 0.013 to 150 g DM m^{-2} , and was also higher in cold years than in warm years. Based on the results of cluster analysis, the copepod community was divided into six groups (A–F). The regional and interannual distributions of each group were distinct. Interannual changes in abundance of the dominant copepod on the Outer shelf and Middle shelf were highly significant ($p < 0.0001$), and their abundances were negatively correlated with temperature and salinity. Interannual changes in copepod community that occurred between cold and warm years are thought to have been caused by differences in the magnitude and timing of the spring phytoplankton bloom between the two regimes.

Abundance and biomass of the chaetognath *Parasagitta elegans* ranged from 30 to 15,180 inds. m⁻² and from 11 to 1,559 mg DM m⁻², respectively. Chaetognath abundance was significantly correlated with the abundance of the dominant copepods ($p < 0.0001$). Differences in cold and warm years may also affect recruitment of walleye pollock. We conclude that on the southeastern Bering Sea shelf, the magnitude and timing of primary production, which is related to climate change, may significantly affect how it is transferred through the food web.

Keywords: Annual variations; Abundance; Biomass; Zooplankton; Community composition; Species diversity

Regional terms: Southeastern Bering Sea, Alaska, USA

1. Introduction

The southeastern Bering Sea shelf is characterized by high biological productivity, has large amounts of phytoplankton, zooplankton, benthos, fishes, seabirds and marine mammals, and is an important fishing ground of walleye pollock (*Theragra chalcogramma*) (Springer, 1992; Springer et al., 1996). Recently, this region has undergone large biological/environmental changes (Jin et al., 2009; Stabeno et al., 2010). In 1997, a coccolithophore (*Emiliania huxleyi*) bloom was observed (Sukhanova and Flint, 1998; Napp and Hunt, 2001; Stockwell et al., 2001), and mass mortality of short-tailed shearwaters (*Puffinus tenuirostris*) occurred near the Pribilof Islands (Baduini et al., 2001). In 1998, a climate regime shift was reported (McFarlane et al., 2000), and the biomass of the jellyfish *Chrysaora melanaster* was observed to peak (Brodeur et al., 2008). To describe the effect of climate regime shifts on lower to higher trophic levels, Hunt et al. (2002) proposed the Oscillation Control Hypothesis (OCH). In 2006, a climate regime shift from a warm period to cold period was reported, and the OCH was revised based on observations during this period (Hunt et al., 2011).

In the Bering Sea ecosystem, mesozooplankton are important because they connect primary production to higher trophic level production. In the mesozooplankton fauna of this region, the dominant taxon is Copepoda, and various studies have been conducted on their biology and ecology. They include studies of seasonal variations of community structure during spring and summer (Smith and Vidal, 1986), interannual variations in their total wet-weight biomass (Sugimoto and Tadokoro,

1997, 1998; Napp et al., 2002; Hunt et al., 2008), seasonal and spatial variations in community structure (Kang et al., 2006), and the effects of El Niño and La Niña on biomass and production (Coyle and Pinchuk, 2002). However, most studies of long-term variations in mesozooplankton have treated only wet-weight biomass data without examining the species caught or separating sub-regions or domains. In studies of species and zooplankton community structure, the study areas have been narrow, and the study periods have been short. Consequently, information on long-term changes in mesozooplankton community and their species composition is extremely scarce.

The second most abundant taxonomic group in the Bering Sea mesozooplankton biomass is chaetognaths (Coyle and Pinchuk, 2002). They are carnivores, prey upon small zooplankton and fish larvae, contribute to vertical fluxes to deep water (Brodeur and Terazaki, 1999), and sometimes account for nearly 50% of the prey of walleye pollock larvae (Brodeur and Wilson, 1996). Thus, chaetognaths are important because of their role linking lower to higher trophic levels. However, little is known about their long-term variation in the Bering Sea.

In the present study, regional and long-term changes in abundance, biomass and community structure of copepods and chaetognaths are evaluated based on samples collected over the southeastern Bering Sea shelf during the summers of 1994–2009. Long-term data on copepods and chaetognaths are compared with hydrographic data, and we discuss possible mechanisms controlling interannual variations in mesozooplankton community structures and species composition.

2. Material and methods

2-1. Field sampling

Sampling was conducted over the southeastern Bering Sea shelf during 1994–2009 between 24 June and 8 August on board the T/S *Oshoro-Maru* of the Faculty of Fisheries, Hokkaido University. For each year, samples were collected within a two week period. Zooplankton samples were collected by vertical hauls of flowmeter-equipped NORPAC nets (45 cm mouth diameter, 0.335 mm mesh; Motoda, 1957) from 150 m depth or near the bottom (where the depth was shallower than 150 m) to the surface. Zooplankton samples were immediately preserved with 5% formaldehyde-seawater buffered with sodium tetraborate. In addition, temperature and salinity were measured with a CTD (Neil Brown, Mark IIIB during 1994–2001 and Seabird SBE-911*plus* during 2002–2009). Water samples from the CTD rosette were filtered through GF/F filters, extracted with acetone or DMF and chlorophyll *a* concentration was measured using a fluorometer (Turner Designs, Inc.).

The number of sampling stations in any one year was 9–49 (total = 96), and the total number of samples was 428 (Fig. 1). Based on bottom depth, the southeastern Bering Sea shelf was divided into three regions: Inner shelf (< 50 m), Middle shelf (50–100 m) and Outer shelf (100–200 m) (Coachman and Charnell, 1979; Coachman, 1986). Areas with depth of 200–1000 m and > 1000 m were classified as Slope and Basin, respectively. Six stations on the Outer shelf and six on the Middle shelf were sampled every year of the study (Fig. 1), so only the data from these twelve stations were used to analyze long-term changes.

2-2. Sample analysis

Based on the biomass of the samples, subsamples were created with a Motoda plankton splitter (Motoda, 1959), and the subsamples were examined under a stereomicroscope.

Copepods were staged and identified to the lowest possible taxonomic level. For copepods, the most important genus in this region (*Calanus*) is reported to be mixture of *C. glacialis* and *C. marshallae* (Nelson et al., 2009). We could not distinguish the two species and treated them as “*Calanus* spp.”. For biomass estimation, copepods were separated into four groups: *Calanus* spp., *Neocalanus* spp., *Eucalanus bungii* and other copepods. One hundred individuals in each group were chosen randomly for measurement of total lengths (TL, μm) by ocular micrometer. Dry mass (DM, μg) was estimated using the following equations derived for organisms from the Oyashio region (Imao, 2005):

$$\text{Calanus and Neocalanus spp. (less lipids)} \quad \log_{10} \text{DM} = 2.48 \log_{10} \text{TL} - 6.00 \quad (1)$$

$$\text{Calanus and Neocalanus spp. (more lipids)} \quad \log_{10} \text{DM} = 3.00 \log_{10} \text{TL} - 7.70 \quad (2)$$

$$\text{Eucalanus bungii} \quad \log_{10} \text{DM} = 3.16 \log_{10} \text{TL} - 9.16 \quad (3)$$

$$\text{Other copepods} \quad \log_{10} \text{DM} = 2.62 \log_{10} \text{TL} - 6.40 \quad (4)$$

For *Calanus* spp. and *Neocalanus* spp., lipid accumulation was separated into two: less lipids (<20% of prosome volume) and more lipids ($\geq$ 20% of prosome volume). Then the equations (1) and (2) were applied for less and more lipid specimens, respectively. The total biomass of copepods (mg DM m^{-2}) was estimated by multiplying the average individual mass (mg DM ind.^{-1}) by the abundance (inds. m^{-2}).

Chaetognaths were identified to species based on the samples collected during 1996–2009. The total length (TL, mm) of the dominant chaetognath, *Parasagitta elegans*, was measured using a ruler for large individuals (TL ≥ 10 mm) and an ocular micrometer for small specimens (TL < 10 mm). Specimens were classified according to McLaren (1969), Sameoto (1973) and Zo (1973): Stage I (juveniles), Stage II (immature) and Stage III (mature). To estimate DM, ash-free dry mass (AFDM, mg) was estimated from the TL using the following equation (Mumm, 1991):

$$\text{AFDM} = 0.0002 \text{ TL}^{2.6924} \quad (5)$$

Dry mass was estimated as $\text{DM} = \text{AFDM} / 0.9$ (Båmstedt, 1986).

2-3. Data analysis

Zooplankton samples were collected day and night. Since the depths of most sampling stations were < 150 m and the sampling was conducted throughout the water column, day-night differences in abundance and biomass were expected to be small. There were no significant differences between day and night abundance or biomass of copepods and chaetognaths for all years ($p = 0.06\text{--}0.93$, *U*-test). Thus, no day-night conversions were done for abundance or biomass.

To examine community structure, we conducted cluster analysis and non-metric multidimensional scaling (NMDS) ordination. Abundance data (X : inds. m^{-2}) of each species were transformed to $\log_{10}(X+1)$ prior to analysis to reduce the bias of abundant species. Rare species which occurred only <2% of the whole stations were eliminated from the data for analysis (cf. Table 1). Similarities between samples

were examined by Bray-Curtis index (Bray and Curtis, 1957) according to the differences in species composition. For grouping the samples, the similarity indices were coupled with hierarchical agglomerative clustering with a complete linkage method. The NMDS ordination was carried out to delineate the sample groups on the two-dimensional map. All of these analyses were carried out using BIOSTAT II software (Sigma Soft).

To evaluate environmental factors that may have influenced sample groups determined by cluster analysis, nonmetric multi-dimensional scaling (NMDS) ordination was carried out to delineate the sample groups in two-dimensional space. We then conducted multiple-regression analysis ($Y = aX_1 + bX_2 + c$) with the hydrographic data (Y : latitude, longitude, depth, sea surface and bottom temperature and sea surface and bottom salinity) are the dependent variables and two-dimensional NMDS as independent variables (thus, axis 1 as X_1 and axis 2 as X_2) (a , b , c are fitted constants). Shannon's diversity index (H') in each group was calculated using the equation:

$$H' = -\sum n_i/N \times \ln n_i/N \quad (6)$$

where n_i is abundance (inds. m⁻²) of the i th species, and N is total copepod abundance (inds. m⁻²) in the group (Shannon and Weaver, 1949).

To evaluate factors affecting the variability of copepod abundance, two-way ANOVA with year and station as independent variables was used. To determine which factors controlled annual changes in copepod abundance, correlation analysis was done between the abundance of dominant copepods and three factors: water temperature

(throughout whole water column, in upper layer and in lower layer), salinity and chlorophyll *a* (whole water column).

For *P. elegans*, cohort analysis was done based on the TL histogram data from the Middle shelf using Microsoft Excel Solver (Aizawa and Takiguchi, 1999).

To evaluate what factors controlled the interannual changes in chaetognath abundance, we tested for correlations between the chaetognath abundance for the Outer shelf and Middle shelf and four factors: water temperature (whole water column, upper and lower layer), salinity (whole water column), chlorophyll *a* (whole water column) and abundance of the dominant copepods.

3. Results

3-1. Hydrography

In all years, a thermocline was present 20–30 m, and the bottom temperature of Middle shelf ranged between 0° and 4°C, but was higher for the Inner and Outer shelves (Fig. 2A). The Middle shelf, bottom temperature was about 4°C during 2001–2005, and lower (0–3°C) during 2007–2009. Owing to Stabeno et al. (2012), there were three regimes in depth-averaged ocean temperature: cold; 1995, 1997, 1999, 2007-2009, average; 1996, 2000, 2006, and warm; 1998, 2001-2005. Our results confirmed this pattern (Fig. 2A).

The peak of summer chlorophyll *a* at each station varied between 0.5 and 25.4 mg m⁻³, and chlorophyll *a* was high above the thermocline and on the Inner shelf (Fig. 2B). The peak of chlorophyll *a* was low (ca. 1 mg m⁻³) during the warm regime

(2001–2005).

3-2. Copepods

Regional and interannual changes in total copepod abundance and biomass were observed (Fig. 3). Copepod abundance ranged from 150 to 834,486 inds. m⁻², and was greatest on the Middle shelf (Fig. 3A). Copepod abundance was high during cold years (1995, 1997, 1999 and 2007–2009), and low in warm years (2001–2005) (Fig. 3A). Copepod biomass ranged from 0.013 to 150 g DM m⁻², and was highest on the Outer and Middle shelves (Fig. 3B). The biomass peak occurred in more oceanic waters than the abundance peak due to the dominance of large-sized oceanic copepods. Copepod biomass and abundance were both high in cold years and low in warm years.

Throughout the study period, 24 calanoid copepod species from 21 genera were identified (Table 1). *Pseudocalanus* spp. was the most numerous, and *Calanus* spp. was dominated the copepod biomass. Large-sized oceanic copepods that diapause at depth (i.e., *E. bungii*, *M. pacifica*, *N. cristatus*, *N. flemingeri* and *N. plumchrus*) were collected on the Outer shelf, on the Slope and in the Basin (Table 1).

Based on abundance of each species, cluster analysis divided the copepod community into six groups (A–F). The groups contained 24–127 stations (Fig. 4A). From multiple-regression analysis, environmental factors that were significantly related with the group separation were latitude, longitude, depth, sea surface temperature, sea surface salinity and bottom salinity. The most important factors were latitude and depth; they determined 70–77% of the variability of the copepod community (Fig. 4B).

Mean copepod abundance was highest in group D and lowest in group E (Fig. 4C). Species diversity was higher in groups A and C ($H' = 1.6$) than in groups B, D, E and F ($H' = 1.0\text{--}1.1$). These group-specific differences in species diversity were caused by their species composition. For groups A–C, *Pseudocalanus* spp. and large, oceanic copepods (*M. pacifica*, *E. bungii*, *Neocalanus* spp.) were abundant (Fig. 4C). For groups D and E, species diversity was lower, and the species compositions were similar; they were comprised of *Pseudocalanus* spp. (55–58%), *Acartia* spp. (24%) and *Calanus* spp. (18–19%). In group F, the proportion of *Calanus* spp. decreased, and the main species were *Pseudocalanus* spp. (69%), and *Acartia* spp. (18%). Endemic species, such as *Epilabidocera amphitrites*, *Eurytemora herdmani* and *Tortanus discaudatus*, were also observed (Fig. 4C, Table 1).

The regional and interannual distributions of each group were clearly separated (Fig. 5A). Composition of group A was greater in cold years (1995, 1997, 1999 and 2007–2009) on the Outer shelf, and that of group B was greater in warm years (1998, 2001–2005) on the Outer shelf, and group C was seen mainly on the Outer shelf and in the Basin, but also on the Middle shelf in 2006–2008 (Fig. 5B). Group D was mainly observed in cold years on the Middle shelf, group E was seen in warm years on the Middle shelf, and group F was centered on the Inner shelf (Fig. 5A).

Two-way analysis of variance (ANOVA) on annual and geographical differences in abundance of the dominant copepods showed significant interannual changes for *Acartia* spp., *M. pacifica* and *Neocalanus* spp. on the Outer shelf (Table 2). Significant station-to-station differences were detected only for *Calanus* spp. on the

243 Middle shelf. All dominant copepods on the Middle shelf (*Acartia* spp., *Calanus* spp.
244 and *Pseudocalanus* spp.) showed highly significant interannual changes in abundance (p
245 < 0.0001 , Table 2).

246 Correlation analysis was used to evaluate which environmental parameters
247 were related to the interannual changes in copepod abundance. Abundances of the
248 three dominant copepods on the Middle shelf, which all showed high interannual
249 variability, were strongly negatively correlated with temperature and salinity (Table 3).

250 3-3. Chaetognaths

251 Two chaetognath species (*P. elegans* and *Eukrohnia hamata*) were collected.
252 *Parasagitta elegans* dominated the chaetognath fauna, composing 84% of the total
253 chaetognath abundance and 89% of the chaetognath abundance on the Middle shelf.
254 For this reason, *E. hamata* was not included in the subsequent analysis of chaetognaths.

255 *Parasagitta elegans* abundance ranged from 30 to 15,180 inds. m^{-2} , and was
256 highest on the Middle shelf (Fig. 6A). Abundance was low during 2000–2004 and
257 high during 1996–1999 and 2005–2009. *Parasagitta elegans* biomass varied from 11
258 to 1,559 mg DM m^{-2} (Fig. 6B). Biomass was low during 2002–2004 and high during
259 1996–1999 and 2005–2009. Differences in the timing of the abundance and biomass
260 peaks might have been caused by interannual changes in *P. elegans* body-size
261 composition.

262 In every year, *P. elegans* TL composition on the Middle shelf had two or three
263 cohorts (Fig. 7). The small-sized cohort was dominated by Stage I (juveniles), the

middle-sized cohort by Stage II (immature), and the large-sized cohort by Stage III (mature) individuals. Stage III was not observed in 2003 or 2004 (Fig. 7). The average TL of the small-sized cohort (Stage I) was significantly larger (3.5–8.5 mm) during 1996–2005 than those (2.5–5.5 mm) during 2006–2009 (Fig. 7) ($p < 0.05$, U -test).

Parasagitta elegans abundance on the Outer shelf was positively correlated with dominant copepod abundance (*Acartia* spp., *Calanus* spp. and *Pseudocalanus* spp.), but not with environmental factors (Table 4). On the Middle shelf, *P. elegans* had a highly significant positive correlation with dominant copepod abundance ($p < 0.0001$) and significant negative correlations with temperature and salinity. Thus on the Middle shelf, *P. elegans* abundance was high when temperature and salinity were low, and low when temperature and salinity were high, which is similar to the pattern observed in copepod abundance (Table 4).

4. Discussion

The copepod community on the Bering Sea shelf showed large interannual differences across the shelf (Fig. 5A). The major environmental factors that were related to the copepod community were latitude and depth (Fig. 4B), and species diversity was highest on the Outer shelf because of the occurrence of the oceanic copepods in addition to the shelf copepods (Fig. 4C). These findings correspond well with the horizontal distribution of copepod community structure reported by Cooney and Coyle (1982). These cross-shelf differences in copepod species may be related to the occurrence of

frontal structures (Coachman and Charnell, 1979; Coachman, 1986). The large number of oceanic species on the Middle shelf in 1998 (Fig. 5C) may be caused by the transportation of oceanic copepods by strong cross-shelf advection (Stabeno et al., 2001) (Fig.1). Stabeno et al. (2012) also reported the transport of oceanic species during warm years.

Long-term changes in zooplankton on the southeastern Bering Sea shelf based upon wet-weight biomass have been reported (Sugimoto and Tadokoro, 1997, 1998; Napp et al., 2002). Recently, Hunt et al. (2008) also reported that the zooplankton wet mass was lower in the early 2000s than in the late 1990s. This corresponds with the interannual changes in copepod abundance and biomass observed in the present study (Fig. 3). In the present study, the copepod community greatly varied between cold years (1995, 1997, 1999 and 2007-2009) and warm years (1998, 2001–2005) on both the Outer and Middle shelves. In warm years, oceanic species decreased on the Outer shelf, but showed no change on the Middle shelf, and the abundance of all copepod species drastically decreased on both the Outer and Middle shelf (Fig. 5). Thus, these findings clearly indicate that both the wet mass and copepod community structure showed interannual changes on the southeastern Bering Sea shelf (Fig. 8).

Physical and biological changes have occurred in the southeastern Bering Sea shelf over the last two decades (Fig. 8). During the 1997 El Niño/Southern Oscillation (ENSO) event, there were anomalous atmospheric conditions over the southeastern Bering Sea shelf (Overland et al., 2001), a warm surface layer, a bloom of the coccolithophorid *Emiliana huxleyi* (Sukhanova and Flint, 1998) and mass mortality of

short-tailed shearwaters (*P. tenuirostris*) (Baduini et al., 2001). Short-tailed shearwaters prey on large zooplankton, especially adult euphausiids (Hunt et al., 1996), and the starvation of shearwaters that occurred during the summer of 1997 may have been due to low densities of adult euphausiids (Stockwell et al., 2001) or difficulty in finding prey patches due to low water transparency caused by the coccolithophore blooms (Lovvorn et al., 2001). Thus, the mass mortality of short-tailed shearwaters may not have been related with the pattern of mesozooplankton abundance and biomass described in this study.

Coccolithophore (*E. huxleyi*) blooms in recent times were first observed in this region in September, 1997 (Vance et al., 1998). Large-scale coccolithophore blooms were also observed in June, 1998 and 2000 (Sukhanova and Flint, 1998; Iida et al., 2002, 2008, 2012). As an ecological consequence of coccolithophores, negative effect for short-tailed shearwaters and their nutritional values for copepods should be considered. Information in the literature regarding ingestion of coccolithophores by copepods is contradictory. For example, Huskin et al. (2000) state that *E. huxleyi* is difficult to digest and has low nutritional value, it is not a preferred food item for *Calanus* species. However, Nejstgaard et al. (1997) reported that about 75% of the carbon consumed by *Calanus finmarchicus* during an *E. huxleyi* bloom came from *E. huxleyi* ($> 30 \times 10^6$ cells L^{-1}), and high abundance of *Calanus* spp. was observed on the Inner shelf during a coccolithophore bloom (Coyle and Pinchuk, 2002). The latter observations suggest that coccolithophore production may have little effect on *Calanus* spp. In fact, years of coccolithophore blooms (1997, 1998 and 2000) did not correspond with years of low

copepod abundance or biomass (Fig. 8B). This discrepancy between coccolithophore blooms and copepod abundance could be explained if coccolithophorids are ingested by *Calanus* species (Nejstgaard et al., 1997) and have some nutritional value. As the other possible cause, turbid waters of coccolithophore bloom may provide copepods refuge from visual predators (Lovvorn et al., 2001).

During the study period, there were cold (1995, 1997, 1999, 2007-2010), average (1996, 2000, 2006) and warm years (1998, 2001-2005) (Fig. 2A). During cold and warm years, the timing and magnitude of the spring bloom differ (Stabeno et al., 2001; Hunt et al., 2002, 2011). During cold years, the sea ice remains during the severe winter storms, and ice-edge blooms occur. In warm years, the sea ice melts before the spring storms, and strong wind mixing delays the start of the bloom until stratification occurs when the thermocline develops. From long-term (1960–2005) simulation modeling, Jin et al. (2009) reported that in cold years ($PDO < -1$), phytoplankton blooms start early, last long and are large, while in warm years ($PDO > 1$), they occur later with short pulses and are small. These differences in bloom timing between cold and warm years have also been recorded in mooring observations in this region (Hunt et al., 2011).

Copepod abundance and biomass on both the Outer shelf and Middle shelf were high during cold years (1995, 1997, 1999 and 2007-2009) and low during warm years except 1998 (2001–2005) (Fig. 3, Table 3). The dominant copepod, *Calanus* spp., has an extended spawning period; it starts well before the spring phytoplankton bloom (February) and ends in May, however the survival rate of early developmental

stages depends on whether they encounter a phytoplankton bloom or not, and recruitment of copepodid stages increases if they encounter early ice associated blooms, which are highly productive (Baier and Napp, 2003). Low temperatures in cold years may also lead to greater abundance of the small copepods *Pseudocalanus* spp. and *Acartia* spp. especially in the Middle shelf (Table 3). Cold temperature may provide the longer growing season and high primary productivity of ice-edge blooms, sufficient food condition for copepods, resulting in increased abundance of both large and small copepods.

In this study, copepod abundance on both the Outer and Middle shelf was negatively correlated with temperature (Table 3). In contrast, on the Inner shelf, positive correlations were reported between spring temperatures and the abundance of three copepods (*Calanus* spp., *Pseudocalanus* spp. and *Acartia* spp.), and between summer temperatures and two copepods (*Pseudocalanus* spp. and *Acartia* spp.) (Coyle and Pinchuk, 2002). This suggests that the response of copepods to temperature may vary between the Outer-Middle shelf (negative, this study), and the Inner shelf (positive, Coyle and Pinchuk, 2002). In this study, differences between cold and warm years on the Middle shelf were observed in copepod abundance, but not in species composition (group D and E in Fig. 4C). This is partly because both large and small copepods were abundant in cold years and less abundant in warm years (Table 3). Coyle et al. (2008) compared the summer zooplankton community on the Middle shelf between a cold (1999) and warm year (2004), and reported that the dominant species shifted from large species in the cold year to small species in the warm year. However, this phenomenon

was not observed during the 16 years of the present study. Our results showed that both large and small-sized copepods showed a clear negative correlation with habitat temperature (Table 3).

Similar to copepods, the biomass of the jellyfish *Chrysaora melanaster* is high in cold years and low in warm years (Brodeur et al., 2008). The increase in cold years may be due to improved feeding conditions for polyps (mass sinking of phytoplankton after ice-edge bloom), and subsequent abundant food (copepods) for pelagic medusae may improve survival and growth (Brodeur et al., 2008). In contrast, oceanic blooms in warm years are less productive and provide less food (sinking phytoplankton) for polyps, as well as less food (copepods) for planktonic medusae, resulting in low growth and survival of polyps and medusae.

Interannual changes in chaetognath abundance and biomass were somewhat similar to the changes in copepods (Fig. 8). This may be an example of bottom-up control on predatory chaetognath abundance by prey (copepods) abundance. Chaetognaths in this region change their food items and size at maturity between cold and warm years; in cold years, they prey on large *Calanus* spp. and mature at large size, while in warm years, they feed on small *Pseudocalanus* spp. and mature at small size (Baier and Terazaki, 2005). The TL of most mature (Stage III) individuals in the present study was > 20 mm in cold years (1997, 1999 and 2007-2009), and < 20 mm in warm years (1998, 2001, 2002 and 2005), and no mature individuals were observed in the warm years 2003 and 2004 (Fig. 7). These facts support the finding of Baier and Terazaki (2005), who reported that sufficient food conditions in cold years result in a

large size at maturity, while food limitation in warm years induces small size at maturity.

For the early life history stages of walleye pollock, an important fisheries resource in this region, food conditions differ between cold and warm years (Napp et al., 2000; Hunt et al., 2011). In cold years, there may be a mismatch between the production of pollock larvae and their prey (Napp et al., 2000), but an early start of the ice-edge bloom and high abundance of lipid-rich prey for age-0 fish (copepods and euphausiids) results in high survival rates and low cannibalism (Hunt et al., 2011; Heintz et al., this issue). In warm years, the late start of the open-water bloom results in a low abundance of lipid-rich prey for the age-0 pollock and chaetognaths. Under this food-limited condition, growth and survival of walleye pollock juveniles decrease, and the proportion of cannibalism may increase. This results in a decrease in recruitment (Hunt et al., 2011, Heintz et al., this volume). Thus, the timing and magnitude of primary production related with climate change affects production through the entire food web structure on the southeastern Bering Sea shelf.

Acknowledgements

We thank Dr. J. R. Bower of Hokkaido University for his comments on the manuscript. All samples and hydrographic data used in this study were collected by the T/S *Oshoro-Marui*. We are grateful to the captain and crew for their help. Some of the chaetognath data were collected by Mr. N. Hagimoto. This study was supported by Grant-in-Aid for Scientific Research (A) 24248032 and Grant-in-Aid for Scientific

416 Research on Innovative Areas 24110005 from the Japan Society for the Promotion of
417 Science (JSPS).

References

- Aizawa, Y., Takiguchi, N., 1999. Consideration of the methods for estimating the age-composition from the length frequency data with MS-Excel. Bull. Jpn. Soc. Fish. Oceanogr. 63, 205–214 (in Japanese).
- Baduini, C. L., Hyrenbach, K. D., Coyle, K. O., Pinchuk, A., Mendenhall, V., Hunt, G. L. Jr., 2001. Mass mortality of short-tailed shearwaters in the south-eastern Bering Sea during summer 1997. Fish. Oceanogr. 10, 117–130.
- Baier, C. T., Napp, J. M., 2003. Climate-induced variability in *Calanus marshallae* populations. J. Plankton Res. 25, 771–782.
- Baier, C. T., Terazaki, M., 2005. Interannual variability in a predator-prey interaction: climate, chaetognaths and copepods in the southeastern Bering Sea. J. Plankton Res. 27, 1113–1125.
- Båmstedt, U., 1986. Chemical composition and energy content, in: Corner, E. D. S., O'Hara, S. C. M. (Eds.), The Biological Chemistry of Marine Copepods. Clarendon Press, Oxford, pp. 1–58.
- Bray, J. R., Curtis, J. T., 1957. An ordination of the upland forest communities of southern Wisconsin. Ecol. Monog. 27, 325–349.
- Brodeur, R. D., Decker, M. B., Ciannelli, L., Purcell, J. E., Bond, N. A., Stabeno, P. J., Acuna, E., Hunt, G. L. Jr., 2008. Rise and fall of jellyfish in the eastern Bering Sea in relation to climate regime shifts. Prog. Oceanogr. 77, 103–111.
- Brodeur, R. D., Terazaki, M., 1999. Springtime abundance of chaetognaths in the shelf region of the northern Gulf of Alaska, with observations on the vertical

440 distribution and feeding of *Sagitta elegans*. Fish. Oceanogr. 8, 93–103.

441 Brodeur, R. D., Wilson, M. T., 1996. A review of the distribution, ecology and
 442 population dynamics of age-0 walleye pollock in the Gulf of Alaska. Fish.
 443 Oceanogr. 5, 148–166.

444 Coachman, L. K., 1986. Circulation, water masses, and fluxes on the southeastern
 445 Bering Sea shelf. Cont. Shelf Res. 5, 23–108.

446 Coachman, L. K., Charnell, R. L., 1979. On lateral water mass interaction-A case
 447 study, Bristol Bay, Alaska. J. Phys. Oceanogr. 6, 278–297.

448 Cooney, R. T., Coyle, K. O., 1982. Trophic implications of cross-shelf copepod
 449 distributions in the southeastern Bering Sea. Mar. Biol. 70, 187–196.

450 Coyle, K. O., Pinchuk, A. I., 2002. Climate-related differences in zooplankton density
 451 and growth on the inner shelf of the southeastern Bering Sea. Prog. Oceanogr.
 452 55, 177–194.

453 Coyle, K. O., Pinchuk, A. I., Eisner, L. B., Napp, J. M., 2008. Zooplankton species
 454 composition, abundance and biomass on the eastern Bering Sea shelf during
 455 summer: The potential role of water-column stability and nutrients in structuring
 456 the zooplankton community. Deep-Sea Res. II 55, 1775–1791.

457 Hunt, G. L. Jr., Allen, B.M., Angliss, R.P., Baker, T., Bond, N., Buck, G., Byrd, G.V.,
 458 Coyle, K.O., Devol, A., Eggers, D.M., Eisner, L., Feely, R., Fitzgerald, S., Fritz,
 459 L.W. Gritsay, E.V., Ladd, C., Lewis, W., Mathis, J., Mordy, C.W., Mueter, F.,
 460 Napp, J., Sherr, E., Shull, D., Staben, P., Stepanenko, M.A., Strom, S.,
 461 Whitley, T.E., 2010. Status and trends of the Bering Sea region, 2003-2008,

462 in: McKinnell, S. M. and Dagg, M. J. (Eds.), Marine Ecosystems of the North
 463 Pacific Ocean, 2003–2008. PICES Special Publication 4, pp. 196–267.

464 Hunt, G. L. Jr., Coyle, K. O., Eisner, L. B., Farley, E. V., Heintz, R. A., Mueter, F.,
 465 Napp, J. F., Overland, J. E., Ressler, P. H., Salo, S., Stabeno, P. J., 2011.
 466 Climate impacts on eastern Bering Sea food webs: a synthesis of new data and
 467 an assessment of the Oscillating Control Hypothesis. ICES J. Mar. Sci. 68,
 468 1230–1243.

469 Hunt, G. L. Jr., Coyle, K. O., Hoffman, S., Decker, M. B., Flint, E. N., 1996. Foraging
 470 ecology of short-tailed shearwaters near the Pribilof Islands, Bering Sea. Mar.
 471 Ecol. Prog. Ser. 141, 1–11.

472 Hunt, G. L. Jr., Stabeno, P. J., Strom, S., Napp, J.M., 2008. Patterns of spatial and
 473 temporal variation in the marine ecosystem of the southeastern Bering Sea, with
 474 special reference to the Pribilof Domain. Deep-Sea Res. II 55, 1919–1944.

475 Hunt, G. L. Jr., Stabeno, P., Walters, G., Sinclair, E., Brodeur, R. D., Napp, J. M., Bond,
 476 N. A., 2002. Climate changes and control of the southeastern Bering Sea
 477 pelagic ecosystem. Deep-Sea Res. II 49, 5821–5853.

478 Huskin, I., Anadón, R., Álvarez-Marqués, F., Harris, R. P., 2000. Ingestion, fecal
 479 pellet and egg production rates of *Calanus helgolandicus* feeding
 480 coccolithophorid versus non-coccolithophorid diets. J. Exp. Mar. Biol. Ecol.
 481 248, 239–254.

482 Iida, T., Mizobata, K., Saitoh, S. I., 2008. Temporal and spatial variability of
 483 phytoplankton in the Bering Sea –From diatom bloom in spring to

484 coccolithophore bloom in summer and fall– Kaiyo Monthly Extra 50, 127–137
 485 (in Japanese).

486 Iida, T., Mizobata, K., Saitoh, S. -I., 2012. Interannual variability of coccolithophore
 487 *Emiliania huxleyi* blooms in response to changes in water column stability in the
 488 eastern Bering Sea. Cont. Shelf Res. 34, 7–17.

489 Iida, T., Saitoh, S. -I., Miyamura, T., Toratani, M., Fukushima, H., Shiga, N., 2002.
 490 Temporal and spatial variability of coccolithophore blooms in the eastern Bering
 491 Sea, 1998–2001. Prog. Oceanogr. 55, 165–177.

492 Imao, F., 2005. Zooplankton community structure and their functional role in carbon
 493 cycle in the Oyashio region, western subarctic Pacific. Master thesis,
 494 Hokkaido University, 42 pp.

495 Jin, M., Deal, C., Wang, J., McRoy, C. P., 2009. Response of lower trophic level
 496 production to long-term climate change in the southeastern Bering Sea. J.
 497 Geophys. Res. 114, C04010, doi: 10.1029/2008JC005105.

498 Kang, Y., Kim, S., Lee, W., 2006. Seasonal and spatial variations of zooplankton in
 499 the central and southeastern Bering Sea during the mid-1990s. Deep-Sea Res. I
 500 53, 795–803.

501 Lovvorn, J. R., Baduini, C. L., Hunt, G. L. Jr., 2001. Modeling underwater visual and
 502 filter feeding by planktivorous shearwaters in unusual sea conditions. Ecology
 503 82, 2342–2356.

504 McFarlane, G. A., King, J. R., Beamish, R. J., 2000. Have there been recent changes
 505 in climate? Ask the fish. Prog. Oceanogr. 47, 147–169.

506 McLaren, I. A., 1969. Population and production ecology of zooplankton in Ogac
 507 Lake, a landlocked Fjord on Baffin Island. J. Fish. Res. Bd. Canada 26,
 508 1485–1559.

509 Miller, C. B., Frost, B. W., Batchelder, H. P., Clemons, M. J., Conway, R. E., 1984. Life
 510 histories of large, grazing copepods in a Subarctic Ocean Gyre: *Neocalanus*
 511 *plumchrus*, *Neocalanus cristatus*, and *Eucalanus bungii* in the Northeast Pacific.
 512 Prog. Oceanogr. 13, 201–243.

513 Motoda, S., 1957. North Pacific standard plankton net. Inf. Bull. Planktol. Japan 4,
 514 13–15.

515 Motoda, S., 1959. Devices of simple plankton apparatus. Mem. Fac. Fish. Hokkaido
 516 Univ. 7, 73–94.

517 Mumm, N., 1991. On the summer distribution of mesozooplankton in the Nansen
 518 Basin, Arctic Ocean. Rep. Polar Res. 92, 1–146.

519 Napp, J. M., Baier, C. T., Brodeur, R. D., Coyle, K. O., Shiga, N., Mier, K., 2002.
 520 Interannual and decadal variability in zooplankton communities of the southeast
 521 Bering Sea shelf. Deep-Sea Res. II 49, 5991–6008.

522 Napp, J. M., Hunt, G. L., 2001. Anomalous conditions in the south-eastern Bering Sea
 523 1997: linkage among climate, weather, ocean, and biology. Fish. Oceanogr. 10,
 524 61–68.

525 Napp, J. M., Kendall Jr., A. W., Schumacher, J. D., 2000. A synthesis of biological and
 526 physical processes affecting the feeding environment of larval walleye pollock
 527 (*Theragra chalcogramma*) in the eastern Bering Sea. Fish. Oceanogr. 9,

528 147-162.

529 Nejstgaard, J. C., Gismervik, I., Solberg, P. T., 1997. Feeding and reproduction by
530 *Calanus finmarchicus*, and microzooplankton grazing during mesocosm blooms
531 of diatoms and the coccolithophore *Emiliana huxleyi*. Mar. Ecol. Prog. Ser.
532 147, 197–217.

533 Nelson, R. J., Carmack, E. C., McLaughlin, F. A., Cooper, G. A., 2009. Penetration of
534 Pacific zooplankton into the western Arctic Ocean tracked with molecular
535 population genetics. Mar. Ecol. Prog. Ser. 381, 129-138.

536 Overland, J.E., Bond, N.A., Adams, J.M., 2001. North Pacific atmospheric and SST
537 anomalies in 1997: links to ENSO? Fish. Oceanogr. 10, 69–80.

538 Sameoto, D. D., 1973. Annual life cycle and production of the chaetognath *Sagitta*
539 *elegans* in Bedford Basin, Nova Scotia. J. Fish. Res. Bd. Canada 30, 333–344.

540 Shannon, C. E., Weaver, W., 1949. The Mathematical Theory of Communication. The
541 University of Illinois Press, Urbana.

542 Smith, S. L., Vidal, J., 1986. Variations in the distribution, abundance, and
543 development of copepods in the southeastern Bering Sea in 1980 and 1981.
544 Cont. Shelf Res. 5, 215–239.

545 Springer, A. M., 1992. A review: Walleye pollock in the North Pacific-how much
546 difference do they really make? Fish. Oceanogr. 1, 80–96.

547 Springer, A. M., McRoy, C. P., Flint, M. V., 1996. The Bering Sea Green Belt:
548 shelf-edge processes and ecosystem production. Fish. Oceanogr. 5, 205–223.

549 Stabeno, P. J., Bond, N. A., Kachel, N. B., Salo, S. A., Schumacher, J. D., 2001. On

550 the temporal variability of the physical environment over the south-eastern
551 Bering Sea. *Fish. Oceanogr.* 10, 81–98.

552 Stabeno, P. J., Napp, J., Mordy, C., Whitledge, T., 2010. Factors influencing physical
553 structure and lower trophic levels of the eastern Bering Sea shelf in 2005: ice,
554 tides and winds. *Prog. Oceanogr.* 85, 180–196.

555 Stabeno, P. J., Kachel, N. B., Moore, S. E., Napp, J. M., Sigler, M., Yamaguchi, A.,
556 Zerbini, A. N., 2012. Comparison of warm and cold years on the southeastern
557 Bering Sea shelf and some implications for the ecosystem. *Deep-Sea Res. II*
558 65-70, 31-45.

559 Stockwell, D. A., Whitledge, T. E., Zeeman, S. I., Coyle, K. O., Napp, J. M., Brodeur, R.
560 D., Pinchuk, A. I., Hunt, G. L. Jr., 2001. Anomalous conditions in the
561 south-eastern Bering Sea, 1997: nutrients, phytoplankton and zooplankton.
562 *Fish. Oceanogr.* 10, 99–116.

563 Sugimoto, T., Tadokoro, K., 1997. Interannual-interdecadal variations in zooplankton
564 biomass, chlorophyll concentration and physical environment in the subarctic
565 Pacific and Bering Sea. *Fish. Oceanogr.* 6, 74–93.

566 Sugimoto, T., Tadokoro, K., 1998. Interdecadal variations of plankton biomass and
567 physical environment in the North Pacific. *Fish. Oceanogr.* 7, 289–299.

568 Sukhanova, I. N., Flint, M. V., 1998. Anomalous blooming of coccolithophorids over
569 the eastern Bering Sea self. *Oceanology* 38, 502–505.

570 Vance, T. C., Schumacher, J. D., Stabeno, P. J., Baier, C. T., Wyllie-Echeverria, T.,
571 Tynan, C. T., Brodeur, R. D., Napp, J. M., Coyle, K. O., Decker, M. B., Hunt, G.

572 L., Jr., Stockwell, D., Whitledge, T. E., Jump, M., Zeeman, S., 1998.
573 Aquamarine waters recorded for first time in Eastern Bering Sea. Eos, Trans.
574 Am. Geophys. Union 79, 121-126.
575 Zo, Z., 1973. Breeding and growth of the chaetognath *Sagitta elegans* in Bedford
576 Basin. Limnol. Oceanogr. 18, 750–756.
577

Figure captions

Fig. 1. Location of the sampling stations in the southeastern Bering Sea during the summers of 1994–2009 (circles). Solid symbols denote stations where sampling was conducted in all years from 1994–2009, providing the data used for annual comparisons. Line indicates 166°W transect where the hydrographic data were analyzed (cf. Fig. 2).

Fig. 2. (A) Temperature and (B) chlorophyll *a* stations along 166°W during summers of 1994–2009.

Fig. 3. Regional variations in (A) copepod abundance and (B) biomass in the southeastern Bering Sea during summers of 1994–2009.

Fig. 4. Copepod assemblages: (A) dendrograms from the cluster analysis based on copepod abundance. Six groups (A–F) were identified with dissimilarity at 20% (dashed line). Numbers in the parentheses indicate quantity of stations included in each group. (B) Nonmetric multi-dimensional scaling plots of the six groups, with arrows and percentages indicating directions of environmental parameters and coefficient of determinations (r^2), respectively. Lat: latitude, Lon: longitude, SSS: sea surface salinity, SST: sea surface temperature, BS: bottom salinity. (C) Mean abundance with standard deviations and species diversity (H') (upper) and taxonomic composition of each group (lower).

Fig. 5. (A) Spatial and temporal distribution of six copepod communities identified from Bray-Curtis dissimilarity analysis. Annual distribution of groups on the (B) Outer shelf and (C) Middle shelf.

600 Fig. 6. Regional variations in *Parasagitta elegans*: (A) abundance and (B) biomass
601 during summers of 1996–2009.

602 Fig. 7. Annual patterns of *Parasagitta elegans* length frequency on Middle shelf
603 during summers of 1996–2009. n: number of measured specimens.

604 Fig. 8. (A) Annual changes in biological/environmental conditions, (B) copepod
605 abundance and biomass and (C) chaetognath (*Parasagitta elegans*) abundance
606 and biomass on Middle shelf of the southeastern Bering Sea during summers of
607 1994–2009. Values are means and error bars indicate standard deviations.
608 For references in (A), 1: Sukhanova and Flint (1998), Iida et al. (2002, 2008), 2:
609 Baduini et al. (2001), 3: McFarlane et al. (2000), 4: Brodeur et al. (2008), Hunt
610 et al. (2010), 5: Stabenho et al. (2012).

Table 1. Calanoid copepod species collected in the southeastern Bering Sea during summers of 1994–2009. Major region of occurrence is also shown for each species. ●: Large oceanic copepods that have a diapause phase in deep layer (Miller et al., 1984). Region abbreviations are; I: Inner shelf, M: Middle shelf, O: Outer shelf, S: Slope and B: Basin. *: Rare species (occurred only <2% of the whole stations) omitted from data for cluster analysis (cf. Fig. 4).

Species	Major region of occurrence
<i>Acartia longiremis</i> (Lilljeborg, 1853)	I, M, O
<i>A. tumida</i> Willey, 1920	M, O
* <i>Aetideopsis rostrata</i> Sars, 1903	B
* <i>Aetideus pacificus</i> Brodsky, 1950	B
<i>Calanus</i> spp.	M, O, S, B
<i>Candacia columbiae</i> Campbell, 1929	S, B
<i>Centropages abdominalis</i> Sato, 1913	I, M
<i>Epilabidocera amphitrites</i> (McMurrich, 1916)	I
● <i>Eucalanus bungii</i> Giesbrecht, 1892	O, S, B
<i>Eurytemora herdmanni</i> Thompson and Scott, 1897	I
<i>Gaetanus intermedius</i> Wolfenden, 1905	S, B
<i>G. simplex</i> Brodsky, 1950	S, B
<i>Heterorhabdus tanneri</i> (Giesbrecht, 1895)	S, B
<i>Metridia pacifica</i> Brodsky, 1950	O, S, B
<i>Microcalanus pygmaeus</i> (Sars, 1900)	M, O
● <i>Neocalanus cristatus</i> (Kröyer, 1848)	O, S, B
● <i>N. flemingeri</i> Miller, 1988	O, S, B
● <i>N. plumchrus</i> (Marukawa, 1921)	O, S, B
<i>Paraeuchaeta elongata</i> (Esterly, 1913)	B
<i>Pleuromamma scutullata</i> Brodsky, 1950	S, B
<i>Pseudocalanus</i> spp.	I, M, O, S, B
* <i>Racovitzanus antarcticus</i> Giesbrecht, 1902	B
<i>Scolecithricella minor</i> (Brady, 1883)	O, S, B
* <i>S. ovata</i> (Farran, 1905)	B
<i>Tortanus discaudatus</i> (Thompson and Scott, 1897)	I
<i>Undinopsis pacificus</i> Brodsky, 1950	O, S

Table 2. Variance analysis (two-way ANOVA) on year-to-year and station-to-station differences in abundance (inds. m⁻²) of dominant copepod taxa on Outer and Middle shelf of the southeastern Bering Sea during summers of 1994–2009.

*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$, ns: not significant.

Domain	Species	<i>F</i> -value <i>df</i> ₁ , <i>df</i> ₂	
		Year _{15,30}	Station _{2,30}
Outer shelf			
	<i>Acartia</i> spp.	2.136*	1.215 ^{ns}
	<i>Calanus</i> spp.	1.259 ^{ns}	3.027 ^{ns}
	<i>Eucalanus bungii</i>	1.402 ^{ns}	2.610 ^{ns}
	<i>Metridia pacifica</i>	2.060*	1.464 ^{ns}
	<i>Neocalanus</i> spp.	5.228***	1.956 ^{ns}
	<i>Pseudocalanus</i> spp.	1.190 ^{ns}	0.194 ^{ns}
Middle shelf			
	<i>Acartia</i> spp.	6.574****	2.798 ^{ns}
	<i>Calanus</i> spp.	6.026****	3.745*
	<i>Pseudocalanus</i> spp.	5.791****	0.195 ^{ns}

Table 3. Correlation coefficients (*r*) between abundance of dominant copepods and various environmental parameters in the southeastern Bering Sea during summers of 1994–2009.

*: *p* < 0.05, **: *p* < 0.01, ***: *p* < 0.001, ****: *p* < 0.0001.

Domain Species	Temperature (n= 91)			Salinity (n= 91)	Chlorophyll <i>a</i> (n=45)
	Whole water column	Upper layer	Lower layer	Whole water column	Whole water column
Outer shelf					
<i>Acartia</i> spp.	0.207 *	0.149	0.186	-0.140	0.067
<i>Calanus</i> spp.	-0.423 ****	-0.101	-0.406 ***	-0.294 **	0.192
<i>Eucalanus bungii</i>	0.138	-0.143	0.307 **	0.517 ****	0.071
<i>Metridia pacifica</i>	-0.231 *	-0.258 *	-0.105	0.076	0.370 *
<i>Neocalanus</i> spp.	-0.285 **	-0.404 ***	-0.108	0.252 *	0.369 *
<i>Pseudocalanus</i> spp.	-0.159	-0.142	-0.044	-0.028	0.341 *
Middle shelf					
<i>Acartia</i> spp.	-0.410 ***	-0.146	-0.431 ****	-0.506 ****	-0.111
<i>Calanus</i> spp.	-0.448 ****	-0.340 **	-0.457 ****	-0.407 ***	-0.126
<i>Pseudocalanus</i> spp.	-0.567 ****	-0.292 **	-0.539 ****	-0.484 ****	-0.037

Table 4. Correlation coefficients (r) between abundance of chaetognath *Parasagitta elegans* and various parameters (water temperature, salinity and copepod abundance) on the Outer shelf and Middle shelf in the southeastern Bering Sea during summers of 1996–2009.

*: $p < 0.05$, **: $p < 0.01$, ***: $p < 0.001$, ****: $p < 0.0001$. —: not applicable.

Parameters	Outer shelf	Middle shelf
Temperature		
Whole water column	-0.137	-0.519 ****
Upper layer	-0.057	-0.301 **
Lower layer	-0.187	-0.547 ****
Salinity		
Whole water column	-0.172	-0.513 ****
Copepods		
<i>Acartia</i> spp.	0.343 **	0.563 ****
<i>Calanus</i> spp.	0.431 ****	0.503 ****
<i>Eucalanus bungii</i>	0.129	—
<i>Metridia pacifica</i>	0.195	—
<i>Neocalanus</i> spp.	0.093	—
<i>Pseudocalanus</i> spp.	0.291 **	0.566 ****

Fig. 1. Ohashi et al.

Fig. 2. Ohashi et al.

Fig. 3. Ohashi et al.

Fig. 4. Ohashi et al.

Fig. 5. Ohashi et al.

Fig. 6. Ohashi et al.

Fig. 7. Ohashi et al.

Fig. 8. Ohashi et al.