Title	Variation in assimilation efficiencies of dominant Neocalanus and Eucalanus copepods in the subarctic Pacific : consequences for population structure models
Author(s)	Abe, Yoshiyuki; Natsuike, Masafumi; Matsuno, Kohei; Terui, Takeshi; Yamaguchi, Atsushi; Imai, Ichiro
Citation	Journal of experimental marine biology and ecology, 449, 321-329 https://doi.org/10.1016/j.jembe.2013.10.023
Issue Date	2013-11
Doc URL	http://hdl.handle.net/2115/54053
Type	article (author version)
File Information	Abe-et-al.pdf

[Instructions for use](#)

Variation in assimilation efficiencies of dominant *Neocalanus* and *Eucalanus* copepods in the
subarctic Pacific: consequences for population structure models

Yoshiyuki Abe^{a*}, Masafumi Natsuike^a, Kohei Matsuno^{a,b}, Takeshi Terui^{b,c}, Atsushi Yamaguchi^a and
Ichiro Imai^a

^a*Graduate School of Fisheries Sciences, Hokkaido University, 3-1-1 Minatomachi, Hakodate,
Hokkaido 041-8611, Japan*

^b*National Institute of Polar Research, 10-3, Midoricho, Tachikawa, Tokyo 190-8518, Japan*

^c*Core Research for Evolutional Science and Technology, Japan Science and Technology Agency, 5
Sanbancho, Chiyoda-ku, Tokyo 102-0075, Japan*

* Corresponding author. Tel.: +81 138 40 5543; fax: +81 138 40 5542.

E-mail address: y.abe@fish.hokudai.ac.jp

Abstract

The assimilation efficiency of zooplankton is an essential parameter required to estimate energy transfer to higher trophic levels in marine ecosystems. However, little information is available for large oceanic copepods, especially the *Neocalanus* and *Eucalanus* species dominant in the subarctic Pacific. In this study, the assimilation efficiencies of the C5 stages of *N. cristatus*, *N. flemingeri* and *E. bungii* were evaluated using eight phytoplankton species as food. The average assimilation efficiencies of *N. cristatus*, *N. flemingeri* and *E. bungii* ranged between 45- 66%, 44 - 66% and 34-65%, respectively. The assimilation efficiency was highly variable depending on the food phytoplankton species. In all species, the assimilation efficiency showed a significant negative relationship with the ash content of the phytoplankton ($r^2 = 0.79 - 0.87$, $p < 0.001$). The assimilation efficiency of large-body sized *N. cristatus* for large-sized diatoms was higher than for the other copepod species. In population models of *N. cristatus*, changes in assimilation efficiency affect the growth and survival rates of the population. The Lagrangian ensemble model (LEM) for *N. cristatus* showed that, for assimilation efficiencies less than 57%, the population could not be maintained. Because variations in assimilation efficiency may have significant effects on the copepod population, their variability should be incorporated into marine ecosystem models in the future.

Keywords: Assimilation efficiency, Copepods, Diatoms, Lagrangian Ensemble Model, *Neocalanus*

31 *cristatus*

1. Introduction

In marine ecosystems, copepods feed on phyto- and microzooplankton and are preyed upon by pelagic fishes, sea birds and whales; thus, they play an important role in energy transfers to higher trophic organisms (Nemoto, 1963; Hunt et al., 1998; Beamish et al., 1999; Ikeda et al., 2008).

In the summer zooplankton community of the subarctic Pacific, large oceanic copepods, *Neocalanus* and *Eucalanus* species, are predominant and form 85- 90% of the total zooplankton biomass (Vinogradov, 1970). Because *Neocalanus* and *Eucalanus* species egest large faecal pellets and perform seasonal vertical migrations, they play an important role in transporting organic material from the surface layer to the deep sea, called the “biological pump” (Kobari et al., 2003, 2008).

Among the various copepod parameters that affect the material flux, the assimilation efficiency is an essential parameter required to estimate the energy transfer to higher trophic levels in marine ecosystems (Conover, 1966a, b). From the 1960s to the present, many studies have been performed on the assimilation efficiencies of copepods. Through these studies, much information has been accumulated, e.g., assimilation efficiency varies with the ash content of the food (Conover, 1966a, b), and the carbon assimilation efficiency is correlated with the concentrations of soluble carbohydrates in diets (Head, 1992). However, our knowledge about assimilation efficiency has mainly come from coastal species (cf. Gottfried and Roman, 1983; Berggreen et al., 1988; Besiktepe and Dam, 2002; Katechakis et al., 2004), and little information is available for large oceanic

copepods, especially the *Neocalanus* and *Eucalanus* species that are dominant in the subarctic Pacific. Even in marine ecosystem models such as NEMURO, a constant value (70%) is applied for the copepod assimilation efficiency (Kishi et al., 2007; Terui and Kishi, 2008; Terui et al., 2012). As mentioned above, a few large oceanic copepods (*Neocalanus* and *Eucalanus* species) are dominant in the zooplankton biomass of the subarctic Pacific; therefore, information on their assimilation efficiency is very important for increasing the accuracy of ecosystem models (e.g., NEMURO) in this region.

The assimilation efficiency of copepods is measured by several methods; the Ratio method, the Radio tracer method and the Calculation method, based on ingestion rate, evacuation rate and faecal pellet volume (Conover, 1966a; Sorokin, 1968; Besiktepe and Dam, 2002). Among these methods, the Radio tracer method is mainly used for minimally metabolised materials, such as heavy metals. For common materials (e.g., carbon and phosphorus), the Radio tracer method cannot be applied because they are readily metabolised, which prevents accurate measurement (Båmstedt et al., 2000). The Calculation method requires the quantitative collection of faecal pellets, so this method is also difficult (Conover and Francis, 1973; Omori and Ikeda, 1984). The Ratio method by Conover (1966a) does not require quantitative collection of faecal pellets, is readily applicable to various animals and is an effective method, even though nearly half a century has passed since it was originally described (Azad et al., 2011; Enríquez-Ocaña et al., 2012; Nelson et al., 2012). Recently,

most studies on assimilation efficiency of copepods have mainly concerned marine heavy metal pollution in coastal small copepods as evaluated using the Radio tracer method (Fisher and Reinfelder, 1991; Hutchins et al., 1995; Wang and Fisher, 1998; Chang and Reinfelder, 2000; Xu and Wang, 2001, 2002; Stewart and Fisher, 2003; Wang et al., 2007; Zheng et al., 2011). However, little information is available on the assimilation efficiencies of common materials (organic material, carbon and nitrogen) by large oceanic copepods.

In this study, the assimilation efficiencies of three large oceanic copepods (*N. cristatus*, *N. flemingeri* and *E. bungii*), which are dominant in the zooplankton biomass in the subarctic Pacific, were measured by applying the Ratio method considering eight phytoplankton species (diatoms, a dinoflagellate and a raphidophycean) as food. Phytoplankton cell size, colony formation, swimming ability and ash contents were also analysed as factors that may affect the assimilation efficiency of copepods. The effects of food carbon concentration on copepod assimilation efficiency were also evaluated. By applying the observed assimilation efficiency of one species of copepod (*N. cristatus*), the effects of changes in the assimilation efficiency on copepod population structure were evaluated using the LEM (Lagrangian ensemble model) population model.

2. Materials and methods

Live specimens of the C5 stages of *N. cristatus*, *N. flemingeri* and *E. bungii* were collected

using vertical hauls of a 80-cm ring net from 0-30 or 0-150 m deep at several stations in the subarctic Pacific from March to July 2011 and May to August 2012. Because adult (*C6*) *Neocalanus* spp. degrade feeding appendages and cease feeding (Miller et al., 1984; Miller, 1988), we used the *C5* stage as experimental specimens. Seawater was collected from 30 m deep using Niskin bottles, filtered through a GF/F filter and used in the subsequent experiments. Ten live specimens were transferred into a 1-L bottle filled with filtered seawater (FSW). Up to 100 specimens of each species were kept at 2°C and then carried to the land laboratory.

To obtain sympatric phytoplankton species, 5 ml of unfiltered seawater was added to a flask containing 300 ml of modified SWM-3 medium (Chen et al., 1969; Imai et al., 1996), then incubated at 15°C under illumination of 100 to 120 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ with a 14 h light and 10 h dark photocycle. Three diatoms (*Chaetoceros* sp., *Ditylum brightwellii* and *Thalassiosira nordenskiöldii*) were isolated from this treatment. Six diatoms (*Attheya septentrionalis*, *Chaetoceros* sp., *D. brightwellii*, *Pauliella taeniata*, *Skeletonema* sp. and *Th. nordenskiöldii*), one dinoflagellate (*Alexandrium tamarense*) and one raphidophycean (*Heterosigma akashiwo*) were incubated under the same conditions as food for the copepods. The carbon contents of *Chaetoceros* sp., *D. brightwellii* and *Skeletonema* sp. were estimated by multiplying 0.43 (for the former two species) or 0.51 (for the latter species) with the ash-free dry weight (AFDW) (Parsons et al., 1961). For *Th. nordenskiöldii* and *A. tamarense*, the carbon contents were obtained by multiplying 0.108 or

0.173 with the dry weight (DW), respectively (Liu and Wang, 2002). Information regarding cell size, carbon and ash contents, colony formation, swimming ability and initial cell density of each phytoplankton is summarised in Table 1.

Before the experiments, no food was added for the copepod specimens for at least one day (24 h). For the experiments, each phytoplankton species was adjusted to a density of 5.0×10^2 - 2.0×10^4 cells ml^{-1} (110 - 2577 $\mu\text{g C L}^{-1}$) (Table 1). For each experiment, 15 individuals of *N. cristatus* or *E. bungii* or 20 individuals of *N. flemingeri* were added to each phytoplankton species in 1-L bottles and incubated for 24 hours under dark conditions at 3°C . Experiments were carried out in triplicate along with one control bottle with no added copepods. During the experiments, the bottles were rotated every 3 h to prevent the phytoplankton from sinking. After the 24-hour experiment, the copepods were transferred to new bottles containing FSW. Faecal pellets were pipetted from the incubation bottles using sterile Pasteur pipettes, placed in petri dishes filled with chilled FSW and rinsed 5-10 times by immersion in FSW to avoid contamination of the remaining phytoplankton cells. Then, the faecal pellets were checked under a stereomicroscope. The faecal pellet volume (FPV , μm^3) was quantified using the following equation:

$$FPV = \frac{4}{3} \pi \left(\frac{FPW}{2} \right)^2 \left(\frac{FPL}{2} \right) \quad (1)$$

where FPW and FPL are faecal pellet width and length in μm , respectively. Both FPL and FPW were measured to a precision of $10 \mu\text{m}$ under a dissecting microscope fitted with an eye-piece

121 micrometer.

122 Previously, GF/F filters were combusted at 480°C for 5 h, then weighed with a
 123 microbalance (Mettler Toledo MT5) to a precision of 1 µg. Food phytoplankton and faecal pellets
 124 were gently filtered with pre-weighed GF/F filter, briefly rinsed with small amount of distilled water
 125 and then dried at 60°C for 5 h. Their DW was then measured with a microbalance. Dried filters
 126 were combusted at 480°C for 5 h, and the ash weight (*ASH*) was measured with a microbalance.
 127 The organic contents of the phytoplankton and faecal pellets were calculated as $DW - ASH$. The
 128 assimilation efficiency was determined by the Ratio method (Conover 1966a, b):

$$129 \quad U' = [(F' - E') / (1 - E') (F')] \times 100 \quad (2)$$

130 where U' is the assimilation efficiency (%), F' is the organic fraction of the food, and E' is the
 131 organic fraction of the faecal pellets.

132 To evaluate differences among species in the assimilation efficiencies and phytoplankton
 133 ash contents, an ANCOVA was applied using the copepod species and phytoplankton ash contents as
 134 independent variables.

135 The relationship between assimilation efficiency and phytoplankton carbon concentration
 136 was analysed as another factor that controls the copepod assimilation efficiency. To express the
 137 nonlinear regression models between assimilation efficiency and carbon concentration, a generalised
 138 additive model (GAM) was applied using the free software “R” and the multivariate smoothing

parameter estimation package “mgcv”.

3. Results

The faecal pellet volumes observed in this study are summarised in Table 2. The faecal pellet volumes varied with phytoplankton species, and their means were 12.23-49.67, 3.77-5.95 and $5.20-21.52 \times 10^6 \mu\text{m}^3$ for *N. cristatus*, *N. flemingeri* and *E. bungii*, respectively (Table 2). These values corresponded well with the predicted values from the prosome length of each copepod species predicted by Mauchline’s equation (1998): 38.88, 5.64 and $16.06 \times 10^6 \mu\text{m}^3$ for *N. cristatus*, *N. flemingeri* and *E. bungii*, respectively (Table 2).

Experiments on the assimilation efficiency of *N. cristatus* were performed with eight phytoplankton species. The assimilation efficiency of *N. cristatus* ranged between 45% and 66% and varied with the phytoplankton species. The highest and lowest assimilation efficiencies were observed for the diatoms *P. taeniata* and *Th. nordenskiöldii*, respectively (Fig. 1).

Many specimens of C5 *N. flemingeri* moulted to C6 during the experiments. If moulting was observed, the whole replicate was thrown out. Because of this limitation, assimilation efficiency data for this species were only obtained for three diatom species. The assimilation efficiency of *N. flemingeri* ranged between 44% and 66% and varied with the phytoplankton species. The highest and lowest values of assimilation efficiency were recorded for the diatoms *Skeletonema*

sp. and *D. brightwellii*, respectively (Fig. 2).

Experiments on *E. bungii* were performed using four phytoplankton species. The assimilation efficiency of *E. bungii* ranged between 34% and 65% and varied with the phytoplankton species. The highest and lowest values of assimilation efficiency were recorded for the diatoms *Skeletonema* sp. and *D. brightwellii*, respectively (Fig. 3).

A common trend for the three copepod species was that the assimilation efficiencies had a significant negative correlation with the ash contents of the food phytoplankton ($r^2 = 0.79 - 0.87$, $p < 0.001$, Fig. 4A-C), and the AFDW content showed no relationship with assimilation efficiency (Fig. 4D-F). From the ANCOVA analysis with assimilation efficiency as the dependent variable and covariate with the ash content of phytoplankton, the effect of the interaction of copepods with ash contents was highly significant (Table 3). Thus, the parallelism of the regression lines for the three copepods (Fig. 4A-C) was rejected (i.e., the slopes of the regression lines varied with species).

The assimilation efficiency was stable for carbon concentrations in the range of 100-830 $\mu\text{g C L}^{-1}$ in the phytoplankton species, and it decreased with increasing carbon concentration above 830 $\mu\text{g C L}^{-1}$ (Fig. 5).

4. Discussion

According to Båmstedt et al. (2000), potential sources of bias in the Ratio method are the

following: (1) non-homogenous food material, (2) food selectivity, (3) sloppy feeding, (4) losses from faecal material, (5) absorbance of inert tracers in the digestive tract and (6) production of non-faecal material mixed with faeces. In the present study, because we applied only one phytoplankton species as food for each experiment, sources (1) and (2) seem to be eliminated. For (3) - (6), we rotated the treated incubation bottles, pipetted faecal pellets by sterile Pasteur pipette, rinsed the pellets 5-10 times within FSW and then checked the faecal pellets under stereomicroscope. Because the FPV observed in this study corresponded well with the predicted values (Table 2), we believe that the effects of coprophagy or coprohexy were negligible, and the biases on the Ratio method were minimised for this study.

The most important result of this study is that the assimilation efficiencies of the three oceanic copepods showed highly significant negative correlations with the ash content of the food phytoplankton (Fig. 4A-C). These findings correspond well with a previous study (Conover, 1966b). Among the phytoplankton species, the highest ash content was observed for diatoms (cf. Fig. 1). As a specialised characteristic of diatoms, their cell walls are made of silica. Although silica is ingested by copepods, 79-90% of the silica is egested as faecal pellets (Tande and Slagstad, 1985; Conover et al., 1986; Cowie and Hedges, 1996). Because copepods do not utilise silica, the highly significant negative relationship between copepod assimilation efficiency and ash content of phytoplankton is a common pattern. The slope (b) of the regression ($Y = a + bX$) between copepod

assimilation efficiency (Y : %) and the ash content of the phytoplankton (X : %) in this study (-0.72) is very close to that found by Conover (1966b) (-0.73). Conover (1966b) mainly used the large oceanic copepod *Calanus hyperboreus* as the experimental species, which is dominant in the northern North Atlantic. *N. cristatus* can be considered as the counterpart species in the North Pacific of *C. hyperboreus* in the North Atlantic (Parsons and Lalli, 1988), and the similar slopes of the regression formula between this study (*N. cristatus*) and Conover (1966b) (*C. hyperboreus*) indicate that they have similar ecological roles.

To clarify the factors that determine the copepod assimilation efficiency, we calculated the adjusted copepod assimilation efficiency (U'_{adj} : %) with phytoplankton ash contents using the following equation (Sokal and Rohlf, 2012):

$$U'_{adj} = U' - b(1 - F') \times 100 \quad (4)$$

where U' is the assimilation efficiency (%), $(1 - F')$ is the ash fraction of the phytoplankton, and b is the fitted constant of the regression for each copepod species (cf. Fig. 4A-C). The relationships between the adjusted copepod assimilation efficiency and each parameter are listed in Table 1 (i.e., phytoplankton cell size, colony formation and movement ability) and were analysed along with copepod body size (Ueda et al., 2008) (Fig. 6). For phytoplankton cell size, both *N. flemingeri* and *E. bungii* had high assimilation rates for small-sized *Skeletonema* sp. (2-21 μ m) and low assimilation rates for large-sized *D. brightwellii* (25-100 μ m). A negative correlation between

assimilation efficiency and phytoplankton cell size was detected ($p < 0.001$) (Fig. 6A). However, changes in assimilation efficiency with the phytoplankton cell size were not detected for *N. cristatus* (Fig. 6A). This is because the assimilation efficiency of the large-sized *D. brightwellii* was higher for *N. cristatus* than for the other two species (Fig. 6C). Thus, the large-sized copepod *N. cristatus* had a high assimilation efficiency for large-sized phytoplankton, so there was no relationship between the assimilation efficiency and phytoplankton cell size for *N. cristatus* (Fig. 6A).

Colony formation and movement ability are closely related among the phytoplankton species treated in this study, i.e., colony-forming species have no movement ability (diatoms), whereas non-colony-forming species have movement ability (dinoflagellates and raphidophyceans) (Table 1). In this study, a significant difference was only detected for *N. cristatus*: the assimilation efficiency was higher for non-colony-forming phytoplankton with movement ability (dinoflagellates and raphidophyceans) than for the colony-forming phytoplankton with no movement ability (diatoms) (Fig. 6B). However, this pattern was not detected for the other two copepods, which was partly because the number of experiments with dinoflagellates and raphidophyceans was extremely limited ($n = 1$) for these species (Fig. 6B). The limitation pattern of *N. cristatus* for this study (assimilation efficiency of dinoflagellates was higher than those of diatoms, $p < 0.01$, Fig. 6B) corresponds well with the results of previous studies (Conover, 1966a; Besiktepe and Dam, 2002).

Concerning the effects of food concentration on assimilation efficiency, Conover (1966b) reported that the assimilation efficiency did not change, regardless of the food carbon concentration. However, it is recognised that the assimilation efficiency decreases with increasing food concentration (Gaudy, 1974; Landry et al., 1984; Besiktepe and Dam, 2002; Thor and Wendt, 2010). The decrease of assimilation efficiency under high food concentrations ($> 830 \mu\text{g C L}^{-1}$) was also the case in this study (Fig. 5). Two factors are considered as possible causes of the decrease of assimilation under high food concentrations: (1) the shortened gut passage time under high food concentration (Besiktepe and Dam, 2002) may prevent sufficient digestion and assimilation (Lehman, 1976; Landry et al., 1984) or (2) changes in the activities of digestive enzymes with food concentration (high under low food concentration) (Hassett and Landry, 1983), where the digestive enzyme activity decreases under high food concentrations, may reduce the assimilation efficiency (Landry et al., 1984). Based on this information, Pahlow and Prome (2010) created an ecosystem model that incorporates the decrease of assimilation efficiency with increasing food concentration. Montagnes and Fenton (2012) also developed an ecosystem model in which assimilation efficiency varied with food concentration and compared it to the model with constant assimilation efficiency. Because the decrease of assimilation efficiency under high food carbon concentrations was observed for an anomalously high food concentration in the experimental condition (Fig. 5), it should be questioned whether this phenomenon would occur under natural food concentrations in the field.

Applying the same carbon contents of food is recommended for future studies of incubation experiments on assimilation efficiency.

Marine ecosystem models, such as NPZD models (e.g., NEMURO), PDM (population dynamics model) and LEM, apply a constant assimilation efficiency (70%) for zooplankton (mainly considering copepods) (Kishi et al., 2007; Terui et al., 2012). This study, however, showed that the assimilation efficiency of large oceanic copepods was lower, in the range of 34%-66%, and varied depending on food ash contents. The low assimilation efficiency observed in this study may be caused by the food applied (phytoplankton). Under natural food conditions, these copepods may prefer to feed on microzooplankton rather than on phytoplankton (cf. Dagg et al., 2009 and references therein). The assimilation efficiency of microzooplankton may be higher than the phytoplankton, especially for the diatoms, which is why the observed assimilation efficiency of this study (34-66%) was lower than the commonly used value (70%).

Next, we tested the effects of changes in assimilation efficiency by applying the LEM for *N. cristatus* (Terui et al., 2012). In this test, the environmental settings (e.g., water temperature and solar radiation) were the same as in the original model, and only the assimilation efficiency (P6 in LEM of Terui et al., 2012) was changed. We ran a model using an assimilation efficiency (as 70% in the original model) between 45% and 66% (i.e., the range experimentally observed for *N. cristatus*, Fig. 1). After a run of 50 years, the results reached a steady state after more than 51 years

and are shown in Fig. 7. Under the 66% assimilation efficiency, *N. cristatus* could maintain the population (Fig. 7B). However, *N. cristatus* could not maintain its population under the lowest 44% assimilation efficiency (Fig. 7C). When the runs changed for every 1%, it was demonstrated that *N. cristatus* could not maintain its population for < 57% assimilation efficiency, and the survival of the population was possible for > 58% assimilation efficiency. In addition to population survival, changes in assimilation efficiency affected development time, i.e., 139 days were required for individuals born on 22 February to reach C5 (solid) under 70% assimilation efficiency, and 150 days were required for same hatch date individuals under 66% assimilation efficiency. Thus, the changes in assimilation efficiency have significant effects on copepod population survival and growth and variations in assimilation efficiency should be incorporated into marine ecosystem models in the future. Because copepod assimilation efficiency is highly significantly correlated with the inorganic content of phytoplankton (Fig. 4), the assimilation efficiency in the model should be estimated using parameters based on the composition of the food phytoplankton taxa.

Acknowledgements

We thank three anonymous reviewers of our manuscript. Their comments were helpful and greatly improved the manuscript. We acknowledge the captains, officers, crew and researchers on board T/S *Oshoro-Maru* from Hokkaido University for their efforts during field sampling. Part

280 of this study was supported by a Grant-in-Aid for Scientific Research (A) 24248032 and a
281 Grant-in-Aid for Scientific Research on Innovative Areas 24110005 from the JSPS.

References

- Azad, A.K., Pearce, C.M., McKinley, R.S., 2011. Effects of diet and temperature on ingestion, absorption, assimilation, gonad yield, and gonad quality of the purple sea urchin (*Strongylocentrotus purpuratus*). *Aquaculture* 317, 187-196.
- Båmstedt, U., Gifford, D.J., Irigoien, X., Atkinson, A., Roman, R., 2000. 8 Feeding. In: Harris, R.P., Wiebe, P.H., Lenz, J., Skjoldal, H.R., Huntley, M. (Eds.), *ICES Zooplankton Methodology Manual*. Academic Press, pp. 297-399.
- Beamish, R.J., Leask, K.D., Ivanov, O.A., Balanov, A.A., Orlov, A.M., Sinclair, B., 1999. The ecology, distribution and abundance of mid water fishes of the Subarctic Pacific gyres. *Prog. Oceanogr.* 43, 399-442.
- Berggreen, U., Hansen, B., Kiørboe, T., 1988. Food size spectra, ingestion and growth of the copepod *Acartia tonsa* during development: implications for determination of copepod production. *Mar. Biol.* 99, 341-352.
- Besiktepe, S., Dam, H.G., 2002. Coupling of ingestion and defecation as a function of diet in the calanoid copepod *Acartia tonsa*. *Mar. Ecol. Prog. Ser.* 229, 151-164.
- Chang, S.I., Reinfelder, J.R., 2000. Bioaccumulation, subcellular distribution, and trophic transfer of copper in a coastal marine diatom. *Environ. Sci. Technol.* 34, 4931-4935.
- Chen, L.C.M., Edelstein, T., McLachlan, J., 1969. *Bonnemaisonia hamifera* Harih in nature and in culture. *J. Phycol.* 5, 211-220.
- Conover, R.J., 1966a. Assimilation of organic matter by zooplankton. *Limnol. Oceanogr.* 11, 338-345.
- Conover, R.J., 1966b. Factors affecting the assimilation of organic matter by zooplankton and the question of superfluous feeding. *Limnol. Oceanogr.* 11, 346-354.
- Conover, R.J., Francis, V., 1973. The use of radioactive isotopes to measure the transfer of materials

- 306 in aquatic food chains. Mar. Biol. 18, 272-283.
- 307 Conover, R.J., Durvasula, R., Roy, S., Wang, R., 1986. Possible loss of chlorophyll-derived pigments
308 during passage through the gut of zooplankton, and some of the consequences. Limnol.
309 Oceanogr. 31, 878-887.
- 310 Cowie, G.L., Hedges, J.I., 1996. Digestion and alteration of the biochemical constituents of a diatom
311 (*Thalassiosira weissflogii*) ingested by an herbivorous zooplankton (*Calanus pacificus*).
312 Limnol. Oceanogr. 41, 581-594.
- 313 Dagg, M., Strom, S., Liu, H., 2009. High feeding rates on large particles by *Neocalanus flemingeri*
314 and *N. plumchrus*, and consequences for phytoplankton community structure in the
315 subarctic Pacific Ocean. Deep-Sea Res. I 56, 716-726.
- 316 Enríquez-Ocaña, L.F., Nieves-S., M., Pina-Valdez, P., Martinez-Cordova, L.R., Medina-Jasso, M.A.,
317 2012. Evaluation of the combined effect of temperature and salinity on the filtration,
318 clearance rate and assimilation efficiency of the mangrove oyster *Crassostrea*
319 *corteziensis* (Hertlein, 1951). Arch. Biol. Sci. 64, 479-488.
- 320 Fisher, N.S., Reinfelder, J.R., 1991. Assimilation of selenium in the marine copepod *Acartia tonsa*
321 studied with a radiotracer ratio method. Mar. Ecol. Prog. Ser. 70, 157-164.
- 322 Gaudy, R., 1974. Feeding four species of pelagic copepods under experimental conditions. Mar. Biol.
323 25, 125-141.
- 324 Gottfried, M., Roman, M.R., 1983. Ingestion and incorporation of coral-mucus detritus by reef
325 zooplankton. Mar. Biol. 72, 211-218.
- 326 Hassett, R.P., Landry, M.R., 1983. Effects of food-level acclimation on digestive enzyme activities
327 and feeding behavior of *Calanus pacificus*. Mar. Biol. 75, 47-55.
- 328 Head, E.J.H., 1992. Comparison of the chemical composition of particulate material and copepod
329 fecal pellets at stations off the coast of Labrador and in the Gulf of St. Lawrence. Mar.

- 330 Biol. 112, 593-600.
- 331 Hunt, G.L., Russell, R.W., Coyle, K.O., Weingartner, T., 1998. Comparative foraging ecology of
 332 planktivorous auklets in relation to ocean physics and prey availability. Mar. Ecol. Prog.
 333 Ser. 167, 241–259.
- 334 Hutchins, D.A., Wang, W., Fisher, N.S., 1995. Copepod grazing and the biogeochemical fate of
 335 diatom iron. Limnol. Oceanogr. 40, 989-994.
- 336 Ikeda, T., Shiga, N., Yamaguchi, A., 2008. Structure, biomass distribution and trophodynamics of the
 337 pelagic ecosystem in the Oyashio region, western subarctic Pacific. J. Oceanogr. 64,
 338 339-354.
- 339 Imai, I., Itakura, S., Matsuyama, Y., Yamaguchi, M., 1996. Selenium requirement for growth of a
 340 novel red tide flagellate *Chattonella verruculosa* (Raphidophyceae) on culture. Fish. Sci.
 341 62, 834-835.
- 342 Katechakis, A., Steibor, H., Sommer, U., Hansen, T., 2004. Feeding selectivities and food niche
 343 separation of *Acartia clausi*, *Penilia avirostris* (Crustacea) and *Doliolum denticulatum*
 344 (Thaliacea) in Blanes Bay (Catalan Sea, NW Mediterranean). J. Plankton Res. 26,
 345 589-603.
- 346 Kishi, M.J., Kashiwai M., Ware D.M., Megrey B.A., Eslinger D.L., Werner F.E. et al., 2007.
 347 NEMURO—a lower trophic level model for the North Pacific marine ecosystem. Ecol.
 348 Model. 202, 12–25.
- 349 Kobari, T., Shinada, A., Tsuda, A., 2003. Functional roles of interzonal migrating mesozooplankton
 350 in the western subarctic Pacific. Prog. Oceanogr. 57, 279-298.
- 351 Kobari, T., Steinberg, D.K., Ueda, A., Tsuda, A., Silver, M.W., Kitamura, M., 2008. Impacts of
 352 ontogenetically migrating copepods on downward carbon flux in the western subarctic
 353 Pacific Ocean. Deep-Sea Res. II 55, 1648-1660.

- Landry, M.R., Hassett, R.P., Fagerness, V., Downs, J., Lorenzen, C.J., 1984. Effect of food acclimation on assimilation efficiency of *Calanus pacificus*. Limnol. Oceanogr. 29, 361-364.
- Lehman, J.T., 1976. The filter-feeder as an optimal forager, and the predicted shapes of feeding curves. Limnol. Oceanogr. 21, 501-516.
- Liu, S., Wang, W.X., 2002. Feeding and reproductive responses of marine copepods in South China Sea to toxic and nontoxic phytoplankton. Mar. Biol. 140, 595–603.
- Mauchline, J., 1998. The biology of calanoid copepods. Adv. Mar. Biol. 33, 1-710.
- Miller, C.B., Frost, B.W., Batchelder, H.P., Clemons, M.J., Conway R.E., 1984. Life histories of large, grazing copepods in a subarctic ocean gyre: *Neocalanus plumchrus*, *Neocalanus cristatus* and *Eucalanus bungii* in the North Pacific. Prog. Oceanogr. 13, 201-243.
- Miller, C.B., 1988. *Neocalanus flemingeri*, a new species of Calanidae (Copepoda: Calanoida) from the subarctic Pacific Ocean, with a comparative redescription of *Neocalanus plumchrus* (Marukawa) 1921. Prog. Oceanogr. 20, 223-273.
- Montagnes, D.J.S., Fenton, A., 2012. Prey-abundance affects zooplankton assimilation efficiency and the outcome of biogeochemical models. Ecol. Model. 243, 1-7.
- Nelson, E.J., MacDonald, B.A., Robinson, S.M.C., 2012. The absorption efficiency of the suspension-feeding sea cucumber, *Cucumaria frondosa*, and its potential as an extractive integrated multi-trophic aquaculture (IMTA) species. Aquaculture 370, 19-25.
- Nemoto, T. 1963. Some aspects of the distribution of *Calanus cristatus* and *C. plumchrus*, in the Bering Sea and its neighbouring waters, with reference to the feeding of baleen whales. Sci. Rep. Whale Res. Inst. 17, 157-170.
- Omori, M., Ikeda, T., 1984. Methods in Marine Zooplankton Ecology. John Wiley and Sons, New York.

- 378 Pahlow, M., Prome, A., 2010. Model of optimal current feeding in zooplankton. Mar. Ecol. Prog. Ser.
379 403, 129-144.
- 380 Parsons, T.R., Stephens, K., Strickland, J.D.H., 1961. On the chemical composition of eleven species
381 of marine phytoplankton. J. Fish. Res. Bd. Can. 18, 1001-1016.
- 382 Parsons, T.R., Lalli, C.M., 1988. Comparative oceanic ecology of the plankton communities of the
383 subarctic Atlantic and Pacific Oceans. Oceanogr. Mar. Biol. Annu. Rev. 26, 317-359.
- 384 Sokal, R.R., Rohlf, F.J., 2012. Biometry: the Principles and Practice of Statistics in Biological
385 Research. 4th Edition. W.H. Freeman, New York.
- 386 Sorokin, J.I., 1968. The use of ^{14}C in the study of the nutrition of aquatic animals. Mitt. Int. Ver.
387 Theor. Angew. Limnol. 16, 40-41.
- 388 Stewart, G.M., Fisher, N.S., 2003. Bioaccumulation of polonium-210 in marine copepods. Limnol.
389 Oceanogr. 48, 2011-2019.
- 390 Tande, K.S., Slagstad, D., 1985. Assimilation efficiency in herbivorous aquatic organisms - The
391 potential of the ratio method using ^{14}C and biogenic silica as markers. Limnol. Oceanogr.
392 30, 1093-1099.
- 393 Terui T., Kishi M.J., 2008. Population dynamics model of Copepoda (*Neocalanus cristatus*) in the
394 northwestern subarctic Pacific. Ecol. Model. 215, 77-88.
- 395 Terui, T., Kishi, M.J., Ueno, H., 2012. Lagrangian ensemble model of Copepoda (*Neocalanus*
396 *cristatus*) in the northwestern subarctic Pacific. J. Oceanogr. 68, 727-741.
- 397 Thor, P., Wendt, I., 2010. Functional response of carbon absorption efficiency in the pelagic calanoid
398 copepod *Acartia tonsa* Dana. Limnol. Oceanogr. 55, 1779-1789.
- 399 Ueda, A., Kobari, T., Steinberg, D.K., 2008. Body length, weight and chemical composition of
400 ontogenetically migrating copepods in the Western Subarctic Gyre of the North Pacific
401 Ocean. Bull. Plankton Soc. Japan 55, 107-114.

- 402 Vinogradov, M.E., 1970. Vertical Distribution of the Oceanic Zooplankton. Israel Program for
 403 Scientific Translations, Jerusalem.
- 404 Wang, W.X., Fisher, N.S., 1998. Accumulation of trace elements in a marine copepod. *Limnol.*
 405 *Oceanogr.* 43, 273-283.
- 406 Wang, M.H., Wang, D.Z., Wang, G.Z., Huang, X.G., Hong, H.S., 2007. Influence of N, P additions
 407 on the transfer of nickel from phytoplankton to copepods. *Environ. Pollution* 148,
 408 679-687.
- 409 Xu, Y., Wang, W., 2001. Individual responses of trace-element assimilation and physiological
 410 turnover by the marine copepod *Calanus sinicus* to changes in food quantity. *Mar. Ecol.*
 411 *Prog. Ser.* 218, 227-238.
- 412 Xu, Y., Wang, W., 2002. The assimilation of detritus-bound metals by the marine copepod *Acartia*
 413 *spinicauda*. *Limnol. Oceanogr.* 47, 604-610.
- 414 Zheng, Y., Dam, H.G., Avery, D.E., 2011. Differential responses of populations of the copepod
 415 *Acartia hudsonica* to toxic and nutritionally insufficient food algae. *Harmful Algae* 10,
 416 723–731.
- 417

Figure captions

Fig. 1. Assimilation efficiency (solid column) of *Neocalanus cristatus* C5 for various food phytoplankton. Experiments were performed in triplicate, and the bars indicate standard deviations. Open circles are the ash contents of the phytoplankton in each experiment. Horizontal lines indicate the means of the assimilation efficiencies. The percentage values in the panels indicate the means of the assimilation efficiencies.

Fig. 2. Assimilation efficiency (solid column) of *Neocalanus flemingeri* C5 for various food phytoplankton. Experiments were performed in triplicate, and the bars indicate standard deviations. Open circles are the ash contents of the phytoplankton in each experiment. Horizontal lines indicate the means of the assimilation efficiencies. The percentage values in the panels indicate the means of the assimilation efficiencies.

Fig. 3. Assimilation efficiency (solid column) of *Eucalanus bungii* C5 for various food phytoplankton. Experiments were performed in triplicate, and the bars indicate standard deviations. Open circles are the ash contents of the phytoplankton in each experiment. Horizontal lines indicate the means of the assimilation efficiencies. The percentage values in the panels indicate the means of the assimilation efficiencies.

Fig. 4. Relationships between assimilation efficiency and ash content of food phytoplankton: (A) *Neocalanus cristatus* C5, (B) *Neocalanus flemingeri* C5 and (C) *Eucalanus bungii* C5. Relationships between assimilation efficiency and ash free dry weight (AFDW) contents of food phytoplankton: (D) *N. cristatus* C5, (E) *N. flemingeri* C5 and (F) *E. bungii* C5. Regression equations are shown for each species.

Fig. 5. The effects of variable phytoplankton carbon concentrations on copepod assimilation efficiency for whole data (A), diatoms (B) and dinoflagellates (C). To express the nonlinear regression model between the assimilation efficiency and the carbon

concentration, a generalised additive model (GAM) was applied using the free software “R” and the multivariate smoothing parameter estimation package “mgcv”.

Fig. 6. Relationships between the adjusted assimilation efficiency and phytoplankton cell size (A), colony formation / movement ability (B) and copepod body size (C). **: $p < 0.01$, ns: not significant. NF: *Neocalanus flemingeri*, EB: *Eucalanus bungii*, NC: *Neocalanus cristatus*.

Fig. 7. Simulated biomass of each stage of *Neocalanus cristatus* by LEM from Terui et al. (2012) applying a modification of the assimilation efficiency to 70% (A), 66% (B) and 45% (C).

Table 1. Data on phytoplankton (cell size, carbon, ash contents, colony formation and movement ability) used as food for copepods in the laboratory experiments. Owing to size, cell density of phytoplankton was changed for assimilation experiments . For carbon and ash contents, values are mean $\pm$ 1 sd.

Taxa / Species	Cell size (μm)	Carbon content (pg C cell ⁻¹)	Ash content (pg Ash cell ⁻¹)	Colony formation	Movement ability	Concentration	
						(cells ml ⁻¹)	(μg C L ⁻¹)
Diatoms							
<i>Attheya septentrionalis</i>	5–10	–	190 ± 99	+	–	5.0 × 10 ³ –1.0 × 10 ⁴	–
<i>Chaetoceros</i> sp.	10–40	249 ± 3	754 ± 105	+	–	1.0 × 10 ³	249
<i>Ditylum brightwellii</i>	25–100	2596 ± 1540	9400 ± 582	+	–	5.0 × 10 ²	1298
<i>Pauliella taeniata</i>	25–30	–	–	+	–	5.0 × 10 ² –1.0 × 10 ⁴	–
<i>Skeletonema</i> sp.	2–21	48 ± 16	105 ± 76	+	–	1.0 × 10 ⁴ –2.0 × 10 ⁴	480–960
<i>Thalassiosira nordenskioeldii</i>	10–50	252 ± 62	1389 ± 540	+	–	1.0 × 10 ³	252
Dinoflagellates							
<i>Alexandrium tamarense</i>	30–40	2577 ± 543	6796 ± 1747	–	+	5.0 × 10 ² –1.0 × 10 ³	1289–2577
Raphidophyceae							
<i>Heterosigma akashiwo</i>	10–20	100	528 ± 85	–	+	5.0 × 10 ³	500

Table 2. The volume of the faecal pellets of *Neocalanus cristatus*, *Neocalanus flemingeri* and *Eucalanus bungii* observed with various phytoplanktons as food. For comparison possible, faecal pellet volumes (*FPV* in μm^3) predicted by prosome length (*PL* in mm) ($\text{Log } FPV = 2.474 \log PL + 5.226$, Mauchline 1998) are shown in the bottom column. Values are mean $\pm$ 1sd.

Taxa / Species	Faecal pellet volume ($10^6 \mu\text{m}^3$)		
	<i>Neocalanus cristatus</i>	<i>Neocalanus flemingeri</i>	<i>Eucalanus bungii</i>
Diatoms			
<i>Attheya septentrionalis</i>	16.06 $\pm$ 14.60		
<i>Chaetoceros</i> sp.	17.16		21.52
<i>Ditylum brightwellii</i>	12.53 $\pm$ 4.44	5.95	10.33 $\pm$ 2.84
<i>Pauliella taeniata</i>	29.38		
<i>Skeletonema</i> sp.	12.23 $\pm$ 4.09		
<i>Thalassiosira nordenskiöldii</i>	49.67	5.20	13.24 $\pm$ 2.64
Dinoflagellates			
<i>Alexandrium tamarense</i>	16.94 $\pm$ 9.80	3.77	5.20 $\pm$ 1.94
Raphidophyceae			
<i>Heterosigma akashiwo</i>	20.98 $\pm$ 1.74		
<i>FPV</i> predicted by <i>PL</i>	38.88	5.64	16.06

Table 3. Result of ANCOVA for the adjusted assimilation efficiencies. For this analysis, copepod species and phytoplankton ash contents applied as independent variables.
df: degree of freedom, SS: sum of squares, ***: $p<0.001$.

Parameter	df	SS	F-value	<i>p</i>
Copepods	2	1385.26	13.93	***
Ash contents	1	31176.81	626.88	***
Copepods × Ash contents	2	7110.58	71.49	***
Error	86	4277.08

Fig. 1 (Abe et al.)

Fig. 2 (Abe et al.)

Fig. 3 (Abe et al.)

○ <i>Attheya septentrionalis</i>	◆ <i>Skeletonema</i> sp.
● <i>Chaetoceros</i> sp.	▼ <i>Thalassiosira nordenskiöldii</i>
☆ <i>Ditylum brightwellii</i>	● <i>Alexandrium tamarense</i>
□ <i>Pauliella taeniata</i>	△ <i>Heterosigma akashiwo</i>

Fig. 4 (Abe et al.)

Fig. 5. Abe et al.

Fig. 6 (Abe et al.)

Fig. 7 (Abe et al.)