



Title	Regulation of multidrug resistance protein 2 (MRP2, ABCC2) expression by statins: Involvement of SREBP-mediated gene regulation
Author(s)	Kobayashi, Masaki; Gouda, Keisuke; Chisaki, Ikumi; Asada, Koji; Ogura, Jiro; Takahashi, Natsuko; Konishi, Toru; Koshida, Yusuke; Sasaki, Shotaro; Yamaguchi, Hiroaki; Iseki, Ken
Citation	International journal of pharmaceutics, 452(1-2), 36-41 https://doi.org/10.1016/j.ijpharm.2013.04.019
Issue Date	2013-08-16
Doc URL	http://hdl.handle.net/2115/54671
Type	article (author version)
File Information	WoS_62834_Kobayashi.pdf



[Instructions for use](#)

1 Regulation of multidrug resistance protein 2 (MRP2, ABCC2) expression by statins:

2 involvement of SREBP-mediated gene regulation

3
4 Masaki Kobayashi^{a, d}, Keisuke Gouda^{a, d}, Ikumi Chisaki^{a, d}, Koji Asada^a, Jiro Ogura^a,
5 Natsuko Takahashi^b, Toru Konishi^a, Yusuke Koshida^a, Shotaro Sasaki^a, Hiroaki
6 Yamaguchi^a, Ken Iseki^{a, c*}

7
8 ^aLaboratory of Clinical Pharmaceutics & Therapeutics, Division of Pharmasciences,
9 Faculty of Pharmaceutical Sciences, Hokkaido University

10 Kita-12-jo, Nishi-6-chome, Kita-ku, Sapporo 060-0812, Japan

11 ^bHokkaido Pharmaceutical University School of Pharmacy, Otaru 047-0264, Japan.

12 ^cDepartment of Pharmacy, Hokkaido University Hospital, Sapporo 060-8648, Japan

13 ^dThese authors contributed equally as first author

14
15 *To whom correspondence should be addressed.

16 Ken Iseki, Ph. D., Laboratory of Clinical Pharmaceutics & Therapeutics, Division of
17 Pharmasciences, Faculty of Pharmaceutical Sciences, Hokkaido University

18 Kita-12-jo, Nishi-6-chome, Kita-ku, Sapporo 060-0812, Japan

19 Tel/Fax: +81-11-706-3770

20 E-mail: ken-i@pharm.hokudai.ac.jp

21

22 **ABSTRACT**

23 Multidrug resistance protein 2 (MRP2, ABCC2) is localized to the apical
24 membrane of hepatocytes and played an important role in the biliary excretion of a
25 broad range of endogenous and xenobiotic compounds and drugs, such as pravastatin.
26 However, the effects of statins on MRP2 in the liver and the precise mechanisms of their
27 actions have been obscure. The goal of this study was to determine the regulatory
28 molecular mechanism for statin-induced MRP2 expression in hepatocytes. *In vitro* and
29 *in vivo* studies suggested that pitavastatin increased MRP2 expression. Pitavastatin
30 promoted liver X receptor (LXR) α/β translocation from the cytosol to nuclei, resulting
31 in LXR activation. Deletion and mutational analysis suggested that the potential a sterol
32 regulatory element (SRE) played a major role in the observed modulation of MRP2
33 expression by pitavastatin. Furthermore pitavastatin increased the protein-DNA
34 complex, and when SRE was mutated, stimulation of the protein-DNA complex by
35 pitavastatin was decreased. It was demonstrated that pitavastatin upregulated MRP2
36 expression by an SREBP regulatory pathway in hepatocytes and that the actions of
37 statins may lead to improve the biliary excretion of MRP2 substrates.

38 **Keywords:** Statin; hepatocytes; multidrug resistance protein 2; sterol regulatory
39 element-binding protein

40

41 **1. Introduction**

42 HMG-CoA reductase inhibitors (statins) reduce low-density lipoprotein (LDL)
43 cholesterol concentration through blockade of the mevalonate pathway and consequent
44 increment of LDL receptor expression in the liver (Goldstein and Brown, 1990). Statins
45 are the most widely used cholesterol-lowering agents for prevention of cardiovascular
46 disease (Havel and Rapaport, 1995). Major functions of hepatocytes include sinusoidal
47 extraction, intracellular metabolism and biliary excretion of endogenous and exogenous
48 lipophilic compounds (Kim, 2002). Xenobiotics, unconjugated bilirubin and hormones
49 are efficiently cleared from sinusoidal blood circulation into the liver by basolateral
50 transporters, including OATP-C (SLCO1B1), and are also conjugated with phase II
51 enzymes such as UDP-glucuronyltransferase (UGT1A1), sulfotransferase and
52 glutathione transferase (Cui et al., 2001; Hagenbuch and Meier, 2004; Kullak-Ublick et
53 al., 2004). After conjugation, the water-soluble metabolites are excreted primarily into
54 bile by ATP-binding cassette (ABC) transporters such as multidrug resistance protein 2
55 (MRP2, ABCC2) (Paulusma and Oude Elferink, 1997). This transporter is localized to
56 the apical membrane of hepatocytes and played an important role in the biliary
57 excretion of a broad range of endogenous and xenobiotic compounds. Pravastatin, one
58 of the statins, is a substrate of MRP2 using a double transporter (OATP1B1 and

59 MRP2)-expressing system (Sasaki et al., 2002). Cerivastatin is also a substrate of MRP2
60 (Matsushima et al., 2005). We previously reported that MRP2 expression is regulated
61 via a cholesterol-sensing nuclear receptor liver X receptor α (LXR α)-dependent
62 pathway (Chisaki et al., 2009). Several studies have shown that statins affect LXR
63 activation in the liver. However, the effects of statins on MRP2 expression are obscure.
64 It is important to investigate the effects of statins on MRP2 expression for considering
65 the biliary excretion of MRP2 substrates.

66 The aim of this study was to determine the effects of statins on MRP2 expression
67 in HepG2 cells and the rat liver.

68

69 **2. Materials and methods**

70 *2.1. Chemicals*

71 Pitavastatin Ca was kindly donated by Kowa (Tokyo, Japan). TO901317 was
72 obtained from Cayman Chemical (Ann Arbor, MI). All other reagents were of the
73 highest grade available and used without further purification.

74

75 *2.2. Cell culture*

76 HepG2 cells were cultured in Dulbecco's modified Eagle's medium (Sigma
77 Aldrich Japan, Tokyo) with 10% fetal bovine serum and 1% penicillin-streptomycin at
78 37°C under 5% CO₂ as described previously (Kobayashi et al., 2008). Rat primary
79 hepatocytes were isolated by the collagenase perfusion technique as described
80 previously with some modifications (Miyazaki et al., 1998). Collagen-coated plates
81 were prepared by using 50 µg/mL collagen solution. The plates were allowed to dry in a
82 laminar flow cabinet for 1 h. Isolated primary hepatocytes were plated onto the
83 collagen-coated plates in William's E medium (Invitrogen, Grand Island, NY)
84 supplemented with 10% fetal bovine serum and 1% penicillin-streptomycin and left to
85 attach for 6 h in an incubator (at 37°C under 5% CO₂). The density of isolated primary
86 hepatocytes was 2.0×10^6 cells/well. A minimum of two animal perfusions were used in

87 the study.

88

89 *2.3. Animals*

90 Male Wistar rats, aged 7 to 8 weeks (265-400 g in weight), were obtained from
91 Jla (Tokyo, Japan). The housing conditions were described previously (Kobayashi et al.,
92 2008). The experimental protocols were reviewed and approved by the Hokkaido
93 University Animal Care Committee in accordance with the “Guide for the Care and Use
94 of Laboratory Animals”. The dosage of statins suspended with methylcellulose was 30
95 mg/kg for rats as described previously (Kobayashi et al., 2011). Rats were starved for 24
96 hours after administration and euthanized.

97

98 *2.4. Quantitative real-time PCR*

99 Quantitative real-time PCR was performed using an Mx3000™ Real-time PCR
100 System with Platinum® SYBR® Green qPCR SuperMix UDG (Invitrogen, Carlsbad,
101 CA) following the manufacturer’s protocol. PCR was performed using human
102 MRP2-specific primers through 40 cycles of 94°C for 15 s, 60°C for 30 s and 72°C for
103 30 s or using human GAPDH-specific primers after pre-incubation at 95°C for 15 min.
104 The primers specific to hMRP2 and hGAPDH were designed on the basis of sequences

105 in the GenBankTM database (accession no.: NM_000392 and NM_002046). The
106 sequences of the specific primers were as follows: the sense sequence was 5'-ACA
107 GAG GCT GGT GGC AAC C-3' and the antisense sequence was 5'-ACC ATT ACC
108 TTG TCA CTG TCC ATG A-3' for hMRP2, and the sense sequence was 5'-AAG GTC
109 ATC CCT GAG CTG AA-3' and the antisense sequence was 5'-TTC TAG ACG GCA
110 GGT CAG GT-3' for hGAPDH. PCR was performed using rat Mrp2-specific primers
111 through 40 cycles of 94°C for 15 s, 52°C for 30 s and 72°C for 30 s or using rat
112 Gapdh-specific primers after pre-incubation at 95°C for 15 min. The primers specific to
113 rMrp2 and rGapdh were designed on the basis of sequences in the GenBankTM
114 database (accession no.: NM_012833 and AF106860). The sequences of the specific
115 primers were as follows: the sense sequence was 5'-TGA TCG GTT TCG TGA AGA
116 GCT-3' and the antisense sequence was 5'-ACG CAC ATT CCC AAC ACA AA-3' for
117 rMrp2, and the sense sequence was 5'-ATG GGA AGC TGG TCA TCA AC-3' and the
118 antisense sequence was 5'-GTG GTT CAC ACC CAT CAC AA-3' for rGapdh. The
119 PCR products were normalized to amplified GAPDH, which was the internal reference
120 gene (housekeeping gene).

121

122 *2.5. Western blot analysis*

123 Western blot analysis was performed as described previously (Kobayashi et al.,
124 2006). HepG2 cells were seeded on 6-well plastic plates. Following cell attachment (24
125 h), various concentrations of statins were added for 24 h. Total protein extracts were
126 prepared from the cells. The cells were scraped and centrifuged at 1,300 g for 1 min at
127 4°C. The pellet was suspended in 1 mL of PBS and centrifuged at 1,300 g for 1 min at
128 4°C. The resulting pellet was suspended in 100 µL of lysis buffer containing 1% Triton
129 X-100, 0.1% SDS and 4.5 M urea. The suspension was allowed to stand for 5 min and
130 was sonicated for 15 min at 4°C. The suspension was then centrifuged at 12,000 g for
131 15 min at 4°C. The liver crude membrane was used for Western blot analysis (Johnson
132 et al., 2006). The protein concentration in the clear supernatant was determined by the
133 method of Lowry et al. (Lowry et al., 1951). The samples were denatured at 100°C for 3
134 min in a loading buffer containing 0.1 M Tris-HCl, 4% SDS, 10% 2-mercaptoethanol,
135 20% glycerol, 0.004% BPB and 9 M urea and separated on 4.5% stacking and 10% SDS
136 polyacrylamide gels. Proteins were transferred electrophoretically onto nitrocellulose
137 membranes at 15 V for 90 min. The membranes were blocked with PBS containing
138 0.05% Tween 20 (PBS/T) and 10% non-fat dry milk for 1 h at room temperature. After
139 being washed with PBS/T, the membranes were incubated with mouse monoclonal
140 antibody to MRP2 (Abcam, Cambridge, UK) and mouse anti-actin monoclonal antibody

141 (Chemicon, Temecula, CA) for 24 h at room temperature and washed three times with
142 PBS/T for 10 min each time. The membranes were subsequently incubated for 1 h at
143 room temperature with horseradish peroxidase-conjugated goat anti-mouse secondary
144 antibody (Santa Cruz Biotechnology, Santa Cruz, CA) or horseradish
145 peroxidase-conjugated goat anti-rabbit secondary antibody (Santa Cruz Biotechnology,
146 Santa Cruz, CA) at a dilution of 1:2000 or 1:4000 and washed three times with PBS/T
147 for 10 min each time. The bands were visualized by enhanced chemiluminescence
148 according to the instructions of the manufacturer (Amersham Biosciences Corp.,
149 Piscataway, NJ).

150

151 *2.6. Immunohistochemistry*

152 HepG2 cells were washed once with PBS and fixed with methanol for 5 min or
153 10% formaldehyde for 15 min. After the fixed cells had been washed three times with
154 PBS, 10% FBS was added and the cells were incubated for 1 h at room temperature.
155 The cells were then treated with LXR α/β (C-19), SREBP-1 (C-20) and SREBP-2
156 (H-164) (Santa Cruz Biotechnology, Santa Cruz, CA) for 24 h at room temperature.
157 Then the cells were washed three times with PBS and treated with donkey anti-goat
158 IgG-FITC or anti-rabbit IgG-FITC secondary antibody (Santa Cruz Biotechnology,

159 Santa Cruz, CA) at a dilution of 1:400. Nuclei were stained with DAPI. The samples
160 were then visualized by using a confocal microscope (Zeiss LSM-510; Carl Zeiss Inc.,
161 Thornwood, NY).

162

163 *2.7. Reporter plasmid construction*

164 The human MRP2 promoter (-1200/+15) was amplified by PCR from human
165 genomic DNA and then subcloned into a pGL3-basic vector (Promega, Madison, WI).
166 Four 5'-deletion fragments (-701/+15, -427/+15 and -85/+15) were generated from
167 -1200/+15 by the PCR amplification method and subcloned into a pGL3-basic vector.
168 Four different forward primers each contained an internal site for Xho I restriction
169 enzyme. The primer sequences were primer-1, 5'-CTA CTC GAG CTC CCA CAT TCT
170 GGA TTT TGA-3'; primer-2, 5'- CAA CTC GAG ATG ATG GCA ACA CTG CAC TC
171 -3'; primer-3, 5'- TAA CTC GAG GGC TCA CAC TGG ATA AGC TAT TTT -3'; and
172 primer-4, 5'-GGT CTC GAG CCC TGT CCC TAG GGC TTT T-3'. In all cases, the
173 sequence of the reverse primer, 5'-ACG AAG CTT ATG ACC TTT CAT CCC AAC
174 CA-3', contained a site for HindIII enzyme. Vectors of the SRE mutant were generated
175 from -1200/+15 by megaprimer PCR methods using forward and reverse primers for the
176 MRP2 promoter and primers with a sense sequence of 5'-TAA CTC GAG GGC TCA

177 CAC TGG ATA AGC TAT TTT ATA CGC TGA CTT CTT CAA AGA A-3' for SRE
178 mutation. All PCR products and deletion constructs for reporter assays were sequenced
179 using an ABI PRISM 3100 Genetic Analyzer with a BigDye Terminator v3.1 cycle
180 sequencing kit according to the manufacturer's instructions.

181

182 *2.8. Transient transfection and luciferase assay*

183 HepG2 cells (1.0×10^5 cells/well) were seeded into 24-well plates and transfected
184 while still in suspension with one of the MRP2 promoter-luciferase constructs using
185 Lipofectin reagent (Invitrogen, Carlsbad, CA). Each well was transfected with 400 ng of
186 pGL3-MRP2 promoter construct encoding a modified firefly luciferase gene (or empty
187 pGL3 vector). After transfection, cells were transferred to a fresh medium supplemented
188 with TO901317 or pitavastatin. Luciferase activity was measured by a luminometer
189 according to the manufacturer's instructions using an assay kit from Promega. Relative
190 firefly luciferase activities were normalized with renilla luciferase activities.

191

192 *2.9. Electrophoretic mobility-shift assay (EMSA)*

193 Probes for the gel shift assays were labeled with [γ - 32 P] ATP and T4
194 polynucleotide kinase. Labeled double-stranded oligonucleotides were prepared by

195 mixing the complementary single-stranded DNA that had been heated to 95 °C for 5
196 min. The labeled probe (30,000 cpm) was incubated with hepatic nuclear extract in a
197 mixture containing 10 mM Tris-HCl (pH 7.5), 50 mM KCl, 1 mM DTT, 1 mM EDTA,
198 5% glycerol, and 0.05 mg/mL poly (dI-dC) for 30 min on ice. The DNA-protein
199 complexes were resolved on a 5% PAGE gel at 100 V for 2 h. The gels were dried and
200 visualized by exposure to x-ray film.

201

202 *2.10. Statistical analyses*

203 Student's t-test was used to determine the significance of differences between two
204 group means. Statistical significance among means of more than two groups was
205 determined by one-way analysis of variance (ANOVA). Statistical significance was
206 defined as $p < 0.05$.

207

208 **3. Results and Discussion**

209 *3.1. Effects of statins on expression of MRP2 in hepatocytes*

210 First, we examined the alterations in MRP2 expression induced by atorvastatin
211 and pitavastatin in HepG2 cells. Pitavastatin up-regulated MRP2 mRNA and protein
212 level (Fig. 1A, C). To determine whether pitavastatin regulates MRP2 expression *in vivo*,
213 statins were given to rats as described previously (Kobayashi et al., 2011). Pitavastatin
214 significantly increased Mrp2 mRNA and protein level in the rat liver (Fig. 1B, D). Next,
215 we examined the location of induced MRP2 molecule by statins in HepG2 cells using
216 immunostaining. Pitavastatin increased MRP2 expression at the plasma membrane
217 compared to that of non-treated or atorvastatin-treated cells (data not shown). **These**
218 **results suggest that pitavastatin increases MRP2 expression in hepatocytes and the**
219 **MRP2 expression alteration may improve the transport function.** Further
220 investigations to measure this function using MRP2 over-expressed cells are in
221 progress.

222

223 *3.2. Involvement of LXR in MRP2 expression in hepatocytes*

224 To clarify the mechanism by which pitavastatin induces alteration in MRP2
225 expression, we focused on liver X receptors (LXRs), members of the nuclear hormone

226 receptor superfamily. We previously reported that LXR α activation induced expression
227 of MRP2 in HepG2 cells and the rat liver (Chisaki et al., 2009). LXRs are members of
228 the nuclear hormone receptor superfamily represented by two subtypes, LXR α and
229 LXR β (Shinar et al., 1994; Willy et al., 1995; Miyata et al., 1996). Previous studies have
230 demonstrated that the effects of statins on ABCA1 regulated by LXR are biphasic:
231 suppression through reduction of oxysterols (ligands for LXR) and enhancement
232 through reduction of geranylgeranyl pyrophosphate (suppression factor of PPARs) in
233 macrophages (Argmann et al., 2005). Therefore, we investigated whether pitavastatin
234 activates LXR in HepG2 cells. We used TO901317, an LXR synthetic agonist, as a
235 positive control in this study. As shown in Fig. 2A, TO901317 promoted LXR α/β
236 translocation from the cytosol to nuclei, and pitavastatin also accelerated LXR α/β
237 transfer to nuclei. Next, an MRP2 promoter-luciferase construct was transiently
238 transfected into HepG2 cells and luciferase activity was measured after the addition of
239 TO901317. There was clearly an increase in luciferase activity from the MRP2
240 promoter construct by LXR activation. To confirm that LXR activation is associated
241 with MRP2 expression, we examined the effect of LXR α siRNA on alteration in MRP2
242 expression induced by pitavastatin. Pitavastatin-induced MRP2 expression alteration
243 was blocked by LXR α siRNA (10 nM) transfected into HepG2 cells (data not shown).

244 Moreover, to determine which region of the MRP2 promoter is responsible for
245 TO901317-induced stimulation of MRP2 promoter activity, a series of MRP2 deletion
246 constructs were transfected into HepG2 cells, and luciferase activity was measured. The
247 full-length promoter construct (-1200/+15) exhibited approximately three-fold
248 activation of promoter activity after TO901317 treatment compared with the control.
249 Similarly, deletion constructs p-701/+15 and p-427/+15 showed an approximately two
250 point five to three-fold increase after TO901317 treatment. On the other hand, deletion
251 of nucleotides -85 to +15 significantly decreased both the basal and TO901317-induced
252 promoter activities (Fig. 2B). These results suggest that the region between -427 and -85
253 is important for basal promoter activity and contains a TO901317-responsive element.

254

255 *3.3. Importance of SREBP in statin-induced MRP2 expression*

256 From deletion analysis, the region between -427 and -85 of the MRP2 promoter
257 was found to be important for basal promoter activity and this region did not contain
258 LXR response elements but contained a sterol regulatory element-binding proteins
259 (SREBPs) binding site (Supplemental Fig. 1A). In the liver, SREBPs regulate the
260 production of lipids for export into the plasma as lipoproteins and into the bile as
261 micelles. An LXR-binding site in the SREBP-1c promoter activates SREBP-1c

262 transcription in the presence of LXR agonists (Repa et al., 2000). Therefore, to clarify
263 the mechanism by which pitavastatin induces MRP2 expression, we focused on the
264 involvement of SREBPs. As shown in Fig. 3A, the protein-DNA complex was increased
265 in a TO901317 concentration-dependent manner and abolished by adding a 30-fold
266 excess of non-radioactive SRE (Fig. 3B). SRE_m sequence was shown in (Supplemental
267 Fig. 1B). Next, we examined the effect of SRE mutation, SRE_m, on the protein-DNA
268 complex. When SRE was mutated, stimulation of the protein-DNA complex by
269 TO901317 was decreased (Fig. 3B). These results suggest that SREBP binds to SRE in
270 the MRP2 promoter and LXR activation promotes SRE binding ability. We examined
271 the effects of statins on the protein-DNA complex in HepG2 cells. As shown in Fig. 4A,
272 pitavastatin increased the protein-DNA complex. When SRE was mutated, stimulation
273 of the protein-DNA complex by pitavastatin was decreased. Moreover, TO901317 and
274 pitavastatin-induced stimulation of MRP2 promoter activity were also decreased
275 (Supplemental Fig. 2). These results suggest that SREBPs play an important role in
276 increment of MRP2 expression level induced by pitavastatin. The mammalian genome
277 encodes three SREBP isoforms, designated SREBP-1a, SREBP-1c, and SREBP-2
278 (Brown and Goldstein, 1997). Therefore, to determine which SREBP isoform is
279 involved in the increase of MRP2 expression induced by pitavastatin, we examined the

280 effects of TO901317 and statins on the expression of SREBPs. TO901317 and
281 pitavastatin promoted SREBP-1 translocation from the cytosol to nuclei. On the other
282 hand, SREBP-2 localized to the nuclei with or without TO901317 and statins (Fig. 4B).
283 These results suggest that pitavastatin induces SREBP-1 activation and increases MRP2
284 expression mediated by the LXR-SREBP-1 pathway. MRP2 is a 190-kDa
285 phosphoglycoprotein localized in the apical membrane of hepatocytes and is involved in
286 the transport of a wide variety of organic anions such as bilirubin glucuronide and
287 glutathione conjugates. MRP2/Mrp2 function plays a pivotal role in homeostasis of
288 bilirubin, as shown in Dubin-Johnson Syndrome (DJS) and transport deficient (TR-)
289 and Eisai hyperbilirubinemic rats (EHBR), all of which are caused by a congenital
290 absence of MRP2/Mrp2 and show hyperbilirubinemia (Kartenbeck et al., 1996;
291 Paulusma et al., 1996; Ito et al., 1997). Therefore, Hayashi et al suggest that that
292 upregulation of MRP2 expression will improve hyperbilirubinemia in patients with DJS
293 (Hayashi et al., 2012). Our results suggest that pitavastatin may lead to improve the
294 biliary excretion of MRP2 substrates, such as bilirubin glucuronide.
295

296 **4. Conclusions**

297 In conclusion, our data demonstrated that pitavastatin upregulated MRP2
298 expression by an LXR-SREBP regulatory pathway in hepatocytes. Our findings suggest
299 that LXR and SREBP can be involved in the up-regulation of MRP2 expression in
300 hepatocytes.

301 **Acknowledgement**

302 This work was supported by Nakatomi Foundation.

303

304 **References**

305 Argmann, C.A., Edwards, J.Y., Sawyez, C.G., O'Neil, C.H., Hegele, R.A., Pickering,
306 J.G., Huff, M.W., 2005. Regulation of macrophage cholesterol efflux through
307 hydroxymethylglutaryl-CoA reductase inhibition: a role for RhoA in ABCA1-mediated
308 cholesterol efflux. *J. Biol. Chem.*, 280, 22212-22221.

309

310 Brown, M.S., Goldstein, J.L., 1997. The SREBP pathway: regulation of cholesterol
311 metabolism by proteolysis of a membrane-bound transcription factor. *Cell*, 89, 331-340.

312

313 Chisaki, I., Kobayashi, M., Itagaki, S., Hirano, T., Iseki, K., 2009. Liver X receptor
314 regulates expression of MRP2 but not that of MDR1 and BCRP in the liver. *Biochim.*
315 *Biophys. Acta*, 1788, 2396-2403.

316

317 Cui, Y., Konig, J., Leier, I., Buchholz, U., Keppler, D., 2001. Hepatic uptake of bilirubin
318 and its conjugates by the human organic anion transporter SLC21A6. *J. Biol. Chem.*,
319 276, 9626-9630.

320

321 Goldstein, J.L., Brown, M.S., 1990. Regulation of the mevalonate pathway. *Nature*, 343,

322 425-430.

323

324 Hagenbuch, B., Meier, P.J., 2004. Organic anion transporting polypeptides of the
325 OATP/SLC21 family: phylogenetic classification as OATP/SLCO superfamily, new
326 nomenclature and molecular/functional properties. *Pflugers Arch*, 447, 653-665.

327

328 Havel, R.J., Rapaport, E., 1995. Management of primary hyperlipidemia. *N. Engl. J.*
329 *Med.*, 332, 1491-1498.

330

331 Hayashi, H., Mizuno, T., Horikawa, R., Nagasaka, H., Yabuki, T., Takikawa, H.,
332 Sugiyama, Y., 2012. 4-Phenylbutyrate modulates ubiquitination of hepatocanalicular
333 MRP2 and reduces serum total bilirubin concentration. *J. Hepatol.*, 56, 1136-1144.

334

335 Ito, K., Suzuki, H., Hirohashi, T., Kume, K., Shimizu, T., Sugiyama, Y., 1997.
336 Molecular cloning of canalicular multispecific organic anion transporter defective in
337 EHBR. *Am. J. Physiol.*, 272, G16-G22.

338

339 Johnson, B.M., Zhang, P., Schuetz, J.D., Brouwer, K.L., 2006. Characterization of

340 transport protein expression in multidrug resistance-associated protein (Mrp) 2-deficient
341 rats. *Drug Metab. Dispos.*, 34, 556-562.

342

343 Kartenbeck, J., Leuschner, U., Mayer, R., Keppler, D., 1996. Absence of the canalicular
344 isoform of the MRP gene-encoded conjugate export pump from the hepatocytes in
345 Dubin-Johnson syndrome. *Hepatology*, 23, 1061-1066.

346

347 Kim, R.B., 2002. Transporters and xenobiotic disposition. *Toxicology*, 181–182,
348 291-297.

349

350 Kobayashi, M., Chisaki, I., Narumi, K., Hidaka, K., Kagawa, T., Itagaki, S., Hirano, T.,
351 Iseki, K., 2008. Association between risk of myopathy and cholesterol-lowering effect:
352 a comparison of all statins. *Life Sci.*, 82, 969-975.

353

354 Kobayashi, M., Gouda, K., Chisaki, I., Ochiai, M., Itagaki, S., Iseki, K., 2011.
355 Regulation mechanism of ABCA1 expression by statins in hepatocytes. *Eur. J.*
356 *Pharmacol.*, 662, 9-14.

357

358 Kobayashi, M., Kagawa, T., Narumi, K., Itagaki, S., Hirano, T., Iseki, K., 2008.
359 Bicarbonate supplementation as a preventive way in statins-induced muscle damage. J.
360 Pharm. Pharm. Sci., 11, 1-8.
361
362 Kobayashi, M., Otsuka, Y., Itagaki, S., Hirano, T., Iseki, K., 2006. Inhibitory effects of
363 statins on human monocarboxylate transporter 4. Int. J. Pharm., 317, 19-25.
364
365 Kullak-Ublick, G.A. Stieger, B., Meier, P.J., 2004. Enterohepatic bile salt transporters in
366 normal physiology and liver disease. Gastroenterology, 126, 322-342.
367
368 Lowry, O.H., Rosebrough, N.J., Farr, A.L., Randall, R.J., 1951. Protein measurement
369 with the folin phenol reagent. J. Biol. Chem., 193, 265-275.
370
371 Matsushima, S., Maeda, K., Kondo, C., Hirano, M., Sasaki, M., Suzuki, H., Sugiyama,
372 Y., 2005. Identification of the hepatic efflux transporters of organic anions using
373 double-transfected Madin-Darby canine kidney II cells expressing human organic
374 anion-transporting polypeptide 1B1 (OATP1B1)/multidrug resistance-associated protein
375 2, OATP1B1/multidrug resistance 1, and OATP1B1/breast cancer resistance protein. J.

376 Pharmacol. Exp. Ther., 314, 1059-1067.

377

378 Miyata, K.S., McCaw, S.E., Patel, H.V., Rachubinski, R.A., Capone, J.P., The orphan
379 nuclear hormone receptor LXR alpha interacts with the peroxisome
380 proliferator-activated receptor and inhibits peroxisome proliferator signaling. J. Biol.
381 Chem., 271, 9189-9192.

382

383 Miyazaki, M., Mars, W.M., Runge, D., Kim, T.H., Bowen, W.C., Michalopoulos, G.K.,
384 1998. Phenobarbital suppresses growth and accelerates restoration of differentiation
385 markers of primary culture rat hepatocytes in the chemically defined hepatocyte growth
386 medium containing hepatocyte growth factor and epidermal growth factor. Exp. Cell.
387 Res., 241, 445-457.

388

389 Paulusma, C.C., Bosma, P.J., Zaman, G.J., Bakker, C.T., Otter, M., Scheffer, G.L.,
390 Scheper, R.J., Borst, P., Oude Elferink, R.P., 1996. Congenital jaundice in rats with a
391 mutation in a multidrug resistance-associated protein gene. Science, 271, 1126–1128.

392

393 Paulusma, C.C., Oude Elferink, R.P., 1997. The canalicular multispecific organic anion

394 transporter and conjugated hyperbilirubinemia in rat and man. *J. Mol. Med.*, 75,
395 420-428.

396

397 Repa, J.J., Liang, G., Ou, J., Bashmakov, Y., Lobaccaro, J.M., Shimomura, I., Shan, B.,
398 Brown, M.S., Goldstein, J.L., Mangelsdorf, D.J., 2000. Regulation of mouse sterol
399 regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors,
400 LXRA and LXRbeta. *Genes Dev.*, 14, 2819-2830.

401

402 Sasaki, M., Suzuki, H., Ito, K., Abe, T., Sugiyama, Y., 2002. Transcellular transport of
403 organic anions across a double-transfected Madin-Darby canine kidney II cell
404 monolayer expressing both human organic anion-transporting polypeptide
405 (OATP2/SLC21A6) and multidrug resistance-associated protein 2 (MRP2/ABCC2). *J.*
406 *Biol. Chem.*, 277, 6497-6503.

407

408 Shinar, D.M., Endo, N., Rutledge, S.J., Vogel, R., Rodan, G.A., Schmidt, A., 1994.
409 NER, a new member of the gene family encoding the human steroid hormone nuclear
410 receptor. *Gene*, 147, 273-276.

411

412 Willy, P.J., Umesono, K., Ong, E.S., Evans, R.M., Heyman, R.A., Mangelsdorf, D.J.,
413 1995. LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Gene*
414 *Dev.*, 9, 1033-1045.
415

416 **Figures Legends**

417 Fig. 1. Effects of statins on MRP2 mRNA levels and protein expression in HepG2 cells
418 (A, C) and the rat liver (B, D).

419 (A) HepG2 cells were treated with various concentrations of statins for 24 h.

420 The bar graphs (n=3-13) are given as means with S.D. of more than two
421 independent experiments.

422 *P < 0.05 compared with vehicle control.

423 (B) Male Wistar rats were administered methylcellulose with or without 30 mg/kg
424 statins for 24 h.

425 The bar graphs (n=3-10) are given as means with S.D. of more than two
426 independent experiments.

427 *P < 0.05 compared with vehicle control.

428 (C) HepG2 cells were treated with 10 μ M statins for 24 h.

429 Data shown are typical results of the three independent experiments.

430 The intensity of Western blot analysis was determined by densitometry using Scion
431 image.

432 (D) Male Wistar rats were administered methylcellulose with or without 30 mg/kg
433 statins for 24 h.

434 Data shown are typical results of the three independent experiments.

435 The intensity of Western blot analysis was determined by densitometry using Scion
436 image.

437

438 Fig. 2. Nuclear translocation of LXR α/β induced by LXR activation and pitavastatin (A)
439 and functional analysis of MRP2 promoter in response to LXR activation (B) in HepG2
440 cells.

441 (A) HepG2 cells were treated with 10 μ M TO901317 and pitavastatin for 24 h.

442 Localization of LXR α/β was determined using an antibody against LXR α/β (green).

443 Nuclei were stained with DAPI (blue). Scale bar shows 10 μ m.

444 Data shown are typical results of three independent experiments.

445 (B) A series of MRP2 deletion promoter constructs were transfected into HepG2 cells
446 for luciferase assay.

447 HepG2 cells were treated with 10 μ M TO901317.

448 The bar graphs (n=4-6) are given as means with S.D. of more than two independent
449 experiments.

450 *P < 0.05 compared with vehicle control. N.S.: not significant.

451

452 Fig. 3. Electrophoretic mobility shift assay of nuclear extract of HepG2 cells binding to
453 human SRE consensus sequence.

454 Double-stranded SRE consensus sequence (A) and SRE sequence of MRP2
455 promoter (B) were labeled with [γ - 32 P]-ATP.

456 These labeled SREs were incubated with nuclear extracts from HepG2 cells.

457 HepG2 cells were treated with various concentrations of TO901317 for 24 h.

458 Data shown are typical results of the more than two independent experiments.

459

460 Fig. 4. Nuclear extract of HepG2 cells binding to human SRE consensus sequence (A),
461 SREBP-1 and 2 nuclear translocation (B) induced by TO901317 and statins in HepG2
462 cells.

463 (A) HepG2 cells were treated with 10 μ M TO901317 and statins for 24 h.

464 Data shown are typical results of the more than two independent experiments.

465 (B) HepG2 cells were treated with 10 μ M TO901317 and statins for 24 h. Localization
466 of SREBP-1 and 2 was determined using an antibody against SREBP-1 and 2
467 (green). Nuclei were stained with DAPI (blue).

468 Data shown are typical results of the more than two independent experiments.

469

470 Supplemental Fig. 1. Sequence of MRP2 promoter (-427 to -85) (A) and SRE mutant

471 that was created to eliminate SREBP binding to the MRP2 promoter construct (B).

472

473 Supplemental Fig. 2. Effects of TO901317 (A) and pitavastatin (B) on SRE and SREm

474 promoter activity in HepG2 cells.

475 HepG2 cells were treated with 10 μ M TO901317 and pitavastatin for 3 and 6 hours.

476 The bar graphs (n=4-9) are given as means with S.D. of more than two independent

477 experiments.

478 *P < 0.05 compared with vehicle control. N.S.: not significant.

Fig. 1





Fig. 2

(A)



(B)





Fig. 3

(A)



(B)






Fig. 4

(A)



(B)



Supplemental Fig. 1

(A)

-427 gctcacactggataagctatfff **ataacctgact** tcttcaaagaaagttt
SREBP
acatcatgtttaaacatgttttagattctatatttttaattaaaaatct
-327 aaggaagaaggatatttcacatttctataaaactctaagatcttgacagcag
aagcgaactgcacatttaggggtgcctgccctctactgatgctgccct
-227 ttgtgggtcatatgtccttaggaaaatgaaagactgtgcactcttgatt
gttgccagctctgtgacatcttcagtggtccttttatgtatggcca
-127 ctctacagaggcctcttgactttgggaactggtgagtctc -85

(B)

SRE ataacctgact
SREm atacgctgact

Supplemental Fig. 2

