



Title	On LF3 (3)
Author(s)	Tsuzuku, Tosihiro
Citation	Journal of the Faculty of Science Hokkaido University. Ser. 1 Mathematics, 22(3-4), 104-107
Issue Date	1972
Doc URL	http://hdl.handle.net/2115/54693
Type	bulletin (article)
File Information	JFSHIU_22_N3-4_104-107.pdf



[Instructions for use](#)

On $LF_3(3)$

Dedicated to Professor Yoshie Katsurada on her 60th birthday

By Tosiro TSUZUKU

Let Ω be a set of $p=4q+1$ symbols where p and q are prime numbers. Let G be an unsolvable transitive permutation group on Ω . Appel and Parker ([1], theorem 2) showed that there is no such a permutation group (G, Ω) for $q \leq 79$ except for the group $(LF_3(3), \mathbf{P})$ with $q=3$, where $LF_3(3)$ denotes the linear fractional group over the field of three elements and \mathbf{P} denotes the projective plane over the field of three elements. In this note we prove the following theorem.

THEOREM. *Let H be the stabilizer of a symbol, say 0, of Ω . If H is imprimitive on $\Omega - \{0\}$, then (G, Ω) is isomorphic to $(LF_3(3), \mathbf{P})$.*

PROOF. By the result of Appel-Parker, we can assume that $q > 79$, and we only need to prove non existence of such a permutation group (G, Ω) for $q > 79$.

Let P be a Sylow p -subgroup of G and let $N(P)$ be the normalizer of P in G . Let N be a minimal normal subgroup of G . Then N is a simple group and transitive on Ω , and G is a product of N and $N(P)$. Hence G/N is cyclic and so N is an unsolvable transitive group on Ω . Hence hereafter we can assume that G is simple.

Let K be the stabilizer of two symbols 0 and 1 of Ω . Then, by a theorem of Witt, $N(K)$ is doubly transitive on $F(K)$, where

$$F(K) = \{\alpha \in \Omega \mid \alpha^a = \alpha \text{ for any } a \in K\},$$

and hence $4qp$ is divisible by $f(f-1)$ where $f = |F(K)|$. From our assumption we have that $f=2$. Since $(H, \Omega - \{0\})$ is imprimitive, K is not a maximal subgroup of H . Let L be a maximal subgroup of H containing K . Since $[H:K]=4q$ and $f=2$ three cases arise: (i) $[H:L]=2$ and $[L:K]=2q$, (ii) $[H:L]=4$ and $[L:K]=q$ and (iii) $[H:L]=q$, and $[L:K]=4$. Let $\Omega = \{\Omega_1, \Omega_2, \dots, \Omega_t\}$, $t[L:K]=4q$, be a complete system of sets of imprimitivity of $(H, \Omega - \{0\})$ corresponding to L , namely, $\Omega - \{0\} = \cup \Omega_i$, $|\Omega_i| = [L:K]$, $\Omega_i \cap \Omega_j = \emptyset$ for $i \neq j$, every element of H induces a permutation on the set $\tilde{\Omega}$ and L is the stabilizer of one symbol of $\tilde{\Omega}$, say Ω_1 . Then we may assume that $\Omega_1 \ni 1$.

Case (i) cannot occur. Assume that $[L : K] = 2q$. Then L is a normal subgroup of H of index 2. Let 1_L be the principal character of L . Then ${}^H 1_L$, the character of H induced by 1_L , is the sum of the principal character 1_H of H and a non trivial linear character ε of H . Since (G, Ω) is doubly transitive, ${}^G 1_H = 1_G + \chi$ where χ is an irreducible character of degree $p-1$. Since ${}^H 1_K = \chi_{1_H}$ (the restriction of χ to H), we have that ${}^G \varepsilon = \chi + \eta$ by the reciprocity theorem of Frobenius, where η is a non principal linear character of G . This is a contradiction.

Case (ii) cannot occur. Assume that $[L : H] = q$. If ${}^H 1_L$ contains a non principal linear character, then we have a contradiction in the similar way as the case (i). Hence we have that

$${}^H 1_L = 1_H + \phi$$

where ϕ is a irreducible character of H of degree 3. Then by the reciprocity theorem of Frobenius we have that

$${}^G 1_L = 1_G + 2\chi_0 + \phi$$

where χ_0 is a irreducible character of G of degree $p-1$ and ϕ is a character of G of degree $2p+1$. By a result of Tuan ([4], Theorem 3), we conclude that ϕ is irreducible or decomposed into two irreducible characters. In the latter case, by a theorem of Brauer [2], the degrees of irreducible characters are p and $p+1$. Let us consider G as a permutation group on G/L , the set of left cosets by L . Then it is known ([5], Proposition 29. 2) that the number of domains of transitivity of $(L, G/L)$ equals to the norm of ${}^G 1_L$. Assume that ϕ is irreducible. Then the norm of ${}^G 1_L$ is six, and hence the number of domains of transitivity of $(L, G/L)$ is six, and it is easy to see that the lengths of those domains are 1, 3, q , $3q$, $3q$ and $9q$. Then, by a theorem of Frame ([3], Theorem B), we have that

$$\frac{(4q)^4 \cdot 3 \cdot q \cdot 3q \cdot 3q \cdot 9q}{(p-1)^4 (2p+1)}$$

is a rational integer. This is a contradiction. Hence ϕ is reducible. Then the number of domains of transitivity of $(L, G/L)$ is seven and the lengths of those domains are 1, 3, q , $3q$, $3q$, $3q$ and $6q$. Then, by a theorem of Frame,

$$\frac{(4p)^5 \cdot 3 \cdot q \cdot 3q \cdot 3q \cdot 3q \cdot 6q}{(p-1)^4 \cdot p(p+1)}$$

is a rational integer. This is also a contradiction.

Case (iii) cannot occur. Assume that $[L : K] = 4$. We can assume that

there is no subgroup T of H such that $H \cong T \cong K$ and $[T:K] \neq 4$. Let M be the kernel of H as a permutation group on $\tilde{\Omega}$, namely,

$$M = \{a \in H \mid \Omega_i^a = \Omega_i \text{ for } i=1, \dots, q\}.$$

If $M \leq K$, then $|M|=1$ and then a Sylow q subgroup Q of H is a normal subgroup of H because H is considered as a Frobenius group of degree q . Then KQ is a subgroup of H and $[KQ:K]=q$. This is a contradiction. If M is not transitive on Ω_1 , then we have that KM is a subgroup of H and $[KM:K]=2$. This is also a contradiction. Hence M is transitive on Ω_1 , also on Ω_i for any index i . Now we assume that H is unsolvable. Then $(H/M, \tilde{\Omega})$ is a unsolvable permutation group of degree q since M is solvable, and hence L is transitive on $\Omega_2 \cup \Omega_3 \cup \dots \cup \Omega_q$ because L is transitive on $\tilde{\Omega} - \{\Omega_1\}$ and M is transitive on M_i for any i . Let N be the kernel of L as a permutation group on Ω_1 . Then N is half-transitive on $\Omega_2 \cup \dots \cup \Omega_q$ and so we have that

$$\begin{aligned} F(N) &= \{\alpha \in \Omega \mid \alpha^a = \alpha \text{ for any } a \in N\} \\ &= \{0\} \cup \Omega_1 \text{ or } \Omega. \end{aligned}$$

If $F(N)=\Omega$, then we have that $q-1 \leq 6$ which is in contradiction with our assumption. Hence $F(N)=\{0\} \cup \Omega_1$. Then, by a theorem of Witt, $[G:N] \equiv 0 \pmod{5}$. This is a contradiction. Hence H is solvable. Since $|FK|=2$, $\Omega_1 - \{1\}$ is an orbit of length 3 of (K, Q) . If $\Omega_1 - \{1\}$ is the unique orbit of length 3 of (K, Q) , then $\{0\} \cup \Omega_1$ is uniquely determined by a (unordered) set $\{0, 1\}$, and then it is easily seen that $\{0\} \cup \Omega_1$ is also uniquely determined by any two symbols of $\{0\} \cup \Omega_1$. Hence we conclude that $p(p-1)/5 \cdot 4$ is a rational integer which is in contradiction with $p(p-1)=4qp$. Therefore the permutation group (K, Ω) has at least two orbits of length 3. For a symbol α of $\Omega - \{0\}$, we set

$$\tilde{\alpha} = \{\Omega_i \mid \Omega_i \cap F(M_\alpha) \neq \emptyset\}$$

where $M_\alpha = \{a \in M \mid \alpha^a = \alpha\}$ and $F(M_\alpha) = \{\beta \in \Omega \mid \beta^x = \beta \text{ for any } x \in M_\alpha\}$. Then we have that $\tilde{\alpha} = \tilde{\beta}$ for $\alpha, \beta \in \Omega_i$ and, if $\tilde{\alpha} \neq \tilde{\beta}$, then $\tilde{\alpha} \cap \tilde{\beta} = \emptyset$ and $|\tilde{\alpha}| = |\tilde{\beta}|$. Hence if $|\tilde{\alpha}| \geq 1$ then $\tilde{\alpha} = \tilde{\Omega}$. Then M is faithful on Ω_1 and we conclude that $|M| \leq 24$. Since $(H/M, \tilde{\Omega})$ is a Frobenius group, $H \triangleright MQ$ where Q is a Sylow q -subgroup of H . Since $q > 24$, $MQ = M \times Q$, namely $H \triangleright Q$. Hence we conclude that KQ is a subgroup of H and $[KQ:K]=q$. This is in contradiction with our assumption. Hence $|\hat{\alpha}|=1$ for any $\alpha \in \Omega - \{0\}$. Then the set of orbits of (M_α, Ω) consists of the following orbits; two orbits of length 1, one orbit of length 3 and other orbits of even length. Therefore this

conclude that (K, Ω) has only one orbit of length 3. This is a contradiction. Thus we have completed the proof of the Theorem.

Department of Mathematics
Hokkaido University

Reference

- [1] APPEL, K. I. and PARKER, E. T.: On unsolvable groups of degree $p=4q+1$, p and q primes, *Canad. J. Math.* 19 (1967), 583-589.
- [2] Brauer, R.: On permutation groups of prime degree and related classes of groups, *Ann. of Math.* 44 (1943), 57-79.
- [3] Frame, J. S.: The double cosets of a finite group, *Bull. Amer. Math. Soc.* 47 (1941), 458-467.
- [4] TUAN, H. F.: On groups whose orders contain a prime number to the first power, *Ann. of Math.* 45 (1944), 110-140.
- [5] WIELANDT, H.: "Finite Permutation Groups" Acad. Press, New York and London, 1964.

(Received August 17, 1971)