Title	Synthesis of pacidamycin analogues via an Ugi-multicomponent reaction
Author(s)	Okamoto, Kazuya; Sakagami, Masahiro; Feng, Fei; Takahashi, Fumiyo; Uotani, Kouichi; Togame, Hiroko; Takemoto, Hiroshi; Ichikawa, Satoshi; Matsuda, Akira
Citation	BIOORGANIC & MEDICINAL CHEMISTRY LETTERS, 22(14), 4810-4815 https://doi.org/10.1016/j.bmcl.2012.05.050
Issue Date	2012-07-15
Doc URL	http://hdl.handle.net/2115/54870
Type	article (author version)
File Information	Matsuda_Akira_2012_Bioorg_Med_Chem_Lett.pdf

[Instructions for use](#)

Synthesis of Pacidamycin Analogues via an Ugi-Multicomponent Reaction

Kazuya Okamoto,^a Masahiro Sakagami,^a Fei Feng,^b Fumiyo Takahashi,^a
Kouichi Uotani,^c Hiroko Togame,^a Hiroshi Takemoto,^a Satoshi
Ichikawa,^{d,*} Akira Matsuda^d

^aShionogi Innovation Center for Drug Discovery, Shionogi & Co., Ltd., Kita-21 Nishi-11, Kita-ku, Sapporo 001-0021, ^bFaculty of Advanced Life Science, Hokkaido University, Kita-21, Nishi-11, Kita-ku, Sapporo, 001-0021, ^cMedicinal Research Laboratories, Shionogi & Co., Ltd., 3-1-1, Futaba-cho, Toyonaka, Osaka 561-0825, ^dFaculty of Pharmaceutical Sciences, Hokkaido University, Kita-12, Nishi-6, Kita-ku, Sapporo 060-0812, Japan

*Corresponding author: Satoshi Ichikawa

Tel (+81)11-706-3229

Fax (+81)11-706-4980

e-mail: ichikawa@pharm.hokudai.ac.jp

Abbreviations

α,β -diaminobutylic acid (DABA),
2-(1*H*-7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
(HATU), phospho-MurNAc-pentapeptide transferase (MraY), *N*-methylmorpholine
(NMM), structure-activity relationship (SAR), Ugi-four component reaction (U-4CR)

Abstract

The second-generation synthesis of 3'-hydroxypacidamycin D (**2**) has been accomplished via an Ugi-four component reaction at a late stage of the synthesis. This approach provided ready access to a range of analogues including diastereomers of the diaminobutylic acid residue and hybrid-type analogues of mureidomycins. Biological evaluations of these analogues indicated that the stereochemistry at the diaminobutylic acid residue has a crucial impact on both the MraY biochemical inhibition and whole-cell antibacterial activity.

The pacidamycins (Figure 1, **1**),¹ first isolated from *Streptomyces coeruleorubidus* AB 1183F-64 in 1989, are members of a class of uridylpeptide antibiotics,² that are composed of the mureidomycins,³ the napsamycins,⁴ and the recently identified sansanmycins.⁵ Sharing a characteristic structural feature, namely a 3'-deoxyuridine which is attached to a tetrapeptide moiety by an enamide linkage. These compounds exhibit selective antibacterial activity against *Pseudomonas aeruginosa*, a common nosocomial pathogen that is intrinsically resistant to a variety of drugs currently used in the clinic. Consequently, immuno-compromised patients are at greater risk of infection. Moreover, the increased rate of acquired multidrug resistance in *P. aeruginosa* complicates anti-*Pseudomonas* chemotherapy, thereby making the development of novel antibacterial agents that are active against multi-drug resistance *P. aeruginosa* all the more urgent.⁶⁻⁸ These uridylpeptide antibiotics are sub-nanomolar inhibitors of their biological target, the phospho-MurNAc-pentapeptide transferase (MraY),^{2,9} which is responsible for the formation of lipid I in the peptidoglycan biosynthesis pathway.^{2,10} Inhibition of the MraY by these antibiotics causes a malfunction of the peptidoglycan production in a manner different from β -lactams such as the carbapenems. Since MraY is an essential enzyme in bacteria² and a novel target, uridylpeptide antibiotics are expected to be highly useful anti-*Pseudomonas* agents. Intrigued by the promising

biological activity, several groups have conducted structure-activity relationship studies.¹¹⁻¹⁵

We have recently accomplished the total synthesis of pacidamycin D (**1**) and its 3'-hydroxyl analogue **2**, which features a stereoselective construction of the *Z*-oxy-acyl enamide architecture by a copper-catalyzed C-N cross-coupling.¹⁶ These compounds exhibited strong *MraY* inhibitory activity (IC_{50} = 22 nM for **1**, 42 nM for **2**) and potent antibacterial activity against *P. aeruginosa* (MIC 8-64 μ g/mL). The existence of the hydroxyl group at the 3'-position of the uridine moiety does not have any impact on either *MraY* inhibition or antibacterial activity. This first generation strategy, however, is a linear synthesis, where the vinyl iodide derivative of uridine was coupled with the tetrapeptide carboxamide moiety. Nonetheless, from a medicinal point of view, the development of a convergent synthetic strategy that would provide analogues more efficiently is preferable. A multi-component reaction would be ideally suited for structure-activity relationship (SAR) studies since it allows for diversification of accessible analogs simply by altering each component. We have successfully applied an Ugi-four component reaction (U-4CR) in pursuing the SAR of the muraymycins, which are antibacterial nucleoside natural products.¹⁷ Herein we describe the second-generation synthesis of **2** via a U-4CR in order to establish a convergent

synthetic strategy appropriate for the SAR study of uridylpeptide antibiotics. The biological evaluation of several analogues is also described.

Our second-generation retrosynthetic analysis of **2** is summarized in Scheme 1. We planned to install the *N*-terminal amino acid residue **3** on **4** in the final stage of the synthesis considering the initial efforts to prepare **2** as described later. We retrosynthetically divided **4** into the urea dipeptide **5**, 2,4-dimethoxybenzylamine **6**, the 2-*N*-methylaminopropionaldehyde derivative **7**, and the α,β -unsaturated isonitrile derivative of uridine **8** for the U-4CR. This strategy allowed us to construct **4** with the non-proteinogenic amino acid α,β -diaminobutylic acid (DABA) residue, with linkages to the urea dipeptide and uridine moieties at the *C*- and *N*-termini. Moreover, the U-4CR strategy has the advantage of efficiently furnishing diastereomers that are useful for the SAR. Considering the nature of the multi-component assemblage, this strategy allows easy access to diversified analogues.

The aldehyde unit **7** was prepared as shown in Scheme 2. Namely, *N*-formyl-L-alanine **9** was reduced by LiAlH_4 , and the resulting *N*-methylaminoalcohol was selectively protected by a Boc group to provide **10** in 98% yield over 2 steps. The primary hydroxyl group of **10** was oxidized by Dess-Martin periodinane to give **7** in 84% yield. The isonitrile **8** was prepared from the *Z*-enamide **12** by the

copper-catalyzed C-N cross-coupling of the *Z*-vinyl iodide **11**¹⁶ and formamide as describe in Scheme 3. The iodide **11** was reacted with formamide using 0.2 equiv. of CuI, 0.4 equiv. of MeNHCH₂CH₂NHMe, Cs₂CO₃ in THF at 70 °C to give the desired *Z*-enamide **12**, which was obtained selectively in 90% yield. The corresponding *E*-isomer was not observed at all during the course of the reaction. Dehydration of **12** was achieved by a treatment with triphosgene and Et₃N in CH₂Cl₂ at -78 °C to provide the isonitrile **8** in 88% yield.

With **7** and **8** in hand, we then examined the assemblage of the four components by the U-4CR (Scheme 4). The assemblage proceeded simply by mixing **5**, **6**, **7**, and **8** in EtOH at room temperature for 48 h, and the desired **13** and its diastereomer at the newly formed stereogenic center were obtained in acceptable yields (33% for **13** and 30% for **14**). These were separated by silica gel column chromatography. Initially we planned to use an aldehyde of the dipeptide derivative such as **21**, which was prepared by reduction of the Weinreb amide derivative of the dipeptide **20**, as the aldehyde component in the U-4CR assemblage (Scheme 5). However, it turned out that the resulting aldehyde **21** easily cyclized to form a rather stable cyclic aminal **22**, and the U-4CR did not proceed at all. Therefore we turned our attention to the stepwise synthetic route toward the target compounds as shown in Scheme 4 in order to avoid the undesired cyclization.

The final assemblage of the remaining amino acid and global deprotection were then investigated. Upon selective removal of the Boc group of **13** (10% TFA, CH₂Cl₂, 0 °C), the liberated secondary amine of **15** was condensed with *N*-Boc-L-Ala **3** (HATU, *N*-methylmorpholine (NMM), CH₂Cl₂) to give the fully protected 3'-hydroxyl pacidamycin D analogue **17** in 79% yield over 2 steps. Finally removal of all the protecting groups (BCl₃, CH₂Cl₂, -78 °C, 5HF·NEt₃, 22% over 2 steps) successfully afforded **2**. The analytical data for **2** in this second generation synthesis were identical to those of the material previously synthesized.^{16a} The deprotection step gave poor yield of the desired **2** because BCl₃ treatment resulted in decomposition of the acid sensitive enamide moiety as well as migration of the 2,4-dimethoxy group to the indole moiety of the Trp residue. Similar problems were also observed for the formation of **19**, **27**, **30**, and **31**, which were described later. The epimer at the DABA residue **19** was also prepared from **14** in a manner similar to that of the synthesis of **2**.

The mureidomycins (Figure 2)³ were first isolated from *Streptomyces flavidoviridens* SANK 60486 and belong to the same class of uridylpeptide antibiotics. However, their amino acids differ from those of the pacidamycins. Mureidomycin C, which has the dipeptide (Gly-L-*m*-Tyr) at the *N*-terminus, showed the most potent

antibacterial activity against strains of *P. aeruginosa* among the uridylpeptide antibiotics with MICs ranging from 0.1 to 1.56 $\mu\text{g/mL}$.^{3b}

We therefore planned to prepare an analogue **27a**, which is a hybrid-type of pacidamycin D and mureidomycin C, by introducing the Gly-L-*m*-Tyr moiety into the intermediates **15** and **16** (Scheme 6). The analogue **27b**, where the L-*m*-Tyr residue at the N-terminus was replaced by L-Tyr, was designed in order to determine the consequences of placing of the hydroxyl group in this position. The glycine analogue **27c** was also prepared. These analogues **27a-c**, as well as **30**, the epimer of **27a** at the DABA residue, were prepared by the same strategy as that for the synthesis of **2** and **19** by using the dipeptide **23**, **24**,¹⁸ or Boc-Gly-OH (**25**). Of particular note was the coupling of the amine derived from **16** and the dipeptide **23**, where extensive racemization occurred to give predominantly the epimer **29** (22% for **28** vs. 66% for **29**). Deprotection of **29** gave **31**, the diastereomer at the L-*m*-Tyr residue of **30**.

The inhibitory activity of the analogues **2**, **19**, **27a-c**, **30**, and **31** on the purified Mray enzyme (*Staphylococcus aureus*) was next examined by fluorescence based Mray assay using UDP-MurNAc-dansylpentapeptide, where the formation of dansylated lipid I was monitored by fluorescence enhancement (excitation at 355 nm, emission at 535 nm),¹⁹ and the results are summarized in Table 1. It turned out that the epimer **19** was much

weaker than the MraY inhibitors **1** or **2** with an IC₅₀ value of 4000 nM. The stereochemistry at the DABA residue was very important for enzyme inhibitory activity. This was also true for the analogues **30** and **31**, which possess the same stereochemistry at the DABA residue (IC₅₀ 650 and 8900 nM, respectively). Analogue **27a** showed similar inhibitory activity (IC₅₀ 22 nM) to that of **1** and **2**, and introducing the Gly-L-*m*-Tyr moiety at the *N*-terminus did not improve but instead retained the activity. However, moving the hydroxyl group to the *para*-position at the *N*-terminal phenyl ring resulted in a large decrease in inhibitory activity with an IC₅₀ value of 4000 nM for **27b**. Therefore the position of the hydroxyl group on the phenyl ring plays a crucial role. Analogue **27c** showed a slight decrease in MraY inhibitory activity. The antibacterial activity of **2**, **19**, **27a-c**, **30**, and **31** was then evaluated against a range of clinically isolated *P. aeruginosa* strains (Table 2).²⁰ The overall antibacterial activity was well correlated to the MraY inhibitory activity. Thus, **27a** showed a similar anti-Pseudomonal activity to that of **1** and **2**, whereas **19**, **27b**, **27c**, **30**, and **31** exhibited no activity up to 32 µg/mL. The exception was the Gly analogue **27c**, which showed no anti-Pseudomonal activity in spite of potent MraY inhibitory activity (Table 1, IC₅₀ 65 nM).

In summary, the second-generation synthesis of **2** has been accomplished via a U-4CR at a late stage of the synthesis. This approach provided ready access to a range of analogues including the diastereomers of the DABA residue and hybrid-type analogues of the mureidomycins. Biological evaluations of these analogues indicated that the stereochemistry at the DABA residue has a significant impact on both the Mray biochemical inhibition and whole-cell antibacterial activity.

References

1. a) Karwowski, J. P.; Jackson, M.; Theriault, R. J.; Chen, R. H.; Barlow, G. J.; Maus, M. L. *J. Antibiot.* **1989**, *42*, 506. b) Chen, R. H.; Buko, A. M.; Whittern, D. N.; McAlpine, J. B. *J. Antibiot.* **1989**, *42*, 512. c) Fernandes, P. B.; Swanson, R. N.; Hardy, D. J.; Hanson, C. W.; Coen, L.; Rasmussen, R. R.; Chen, R. H. *J. Antibiot.* **1989**, *42*, 521. d) Fronko, R. M.; Lee, J. C.; Galazzo, J. G.; Chamberland, S.; Malouin, F.; Lee, M. D. *J. Antibiot.* **2000**, *53*, 1405.
2. a) Winn, M.; Goss, R. J. M.; Kimura, K.-I.; Bugg, T. D. H. *Nat. Prod. Rep.* **2010**, *27*, 279. b) Bugg, T. D. H.; Lloyd, A. J.; Roper, D. I. *Infect. Dis. Drug Targets* **2006**, *6*, 85. c) Kimura, K.-I.; Bugg, T. D. H. *Nat. Prod. Rep.* **2003**, *20*, 252.
3. a) Inukai, M.; Isono, F.; Takahashi, R.; Enokita, S.; Sakaida, Y.; Haneishi, T. *J. Antibiot.* **1989**, *42*, 662. b) Isono, F.; Katayama, T.; Inukai, M.; Haneishi, T. *J.*

- Antibiot.* **1989**, 42, 674. c) Isono, F.; Inukai, M.; Takahashi, S.; Haneishi, T.; Kinoshita, T.; Kuwano, H. *J. Antibiot.* **1989**, 42, 667. d) Isono, F.; Sakaida, Y.; Takahashi, S.; Kinoshita, T.; Nakamura, T.; Inukai, M. *J. Antibiot.* **1993**, 46, 1203.
4. Chatterjee, S.; Nadkarni, S. R.; Vijayakumar, E. K. S.; Patel, M. V.; Ganguli, B. N. *J. Antibiot.* **1994**, 47, 595.
5. a) Xie, X. Y.; Chen, R. X.; Si, S. Y.; Sun, C. H.; Xu, H. Z. *J. Antibiot.* **2007**, 60, 158.
b) Xie, X. Y.; Xu, H. Z.; Si, S. Y.; Sun, C. H.; Chen, R. X. *J. Antibiot.* **2008**, 61, 237.
6. Tam, V. H.; Chang, K. T.; Abdelraouf, K.; Brioso, C. G.; Ameka, M.; McCaskey, L. A.; Weston, J. S.; Caeiro, J. P.; and Garey, K. W. *Antimicrob. Agents Chemother.* **2010**, 54, 1160.
7. Page, M. G.; Heim, J. *Curr. Opin. Pharmacol.* **2009**, 9, 558.
8. Scheld, M.; Spellberg, B.; Bartlett, J. *Clin. Infect. Dis.* **2009**, 48, 1.
9. a) Isono, F.; Inukai, M. *Antimicrob. Agents Chemother.* **1991**, 35, 234. b) Inukai, M.; Isono, F.; Takatsuki, A. *Antimicrob. Agents Chemother.* **1993**, 37, 980.
10. a) Bouhss, A.; Mengin-Lecreulx, D.; Le Beller, D.; Van Heijenoort, J. *Mol. Microbiol.* **1999**, 34, 576. b) Bouhss, A.; Trunkfield, A. E.; Bugg, T. D.; Mengin-Lecreulx, D. *FEMS Microbiol. Rev.* **2008**, 32, 208. c) Al-Dabbagh, B.;

- Henry, X.; El Ghachi, M.; Auger, G.; Blanot, D.; Parquet, C.; Mengin-Lecreulx, D.; Bouhss, A. *Biochemistry* **2008**, *47*, 8919.
11. a) Boojamra, C. G.; Lemoine, R. C.; Lee, J. C.; Léger, R.; Stein, K. A.; Vernier, N. G.; Magon, A.; Lomovskaya, O.; Martin, P. K.; Chamberland, S.; Lee, M. D.; Hecker, S. J.; Lee, V. J. *J. Am. Chem. Soc.* **2001**, *123*, 870. b) Lemoine, R. C.; Magon, A.; Hecker, S. *Bioorg. Med. Chem. Lett.* **2002**, *12*, 1121. c) Boojamra, C. G.; Lemoine, R. C.; Blais, J.; Vernier, N. G.; Stein, K. A.; Magon, A.; Chamberland, S.; Hecker, S. J.; Lee, V. J. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 3305.
12. Gruschow, S.; Rackham, E. J.; Elkins, B.; Newill, P.; Hill, L. M.; Goss, R. J. M. *ChemBioChem* **2009**, *10*, 355.
13. a) Gentle, C. A.; Harrison, S. A.; Inukai, M.; Bugg, T. D. H. *J. Chem. Soc. Perkin Trans. 1* **1999**, 1287. b) Howard, N. I.; Bugg, T. D. H. *Bioorg. Med. Chem.* **2003**, *11*, 3083.
14. Bozzoli, A.; Kazmierski, W.; Kennedy, G.; Pasquarello, A.; Pecunioso, A. *Bioorg. Med. Chem. Lett.* **2000**, *10*, 2759.
15. Sun, D. Q.; Jones, V.; Carson, E. I.; Lee, R. E. B.; Scherman, M. S.; McNeil, M. R.; Lee, R. E. *Bioorg. Med. Chem. Lett.* **2007**, *17*, 6899.

16. a) Okamoto, K.; Sakagami, M.; Feng, F.; Togame, H.; Takemoto, H.; Ichikawa, S.; Matsuda A. *J. Org. Chem.* **2012**, *77*, 1367. b) Okamoto, K.; Sakagami, M.; Feng, F.; Togame, H.; Takemoto, H.; Ichikawa, S.; Matsuda A. *Org. Lett.* **2011**, *13*, 5240.
17. a) Tanino, T.; Al-Dabbagh, B.; Mengin-Lecreulx, D.; Bouhss, A.; Oyama, H.; Ichikawa, S.; Matsuda, A. *J. Med. Chem.* **2011**, *54*, 8421. b) Tanino, T.; Ichikawa, S.; Al-Dabbagh, B.; Bouhss, A.; Matsuda, A. *ACS Med. Chem. Lett.* **2010**, *1*, 258. c) Tanino, T.; Ichikawa, S.; Shiro, M.; Matsuda, A. *J. Org. Chem.* **2010**, *75*, 1366.
18. The dipeptides **23** and **24** were prepared by peptide coupling of either L-Tyr-OBn or L-*m*-Tyr-OBn with **25**, protection of the phenol with the TBS group, and hydrogenolysis of the Bn group.
19. a) Bouhss, A.; Crouvoisier, M.; Blanot, D.; Mengin-Lecreulx D. *J. Biol. Chem.* **2004**, *279*, 29974. b) Ahmed, A.; Bararia, D.; Vinayak, S.; Yameen, M.; Biswas, S.; Dev, V.; Kumar, A.; Ansari, M. A.; Sharma, Y. D. *Antimicrob. Agents Chemother.* **2004**, *48*, 879.
20. Maki, H.; Miura, K.; Yamano, Y. *Antimicrob. Agents Chemother.* **2001**, *45*, 1823.

Figure 1. Structure of Pacidamycin D and Its 3'-Hydroxy analogue 2

Figure 2. Structure of Mureidomycins

Scheme 1. Retrosynthetic Analysis of 2 Disconnected by Ugi-Four Component Reaction

Scheme 2. Preparation of Aldehyde Unit 7

Scheme 3. Preparation of Isonitrile Unit 8

Scheme 4. Total Synthesis of 3'-Hydroxypacidamycin D (2) and Its Epimer 19

Scheme 5. Attempt to Use Dipeptide Aldehyde 21 in U-4CR

Scheme 6. Synthesis of Analogues 27a-c, 30 and 31

Table 1. MraY Inhibitory Activity of 3'-Hydroxypacidamycin Analogues

	1	2	19	27a	27b	27c	30	31
IC ₅₀ (nM) ^a	22	42	4000	22	4000	65	650	8900

^aThe inhibitory activity of compounds on the purified MraY enzyme (*Staphylococcus aureus*) was then examined by fluorescence based MraY assay using UDP-MurNAc-dansylpentapeptide, where the formation of dansylated lipid I was monitored by fluorescence enhancement (excitation at 355 nm, emission at 535 nm).

Table 2. Anti-Pseudomonal Activity of 3'-Hydroxypacidamycin Analogues

compound	MIC ($\mu\text{g/mL}$) ^a			
	<i>P. aeruginosa</i> PAO1	<i>P. aeruginosa</i> YY165 (ΔmexB)	<i>P. aeruginosa</i> ATCC 25619	<i>P. aeruginosa</i> SR 27156
1	64	16	16	16
2	32	8	16	16
19	>32	>32	>32	>32
27a	>32	16	32	16
27b	>32	>32	>32	>32
27c	>32	>32	>32	>32
30	>32	>32	>32	>32
31	>32	>32	>32	>32

^aMICs were determined by a microdilution broth method as recommended by the NCCLS with cation-adjusted Mueller-Hinton broth (CA-MHB). Serial two-fold dilutions of each compound were made in appropriate broth, and the plates were inoculated with 5×10^4 CFU of each strain in a volume of 0.1 mL. Plates were incubated at 35 °C for 20 h and then MICs were scored.

Captions

Figure 1. Structure of Pacidamycin D and Its 3'-Hydroxy analogue **2**

Figure 2. Structure of the Mureidomycins

Scheme 1. Retrosynthetic Analysis of **2** Disconnected by an Ugi-Four Component Reaction

Scheme 2. Preparation of the Aldehyde Unit **7**

Scheme 3. Preparation of the Isonitrile Unit **8**

Scheme 4. Total Synthesis of 3'-Hydroxypacidamycin D (**2**) and Its Epimer **19**

Scheme 5. Attempt to Use the Dipeptide Aldehyde **21** in the U-4CR

Scheme 6. Synthesis of Analogues **27a-c**, **30** and **31**

Table 1. MraY Inhibitory Activity of 3'-Hydroxypacidamycin Analogues

Table 2. Anti-Pseudomonal Activity of 3'-Hydroxypacidamycin Analogues